

Research Paper

Effect of Short-time Exposure of Local Extremely Low-Frequency Magnetic Fields on Sleepiness in Male Rats



Elnaz Azizi¹, Fatemeh Ayoobi¹, Ali Shamsizadeh², Amir Moghadam-Ahmadi¹, Seyed Ali Shafiei^{3*}, Alireza Khoshdel⁴, Mohammad Reza Mirzaei⁵

1. Non-Communicable Diseases Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

2. Physiology-Pharmacology Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.

3. Neuroscience Research Center, Qom University of Medical Sciences, Qom, Iran.

4. Department of Biochemistry, School of Medicine, Semnan University of Medical Sciences, Semnan, Iran.

5. Molecular Medicine Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran.



Citation Azizi, E., Ayoobi, F., Shamsizadeh, A., Moghadam-Ahmadi, A., Shafiei, S. A., Khoshdel, A. R., and Mirzaei, M. R. (2012). Effect of Short-Time Exposure of Local Extremely Low-Frequency Magnetic Fields on Sleepiness in Male Rats. *Basic and Clinical Neuroscience*, 13(4), 519-530. <http://dx.doi.org/10.32598/bcn.2022.2610.1>

doi <http://dx.doi.org/10.32598/bcn.2022.2610.1>

**Article info:**

Received: 02 Jun 2020

First Revision: 13 Mar 2021

Accepted: 10 Nov 2021

Available Online: 01 Jul 2022

Keywords:

Magnetic field, Extremely low-frequency magnetic fields (ELF-MFs), Oxalic acid, Anxiety, Sleepiness

ABSTRACT

Introduction: Lack of high-quality sleep causes severe side effects like anxiety and changes in plasma concentration of oxalate. The current study investigated the impact of local extremely low-frequency magnetic fields (ELF-MFs) on inducing sleep (sleepiness) and anxiety in male rats.

Methods: In this experimental study, 40 male rats were divided into four groups (n=10 for each group). The ELF-MF exposure (0, 10, and 18 Hz) was applied with an intensity of 200 μ T for three days (10 min/d). The sham-treated animal did not receive ELF-MF. Serum levels of oxalic acid (OA) and sleepiness were measured before and after the last exposure to ELF-MF or sham. Anxiety, sleepiness, and OA were measured using the elevated plus maze, open-field test (OFT), and ELISA test.

Results: A comparison of oxalate levels before and after exposure to ELF-MF revealed that ELF-MF (10 Hz) decreased the serum level of oxalate (P<0.05). Comparing open/closed arm entry (in an elevated plus maze) between before and after exposure to ELF-MF revealed significant differences. Also, frequency, velocity, and distance moved were decreased in the open-field test.

Conclusion: Results of the present study demonstrated that ELF-MF with short-time exposure may modulate the metabolism of OA and may modulate anxiety-like behavior or kind of induction of sleepiness in male rats.

*** Corresponding Author:**

Seyed Ali Shafiei, PhD.

Address: Neuroscience Research Center, Qom University of Medical Sciences, Qom, Iran.

Tel: +98 (25) 31971040

E-mail: sashafiei@muq.ac.ir ; salishafiei@yahoo.com

Highlights

- Oxalate acid concentration may reduce after short time ELF-MF exposure.
- Locomotor activity in male rats may decrease after the ELF-MF exposure.
- Short time ELF-MF exposure may induce sleepiness in male rats that may be used to treat sleep disorders.

Plain Language Summary

It is necessary for a person to have good sleep to feel happy during the day. The usual way to treat the patient's sleep disorders is drug therapy, but there are some non-pharmacological treatments such as cognitive behavioral therapy and proper diet. In this study we decided to evaluate the effect of ELF-MFs on sleep induction (sleepiness) in male rats by assessing behavioral tests and measuring oxalate acid density. The results showed that the activity of rats and oxalate acid concentration reduced after ELF-MF exposure. This was consistent with results of the plus maze test and the reduction of velocity, frequency and in the open-field test can be attributed to sleepiness. The results of this research showed that ELF-MF with short time exposure may modulate the anxiety-like behavior or kind of induction of sleepiness in male rats. This effect may be used to treat sleep disorders and requires further human studies.

1. Introduction

A person must sleep well to feel happy during the day (Sadock, Flaherty, & Sadock, 2011). Insomnia is a general health disorder associated with occupational problems, interpersonal problems, driving accidents, educational problems, difficulty working night shifts, and anxiety (Bélanger, Morin, Langlois, & Ladouceur, 2004; Husby & Lingjaerde, 1990; Lader, 1999; Okada et al., 2000). According to studies, insomnia may develop at any age, although it is more common in aging (Friedman, 2006). Reduced sleep duration in humans has been associated with risk for metabolic disorders, diabetes, obesity, weight gain, and cardiovascular disease (Ayas et al., 2003; Di Milia, Vandelanotte, & Duncan, 2013; Schmid, Hallschmid, & Schultes, 2015; Singh, Drake, Roehrs, Hudgel, & Roth, 2005). The usual way to treat a patient's sleep disorders is drug therapy, but there are some non-pharmacological treatments such as cognitive behavioral therapy and proper diet (Morin et al., 2006).

Some studies showed that exposure to extremely low-frequency magnetic fields (ELF-MFs) might affect cognitive functions (Akbarnejad et al., 2018; Shafiei, Firoozabadi, Rasoulzadeh Tabatabaie, & Ghabaee, 2014; Sharma et al., 2017). Nevertheless, there are conflicting reports on the influence of MFs on sleep quality. For example, electromagnetic fields may affect sleep and cognition tasks in the rat (Dyche, Anch, Fogler, Barnett, & Thomas, 2012; Hassanshahi et al., 2017).

Chronic exposure to pulsed ELF-MFs can alleviate the sleep disorders of patients with insomnia (Pelka Jaelnicke, C., & Gruenwald, 2001) and affect sleep quality (Altpeter et al., 2006). On the other hand, Schmid et al. and Tworoger et al. did not report any significant effect on the sleep quality of young women (Schmid et al., 2012; Tworoger et al., 2004), even in diabetic patients with sleep disorders (Wrobel et al., 2008) and sleeping patterns of healthy people (Wagner et al., 2000). However, our latest studies showed that exposure to ELF-MFs can change EEGs recorded from some areas of the scalp and may even influence reaction time in young, healthy women (Ayoobi, Shamsizadeh, & Shafiei, 2017; Shafiei, Firoozabadi, Rasoulzadeh Tabatabaie, & Ghabaee, 2012). A good night's sleep after ELF-MF exposure had been reported by some participants who have sleep problems. However, the evaluation of this issue was not the focus of this research (Amiri Fallah, Firoozabadi, Shafiei, & Assadi, 2011; Amirifalah, Mohammad, Firoozabadi, & Shafiei Darabi, 2013). These studies have shown that local ELF-MFs may become a helpful therapeutic tool in treating some psychological symptoms (Shafiei & Firoozabadi, 2014).

Then again, the oxalic acid level is a good marker of sleep deprivation (Weljie et al., 2015). The primary sources of blood oxalate are a vegetarian diet, ascorbic acid decomposition, and endogenous synthetic pathways in the liver and erythrocytes (Hodgkinson & Zarembski, 1968). Endogenous substrates for oxalate synthesis include glycolate in peroxisome and hydroxyproline, and sleep deprivation for several days is related to hydroxy-

proline levels in mice (Koehl et al., 2006). Acute sleep deprivation eliminated oxalate from plasma via increased urinary excretion (Weljie et al., 2015).

Therefore, due to the importance of inducing sleep and the safety of extremely weak magnetic fields at very low frequencies, we decided to evaluate the effect of ELF-MFs on sleep induction (sleepiness) in male rats by assessing behavioral tests and measuring oxalate acid density.

2. Material And Methods

Study subjects

This study was obtained from the Central Animal House, Rafsanjan University of Medical Sciences, Iran. Forty male rats of the Wistar strain (160-210 g) were distributed into four experimental groups (n=40: three exposed groups [0, 10, and 18 Hz] n=10/group; sham exposed n=10) and were individually housed before and during experimental protocols. The animals were exposed to a stable temperature (22°C-24°C) and relative humidity of 45%-55%. They were kept under a cycle of 12h light and 12h darkness for at least 2 weeks in our laboratory before starting experiments. Rodents were kept awake at night and asleep during the day. In order to adjust the sleep-wake cycle of rats, we reversed 12h of light and 12h of darkness for these animals in the laboratory so that they have time to wake up during the day and we could do convenient experiments. Animals were fed daily, and cages cleaned.

ELF-MFs exposure

Helmholtz coils with 40 cm diameter separated by a distance of 20 cm were used to generate a nearly uniform magnetic field. A dodecagon made of acrylic with 25 cm diameter and 15 cm height was used and located between two Helmholtz coils. The animal was placed in this box, and a magnetic field with an intensity of 200 μ T was applied for 10 min/d for three days. ELF-MFs exposure was performed at 9 AM. The three experimental groups (0, 10, and 18 Hz) were differentiated in terms of the transmitted frequency, and in group four, all stages were performed like those of the previous groups, except that the set was turned off and there was no magnetic field (Figure 1). Food and water were unavailable during exposures.

Open-field test

Anxiety-like behaviors and locomotor activity were measured with an open-field and behavioral tracking

analysis system; we used a box 50cm in height, and 70cm in diameter, with a video camera, installed 2.5m above the apparatus. Each rat activity was digitally monitored and then analyzed using EthoVision XT software, version 7.1, Noldus Information Technology, Wageningen, The Netherlands. The interior area of the box was divided into inner and outer zones. First, the rat was placed in the center of the box, and its activity was recorded for 5 min. Then the following behavioral parameters were scored: velocity (cm/s), frequency, and total distance moved (cm) in the inner zone. After removing the rat from the open field, the experimental chamber was cleaned with diluted ethanol (10%) and dried (Crawley, 1999).

Elevated plus maze test

we used the elevated plus maze test to measure the rat's movement and anxiety. It comprised 2 open arms (50 \times 10 cm) and 2 enclosed arms (50 \times 10 \times 40 cm) that extend from a common central platform (10 \times 10 cm). A wooden apparatus was elevated 50 cm above floor level. Testing was conducted between 9.00 am and 1.00 pm in a quiet room. The rats were placed in the center of the maze facing an open arm individually, and 5 min were allowed for free exploration. All sessions were filmed. The floor was cleaned after each test with distillate water. The animals were tested before the first ELF-MF exposure and after the last ELF-MF exposure. Measurements were made from the time spent in the maze's central, open and closed parts. In the absence of anxiety, the animal moves more and has a great deal of activity in the open arm. However, if the rat experiences anxiety and sleepiness, it will remain in the closed arm (Rodgers & Dalvi, 1997).

Oxalic acid (OA) measurement

The total serum metabolic oxalic acid (OA) was measured by oxalate assay kit (Colorimetric-KA4532, Abnova, Taiwan) to evaluate how systemic metabolism is impacted by sleep restriction. Blood was taken in the morning one day before the first exposure to ELF-MF (before behavioral tests) and one day after the last exposure to ELF-MF (after behavioral tests). The kit uses a double-antibody sandwich enzyme-linked immunosorbent assay (ELISA) to assay the level of OA in samples. OA is added to a monoclonal antibody enzyme well which is pre-coated with OA monoclonal antibody; this is subject to incubation, and then OA antibodies labeled with biotin are added and combined with streptavidin-HRP to form the immune complex. Next, incubation is carried out, and the material is re-



Figure 1. Schematic view of Helmholtz coils

NEURSCIENCE

washed to remove the uncombined enzyme. To close, chromogen solution A and B is added, and the color of the liquid changes into blue, and with the effect of the acid, the color finally becomes yellow. The chroma of color and the concentration of the substance OA in the sample were positively correlated.

Procedure

Figure 2 represents all of the protocols on different days. This protocol was implemented at different frequencies (0, 10, and 18Hz) for these groups and the sham group, in which the signal generator did not produce any electrical signals, and thus there were no ELF-MFs in the sham group.

Statistical analysis

The obtained data were analyzed by SPSS software version 20.0. The quantitative data were reported as Mean \pm SD and qualitative data in percentages. A paired t-test was used to compare open-field scores, elevated plus maze, ELF-MF, and mean oxalic acid before and

after the last exposure. Differences among groups were compared using 1-way ANOVA across all groups. The significance level in all tests was $P < 0.05$.

3. Results

ELF-MF exposure and anxiety

The average presence of rats in the open arm decreased after the last ELF-MF exposure, and there were significant differences in the ELF-MF exposure 10Hz ($P=0.03$) and ELF-MF exposure 18Hz groups ($P=0.007$). On the other hand, the average presence of rats in the closed arm increased significantly after the last ELF-MF exposure in the ELF-MF exposure 18Hz group ($P=0.01$) (Figure 3).

ELF-MF exposure and locomotor activity

The open-field test was used to determine anxiety-like behaviors and locomotor activity in different groups before and after ELF-MF exposure. For locomotor activity, the traveled distance of movements decreased after the last ELF-MF exposure, and the traveled distances were




Figure 2. Schematic view of the protocol of study on different days

NEURSCIENCE




Figure 3. Elevated plus maze test in all groups

There was a significant effect between the groups before the first and after the last ELF-MF exposures.

* $P < 0.05$; ** $P < 0.01$.

significant between before the first and after the last ELF-MF exposure in the direct current (DC) magnetic field exposure, in which frequency was 0Hz ($P=0.02$), ELF-MF exposure 10Hz ($P=0.002$) and ELF-MF exposure 18Hz groups ($P=0.000$). Also, the velocity decreased after the last ELF-MF exposure and was significant in ELF-MF exposure 0Hz ($P=0.02$), ELF-MF exposure 10Hz ($P=0.002$), and ELF-MF exposure 18Hz groups ($P=0.000$). Frequency decreased after the last ELF-MF exposure, and it was significant in all groups ($P=0.01$) except the sham group ($P=0.48$) (Figure 4).

ELF-MF exposure and OA density

The serum levels of OA were measured by ELISA. The mean serum level of OA was 2.10 ± 0.56 nmol/ μ L before ELF-MF exposure, and it was significantly higher as compared after ELF-MF exposure (1.28 ± 0.47) in the wave 10Hz group ($P < 0.05$). No significant difference was observed among the other groups before and after ELF-MF exposure (Figure 5). All results are summarized in Table 1.

4. Discussion

Our previous human experiments suggested that exposure to 200μ T MFs with extremely low-frequency may improve sleep disturbance (insomnia) caused by anxiety (Amirifallah et al., 2011; Amirifallah et al., 2013). This study used magnetic fields at very low frequencies

to evaluate the impact of DC (0Hz), 10Hz, 18Hz, and 200μ T MF on sleepiness induction rates.

We suggest that the reduction in the frequency, velocity, and distance moved in open-field tests after exposure (Figure 4) and the decrease in the average presence of rats in the open arm (Figure 3) were due to a kind of induction of sleepiness. Therefore, the findings of this study are consistent with those of our previous studies, confirming our hypothesis that the ELF-MFs can induce sleepiness (Table 1).

Ayoobi et al. examined the effect of ELF-MF on sleepiness induction in healthy students, especially females. They used 200μ T MF (3 min duration) at three frequencies (18, 14, and 10Hz), applied to the skull in areas C4, CZ, and C3, respectively, on 32 males and 31 females. Their study showed that exposure to local ELF-MFs might influence minimum reaction time in young, healthy females (Ayoobi et al., 2017). Their findings are consistent with the results of this study.

However, some studies, like Kurokawa et al., had inconsistent findings. They evaluated the effects of continuous exposure to 50Hz, 20μ T MF on 20 subjects for 55 min. The reaction time in time perception tasks did not change with ELF-MF exposure (Kurokawa, Nitta, Imai, & Kabuto, 2003). In this study, the magnetic field used was 10 times less than the magnetic field in this study.




Figure 4. Open-field tests in all groups

There was a significant effect among the groups before the first and after the last ELF-MF exposures.

* P<0.05; ** P<0.01; *** P<0.001.

Table 1. Comparing all tests before the first and after the last ELF-MF exposures

| Tests | | | Mean±SD | | | |
|----------------------------------|---------------------|-----------|----------------|-----------------|------------------|--------------------|
| | | | Sham | Exposure 0 | Exposure 10 | Exposure 18 |
| Elevated plus maze test (s) | Open arm | Before | 171.22±35.68 | 124.65±55.49 | 156.67±28.66 | 204.44±27.28 |
| | | After | 130.71±69.58 | 105.14±80.88 | 97.81±42.18 * | 148.10±79.78 ** |
| | Close arm | Before | 187.33±51.35 | 190.92±58.24 | 198.00±34.84 | 123.00±31.56 |
| | | After | 205.22±78.35 | 217.51±81.17 | 229.71±53.21 | 175.71±71.53 * |
| Open-field test | Distance moved (cm) | Before | 1264.41±375.82 | 1233.51±257.88 | 1463.85±258.55 | 1683.34±174.31 |
| | | After | 940.35±377.01 | 877.33±169.85 * | 991.16±240.35 ** | 1179.85±154.59 *** |
| | Velocity (cm/s) | Before | 4.24±1.25 | 4.11±0.86 | 4.88±0.86 | 5.61±0.58 |
| | | After | 3.13±1.26 | 2.92±0.57 | 3.30±0.80 ** | 3.93±0.52 *** |
| | Frequency | Before | 15.00±10.54 | 20.00±8.88 | 22.57±7.37 | 26.66±6.12 |
| | | After | 11.00±8.49 | 9.29±6.75 * | 12.00±8.14 * | 16.11±5.58 * |
| Oxalic acid (OA) Density (n/mol) | Before | 1.51±0.24 | 1.96±0.20 | 2.10±0.56 | 1.89±0.53 | |
| | After | 1.52±0.36 | 1.30±0.41** | 1.28±0.48*** | 1.42±0.22 | |

* Significant difference between before the first and after the last ELF-MF exposures.

* P<0.05; ** P<0.01; *** P<0.001.

NEUROSCIENCE

Generally, reduced time spent in the central area in the open-field test (Prut & Belzung, 2003) and reduced time spent in the open arms plus maze (Lister, 1987) have been established as an index of increased anxiety behavior. Some animal studies indicated that high in-

tensity and chronic exposure to ELF-MF might induce anxiety-like behavior in rodents (Kitaoka, Kitamura, Aoi, Shimizu, & Yoshizaki, 2013; Liu, Wang, He, & Ye, 2008), and in this study, we expect them to be sleepy because of short exposure.




Figure 5. Comparing serum levels of oxalic acid before the first and after the last of the ELF-MF exposure groups

Serum OA levels in wave DC and wave 10Hz group after ELF-MF exposure were significantly less than before the ELF-MF exposure. No significant difference was observed among the other groups. ** P<0.01, *** P<0.001.

NEUROSCIENCE

Hi, et al. conducted an experimental study and showed that 4 h/d of chronic exposure to ELF-MF significantly affects adult rats' anxiety level and spatial memory. However, 1 h/d exposure to ELF-MFs had no effect in any test (He et al., 2011). ELF-MF exposure of rats 4 h/day increased the anxiety-like behaviors in the open-field test and the elevated plus maze test without altering locomotor activity but had no observed effect in the light/dark box test. Moreover, 1 h/d of ELF-MF exposure did not affect any test. Increased exposure to ELF-EMF may directly and significantly affect anxiety (Hosseinabadi, Khanjani, Ebrahimi, Haji, & Abdollahfard, 2019).

Generally, in a short time exposure, evidence shows that ELF-MF exposure had no significant effects on anxiety induction. Furthermore, the results of the meta-analyses have shown that exposure to ELF-MFs has no considerable effect on cognitive functions (Barth, Ponocny, Ponocny-Seliger, Vana, & Winker, 2010).

Ouyang et al. measured plasma concentrations of oxalic acid in free-ranging wild birds' blood as a sleep restriction biomarker based on earlier studies (Ouyang et al., 2017; Weljie et al., 2015). The findings showed that oxalic acid levels reduced strongly in birds with their highest activity levels at night. They assumed it is likely that night activity may cause sleeplessness and, as a result, a decrease in the concentration of oxalic acid (Ouyang et al., 2017).

Our results suggest that the activity of rats and oxalate acid concentration reduced after ELF-MF exposure (Figure 3). ELF-MF is assumed to cause severe anxiety in rats, resulting in oxalate decrease (Table 1). These results are consistent with the results of the plus maze test (Figure 1) and the reduction of velocity frequency in the open-field test (Figure 2). However, such a hypothesis is incompatible with the evidence obtained in our previous human research, which showed that healthy female participants experienced good sleep after the ELF-MF exposure. The result of Figure 2 (plus maze) can be attributed to sleepiness.

In 2011, Akhtary et al. analyzed the effects of ELF-MFs on learning, memory, and pseudo-anxiety behaviors in giant white laboratory mice. Field groups lived for 28 days in special fiberglass cages with a magnetic field of 10 or 100 μ T. The anxiety was measured by the elevated plus maze test. The groups showed no significant difference (Akhtary, Rashidy-Pour, Vafaei, & Jaddi, 2011). It can be concluded that a 10 μ T field with chronic conditions could not induce anxiety. These findings are similar to the results of the present study.

In another study, 60 diabetic patients were divided into two groups, and 32 were subjected to a magnetic field with a frequency of 180Hz to 195Hz with an intensity of about 100 μ T for 20 min, 5 times per week. The results showed that ELF-MFs did not affect sleep disorders and serum level oxalate (Wrobel et al., 2008).

Monazzam et al. reviewed the quality of sleep and general health of employees exposed to ELF-MFs in a petrochemical complex. They examined the relationship between the levels of ELF, sleep quality, and the general health of 40 EMF-exposed workers and 22 controls. The results showed that 28% of the subjects were in poor health, and 61% had a sleep disorder. No significant difference ($P < 0.01$) was found among the case and control groups regarding sleep quality and general health (Monazzam et al., 2014).

In 2011, the effects of ELF-MFs on mouse anxiety levels were investigated. A magnetic field of 50Hz was used with two groups of animals for 1 and 4 hr. Pseudo-anxiety behaviors were evaluated using an open-field experiment and evaluated plus maze. Mice from the 4-h exposure had more pseudo-anxiety behaviors and static motor activities followed by increased anxiety (He et al., 2011). This difference can be due to the difference in the exposure time and frequency of the magnetic field in these two studies.

The limitations of our study were not measuring melatonin levels. Some reports suggest that melatonin production is slightly increased in exposed animals (Altpeter et al., 2006; Dyche et al., 2012). There are many studies in this field, but when we examined them, there was a significant discrepancy and even a negative one.

5. Conclusion

This study showed that ELF-MF with short-time exposure might modulate the anxiety-like behavior or kind of induction of sleepiness in male rats. This Effect may be used to treat sleep disorders and requires further study.

Ethical Considerations

Compliance with ethical guidelines

The research was approved by the Ethical Committee of Rafsanjan University of Medical Science (RUMS).

Funding

This study was supported by Rafsanjan University of Medical Science (RUMS).

Authors' contributions

Conceptualization and Supervision: Fatemeh Ayoobi, Ali Shamsizadeh, Amir Moghadam-Ahmadi, Seyed Ali Shafiei and Alireza Khoshdel; Methodology: Fatemeh Ayoobi, Ali Shamsizadeh, Amir Moghadam-Ahmadi, Seyed Ali Shafiei; Investigation, Writing—original draft, and Writing—review & editing: All authors; Data collection: Elnaz Azizi, Fatemeh Ayoobi; Data analysis: Elnaz Azizi, Fatemeh Ayoobi, Ali Shamsizadeh, Amir Moghadam-Ahmadi, Seyed Ali Shafiei, and Alireza Khoshdel; Funding acquisition and Resources: All authors.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgments

We especially thank S. Skies for editing this manuscript.

References

- Akbarnejad, Z., Esmailpour, K., Shabani, M., Asadi-Shekaari, M., Saeedi goraghani, M., & Ahmadi-Zeidabadi, M. (2018). Spatial memory recovery in Alzheimer's rat model by electromagnetic field exposure. *International Journal of Neuroscience*, 128(8), 691-696. [DOI:10.1080/00207454.2017.1411353]
- Akhtary, Z., Rashidy-Pour, A., Vafaei, A. A., & Jadidi, M. (2011). [Effects of extremely low-frequency electromagnetic fields on learning, memory, and anxiety-like behaviors in rats (Persian)]. *Koomesh*, 12(4), 435-446. <http://koomesh-journal.semums.ac.ir/article-1-1367-fa.html>
- Altpeter, E. S., Roosli, M., Battaglia, M., Pfluger, D., Minder, C. E., & Abelin, T. (2006). Effect of short-wave (6-22 MHz) magnetic fields on sleep quality and melatonin cycle in humans: The Schwarzenburg shut-down study. *Bioelectromagnetics*, 27(2), 142-150. [DOI:10.1002/bem.20183]
- Amirifallah, Z., Firoozabadi, M., Shafiei, A., & Assadi, A. (2011). Scrutiny of brain signals variations in regions Cz, C3, and C4 under local exposure of extremely low frequency and weak-pulsed magnetic field to promote neurofeedback systems. *Physiology and Pharmacology*, 15(1), 144-163. [Link]
- Amirifalah, Z., Mohammad P. Firoozabadi, S., & Shafiei Darabi, S. A. (2013). Local exposure of brain central areas to a pulsed ELF magnetic field for a purposeful change in EEG. *Clinical EEG and Neuroscience*, 44(1), 44-52. [DOI:10.1177/1550059412460164]
- Ayas, N. T., White, D. P., Manson, J. E., Stampfer, M. J., Speizer, F. E., Malhotra, A., & Hu, F. B. (2003). A prospective study of sleep duration and coronary heart disease in women. *Archives of Internal Medicine*, 163(2), 205-209. [DOI:10.1001/archinte.163.2.205]
- Ayoobi, F., Shamsizadeh, A., & Shafiei, S. A. (2017). The effect of local extremely low-frequency magnetic field on student sleepiness. *Neurological Research*, 39(12), 1080-1085. [DOI:10.1080/01616412.2017.1380931]
- Barth, A., Ponocny, I., Ponocny-Seliger, E., Vana, N., & Winkler, R. (2010). Effects of extremely low-frequency magnetic field exposure on cognitive functions: Results of a meta-analysis. *Bioelectromagnetics*, 31(3), 173-179. [DOI:10.1002/bem.20543]
- Bélanger, L., Morin, C. M., Langlois, F., & Ladouceur, R. (2004). Insomnia and generalized anxiety disorder: Effects of cognitive behavior therapy for gad on insomnia symptoms. *Journal of Anxiety Disorders*, 18(4), 561-571. [DOI:10.1016/S0887-6185(03)00031-8]
- Crawley, J. N. (1999). Behavioral phenotyping of transgenic and knockout mice: Experimental design and evaluation of general health, sensory functions, motor abilities, and specific behavioral tests. *Brain Research*, 835(1), 18-26. [DOI:10.1016/S0006-8993(98)01258-X]
- Di Milia, L., Vandelanotte, C., & Duncan, M. J. (2013). The association between short sleep and obesity after controlling for demographic, lifestyle, work, and health-related factors. *Sleep Medicine*, 14(4), 319-323. [DOI:10.1016/j.sleep.2012.12.007]
- Dyche, J., Anch, A. M., Fogler, K. A., Barnett, D. W., & Thomas, C. (2012). Effects of power frequency electromagnetic fields on melatonin and sleep in the rat. *Emerging Health Threats Journal*, 5(1), 10904. [DOI:10.3402/ehjt.v5i0.10904]
- Friedman, L. (2006). Insomnia and aging. *Sleep: A comprehensive handbook*, 611-614. [DOI:10.1002/0471751723.ch79]
- Hassanshahi, A., Shafeie, S. A., Fatemi, I., Hassanshahi, E., Allahtavakoli, M., Shabani, et al. The effect of Wi-Fi electromagnetic waves in unimodal and multimodal object recognition tasks in male rats [Article in Press]. *Neurological Sciences*, 1-8. [DOI:10.1007/s10072-017-2920-y]
- He, L.-H., Shi, H.-M., Liu, T.-T., Xu, Y.-C., Ye, K.-P., & Sheng, W. (2011). Effects of extremely low-frequency magnetic field on adult rats' anxiety level and spatial memory. *Chinese Medical Journal*, 124(20), 3362-3366.
- Hodgkinson, A., & Zaremski, P. (1968). Oxalic acid metabolism in man: A review. *Calcified Tissue Research*, 2(1), 115-132. [DOI:10.1007/BF02279201]
- Hosseinabadi, M. B., Khanjani, N., Ebrahimi, M. H., Haji, B., & Abdolahfard, M. (2019). The effect of chronic exposure to extremely low-frequency electromagnetic fields on sleep quality, stress, depression, and anxiety. *Electromagnetic Biology and Medicine*, 38(1), 96-101. [DOI:10.1080/15368378.2018.1545665]
- Husby, R., & Lingjaerde, O. (1990). Prevalence of reported sleeplessness in northern Norway in relation to sex, age, and season. *Acta Psychiatrica Scandinavica*, 81(6), 542-547. [DOI:10.1111/j.1600-0447.1990.tb05009.x]
- Kitaoka, K., Kitamura, M., Aoi, S., Shimizu, N., & Yoshizaki, K. (2013). Chronic exposure to an extremely low-frequency magnetic field induces depression-like behavior and corti-

- costerone secretion without enhancement of the hypothalamic-pituitary-adrenal axis in mice. *Bioelectromagnetics*, 34(1), 43-51. [DOI:10.1002/bem.21743]
- Koehl, M., Battle, S., & Meerlo, P. (2006). Sex differences in sleep: The response to sleep deprivation and restraint stress in mice. *Sleep*, 29(9), 1224-1231. [DOI:10.1093/sleep/29.9.1224]
- Kurokawa, Y., Nitta, H., Imai, H., & Kabuto, M. (2003). No influence of short-term exposure to 50-Hz magnetic fields on cognitive performance function in humans. *International Archives of Occupational and Environmental Health*, 76(6), 437-442. [DOI:10.1007/s00420-003-0445-6]
- Lader, M. H. (1999). Limitations on the use of benzodiazepines in anxiety and insomnia: Are they justified? *European Neuropsychopharmacology*, 9, S399-S405. [DOI:10.1016/S0924-977X(99)00051-6]
- Lister, R. G. (1987). The use of a plus-maze to measure anxiety in the mouse. *Psychopharmacology*, 92(2), 180-185. [DOI:10.1007/BF00177912]
- Liu, T. T., Wang, S., He, L. H., & Ye, K. P. (2008). Anxiogenic effect of chronic exposure to the extremely low-frequency magnetic field in adult rats. *Neuroscience Letters*, 434(1), 12-17. [DOI:10.1016/j.neulet.2008.01.019]
- Monazzam, M. R., Hosseini, M., Matin, L. F., Aghaei, H. A., Khosroabadi, H., & Hesami, A. (2014). Sleep quality and general health status of employees exposed to extremely low-frequency magnetic fields in a petrochemical complex. *Journal of Environmental Health Science and Engineering*, 12(1), 78. [DOI:10.1186/2052-336X-12-78]
- Morin, C. M., Bootzin, R. R., Buysse, D. J., Edinger, J. D., Espie, C. A., & Lichstein, K. L. (2006). Psychological and behavioral treatment of insomnia: Update of the recent evidence (1998-2004). *Sleep*, 29(11), 1398-1414. [DOI:10.1093/sleep/29.11.1398]
- Okada, T., Sugishita, T., Murakami, T., Murai, H., Saikusa, T., Horino, T., et al. (2000). Effect of the defatted rice germ enriched with GABA for sleeplessness, depression, autonomic disorder by oral administration. *Nippon Shokuhin Kagaku Kogaku Kaishi = Journal of the Japanese Society for Food Science and Technology*, 47(8), 596-603. [DOI:10.3136/nshkk.47.596]
- Ouyang, J. Q., de Jong, M., van Grunsven, R. H. A., Matson, K. D., Haussmann, M. F., Meerlo, P., et al. (2017). Restless roots: Light pollution affects a free-living songbird's behavior, sleep, and physiology. *Global Change Biology*, 23(11), 4987-4994. [DOI:10.1111/gcb.13756]
- Pelka, R. B., Jaenicke, C., & Gruenewald, J. (2001). A double-blind, placebo-controlled study is an impulse magnetic-field therapy for migraine and other headaches. *Advances in Therapy*, 18(3), 101-109. [DOI:10.1007/BF02850298]
- Prut, L., & Belzung, C. (2003). The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: A review. *European Journal of Pharmacology*, 463(1-3), 3-33. [DOI:10.1016/S0014-2999(03)01272-X]
- Rodgers, R., & Dalvi, A. (1997). Anxiety, defense, and the elevated plus-maze. *Neuroscience & Biobehavioral Reviews*, 21(6), 801-810. [DOI:10.1016/S0149-7634(96)00058-9]
- Sadock, V. A., Flaherty, A. W., & Sadock, B. J. (2011). Kaplan and Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry. Wolters Kluwer Health. [Link]
- Schmid, M. R., Murbach, M., Lustenberger, C., Maire, M., Kuster, N., Achermann, P., et al. (2012). Sleep EEG alterations: Effects of pulsed magnetic fields versus pulse-modulated radio frequency electromagnetic fields. *Journal of Sleep Research*. 21(6), 620-629. [DOI:10.1111/j.1365-2869.2012.01025.x]
- Schmid, S. M., Hallschmid, M., & Schultes, B. (2015). The metabolic burden of sleep loss. *The Lancet Diabetes & Endocrinology*, 3(1), 52-62. [DOI:10.1016/S2213-8587(14)70012-9]
- Shafiei, S. A., Firoozabadi, S., Rasoulzadeh Tabatabaie, K., & Ghabaee, M. (2012). Study of the frequency parameters of EEG influenced by zone-dependent local ELF-MF exposure on the human head. *Electromagnetic Biology and Medicine*, 31(2), 112-121. [DOI:10.3109/15368378.2011.624658]
- Shafiei, S. A., Firoozabadi, S. M., Tabatabaie, K. R., & Ghabaee, M. (2012). Evaluating the changes in alpha-1 band due to exposure to magnetic field. *Iranian Journal of Medical Physics*, 9(2), 141-152. [10.22038/IJMP.2012.320]
- Shafiei, S. A., Firoozabadi, S. M., Tabatabaie, K. R., & Ghabaee, M. (2014). Investigation of EEG changes during exposure to the extremely low-frequency magnetic field to conduct brain signals. *Neurological Sciences*, 35(11), 1715-1721. [DOI:10.1007/s10072-014-1819-0]
- Sharma, A., Kesari, K.K., Verma, H.N., Sisodia, R. (2017). Neurophysiological and behavioral dysfunctions after electromagnetic field exposure: A dose response relationship. In: Kesari, K. (eds) Perspectives in Environmental Toxicology. Environmental Science and Engineering. Springer, Cham. [DOI:10.1007/978-3-319-46248-6_1]
- Singh, M., Drake, C. L., Roehrs, T., Hudgel, D. W., & Roth, T. (2005). The association between obesity and short sleep duration: a population-based study. *Journal of Clinical Sleep Medicine*, 1(04), 357-363. [DOI:10.5664/jcs.m.26361]
- Tworoger, S. S., Davis, S., Emerson, S. S., Mirick, D. K., Lentz, M. J., & McTiernan, A. (2004). Effect of a nighttime magnetic field exposure on sleep patterns in young women. *American Journal of Epidemiology*, 160(3), 224-229. [DOI:10.1093/aje/kwh215]
- Wagner, P., Roschke, J., Mann, K., Fell, J., Hiller, W., Frank, C., et al. (2000). Human sleep EEG under the influence of pulsed radiofrequency electromagnetic fields. Results from polysomnography using submaximal high power flux densities. *Neuropsychobiology*, 42(4), 207-212. [DOI:10.1159/000026695]
- Research Support, Non-U.S. Gov't]. *Neuropsychobiology*, 42(4), 207-212. [DOI:26695]
- Weljie, A. M., Meerlo, P., Goel, N., Sengupta, A., Kayser, M. S., Abel, T., et al. (2015). Oxalic acid and diacylglycerol 36:3 are cross-species markers of sleep debt. *Proceedings of the National Academy of Sciences of the United States of America*, 112(8), 2569-2574. [DOI:10.1073/pnas.1417432112]
- Wrobel, M., Szyborska-Kajane, A., Wystrychowski, G., Bioniszewicz, T., Sieroń-Stoły, K., Sieroń, A., et al. (2008).

Impact of low frequency pulsed magnetic fields on pain intensity, quality of life and sleep disturbances in patients with painful diabetic polyneuropathy. *Diabetes & Metabolism*, 34(4), 349-354. [[DOI:10.1016/j.diabet.2008.02.003](https://doi.org/10.1016/j.diabet.2008.02.003)]

This Page Intentionally Left Blank