

MnSOD downregulation induced by extremely low 0.1 mGy single and fractionated X-rays and microgravity treatment in human neuroblastoma cell line, NB-1

Hiroko P. Indo,^{1,†} Tsukasa Tomiyoshi,^{1,†} Shigeaki Suenaga,¹ Kazuo Tomita,¹ Hiromi Suzuki,^{2,3} Daisuke Masuda,^{2,4} Masahiro Terada,^{2,5} Noriaki Ishioka,^{2,6} Oleg Gusev,^{2,6,7,8} Richard Cornette,⁸ Takashi Okuda,⁸ Chiaki Mukai⁹ and Hideyuki J. Majima^{1,2,*}

¹Department of Oncology and ²Department of Space Environmental Medicine, Kagoshima University Graduate School of Medical and Dental Sciences, 8-35-1 Sakuragaoka, Kagoshima 890-8544, Japan

³Life Science Research Group, Department of Science and Applications, Japan Space Forum, 3-2-1 Surugadai, Chiyoda, Tokyo 100-0004, Japan

⁴Utilization & Engineering Department, Japan Manned Space Systems Corporation, 2-1-6 Tsukuba, Ibaraki 305-0047, Japan

⁵Space Biosciences Division, NASA Ames Research Center, Moffett Field, California 94035, USA

⁶Department of Space Biology and Microgravity Sciences, Institute of Space and Astronautical Science and ⁹Center for Applied Space Medicine and Human Research, Japan Aerospace Exploration Agency, 2-1-1 Sengen, Tsukuba, Ibaraki 305-8505, Japan

⁷Department of Invertebrates Zoology and Functional Morphology, Institute of Fundamental Medicine and Biology, Kazan Federal University 420008, Kremenskaya str., 17 Kazan 420-008, Russia

⁸Anhydrobiosis Research Unit, National Institute of Agrobiological Sciences, 1-2 Ohwashi, Tsukuba, Ibaraki 305-8634, Japan

(Received 6 February, 2015; Accepted 16 February, 2015; Published online 11 July, 2015)

A human neuroblastoma cell line, NB-1, was treated with 24 h of microgravity simulation by clinostat, or irradiated with extremely small X-ray doses of 0.1 or 1.0 mGy using single and 10 times fractionation regimes with 1 and 2 h time-intervals. A quantitative real-time reverse transcription polymerase chain reaction (qRT-PCR) examination was performed for apoptosis related factors (*BAX*, *CYTC*, *APAF1*, *VDAC1-3*, *CASP3*, *CASP8*, *CASP9* *P53*, *AIF*, *ANT1* and *2*, *BCL2*, *MnSOD*, autophagy related *BECN* and necrosis related *CYP-40*). The qRT-PCR results revealed that microgravity did not result in significant changes except for an upregulation of proapoptotic *VDAC2*, and downregulations of proapoptotic *CASP9* and antiapoptotic *MnSOD*. After 0.1 mGy fractionation irradiation, there was increased expression of proapoptotic *APAF1* and downregulation of proapoptotic *CYTC*, *VDAC2*, *VDAC3*, *CASP8*, *AIF*, *ANT1*, and *ANT2*, as well as an increase in expression of antiapoptotic *BCL2*. There was also a decrease in *MnSOD* expression with 0.1 mGy fractionation irradiation. These results suggest that microgravity and low-dose radiation may decrease apoptosis but may potentially increase oxidative stress.

Key Words: space radiation, microgravity, extremely small dose, neuron, NB-1

To ensure the long-term safety and normal expected life span of astronauts exposed to space-related environmental stresses and to help clarify the risks associated with long-term manned space missions, basic data regarding biological responses to such stressors need to be collected. Studies of these biological responses may contribute to a better understanding of the longer-term effects (e.g., carcinogenicity, neurotoxicity, lifespan changes) of exposure to conditions in outer space. Recently accumulated data have revealed that astronauts living on the International Space Station (ISS) are exposed to 0.2–1.0 mSv of space radiation on a daily basis.⁽¹⁾ Thus, several days spent in space will expose an individual to more radiation exposure than one year spent exposed to natural space radiation on Earth (~1.0 mSv/year). The bio-

logical effects of long-term, low-dose exposure to space radiation, including intracellular DNA and mitochondrial damage, may increase the risk of developing a number of disorders.^(2,3) Such radiation effects are important factors to be considered in planning extended manned space missions.⁽³⁻⁵⁾ Similarly, microgravity may influence cellular metabolism and function, including gene expression and signal transduction, and may result in cell death.⁽⁶⁾

Biological experiments involving radiation are aimed at examining dose-effect relationships in which a minimum dose yields a significant irradiation effect. The minimum dose is dependent on study endpoints, and the smaller the minimum dose, the more sensitive the endpoint. Studies of chromosomal aberration and micronucleus formation,⁽⁷⁻¹²⁾ as well as mutation and transformation assays,^(13,14) are sensitive approaches that utilize small doses of approximately 0.1 Gy (100 mGy). However, it remains controversial whether or not the dose-effect relationship is linear at lower doses, according to linear non-threshold (LNT), or non-LNT theory.⁽¹⁵⁾ It is likely that the dose-effect relationship (whether LNT or non-LNT) is partially dependent on the endpoints used in a particular study. To verify the LNT dose-effect relationship, it would be necessary to conduct thousands of experiments.⁽¹⁵⁾ Moreover, examinations at doses smaller than 0.1 Gy (100 mGy) would be difficult due to the lack of a model and/or endpoints. Transcriptomics is an effective tool for examining grouped signaling pathways such as apoptosis, and for investigating changes in the expression patterns of mitochondrial or protein-synthesis genes.

In the present study, we performed a quantitative real-time reverse transcription polymerase chain reaction (qRT-PCR) analysis in order to determine radiation and microgravity-related effects on gene expression in the human neuroblastoma cell line NB-1. The cells were exposed to single, low doses of 0.1 or

[†]These authors have contributed equally to this work.

*To whom correspondence should be addressed.

E-mail: hmajima@dent.kagoshima-u.ac.jp

1.0 mGy, or 10 times fractionated 0.1 or 1.0 mGy X-ray irradiation at 1 or 2 h time intervals. In addition, the cells were separately exposed to clinostat microgravity simulation conditions for 24 h.

Materials and Methods

Cells. The human neuroblastoma cell line NB-1 (RBC1953),⁽¹⁶⁾ purchased from the Riken Bioresource Center Cell Bank (Ibaraki, Japan), was used in this study. The *p53* gene of these cells has a G215C polymorphism, a common polymorphism substituting an arginine for a proline at codon 72.⁽¹⁷⁾ Cells were maintained in Dulbecco's modified Eagle's medium (DMEM; Sigma-Aldrich Inc., St. Louis, MO) supplemented with 10% fetal bovine serum (FBS; Hyclone, Logan, UT) at 37°C in humidified air containing 5% CO₂.

X-irradiation. Each flask of cells was irradiated with a dose of 0.1 or 1.0 mGy at room temperature using a Hitachi MBR-1505R X-ray generator (Hitachi, Tokyo, Japan). The machine was operated at 120 kVp and at either 0.2 mA with a filter of Al 1.0 mm + Cu 1.7 mm to achieve a dose of 0.1 mGy, or at 0.2 mA with a filter of Al 1.0 mm + Cu 0.2 mm to achieve a dose of 1.0 mGy. The dose rates were 0.59 mGy/min (9.84 nGy/s) and 3.69 mGy/min (61.5 nGy/s) for doses of 0.1 and 1.0 mGy, respectively, at a focus-surface distance of 60 cm in all cases. The doses were measured by Radiation Monitor (10x5-60, 9015; Radical Corporation, Monrovia, CA).

Clinostat treatment. Each flask of cells was subjected to microgravity treatment using a clinostat (Portable Microgravity Simulator PMS-1; Advanced Engineering Services Co., Ltd.). The rotation rates around the X- and Y-axes were 4.8 and 7.2 rpm, respectively. The machine operates to give theoretically 0 g to cells for 24 h.

Isolation of the total RNA. Total RNA was isolated using ISOGEN from cultured cells as recommended by the manufacturer (Nippon gene Toyama, Japan). Briefly, after washing cells by PBS 3 times, 1 ml of ISOGEN was added to cells and collected into 1.5 ml tubes. Two hundred μ l of chloroform was added to the samples and vortexed. After centrifugation at 12,000 g for 15 min at 4°C, the aqueous phase was transferred to new 1.5 ml tubes and 0.5 ml of 2-propanol was added. After incubation for 5 min at R.T., centrifugation was performed at 12,000 g for 10 min at 4°C to precipitate RNA. The pellets were washed with 70% of EtOH and after centrifugation 7,500 g for 5 min at 4°C, pellets were dried briefly and dissolved in TE. The RNA quality was checked by measuring 260 nm absorbance and electrophoresis. All cDNAs were prepared by reverse transcription of 1 μ g total RNA using oligo dT(20) primer (0.4 μ M/50 μ l final volume), and ReverTra Ace (TOYOBO) as recommended by the manufacturer. An equivalent volume of 0.1 μ l of cDNA solution was used for quantification of specific cDNA by qRT-PCR.

SYBR Green-based qRT-PCR. Total RNA was isolated from NB-1 cells and cDNA was synthesized as described above. The sequences of primers were compared to those from the available human genome and the EST database (<http://blast.genome.jp/>) in order to select primers that would produce a single amplification product. The forward and reverse primers are listed in Table 1. The qRT-PCR assays were performed on an ABI prism 7000 Sequence Detection System (Applied Biosystems, Foster City, CA) using the QuantiTect SYBR Green PCR Kit (Qiagen, Valencia, CA). Four sets of PCR profiles were used, depending on the primer sets (conditions A–D, Table 1). The experimental conditions were those recommended by the manufacturer, except for the annealing temperature, annealing time, elongation temperature, and elongation time; specifically, the differences were as follows: condition A, annealing and elongation 60°C 1 min; condition B, annealing 55°C 30 s and elongation 72°C 40 s; condition C, annealing 62°C 5 s and elongation 72°C 30 s; and condition D, annealing 55°C 5 s and elongation 72°C 30 s. All PCR assays

were performed for 40 cycles. The size of single amplification products was further verified by gel electrophoresis. All data were normalized to an internal standard (glyceraldehyde-3-phosphate dehydrogenase; GAPDH). Three triplicate samples were used in an assay of qRT-PCR and repeated for three times. The average values were calculated and the bar was expressed as SD.

The NB-1 cells were irradiated with X-rays or were subjected to microgravity for 24 h prior to qRT-PCR gene expression analysis of cell death-related genes. The list of the genes is shown in Table 2. To examine the activation of apoptosis, the following genes were targeted: Bax (*BAX*); cytochrome *c* (*CYTC*); apoptotic protease activating factor 1 (*APAF1*); voltage-dependent anion channels (*VDACs*) 1, 2 and 3 (*VDAC1*, *2 and 3*); caspases 3, 8 and 9, (*CASP3*, *8 and 9*); p53 (*P53*); apoptosis-inducing factor (*AIF*); and adenine nucleotide translocators (*ANT*) 1 and 2 (*ANT1 and 2*). In addition, three more gene groups were targeted to examine the following. Apoptosis suppression: B-cell CLL/lymphoma 2 (*BCL2*) and manganese superoxide dismutase (*MnSOD*), autophagy related gene, Beclin1 (*BECN1*), and necrosis related gene, cyclophilin 40 (*CYP-40*).

Cells were exposed to single doses of radiation of 0.1 and 1 mGy, and analyses were performed 30 min and 2 h after irradiation. To evaluate the effects of long-term, low-dose exposure, cells were irradiated with doses of 0.1 and 1 mGy 10 times in 1 h-intervals ($\times 10$ [1 h]) and analyses were performed immediately, 30 min, and 2 h after irradiation. Cells were also irradiated at 2 h-intervals ($\times 10$ [2 h]) and analyses were performed immediately, 30 min, and 2 h after irradiation. The effects of microgravity, which was applied over a 24 h period using the clinostat, were also evaluated immediately after the procedure. Analyses of cell death-related genes were subsequently conducted.

Statistical analysis. Statistical analyses were performed using Student's *t* test. All *p* values less than 0.05 were considered to be statistically significant. Data are presented as the mean \pm SD. Calculations were carried out using Microsoft Excel (Microsoft Corporation, Redmond, WA) on a MacBook Air (Apple Inc. Cupertino, CA).

Results

NB-1 cells were either subjected to microgravity for 24 h or irradiated with extremely low 0.1 and 1.0 mGy single or 0.1 mGy 10 times fractionated X-ray irradiation. The expression levels of cell death-related genes were then analyzed by qRT-PCR assay (Table 2). As shown in Table 3, microgravity treatment resulted in increased expression of *VDAC2* and decreased expression of *MnSOD*, while *CASP9* was also downregulated. Other genes investigated did not change significantly after a 24 h period of exposure of the cells to microgravity. On the other hand, 0.1 mGy single irradiation significantly decreased the expression of *CYTC*, *APAF1*, *VDAC1–3*, *CASP3*, *P53*, *AIF*, *ANT1,2*, *BCL2*, *MnSOD*, autophagy related *BECN1* and necrosis related *CYP-40*. 1.0 mGy single irradiation significantly downregulated *BAX*, *VDAC2*, *ANT1 and 2*, *BECN1* and *CYP-40*. 0.1 mGy fractionated irradiation increased the expression of *APAF1* and *BCL2* at both 1 and 2 h intervals. The expression of other genes (*CYTC*, *VDAC2 and 3*, *AIF*, *ANT1*, *MnSOD*, and *CYP-40*) was significantly downregulated at both 1 and 2 h intervals. For three of the genes (*CASP8*, *ANT2* and *BECN1*) expression was decreased at only the 1 h intervals, with no changes observed in the 2 h interval regime. A significantly downregulation of *BAX* was shown at the 2 h interval regime.

Discussion

The effects of 0.1 and 1.0 mGy irradiation and 24 h microgravity treatment on gene expression were examined using a human neuroblastoma cell line, NB-1, with active neuronal func-

Table 1. Primer sequences for genes examined in this study

Gene Group	Primer name	Primer sequence	PCR conditions
Apoptosis activation	BAX-F	5'-TTGGGTGAGACTCCTCAAGCCTCC-3'	C
	BAX-R	5'-TCTGAAGATGGGGAGAGGGCACCA-3'	
	CYTC-F	5'-GGAGAGGATACACTGATGGAGTATTTGGA-3'	A
	CYTC-R	5'-GAGATAAGCTATTAAGTCTGCCCTTCTTC-3'	
	APAF1-F	5'-ATATTAAGTGGTGGAAACGTTGCTACTGGG-3'	A
	APAF1-R	5'-AGTCACATATGTTTGAAGTCAGGGGACAC-3'	
	VDAC1-F	5'-GGATACACTCAGACTCTTAAAGCCAGG-3'	A
	VDAC1-R	5'-ATGCTTGAAATTCAGTTCCTAGACCAAGC-3'	
	VDAC2-F	5'-TTGGATCCCACTGCTCCATTCTGCAAA-3'	D
	VDAC2-R	5'-ATTAATGCTCTCCCATCTACCAGAGCAGA-3'	
	VDAC3-F	5'-AAATAATGCCAGCCTGATTGGACTGGGT-3'	A
	VDAC3-R	5'-TTCAAATCCAAGCCAACCTTGACCT-3'	
	CASP3-F	5'-CAGAGGGGATCGTTGTAGAAGTCT-3'	B
	CASP3-R	5'-CGGCCTCCACTGGTATTTTATGAC-3'	
	CASP8-F	5'-AGAGCGATGCTCTCGAGGCGATGATATT-3'	A
	CASP8-R	5'-AAGTAGGCTGAGGCATCTGTTCCCAT-3'	
	CASP9-F	5'-CAAGAGTGGCTCCTGTACGTTGAGA-3'	A
	CASP9-R	5'-CTGTTTATAAATCCCTTTCACCGAAACAGC-3'	
	P53-F	5'-TCAGTCTGAGTCAGGCCCTTCTGT-3'	B
	P53-R	5'-AGGCCTTGGAACTCAAGGATGCCC-3'	
AIF-F	5'-CAAAGGTGTCATCTTACCTCAGGGAC-3'	A	
AIF-R	5'-CTCACCGTCCTTAATGATCTTCTTGTCT-3'		
ANT1-F	5'-TTCTTAGGGGGTGTGGATCGGCATAA-3'	A	
ANT1-R	5'-CACATCAGCAGCCAACCTGGTCTTA-3'		
ANT2-F	5'-ACTGTCAGTCTGTTGCCGGTTGA-3'	A	
ANT2-R	5'-AAAAGCTTTGCCTCCTTCATCACGAGCA-3'		
Apoptosis suppression	BCL2-F	5'-GGAGGCTGGGATGCCTTTGTGGAA-3'	C
	BCL2-R	5'-TAGGCACCCAGGGTGATGCAAGCT-3'	
	MnSOD-F	5'-TTCTGGACAAACCTCAGCCCTAACGGT-3'	A
MnSOD-R	5'-AACAGATGCAGCCGTGAGCTTCTCCTAAA-3'		
Autophagy	BECN1-F	5'-AGTGGCGGCTCCTATTCCATCAAAC-3'	A
	BECN1-R	5'-AAGACCCCACTTAAGATTGCTGAGCATGAA-3'	
Necrosis	CYP-40-F	5'-CAGAGCTCAAGGGATGGCAAGGATTAAG-3'	A
	CYP-40-R	5'-GACTTTCAGCAATTCTGCCTGGATAGCTTT-3'	
Internal control	GAPDH-F	5'-GGGCAAGGTCATCCCTGAGCTGAA-3'	A, B, C, D
	GAPDH-R	5'-TCTAGACGGCAGGTCAGGTCCACC-3'	
Condition A	Annealing and elongation: 60°C 1 min		
Condition B	Annealing: 55°C 30 s, elongation: 72°C 40 s		
Condition C	Annealing: 62°C 5 s, elongation: 72°C 30 s		
Condition D	Annealing: 55°C 5 s, elongation: 72°C 30 s		

tion.^(16,17) The radiation doses used in this study (0.1 or 1.0 mGy) are considered extremely low-dose. The surviving fraction after 2 Gy (2,000 mGy), which is the daily dose in conventional radiation therapy, ranges from 0.35–0.62.^(18,19) In conventional radiation therapy, patients receive 2 Gy × 30 times in 6 weeks, totally 60 Gy to treat cancer. Therefore, the endpoint to evaluate such extremely dose, 0.1 and 1.0 mGy is limited to molecular methods, such as qRT-PCR.

Experimental microgravity data. DNA microarray technology is a powerful tool for identifying “space genes” that play key roles in cellular responses to microgravity.^(20–23) In animals, the hindlimb suspension model simulates space flight models and is well tolerated by the animals with minimal evidence of stress.⁽²⁴⁾ Frigeri and co-workers investigated the effects of microgravity on gene expression in the mouse brain by using 2-week-old hindlimb-unloaded mice.⁽²³⁾ They found that the TIC class of genes (i.e., the class responsible for the transport of small molecules and ions into cells) had the highest percentage of upregulation, including that of *VDAC1*, whereas the most commonly downregulated genes were

those of the JAE class [cell junction, adhesion, and extracellular matrix (ECM)].⁽²³⁾ Schatten *et al.*⁽²⁵⁾ reported that space flight and clinorotation causes apoptosis. However, this study showed relatively no change of apoptosis elements. In this study, microgravity conditions increased expression of apoptotic *VDAC2*⁽²⁶⁾ and decreased expression of antiapoptotic *MnSOD*,^(27,28) while *CASP9* was downregulated (Table 2 and 3), suggesting that these cells may be more susceptible to further oxidative stress.

Wang and Good⁽²⁹⁾ demonstrated the similarity between old age and prolonged exposure to microgravity (i.e., in terms of heart, muscle, and bone atrophy), and emphasized that determining the underlying signal transduction pathways (e.g., the control mechanisms for the processes of proliferation, differentiation, and apoptosis) remains important for creating therapeutic modalities to retard, reduce, and prevent tissue atrophy, both in space and on earth. Experiments examining the differentiation of neuron-like PC12 and SH-SY5Y cells revealed less neurite extension under conditions of microgravity created by a rotating wall bioreactor (RWB) than that observed in static cultures, suggesting that less-

Table 2. Cell death-related genes analyzed by qRT-PCR

Gene group related to apoptosis activation	
Gene	Description
<i>BAX</i>	Bax
<i>CYTC</i>	Cytochrome c
<i>APAF1</i>	Apoptotic protease activating factor 1
<i>VDAC1</i>	Voltage-dependent anion channel 1
<i>VDAC2</i>	Voltage-dependent anion channel 2
<i>VDAC3</i>	Voltage-dependent anion channel 3
<i>CASP3</i>	Caspase 3
<i>CASP8</i>	Caspase 8
<i>CASP9</i>	Caspase 9
<i>P53</i>	p53
<i>AIF</i>	Apoptosis-inducing factor
<i>ANT1</i>	Adenine nucleotide translocator 1
<i>ANT2</i>	Adenine nucleotide translocator 2
Gene group related to apoptosis suppression	
Gene	Description
<i>BCL2</i>	B-cell CLL/lymphoma 2
<i>MnSOD</i>	Manganese superoxide dismutase (SOD2)
Gene group related to autophagy	
Gene	Description
<i>BECN1</i>	Beclin1
Gene group related to necrosis	
Gene	Description
<i>CYP-40</i>	Cyclophilin 40 (Cyclophilin D)

than-optimal differentiation occurs under conditions of simulated microgravity relative to normal gravity.⁽²⁹⁾

Stein *et al.*⁽³⁰⁾ found that energy intake, as well as protein synthesis, decreased in a correlated manner during extended (>3 months) flights on the Russian space station, MIR. Post-flight energy intake and protein synthesis returned to, but did not exceed, the pre-flight levels. Stein and Leskiw further examined oxidative damage, urinary excretion of F2 isoprostane (isoprostane), 8-isoprostaglandin F2a (PG), and 8-oxo-7,8-dihydro-2 deoxyguanosine (8-OHdG), during and after space flight.⁽³¹⁾ Samples were obtained from MIR space flight participants between 88 and 186 days in orbit. They concluded that oxidative damage was decreased in-flight, and oxidative damage was increased post-flight.⁽³¹⁾ While the putative increase in oxidative stress associated with space flight may persist after a return to normal gravity, this effect appears to be more pronounced after an extended period of time spent in space.⁽³²⁾ However, Hollander *et al.*⁽³³⁾ reported that space flight itself was associated with the downregulation of antioxidant defense systems in the rat liver. Space flight (STS-63 for 8 days, $n = 6$, male Sprague-Dawley rats) significantly reduced catalase, GSH reductase, and GSH sulfur-transferase activity in the liver. The relative levels of mRNA for Cu-Zn SOD and catalase were also significantly reduced by space flight. In addition, space flight dramatically reduced hepatic GSH, glutathionedisulfide, and total GSH contents, and these reductions were accompanied by reduced γ -glutamyl transpeptidase activity. Furthermore, space-flight rats exhibited an increase in the liver malondialdehyde concentration. Taken together, these results indicate that space flight can induce a downregulation of the antioxidant defense capacity and promote oxidative stress in the liver.⁽³³⁾ Degan *et al.*⁽³⁴⁾ reported that exposure to simulated microgravity reduced the intracellular ATP concentration in freshly drawn lymphocytes and lymphoblastoid cells (LB and COR3) by about 40–50% of the level found under normal growth conditions. Poly(ADP-ribose) polymerase (PARP) activity increased, indicating that cells exposed to reduced gravity

Table 3. Results of gene expression analysis by qRT-PCR

Apoptosis activation	<i>BAX</i>	<i>CYTC</i>	<i>APAF1</i>	<i>VDAC1</i>	<i>VDAC2</i>	<i>VDAC3</i>	<i>CASP3</i>
Microgravity	0.94 ± 0.31	1.22 ± 0.31	1.04 ± 0.27	1.03 ± 0.30	1.25 ± 0.10 [¶]	1.02 ± 0.13	1.06 ± 0.16
0.1 mGy × 1–0.5 h	0.99 ± 0.15	0.70 ± 0.12 [§]	0.63 ± 0.16 [§]	0.57 ± 0.18 [§]	0.58 ± 0.13 [§]	0.51 ± 0.07 [§]	0.79 ± 0.07 [§]
0.1 mGy × 1–2 h	0.83 ± 0.33	0.66 ± 0.14 [§]	0.67 ± 0.23 [§]	0.63 ± 0.12 [§]	0.45 ± 0.09 [§]	0.50 ± 0.11 [§]	0.89 ± 0.13
1 mGy × 1–0.5 h	0.65 ± 0.19	0.88 ± 0.26	0.95 ± 0.51	0.96 ± 0.23	0.20 ± 0.05 [§]	1.03 ± 0.17	0.81 ± 0.14
1 mGy × 1–2 h	0.55 ± 0.12 [§]	1.02 ± 0.28	0.78 ± 0.27	0.85 ± 0.16	0.76 ± 0.26	0.98 ± 0.21	0.85 ± 0.11
0.1 mGy-1 h × 10–0 h	0.93 ± 0.21	0.78 ± 0.15	1.35 ± 0.34 [¶]	0.80 ± 0.17	0.42 ± 0.18 [§]	0.37 ± 0.11 [§]	0.82 ± 0.20
0.1 mGy-1 h × 10–0.5 h	0.91 ± 0.16	1.03 ± 0.24	1.49 ± 0.33 [¶]	1.18 ± 0.30	0.33 ± 0.08 [§]	0.53 ± 0.04 [§]	0.87 ± 0.14
0.1 mGy-1 h × 10–2 h	1.04 ± 0.40	0.71 ± 0.19 [§]	1.14 ± 0.19	0.87 ± 0.18	0.45 ± 0.22 [§]	0.55 ± 0.06 [§]	0.89 ± 0.18
0.1 mGy-2 h × 10–0 h	0.72 ± 0.27	1.00 ± 0.18	1.26 ± 0.26 [¶]	1.11 ± 0.13	0.74 ± 0.18 [§]	0.72 ± 0.09 [§]	0.83 ± 0.13
0.1 mGy-2 h × 10–0.5 h	1.09 ± 0.25	0.94 ± 0.15	1.66 ± 0.41 [¶]	0.94 ± 0.14	0.47 ± 0.21 [§]	0.64 ± 0.06 [§]	0.92 ± 0.21
0.1 mGy-2 h × 10–2 h	0.70 ± 0.16 [§]	0.71 ± 0.13 [§]	1.17 ± 0.29	0.93 ± 0.09	0.54 ± 0.34 [§]	0.63 ± 0.06 [§]	0.86 ± 0.13
Apoptosis activation	<i>CASP8</i>	<i>CASP9</i>	<i>P53</i>	<i>AIF</i>	<i>ANT1</i>	<i>ANT2</i>	
Microgravity	0.65 ± 0.20	0.73 ± 0.36 [§]	0.94 ± 0.17	0.91 ± 0.22	1.10 ± 0.34	1.04 ± 0.14	
0.1 mGy × 1–0.5 h	0.98 ± 0.37	0.91 ± 0.16	0.65 ± 0.11 [§]	0.66 ± 0.14 [§]	0.74 ± 0.18 [§]	0.76 ± 0.18 [§]	
0.1 mGy × 1–2 h	1.10 ± 0.43	0.88 ± 0.19	0.71 ± 0.16 [§]	0.69 ± 0.14 [§]	0.79 ± 0.22 [§]	0.68 ± 0.16 [§]	
1 mGy × 1–0.5 h	1.00 ± 0.40	1.00 ± 0.45	1.10 ± 0.19	0.97 ± 0.28	1.06 ± 0.31	0.81 ± 0.12 [§]	
1 mGy × 1–2 h	0.83 ± 0.14	1.03 ± 0.43	1.22 ± 0.42	0.92 ± 0.13	0.77 ± 0.16 [§]	0.75 ± 0.28 [§]	
0.1 mGy-1 h × 10–0 h	0.76 ± 0.33 [§]	0.84 ± 0.38	0.79 ± 0.21 [§]	0.66 ± 0.14 [§]	0.75 ± 0.17 [§]	0.87 ± 0.10 [§]	
0.1 mGy-1 h × 10–0.5 h	0.85 ± 0.35 [§]	0.79 ± 0.24	0.87 ± 0.17	0.77 ± 0.21 [§]	0.78 ± 0.18 [§]	0.79 ± 0.22 [§]	
0.1 mGy-1 h × 10–2 h	0.65 ± 0.13 [§]	0.86 ± 0.45	0.92 ± 0.18	0.87 ± 0.28 [§]	0.83 ± 0.23 [§]	0.84 ± 0.17	
0.1 mGy-2 h × 10–0 h	1.15 ± 0.16	0.87 ± 0.32	1.01 ± 0.32	0.71 ± 0.13 [§]	0.82 ± 0.16 [§]	0.99 ± 0.23	
0.1 mGy-2 h × 10–0.5 h	0.93 ± 0.13	0.85 ± 0.25	1.13 ± 0.13	0.82 ± 0.12 [§]	0.77 ± 0.10 [§]	0.95 ± 0.19	
0.1 mGy-2 h × 10–2 h	1.03 ± 0.11	0.86 ± 0.40	0.99 ± 0.15	0.95 ± 0.11	0.82 ± 0.19	0.86 ± 0.20	

[§]Decreased significantly by *t* test ($p < 0.05$). [¶]Increased significantly by *t* test ($p < 0.05$).

are stressed. Maillet *et al.*⁽³⁵⁾ reviewed microgravity in terms of an accelerated model of nutritional disturbances, and concluded that adopting an integrated approach will be essential for optimizing the health of astronauts. They also found that current dietary approaches for adults exposed to normal gravity are inadequate for preventing significant changes in the nutritional status of astronauts, and they suggested areas for further research in both fields of ground and space medicine. It is noted that since space experiments consist of microgravity and space radiation conditions, these findings might be caused by space radiation and not by microgravity.

Experimental data from studies of exposure to time spent in space and extremely low-dose radiation. In this study, 0.1 mGy single dose irradiation resulted in significant downregulation of *CYTC*, *APAF1*, *VDAC1-3*, *CASP3*, *P53*, *AIF*, *ANT1 and 2*, *BCL2*, *MnSOD*, autophagy related *BECN1* and necrosis related *CYP-40* (Table 3 and 4). However, 1.0 mGy single irradiation revealed significant downregulation in *BAX*, *VDAC2*, *ANT1 and 2*, autophagy related *BECN1*⁽³⁶⁾ and necrosis related *CYP-40*⁽³⁷⁾ (Table 3 and 4), suggesting the effects of 0.1 and 1.0 Gy single dose irradiation differentially affect the cells. In the extremely low 0.1 mGy dose fractionated irradiation group, proapoptotic *APAF1* significantly increased. On the other hand, antiapoptotic *BCL2*⁽³⁸⁾ significantly increased, and downregulation of proapoptotic *CYTC*, *VDAC2 and 3*, *CASP8*, *AIF*, *ANT1 and 2* was observed. These results demonstrate changes in the expression profiles of apoptosis-related genes in NB-1 cells subjected to low-dose radiation, with a shift towards an inhibition of apoptosis. Radiation-induced apoptosis is considered to be p53-dependent.⁽³⁹⁻⁴⁸⁾ In this study, the level of *P53* did not show significant change. In addition, the levels of *CASP3* and *CASP9* did not show change. Taken together, these results suggest that long-term, low-dose radiation exposure may inhibit apoptosis, and that non-apoptotic death pathways may also not be affected by such exposure due to downregulation of *BECN1* and *CYP-40* (Table 4). However, the level of *MnSOD*, which suppresses apoptosis, was downregulated in the fractionated regimes, suggesting the possibility of apoptosis. These results may indicate extremely low dose 0.1 mGy X-irradiation fractionation regimes do not promote cell death, but may make cells susceptible to further oxidative stress due to *MnSOD* downregulation.

Few studies have investigated the effects of extremely low-dose irradiation. Suzuki and coworkers found that extremely low-dose X-ray irradiation (20–50 mGy) stimulated the proliferation of human cultured cells and the phosphorylation of extracellular signal-regulated kinase.⁽⁴⁹⁾ Ding and coworkers applied cDNA microarray analyses to G1-arrested normal human skin fibroblasts subjected to X-ray irradiation (low dose, 20 mGy; high dose, 4 Gy).⁽⁵⁰⁾ The predominant cell functions affected by low-dose

radiation were those involved in cell-cell signaling, signal transduction, cell development, and DNA damage responses. At high-doses, the responding genes were involved in apoptosis and cell proliferation. Interestingly, several genes (i.e., the cytoskeleton components anillin (*ANLN*; actin binding protein), and keratin 15 (*KRT15*) and the cell-cell signaling genes Growth factor receptor-bound protein 2 (GRB2)-related adapter protein 2 (*GRAP2*) and G-protein-coupled receptor 51 (*GPR51*) responded to low-dose radiation, but not to high-dose radiation. Pathways that are specifically activated by low-dose radiation were also evident. The researchers concluded that these quantitative and qualitative differences in gene expression may help explain the non-linear correlation of biological effects of ionizing radiation ranging from low doses to high doses.⁽⁵⁰⁾ Moreover, Ogura and coworkers found that the mutation frequency in a 500 μ Gy-irradiated group of *Drosophila melanogaster* was significantly lower than that of the control group, whereas in the 10 Gy-irradiation group, the mutation frequency was significantly higher than that of the control group.⁽⁵¹⁾ They suggested the apparent upregulation of a positive regulator of apoptosis immediately after irradiation with 500 μ Gy, furthermore suggesting that the linear non-threshold (LNT) model for stochastic effects of ionizing radiation was not applicable. Thus, their results will be relevant to further discussions of the LNT model.

Kiefer⁽⁵²⁾ addressed some pertinent questions related to the assessment of radiation risk to humans in space. The effects remain to be clarified for low-dose rates of γ -rays, protons, and heavy particles, which dominate in the case of space-acquired radiation. It will be necessary to conduct experiments in space in order to determine the combined effects of space radiation and microgravity. Ohnishi and coworkers⁽⁵³⁾ discussed cellular risk as well as molecular events following exposure to space radiation, and they have advocated for prioritizing space-radiation research. Gridley *et al.*⁽⁵⁴⁾ conducted a RT-PCR assay to investigate the effects of pre-exposure (“Pre”) of low-dose/low-dose-rate photons (⁵⁷Co, total of 0.049 Gy at 0.24 mGy/h) on gene expression in the 2 Gy whole-body irradiated C57BL/6 mouse liver. There were various genes downregulated in all groups studied (13 in the “2 Gy-alone”, 16 in the “Pre”, and 16 in the “Pre + 2 Gy” groups). These results suggest that exposure to only 0.049 Gy at 0.24 mGy/h can alter gene expression.

Space radiation may increase oxidative stress by reducing levels of antioxidants. Manda *et al.*⁽⁵⁵⁾ tested the effects of high-linear energy transfer (LET) ⁵⁶Fe beams (500 MeV/nucleon, 1.5 Gy) on memory impairment and apoptosis using male C57BL mice. High-LET radiation substantially impaired the reference memory of mice 30 days after irradiation, whereas no significant effect was observed on the motor activities of the mice. The memory dysfunction caused by irradiation was attenuated by α -lipoic acid.

Table 4. Results of gene expression analysis by qRT-PCR (continued)

Apoptosis suppression	<i>BCL2</i>	<i>MnSOD</i>	Autophagy	<i>BECN1</i>	Necrosis	<i>CYP-40</i>
Microgravity	1.07 ± 0.17	0.67 ± 0.18 [§]	Microgravity	0.87 ± 0.11	Microgravity	0.88 ± 0.14
0.1 mGy × 1–0.5 h	0.75 ± 0.19 [§]	0.61 ± 0.07 [§]	0.1 mGy × 1–0.5 h	0.83 ± 0.19 [§]	0.1 mGy × 1–0.5 h	0.74 ± 0.10 [§]
0.1 mGy × 1–2 h	1.04 ± 0.32	0.64 ± 0.18 [§]	0.1 mGy × 1–2 h	0.80 ± 0.20 [§]	0.1 mGy × 1–2 h	0.72 ± 0.11 [§]
1 mGy × 1–0.5 h	1.04 ± 0.23	0.94 ± 0.25	1 mGy × 1–0.5 h	0.73 ± 0.16 [§]	1 mGy × 1–0.5 h	0.86 ± 0.10 [§]
1 mGy × 1–2 h	0.85 ± 0.29	1.06 ± 0.33	1 mGy × 1–2 h	0.84 ± 0.12 [§]	1 mGy × 1–2 h	0.81 ± 0.09 [§]
0.1 mGy-1 h × 10–0 h	1.37 ± 0.19 [¶]	0.50 ± 0.13 [§]	0.1 mGy-1 h × 10–0 h	0.82 ± 0.11 [§]	0.1 mGy-1 h × 10–0 h	0.74 ± 0.13 [§]
0.1 mGy-1 h × 10–0.5 h	2.38 ± 0.39 [¶]	0.65 ± 0.10 [§]	0.1 mGy-1 h × 10–0.5 h	0.62 ± 0.09 [§]	0.1 mGy-1 h × 10–0.5 h	0.78 ± 0.16
0.1 mGy-1 h × 10–2 h	1.33 ± 0.21 [¶]	0.79 ± 0.15 [§]	0.1 mGy-1 h × 10–2 h	0.83 ± 0.15 [§]	0.1 mGy-1 h × 10–2 h	0.56 ± 0.17 [§]
0.1 mGy-2 h × 10–0 h	1.56 ± 0.36 [¶]	0.75 ± 0.14 [§]	0.1 mGy-2 h × 10–0 h	0.96 ± 0.08	0.1 mGy-2 h × 10–0 h	0.73 ± 0.34 [§]
0.1 mGy-2 h × 10–0.5 h	1.72 ± 0.49 [¶]	0.80 ± 0.13 [§]	0.1 mGy-2 h × 10–0.5 h	0.97 ± 0.06	0.1 mGy-2 h × 10–0.5 h	0.71 ± 0.30 [§]
0.1 mGy-2 h × 10–2 h	1.57 ± 0.18 [¶]	0.82 ± 0.11	0.1 mGy-2 h × 10–2 h	1.07 ± 0.14	0.1 mGy-2 h × 10–2 h	0.75 ± 0.36 [§]

[§]Decreased significantly by *t* test ($p < 0.05$). [¶]Increased significantly by *t* test ($p < 0.05$).

Radiation-induced apoptotic and necrotic cell death of granule and Purkinje cells was also significantly inhibited by pretreatment with α -lipoic acid.⁽⁵⁵⁾ The effects of single whole-body irradiation of 3 Gy by γ -rays, 3 Gy by 1,000 MeV protons, or 50 cGy (500 mGy) by 1 GeV/nucleon iron ions were examined in terms of the total antioxidant status (TAS) of male CBA mice, and the protective effects of diet supplementation with Bowman-Birk Inhibitor Concentrate (BBIC), L-selenomethionine (L-SeM), or a combination of N-acetyl cysteine, sodium ascorbate, coenzyme Q10 (CoQ10), α -lipoic acid, and vitamin E succinate were also examined.^(56,57) Guan *et al.*⁽⁵⁷⁾ found that BBIC, L-SeM, and certain antioxidant combinations that include CoQ10 exhibit protective effects, suggesting that antioxidants act as protectors against adverse biological effects induced by space radiation. Our qRT-PCR results regarding changes in cell death-related gene expression revealed that single doses of 0.1 mGy, as well as fractionated dose regimes, downregulated apoptosis-related genes (*CYTC*, *VDAC2 and 3*, *AIF*, *ANT1 and 2*). However, downregulation of antioxidant gene (*MnSOD*) and increased apoptosis related gene (*APAF1*) expression may indicate increased risk by extremely low dose irradiation.

Conclusion

The qRT-PCR examination of cell death-related gene expression changes revealed that microgravity treatment increased apoptosis related gene *VDAC2*, but reduced the expression levels of apoptosis-related gene *CASP9* and the antioxidant gene *MnSOD* using a neuroblastoma cell line NB-1. After 0.1 mGy

single irradiation, most of the genes examined were down-regulated, while 1.0 mGy single irradiation revealed significant downregulation in only six genes (*BAX*, *VDAC2*, *ANT1 and 2*, *BECN1* and *CYP-40*), suggesting 0.1 and 1.0 mGy radiation have different effects on cells. After 0.1 mGy fractionated irradiation, apoptotic *APAF1* and anti-apoptotic *BCL2* were upregulated, and apoptosis related *CYTC*, *VDAC2 and 3*, *CASP8*, *AIF*, *ANT1* genes were significantly downregulated. These results suggest decreased risk of apoptosis. However, downregulation of *MnSOD* was also observed. These results may indicate that space environment including microgravity and space radiation may increase oxidative stress by reducing *MnSOD*.

Acknowledgments

The authors would like to express their appreciation to Dr. Aaron K. Holley for the helpful discussion. This work was supported in part by a Grant-in-Aid for Scientific Research (C) (No. 22592093 to H.J.M. and No. 23592774 to H.P.I.) and the Strategic Promotion Program for Basic Nuclear Research (to H.J.M.) of the Ministry of Education, Culture, Sports, Science and Technology of Japan; and the Program for Promotion of Basic and Applied Researches for Innovations in Bio-oriented Industry (BRAINI) (to H.J.M.) by the Ministry of Agriculture, Forestry and Fisheries of Japan.

Conflict of Interest

No potential conflicts of interest were disclosed.

References

- Yasuda H. Effective dose measured with a life size human phantom in a low Earth orbit mission. *J Radiat Res* 2009; **50**: 89–96.
- Majima HJ, Indo HP, Tomita K, *et al.* Bio-assessment of risk in long-term manned space exploration—cell death factors in space radiation and/or microgravity: a review—. *Biol Sci Space* 2009; **23**: 43–53.
- Kennedy AR. Biological effects of space radiation and development of effective countermeasures. *Life Sci Space Res (Amst)* 2014; **1**: 10–43.
- Cucinotta FA. Review of NASA approach to space radiation risk assessments for Mars exploration. *Health Phys* 2015; **108**: 131–142.
- Sridharan DM, Asaithamby A, Bailey SM, *et al.* Understanding cancer development processes after HZE-particle exposure: roles of ROS, DNA damage repair and inflammation. *Radiat Res* 2015; **183**: 1–26.
- Yatagai F, Ishioka N. Are biological effects of space radiation really altered under the microgravity environment? *Life Sci Space Res* 2014; **3**: 76–89.
- Suzuki M, Watanabe M, Kanai T, *et al.* LET dependence of cell death, mutation induction and chromatin damage in human cells irradiated with accelerated carbon ions. *Adv Space Res* 1996; **18**: 127–136.
- Suzuki M, Kase Y, Kanai T, Ando K. Correlation between cell death and induction of non-rejoining PCC breaks by carbon-ion beams. *Adv Space Res* 1998; **22**: 561–568.
- Durante M, Yamada S, Ando K, *et al.* X-rays vs. carbon-ion tumor therapy: cytogenetic damage in lymphocytes. *Int J Radiat Oncol Biol Phys* 2000; **47**: 793–798.
- Hada M, Chappell LJ, Wang M, George KA, Cucinotta FA. Induction of chromosomal aberrations at fluences of less than one HZE particle per cell nucleus. *Radiat Res* 2014; **182**: 368–379.
- Ono K, Masunaga S, Akaboshi M, Akuta K. Estimation of the initial slope of the cell survival curve after irradiation from micronucleus frequency in cytokinesis-blocked cells. *Radiat Res* 1994; **138** (1 Suppl): S101–S104.
- Balajee AS, Bertucci A, Taveras M, Brenner DJ. Multicolour FISH analysis of ionising radiation induced micronucleus formation in human lymphocytes. *Mutagenesis* 2014; **29**: 447–455.
- Drooger JC, Hoening MJ, Seynaeve CM, *et al.* Diagnostic and therapeutic ionizing radiation and the risk of a first and second primary breast cancer, with special attention for BRCA1 and BRCA2 mutation carriers: A critical review of the literature. *Cancer Treat Rev* 2015; **41**: 187–196.
- Miller RC, Randers-Pehrson G, Geard CR, Hall EJ, Brenner DJ. The oncogenic transforming potential of the passage of single alpha particles through mammalian cell nuclei. *Proc Natl Acad Sci U S A* 1999; **96**: 19–22.
- Valentin J. Low-dose extrapolation of radiation-related cancer risk. *Ann ICRP* 2005; **35**: 1–140.
- Kunimoto M. Possible involvement of the 440 kDa isoform of ankyrinB in neurogenesis in human neuroblastoma NB-1 cells. *FEBS Lett* 1995; **357**: 217–220.
- Klug SJ, Rensing M, Koenig J, *et al.* TP53 codon 72 polymorphism and cervical cancer: a pooled analysis of individual data from 49 studies. *Lancet Oncol* 2009; **10**: 772–784.
- Majima HJ, Suzuki M, Ando K, *et al.* Radiation biology. Theoretical strategy: how to make treatment planning for heavy ion-beam therapy? *Jpn J Med Phys* 1999; **19**: 37–47.
- Motoori S, Majima HJ, Ebara M, *et al.* Overexpression of mitochondrial manganese superoxide dismutase protects against radiation-induced cell death in the human hepatocellular carcinoma cell line HLE. *Cancer Res* 2001; **61**: 5382–5388.
- Wang E. Age-dependent atrophy and microgravity travel: what do they have in common?. *FASEB J* 1999; **13** Suppl: S167–S174.
- Ward NE, Pellis NR, Risin SA, Risin D. Gene expression alterations in activated human T-cells induced by modeled microgravity. *J Cell Biochem* 2006; **99**: 1187–1202.
- Patel MJ, Liu W, Sykes MC, *et al.* Identification of mechanosensitive genes in osteoblasts by comparative microarray studies using the rotating wall vessel and the random positioning machine. *J Cell Biochem* 2007; **101**: 587–599.
- Frigeri A, Iacobas DA, Iacobas S, *et al.* Effect of microgravity on gene expression in mouse brain. *Exp Brain Res* 2008; **191**: 289–300.
- Ziambaras K, Civitelli R, Papavasiliou SS. Weightlessness and skeleton homeostasis. *Hormones (Athens)* 2005; **4**: 18–27.
- Schatten H, Lewis ML, Chakrabarti A. Spaceflight and clinorotation cause cytoskeleton and mitochondria changes and increases in apoptosis in cultured cells. *Acta Astronaut* 2001; **49**: 399–418.
- Ma SB, Nguyen TN, Tan I, *et al.* Bax targets mitochondria by distinct mechanisms before or during apoptotic cell death: a requirement for VDAC2 or Bak for efficient Bax apoptotic function. *Cell Death Differ* 2014; **21**: 1925–1935.

- 27 Majima HJ, Oberley TD, Furukawa K, *et al.* Prevention of mitochondrial injury by manganese superoxide dismutase reveals a primary mechanism for alkaline-induced cell death. *J Biol Chem* 1998; **273**: 8217–8224.
- 28 Indo HP, Yen H-C, Nakanishi I, *et al.* A mitochondrial superoxide theory for oxidative stress diseases and aging. *J Clin Biochem Nutr* 2015; **56**: 1–7.
- 29 Wang SS, Good TA. Effect of culture in a rotating wall bioreactor on the physiology of differentiated neuron-like PC12 and SH-SY5Y cells. *J Cell Biochem* 2001; **83**: 574–584.
- 30 Stein TP, Leskiw MJ, Schluter MD, Donaldson MR, Larina I. Protein kinetics during and after long-duration spaceflight on MIR. *Am J Physiol* 1999; **276** (6 Pt 1): E1014–E1021.
- 31 Stein TP, Leskiw MJ. Oxidant damage during and after spaceflight. *Am J Physiol Endocrinol Metab* 2000; **278**: E375–E382.
- 32 Stein TP. Space flight and oxidative stress. *Nutrition* 2002; **18**: 867–871.
- 33 Hollander J, Gore M, Fiebig R, *et al.* Spaceflight downregulates antioxidant defense systems in rat liver. *Free Radic Biol Med* 1998; **24**: 385–390.
- 34 Degan P, Sancandi M, Zunino A, *et al.* Exposure of human lymphocytes and lymphoblastoid cells to simulated microgravity strongly affects energy metabolism and DNA repair. *J Cell Biochem* 2005; **94**: 460–469.
- 35 Maillot A, Beaufre B, Di Nardo P, Elia M, Pichard C. Weightlessness as an accelerated model of nutritional disturbances. *Curr Opin Clin Nutr Metab Care* 2001; **4**: 301–306.
- 36 Liang XH, Jackson S, Seaman M, *et al.* Induction of autophagy and inhibition of tumorigenesis by beclin 1. *Nature* 1999; **402**: 672–676.
- 37 Nakagawa T, Shimizu S, Watanabe T, *et al.* Cyclophilin D-dependent mitochondrial permeability transition regulates some necrotic but not apoptotic cell death. *Nature* 2005; **434**: 652–658.
- 38 Czabotar PE, Lessene G, Strasser A, Adams JM. Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol* 2014; **15**: 49–63.
- 39 Kharbanda S, Yuan ZM, Weichselbaum R, Kufe D. Determination of cell fate by c-Abl activation in the response to DNA damage. *Oncogene* 1998; **17**: 3309–3318.
- 40 Lakin ND, Jackson SP. Regulation of p53 in response to DNA damage. *Oncogene* 1999; **18**: 7644–7655.
- 41 Rosen EM, Fan S, Goldberg ID, Rockwell S. Biological basis of radiation sensitivity. Part 2: Cellular and molecular determinants of radiosensitivity. *Oncology (Williston Park)* 2000; **14**: 741–757.
- 42 Ding HF, Fisher DE. p53, caspase 8, and regulation of apoptosis after ionizing radiation. *J Pediatr Hematol Oncol* 2001; **23**: 185–188.
- 43 Pruschy M, Rocha S, Zaugg K, *et al.* Key targets for the execution of radiation-induced tumor cell apoptosis: the role of p53 and caspases. *Int J Radiat Oncol Biol Phys* 2001; **49**: 561–567.
- 44 Wahl GM, Carr AM. The evolution of diverse biological responses to DNA damage: insights from yeast and p53. *Nat Cell Biol* 2001; **3**: E277–E286.
- 45 Fei P, El-Deiry WS. P53 and radiation responses. *Oncogene* 2003; **22**: 5774–5783.
- 46 Houtgraaf JH, Versmissen J, van der Giessen WJ. A concise review of DNA damage checkpoints and repair in mammalian cells. *Cardiovasc Revasc Med* 2006; **7**: 165–172.
- 47 Lindsay KJ, Coates PJ, Lorimore SA, Wright EG. The genetic basis of tissue responses to ionizing radiation. *Br J Radiol* 2007; **80**: S2–S6.
- 48 Postiglione I, Chiaviello A, Palumbo G. Twilight effects of low doses of ionizing radiation on cellular systems: a bird's eye view on current concepts and research. *Med Oncol* 2010; **27**: 495–509.
- 49 Suzuki K, Kodama S, Watanabe M. Extremely low-dose ionizing radiation causes activation of mitogen-activated protein kinase pathway and enhances proliferation of normal human diploid cells. *Cancer Res* 2001; **61**: 5396–5401.
- 50 Ding LH, Shingyoji M, Chen F, *et al.* Gene expression profiles of normal human fibroblasts after exposure to ionizing radiation: a comparative study of low and high doses. *Radiat Res* 2005; **164**: 17–26.
- 51 Ogura K, Magae J, Kawakami Y, Koana T. Reduction in mutation frequency by very low-dose gamma irradiation of *Drosophila melanogaster* germ cells. *Radiat Res* 2009; **171**: 1–8.
- 52 Kiefer J. Radiation risk in manned space flights. *Mutat Res* 1999; **430**: 307–313.
- 53 Ohnishi T, Takahashi A, Ohnishi K. Studies about space radiation promote new fields in radiation biology. *J Radiat Res* 2002; **43 Suppl**: S7–S12.
- 54 Gridley DS, Coutrakon GB, Rizvi A, *et al.* Low-dose photons modify liver response to simulated solar particle event protons. *Radiat Res* 2008; **169**: 280–287.
- 55 Manda K, Ueno M, Anzai K. Memory impairment, oxidative damage and apoptosis induced by space radiation: ameliorative potential of alpha-lipoic acid. *Behav Brain Res* 2008; **187**: 387–395.
- 56 Kennedy AR, Guan J, Ware JH. Countermeasures against space radiation induced oxidative stress in mice. *Radiat Environ Biophys* 2007; **46**: 201–203.
- 57 Guan J, Stewart J, Ware JH, Zhou Z, Donahue JJ, Kennedy AR. Effects of dietary supplements on the space radiation-induced reduction in total antioxidant status in CBA mice. *Radiat Res* 2006; **165**: 373–378.