

Hypothermic activity of acetaminophen; involvement of GABA_A receptor, theoretical and experimental studies

Nematollah Ahangar¹, Zohreh Esam¹, Ahmadreza Bekhradnia¹, Mohammad Ali Ebrahimzadeh^{1*}

¹ Pharmaceutical Sciences Research Center, School of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

ARTICLE INFO

Article type:

Original article

Article history:

Received: Apr 18, 2015

Accepted: Feb 4, 2016

Keywords:

Abinitio study
Acetaminophen
Benzodiazepine receptor
Hypothermia
GABA_A

ABSTRACT

Objective(s): The mechanism of hypothermia action of acetaminophen (APAP) remains unclear even 125 years after its synthesis. Acetaminophen produces hypothermia. The mechanism of this reduction in core body temperature is not clear but evidence shows that it is not dependent on opioid and cannabinoid receptors. Because of strong documents about the roles of GABA and benzodiazepine receptors in hypothermic activity of some drugs such as diazepam, we determined if these receptors also contribute to the hypothermic effect of APAP.

Materials and Methods: Diazepam (5 mg/kg, IP) was used for induction of hypothermia. Flumazenil (10 mg/kg, IP) or picrotoxin (2 mg/kg, IP) used for reversal of this effect. Rats injected with APAP (100, 200 or 300 mg/kg, IP). Baseline temperature measurements were taken with a digital thermometer via rectum. To evaluate the structural correlation between APAP and benzodiazepine receptor ligands, numerous models are selected and studied at HF/6-31G* level of theory. Relative energies, enthalpies and Gibbs free energies were calculated for all selected drugs.

Results: Diazepam induced hypothermia was reversed by flumazenil or picrotoxin. Rats injected with APAP displayed dose- and time-related hypothermia. For combined administration, the hypothermic effect of APAP (200 mg/kg) was strongly reduced by pretreatment with picrotoxin or flumazenil $P < 0.0001$ and $P < 0.01$, respectively. Selective structural data, bond length, dihedral angles, and related distance in pharmacophore of APAP and BZD_R models were the same. Some significant structural analogues were obtained between these drugs.

Conclusion: Results suggest hypothermic action of acetaminophen may be mediated by its effect at GABA_A benzodiazepine receptor.

► Please cite this article as:

Ahangar N, Esam Z, Bekhradnia AR, Ebrahimzadeh MA. Hypothermic activity of acetaminophen; involvement of GABA_A receptor, Theoretical and experimental studies. Iran J Basic Med Sci 2016; 19:470-475.

Introduction

Acetaminophen (APAP) is widely used as an analgesic and antipyretic drug (1). It also induces hypothermia in non-febrile humans, mice and rats (2-5). APAP is often classified as a nonsteroidal anti-inflammatory drug (NSAID), but differs from other NSAIDs because it does not reduce inflammation (by inhibiting COX-2) or cause ulceration and bleeding of the stomach mucosa (by inhibiting COX-1) (6). In addition, unlike the aspirin-like drugs, APAP does not inhibit platelet aggregation or increase bleeding time (6). The mechanism of APAP hypothermia, especially in regard to the receptors, is poorly understood. APAP induced hypothermia in mice is accompanied by a reduction in brain levels of prostaglandin E₂ (PGE₂). COX-3 has been proposed to account for this hypothermic effect (5, 7, 8). However, there is no consistent evidence in the literature that central PGE₂ participates in the control of normal body temperature

in any species, and a recent study demonstrated that APAP produces hypothermia in mice by a COX-3-independent mechanism (5).

Benzodiazepine receptors (BZD_R) are an integral part of GABA_A receptor/chloride ionophore complex, which is involved in regulating the coupling between GABA_A receptors and opening of the chloride ion channels (9). Animal studies have shown that diazepam and other benzodiazepines induce hypothermia and reducing fever and pain (just like APAP) and these effects are mediated by benzodiazepine receptors (10, 11). These findings led to the suggestion that the mechanism of induction of hypothermia by acetaminophen is not fully understood until now. It seems another mechanism may also play a role in induction of hypothermia by acetaminophen, perhaps it is unrelated to COX inhibition. The aim of the present study was to find a relationship between GABAergic system and induction of hypothermia by APAP.

*Corresponding author: Mohammad Ali Ebrahimzadeh. Pharmaceutical Sciences Research Center, School of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran. Tel: +98-1133543081-3; Fax: +98-1133543084; email: zadeh20@yahoo.com

Materials and Methods

Drugs

All chemicals used were of the highest purity. APAP (Sigma-Aldrich, USA) was sonicated and dissolved in 20% 1,2-propanediol (Merck, Germany). Picrotoxin (Sigma-Aldrich, USA) and flumazenil (Sigma-Aldrich, USA) were sonicated and suspended with CMC (1%, Synofarm, Germany) and Tween 80 (5%, Merck, Germany) and were administered 15 and 10 minutes before each acquisition session. Diazepam (Sigma-Aldrich, USA) was sonicated and dissolved in normal saline. Drugs were prepared freshly on test days and administered intraperitoneally in a volume of 5 ml/kg.

Experimental part of study

All experiments were carried out on male Sprague–Dawley rats (weight range 200–250 g). The rats housed in a colony room under controlled conditions of temperature ($23 \pm 1^\circ\text{C}$) and light (dark cycle of 12:12). Food and water were freely available throughout the experimental protocol. The experiments were performed between 10 a.m. and 2 p.m. On the day of experiments, rats were placed individually in glass boxes (14 cm width, 24 cm length) with ability of free movement (unrestrained), and allowed to rest for 1 hr before drug injection. Seven rats were used in each experiment. Baseline temperature (rectal temperature) was measured with a digital thermometer (Microlife®/M.I Switzerland), lubricated and inserted to a depth of 4 cm. The data are presented as the changes in rectal temperature from the basal values. Basal values are those for the temperature (Time 0) taken immediately before the first drug/solvent administration (1). After each injection, core body temperature were measured every 15 min until 60th min. The animals were preconditioned to the temperature probe by taking temperature measurements 3 days before the experiment and twice on the day of the experiment before drug administration to reduce handling-induced temperature changes associated with stress. Rats were randomly divided into 13 groups consisting of 7 rats each.

Part 1: Picrotoxin (2 mg/kg, IP).

Part 2: Vehicle of acetaminophen; acetaminophen (100 mg/kg, IP); acetaminophen (200 mg/kg, IP); acetaminophen (300 mg/kg, IP).

Part 3: Diazepam (5 mg/kg, IP); vehicle of diazepam (5 ml/kg); diazepam (5 mg/kg) + vehicle of flumazenil (5 ml/kg); diazepam (5 mg/kg) + flumazenil (10 mg/kg).

Based on results obtained from initial experiment (Figure 1), we selected a fixed, submaximal dose of 200 mg/kg of APAP for our combination experiments.

Part 4: Vehicle of acetaminophen (5 ml/kg); acetaminophen (200 mg/kg, IP); acetaminophen (200 mg/kg) + flumazenil (10 mg/kg); aceta-

minophen (200 mg/kg) + picrotoxin (2 mg/kg).

Single intraperitoneally injection of drugs was given in each group.

Theoretical part of study

Complete geometry optimizations were performed on the APAP, diazepam, lorazepam, temazepam, flurazepam (BZD_R agonists) and flumazenil (BZD_R antagonist) molecules. All structures were optimized at the HF level of theory with the 6-31G* basis set. The frequency calculations were done at the same level to determine the nature of the optimized structures to obtain zero-point energies (un-scaled) and thermochemical corrections. All calculations were carried out using the Gaussian 98 program. Structures, thermal energies (*E*), thermal enthalpies (*H*), thermal Gibbs free energies (*G*) for each drug are calculated at HF level of theory, using 6-31G* basis sets.

Data analysis

The statistical significance of difference between experimental groups at each point was calculated by student's t-test. Differences with $P < 0.05$ were considered statistically significant.

Results

Results of experimental part of study

Effects of APAP on the rat core body temperature

APAP at the doses of 100, 200 and 300 mg/kg, IP, produced a dose- and time dependent hypothermic effect, measured as reduction in the rectal temperature (Figure 1). Acetaminophen caused hypothermia that was maximum (about 3 °C below normal) with a dose of 300 mg/kg. Compared with the vehicle control group, acetaminophen after 1 hr produced a -0.5 ± 0.51 , -2.77 ± 0.48 and -2.98 ± 0.45 °C fall in basal body temperature at doses of 100, 200, and 300 mg/kg, respectively. The mean initial body temperature in the vehicle group was 36.04 ± 0.27 °C.




Figure 1. Time-course of the change in core body temperature after intraperitoneally injection of APAP in rats (n=7). Each point represents the mean change in core body temperature. Zero on the abscissa represents body temperature at time of APAP injection. A dose-related fall in temperature was obtained after administration of 100, 200 and 300 mg/kg APAP with a maximum effect at 300 mg/kg. * $P < 0.05$ and ** $P < 0.01$ in comparison with control group




Figure 2. Each point here represents the mean change in core body temperature of 7 rats after intraperitoneally injection of diazepam (5 mg/kg), normal saline (5 ml/kg), vehicle of flumazenil (5 ml/kg) + diazepam (5 mg/kg) and flumazenil (10 mg/kg) + diazepam (5 mg/kg). Zero on the abscissa indicates body temperature at time of saline or flumazenil (BZR-antagonist) administration. ** $P < 0.01$ and **** $P < 0.0001$ in comparison with control groups

Effect of flumazenil (a BZDR-antagonist) on diazepam induced hypothermia

As shown in Figure 2, pretreatment of animals with Flumazenil (single dose 10 mg/kg, IP) reduced the hypothermic effect of diazepam (single dose of 5 mg/kg).

Effects of picrotoxin and flumazenil (GABA_A and BZDR antagonists) on APAP induced hypothermia

Pretreatment of animals with picrotoxin (single dose 2 mg/kg, IP) reduced the hypothermic effect of APAP (single dose 200 mg/kg, IP). Pretreatment of rats with flumazenil (single dose 10 mg/kg, IP) also decreased APAP hypothermia (Figure 3).

Effects of picrotoxin on the rat core body temperature

Unlike the rats injected with picrotoxin (single dose 2 mg/kg, IP) and APAP (single dose 200 mg/kg), single administration of picrotoxin (2 mg/kg) induced convulsion in all treated rats. The results of this administration on basal body temperature are shown in Table 1.

Results of theoretical part of study

Using HF/6-31G*, level of theory, relative energies (E), enthalpies (H), and Gibbs free energies (G) (kcal/mol) are calculated for all selected drugs (Figure 4, Table 2). Some selective structural data, bond length, dihedral angles, and related distance in pharmacophore of APAP and BZDR models were calculated and some significant structural analogues were obtained between these drugs (Figure 4, Table 3).




Figure 3. Effect of picrotoxin (GABA_A antagonist) and flumazenil (BZDR antagonist) on hypothermic effect of APAP. Each tow new point represents the mean change in core body temperature of 7 rats. APAP-treated group; Pretreatment with picrotoxin (2 mg/kg) or flumazenil (10 mg/kg) + APAP (200 mg/kg). * $P < 0.05$ and ** $P < 0.01$ in comparison with control group

Table 1. Effect of injection of picrotoxin (single dose 2 mg/kg, IP) on core body temperature

| T ₀ | T ₃₀ | T ₆₀ | Mean change ₃₀ | Mean change ₆₀ |
|----------------|-----------------|-----------------|---------------------------|---------------------------|
| 37.26 ± 0.14 | 36.7 ± 0.14 | 36.6 ± 0.32 | - 0.23 ± 0.14 | - 0.47 ± 0.14 |

Data are represented as mean change in core body temperature ± SEM of 7 rats in this group. Basal body temperatures (T₀), immediately before administration of picrotoxin (2 mg/kg, IP) and 30 (T₃₀) and 60 (T₆₀) min after injection

Table 2. Energies (E), enthalpies (H), and Gibbs free energies (G) (kcal/mol) for APAP and selected models of BZDR calculated at HF/6-31G*


| Ligand | E (kcal/mol) | H (kcal/mol) | G (kcal/mol) |
|------------|--------------|--------------|--------------|
| Diazepam | -788241.16 | -788240.04 | -788310.85 |
| Lorazepam | -1098716.27 | -1098715.15 | -1098789.79 |
| Flurazepam | -1007050.83 | -1007049.71 | -1007144.66 |
| Temazepam | -835201.30 | -835200.18 | -835273.87 |
| Flumazenil | -845057.24 | -845056.12 | -845138.08 |
| APAP | -321300.57 | -321299.45 | -321353.87 |

Discussion

Acetaminophen (APAP) is a widely used as an antipyretic analgesic and reducing fever caused by bacterial or viral infections and/ or by clinical trauma (1). APAP demonstrated a dose-dependent hypothermic action (1, 5), thus it can improve functional outcome after acute ischemic stroke (2) or other conditions such as status epilepticus, cardiac arrest, convulsion, epilepsy in which controlling the hypothermia is useful and shows therapeutic effectiveness (12, 13). Some studies suggest that this hypothermic action is mediated by inhibition of COX-3 in CNS, in the anterior hypothalamus (7), but more documents will be required to improve this mechanism.

Table 3. Comparison of distances (Å) and angles (°) between the aromatic ring centers and carbonyl group atoms in APAP and selected models of BZD_R calculated at HF/6-31G*

| Ligand | Ar- O ₄ | Ar- C=O | Ar- C ₃ | Angles (1,2,3)° | Dihedral angles (1,2,3,4)° |
|------------|--------------------|---------|--------------------|-----------------|----------------------------|
| Diazepam | 4.876 | 1.195 | 3.748 | 122.67 | 13.47 |
| Lorazepam | 4.937 | 1.192 | 3.838 | 129.62 | 3.37 |
| Flurazepam | 4.913 | 1.197 | 3.776 | 123.54 | 2.6 |
| Temazepam | 4.900 | 1.195 | 3.770 | 124.12 | 8.61 |
| Flumazenil | 4.924 | 1.360 | 3.787 | 126.06 | 3.74 |
| APAP | 4.041 | 1.198 | 3.843 | 129.08 | 0 |

**Figure 4.** Structural analogues of APAP and selected benzodiazepine receptor ligands

APAP penetrates the brain following peripheral administration, and some evidence suggest a centrally acting mechanism for its action (4, 14). But really the mechanism of action of APAP remains unclear even 125 years after its original synthesis (5). The principal inhibitory neurotransmitter in mammalian CNS, γ -amino butyric acid (GABA), exerts its physiological roles by binding to three major classes of receptors, the GABA_A, GABA_B and GABA_C. The GABA_A receptor belongs to Cysloop super family of ligand-gated ion channels (LGIC) (15). Mammalian GABA_A receptors have been shown to be heteropentameric assemblies of protein subunits, where the most abundant receptor-ion channels have the composition (α_n)₂ (β_n)₂ (γ) (16). This receptor has received the greatest attention in terms of research because of its importance as a biological target for clinically important drugs such as barbiturates, neurosteroids, anesthetics, ethanol, and benzodiazepines (BZDs) (17, 18). Classically, BZD agonists (anxiolytic, anticonvulsant, muscle relaxant, sedative-hypnotic, reducing pain, fever and

basal body temperature) act as positive modulators at BZD/GABA_A receptor (BZD_R) by increasing the frequency of chloride channel openings (19). The allosteric BZD_R has been proposed to reside at the interface between α and γ subunits (16, 20). Exclusive of γ and even β subunits which are respectively necessary for the GABA_A receptor to be able to respond to low concentrations of BZDs and suggested to be responsible for Etomidate-induced hypothermia (21). α subunit ($\alpha_{1,2,3,5}$) is predominant subunit in CNS and is necessary for binding of BZDs. A significant structural analogue exists between APAP and specific ligands of this subunit (TPA-023, TP-003, N-(indol-3-ylglyoxylyl) piperidines (22).

Acetaminophen reduces the core temperatures of febrile and non-febrile in different ways. The hypothermia induced by APAP (Figure 1) has been reported previously in human, mice and rats (1, 2, 5). It has been demonstrated that APAP produces hypothermia independent of opioid or cannabinoid CB1 or NOP receptor activation, in contrary to its antinociceptive effect, which requires opioid and cannabinoid CB1 receptor activation (5). The aim of

the present study was to investigate the mechanism of this hypothermic action in rat. The results suggest that the hypothermic action of acetaminophen (velocity and peak hypothermia respectively in 30 and 60 min after APAP administration) is antagonized by the GABA_A receptor antagonist, picrotoxin and benzodiazepine receptor antagonist, flumazenil (Figure 3). Flumazenil did not alter body temperature (23). Theoretical part of study has been presented in Table 3. Data suggests that structural similarity exist between APAP and BZD_R ligands (diazepam, lorazepam, flurazepam, temazepam and flumazenil).

The present results are in part consistent with some of previous findings. It has been reported that in the presence of APAP, enoxacin (a quinolone antimicrobial agent) did not decrease GABA responses (24), whilst quinolones such as PNU-101017, a novel, imidazoquinoline amide and benzodiazepine receptor partial agonist that has high affinity for the GABA_A receptor subtypes containing the α_1 and α_3 or α_5 subunits, have shown related to GABA_A receptor (25). Administration of flumazenil antagonized the analgesic effect exerted by acetaminophen (26), and also prevented the lipopolysaccharide anxiety sensitization (27). The latter, can correlate APAP as the most proposed antipyretic drug to GABA_A receptor indirectly. Furthermore, study of analgesic effects of APAP and morphine in a mouse model of bone cancer pain supposed that APAP potentiates opioid inhibition of GABAergic synaptic transmission (28). These results also correlate APAP to GABAergic system too. Body temperature may be a useful test for discriminating between full and partial agonists at the BZD_R (29). It is possible to consider APAP as a partial agonist at benzodiazepine receptor level. This finding will be a new way for understanding the mechanism of action of APAP.

Conclusion

These results support the view that hypothermic activity of acetaminophen is mediated by its effect at GABA_A BZD_R. However, further studies will clarify this hypothesis.

Acknowledgment

This research was supported by a grant from the Research Council of Mazandaran University of Medical Sciences.

Conflict of interest

The authors declare that there are no conflicts of interest.

References

1. Ayoub S, Botting R. Acetaminophen-induced hypothermia in mice is mediated by a prostaglandin

endoperoxide synthase 1 gene-derived protein. *Proc Natl Acad Sci* 2004; 101:11165-11169.

2. Dippel DW, van Breda EJ, van der Worp HB, van Gemert HM, Meijer RJ, Kapelle LJ, *et al*. Effect of paracetamol (acetaminophen) and ibuprofen on body temperature in acute ischemic stroke PISA, a phase II double blind, randomized, placebo controlled trial. *BMC Cardiovasc Disord* 2003; 3:2-9.

3. Denes E, Amaniou M, Rogez JP, Weinbreck P, Merle L. Acetaminophen-induced hypothermia, an AIDS related side-effect? About 4 cases. *Ann Med Intern* 2002; 153:411-413.

4. Li S, Dou W, Tang Y, Goorha S, Ballou LR, Blatteis CM. Acetaminophen: antipyretic or hypothermic in mice? In either case, PGHS-1b (COX-3) is irrelevant. *Prostaglandins Other Lipid Mediat* 2008; 85:89-99.

5. Corley G, Rawls S. Opioid, cannabinoid CB1 and NOP receptors do not mediate APAP-induced hypothermia in rats. *Pharmacol Biochem Behav* 2009; 92:503-507.

6. Hilal-Dandan R, Brunton L. *Goodman & Gilman's Manual of Pharmacology and Therapeutics*. New York: McGraw-Hill; 2014. p. 599-600.

7. Botting R, Ayoub S. COX-3 and the mechanism of action of paracetamol/acetaminophen. *Prostaglandins Leukot Essent Fatty Acids* 2005; 72: 85-87.

8. Chandrasekharan NV, Dai H, Roos KL, Evanson NK, Tomsik J, Elton TS, *et al*. COX-3, a cyclooxygenase-1 variant inhibited by acetaminophen and other analgesic/antipyretic drugs: cloning, structure and expression. *Proc Natl Acad Sci* 2002; 99:13926-13931.

9. Bailey DJ, Tetzlaff JE, Cook JM, He X, Helmstetter FJ. Effects of hippocampal injections of a novel ligand selective for the alpha 5 beta 2 gamma 2 subunits of the GABA/ benzodiazepine receptor on Pavlovian conditioning. *Neurobiol Learn Mem* 2002; 78:1-10.

10. Sarnowska A, Beresewicz M, Zabłocka B, Domańska-Janik K. Diazepam neuroprotection in excitotoxic and oxidative stress involves a mitochondrial mechanism additional to the GABA_A and hypothermic effects. *Neurochem Int* 2009; 55:164-173.

11. Vinkers CH, Klanker M, Groenink L, Korte SM, Cook JM, Van Linn ML, Hopkins SC, Olivier B. Dissociating anxiolytic and sedative effects of GABAergic drugs using temperature and locomotor responses to acute stress. *Psychopharmacology (Berl)* 2009; 204:299-311.

12. Schmitt FC, Buchheim K, Meierkord H. Anticonvulsant properties of hypothermia in experimental status epilepticus. *Neurobiol Dis* 2006; 23:689-696.

13. Inamasu J, Nakatsukasa M, Suzuki M. Therapeutic hypothermia for out-of-hospital cardiac arrest: An update for neurosurgeons. *World Neurosurg* 2010; 74:120-128.

14. Massey TE, Walker RM, McElligott TF, Raczy WJ. Acetaminophen induced hypothermia in mice: Evidence for a central action of the parent compound. *Toxicology* 1982; 25:187-200.

15. Trudell JR, Bertaccini E. Comparative modeling of a GABA_A alpha1 receptor using three crystal structures as templates. *J Mol Graphics Model* 2004; 23:39-49.

16. Olsen R, Sieghart W. GABA_A receptors: Subtypes provide diversity of function and pharmacology. *Neuropharmacology* 2009; 56:141-148.
17. Olsen RW. Allosteric ligands and their binding sites define γ -aminobutyric acid (GABA) type A receptor subtypes. *Adv Pharmacol* 2015; 73:167-202.
18. Sieghart W. Allosteric modulation of GABA_A receptors via multiple drug-binding sites. *Adv Pharmacol* 2015; 72:53-96.
19. Chebib M, Johnston GA. GABA-activated ligand gated ion channels: medicinal chemistry and molecular biology. *J Med Chem* 2000; 43:1427-1447.
20. Jembrek MJ, Vlainic J. GABA receptors: pharmacological potential and pitfalls. *Curr Pharm Des* 2015; 21:4943-4959.
21. Cirone J, Rosahl TW, Reynolds DS. Gamma-aminobutyric acid type A receptor beta 2 subunit mediates the hypothermic effect of etomidate in mice. *Anesthesiology* 2004; 100:1438-1445.
22. Gray JA, Roth B. Molecular targets for treating cognitive dysfunction in schizophrenia. *Schizophr Bul* 2007; 33:1100-1119.
23. Jackson HC, Nutt DJ. Strain differences in sensitivity to the hypothermic effects of benzodiazepine receptor ligands in mice. *Psychopharmacology* 1992; 109:365-368.
24. Yakushiji T, Shirasaki T, Akaike N. Non-competitive inhibition of GABA_A responses by a new class of quinolones and non-steroidal antiinflammatories in dissociated frog sensory neurones. *Br J Pharmacol* 1992; 105:13-18.
25. Hall E, Andrus P, Fleck T. Neuroprotective properties of the benzodiazepine receptor, partial agonist PNU-101017 in the gerbil forebrain ischemia model. *J Cereb Blood Flow Metab* 1997; 17:875-883.
26. Madenoglu H, Kaçmaz M, Aksu R. Effects of naloxone and flumazenil on antinociceptive action of acetaminophen in rats. *Curr Ther Res* 2010; 71:111-117.
27. Breese GR, Knapp DJ, Overstreet DE. Repeated lipopolysaccharide (LPS) or cytokine treatments sensitize ethanol withdrawal induced anxiety like behavior. *Neuropsychopharmacology* 2008; 33:867-876.
28. Saito O, Aoe T, Yamamoto T. Analgesic effects of nonsteroidal antiinflammatory drugs, acetaminophen and morphine in a mouse model of bone cancer pain. *J Anesth* 2005; 19:218-224.
29. Jackson H, Nutt D. Body temperature discriminates between full and partial benzodiazepine receptor agonists. *Eur J Pharmacol* 1990; 185:243-246.