



POSTER PRESENTATION

Open Access

# 12/15-lipoxygenase orchestrates the clearance of apoptotic cells and maintains immunologic tolerance

Stefan Uderhardt<sup>1</sup>, Martin Herrmann<sup>1</sup>, Olga Oskolkova<sup>2</sup>, Reinhard Voll<sup>1</sup>, Falk Nimmerjahn<sup>3</sup>, Valery Bochkov<sup>2</sup>, Georg Schett<sup>1</sup>, Gerhard Kroenke<sup>1\*</sup>

From 6th European Workshop on Immune-Mediated Inflammatory Diseases  
Nice, France. 23-25 November 2011

The coordinated and non-inflammatory phagocytosis of apoptotic cells is crucial to maintain immunological tolerance. During inflammation, however, ingestion of apoptotic material by inflammatory phagocytes can provoke a break in self-tolerance. Hence, and though poorly understood, mechanisms governing the sorting of apoptotic cells into distinct and differentially-activated subsets of phagocytes are essential to prevent autoimmunity.

Here we identify the enzyme 12/15-lipoxygenase (12/15-LO) as a crucial factor orchestrating the clearance of apoptotic cells under inflammatory conditions. During peritonitis, the ingestion of apoptotic cells is confined to a distinct population of alternatively-activated, 12/15-LO-expressing resident macrophages (resMΦ). Deletion of 12/15-LO changed this pattern and neighbouring inflammatory macrophages (infMΦ) start to engulf apoptotic cells.

We hypothesized that resMΦ exert a paracrine and 12/15-LO-mediated inhibitory activity on the phagocytic capacity of infMΦ and identified oxidation products of phosphatidyl ethanolamine (oxPE) as the corresponding mediators. oxPE is generated by the action of 12/15-LO in resMΦ and consequently exposed on the macrophages' surfaces. Here, oxPE binds, blocks and scavenges soluble MFG-E8 protein and thereby selectively blocks the major pathway involved in the uptake of apoptotic cells into infMΦ. In turn, we observed a break in self-tolerance and lupus-like autoimmune disease in aged 12/15-LO-deficient mice. These mice displayed spontaneous production of autoantibodies and glomerulonephritis,

which both exacerbated after apoptotic challenge in the pristane-induced model of experimental murine lupus.

Our data point towards a so far unrecognized role for enzymatic lipid oxidation during the maintenance of self-tolerance and identify a mechanism, which orchestrates the cell- and context-specific uptake of antigens by different subsets of phagocytes, imposing a new paradigm in our understanding of the clearance of apoptotic cells.

#### Author details

<sup>1</sup>Internal Medicine 3, University of Erlangen, Germany. <sup>2</sup>Vascular Biology, Medical University of Vienna, Vienna, Austria. <sup>3</sup>Dept. of Genetics, University of Erlangen, Germany.

Published: 23 November 2011

doi:10.1186/1479-5876-9-S2-P5

Cite this article as: Uderhardt et al.: 12/15-lipoxygenase orchestrates the clearance of apoptotic cells and maintains immunologic tolerance. *Journal of Translational Medicine* 2011 **9**(Suppl 2):P5.

#### Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Internal Medicine 3, University of Erlangen, Germany  
Full list of author information is available at the end of the article