

REVIEW

Open Access



Chimeric antigen receptor-engineered natural killer cells for cancer immunotherapy

Ahmet Yilmaz^{1†}, Hanwei Cui^{1†}, Michael A. Caligiuri^{2,3,4,5†} and Jianhua Yu^{2,3,4,5*†}

Abstract

Natural killer (NK) cells are a critical component of the innate immune system. Chimeric antigen receptors (CARs) redirect NK cells toward tumor cells carrying corresponding antigens, creating major opportunities in the fight against cancer. CAR NK cells have the potential for use as universal CAR cells without the need for human leukocyte antigen matching or prior exposure to tumor-associated antigens. Exciting data from recent clinical trials have renewed interest in the field of cancer immunotherapy due to the potential of CAR NK cells in the production of “off-the-shelf” anti-cancer immunotherapeutic products. Here, we provide an up-to-date comprehensive overview of the recent advancements in key areas of CAR NK cell research and identify under-investigated research areas. We summarize improvements in CAR design and structure, advantages and disadvantages of using CAR NK cells as an alternative to CAR T cell therapy, and list sources to obtain NK cells. In addition, we provide a list of tumor-associated antigens targeted by CAR NK cells and detail challenges in expanding and transducing NK cells for CAR production. We additionally discuss barriers to effective treatment and suggest solutions to improve CAR NK cell function, proliferation, persistence, therapeutic effectiveness, and safety in solid and liquid tumors.

Keywords: Chimeric antigen receptor, Natural killer cells, T cells, Cancer immunotherapy

Background

Natural killer (NK) cells are large granular lymphocytes with surface markers CD3⁻CD56⁺NKp46⁺ in humans. They have been recognized as a subset of innate lymphoid cells (ILCs) that lack antigen receptors with recombination activating gene (RAG)-dependent rearrangement [1, 2]. There are three groups of ILCs (Groups 1–3), each with distinct functions and expression profiles of key transcription factors. NK cells are included in Group 1 ILCs, which also include innate lymphoid cell 1 (ILC1). Both NK cells and ILC1 depend on the transcription factor T-bet for their proper function and development, and secrete interferon- γ (IFN- γ) rather than type 2 cytokines (IL-5 and IL-13), IL-17, or IL-22 [3]. The development

of NK cells involves bone marrow and extra-medullary sites such as tonsils, uterus, and liver [4–6]. The NK cell developmental process is initiated in the bone marrow microenvironment where hematopoietic stem cells differentiate into common lymphoid progenitors (CLPs) which can give rise to all subtypes of lymphoid cells. CLPs first develop into NK progenitor cells, and then differentiate into immature and finally mature or resident NK cells. These developmental processes may occur in different organs or tissues [7–9].

NK cells are able to recognize and kill “stressed cells” including cells infected with viruses, allogeneic cells, and tumor cells without prior sensitization in a non-HLA (human leukocyte antigen)-restricted manner. The effector functions in NK cells are regulated through a balance of signals delivered by activating and inhibitory receptors as well as cytokines such as IL-2, IL-12, IL-15, or IL-18 [10]. The inhibitory receptors, including inhibitory killer immunoglobulin receptors (KIRs) and NKG2A/CD94 in humans, keep NK cells in a quiescent state through the

*Correspondence: Jiayu@coh.org

[†]All authors contributed equally to this work

² Department of Hematology and Hematopoietic Cell Transplantation, City of Hope National Medical Center, 1500 E. Duarte Road, KCRB, Bldg. 158, 3rd Floor, Room 3017, Los Angeles, CA 91010, USA

Full list of author information is available at the end of the article



recognition of constitutively expressed self-MHC class I molecules on healthy cells. A variety of activating receptors are expressed on NK cells, exemplified by NKG2D, CD16, natural cytotoxicity receptors (NCRs), and activating KIRs. The activating receptors contribute to NK cell activation via recognition of host-derived or pathogen-encoded ligands or antibodies [11, 12].

NK cells play critical roles against tumor initiation and metastasis [13, 14]. Their activation by tumor cells, which depends on the type of ligands on the surface of the tumor cells, can be accomplished via several mechanisms. The MHC-I molecules on target cells, which inhibit NK cell function through binding to inhibitory receptors on NK cells, are often reduced or lost in tumors (missing-self recognition). Activation of NK cells can also be triggered by upregulation of ligands on tumor cells by certain therapeutic agents [15–17].

Activated NK cells are capable of eliminating tumor cells through direct cell cytotoxicity and/or production of pro-inflammatory cytokines. They can directly release perforin and granzymes to kill tumor cells. They can also mediate antibody-dependent cellular cytotoxicity (ADCC) via the membrane receptor CD16, or apoptotic pathways mediated by Fas ligand (FasL) or TNF-related apoptosis-inducing ligand (TRAIL) [18]. In addition, cytokines and chemokines such as IFN- γ and granulocyte macrophage colony stimulating factor (GM-CSF) produced by NK cells can modulate immune responses of other cells. For example, IFN- γ helps in activation of CD8+ T cells, monocytes, and macrophages [4, 19, 20].

Although NK cells are able to recognize and lyse tumor cells, the tumor cells develop their own mechanisms to evade recognition by NK cells or repress NK cell function. For example, tumor cells can produce immunosuppressive cytokines such as IL-10 or transforming growth factor-beta (TGF- β) to repress NK cell function [21, 22]. They may also reduce expression of tumor-associated antigens (TAAs) or elevate expression of MHC class I-related molecules to inactivate NK cells [23]. Engineering NK cells to express a chimeric antigen receptor (CAR) may overcome immune evasion. Although several approaches including CAR T cells, checkpoint inhibitors, antibodies, antibody–drug conjugates, and tumor vaccinations are being investigated [24, 25], development of “off-the-shelf” CAR-modified NK cells is an especially promising and unique approach in developing sustainable anti-cancer immunotherapeutic products.

Chimeric antigen receptors (CARs): structure and function

A CAR is a functional hybrid antigen receptor usually combining variable regions of an antibody with the constant region of a T cell receptor (TCR), grafting an immune cell with the specificity for TAA expressed on

tumor cells [26]. A typical CAR is composed of a signal peptide, single-chain antibody variable fragment (scFv), and transmembrane and intracellular signaling domains. A linker combines the variable regions of heavy and light chains of an antibody molecule. A hinge, also called spacer, connects the scFv to the transmembrane domain. scFv is derived from a tumor-specific antibody and can target any antigen expressed on tumor cell surface [27]. In some CARs, a ligand of a tumor antigen is used [28, 29]. An intracellular signaling domain is derived from immunoreceptor tyrosine-based activation motifs (ITAMs) of TCR or a cytoplasmic domain of other activating receptors. Phosphorylation of ITAMs recruits adapter molecules, stimulates downstream pathways, and activates immune cells [30, 73].

The first-generation CARs contained only the CD3 ζ chain, the common signal transducing subunit of TCR, or Fc γ R subunits as the intracellular signaling domain [31–33]. They were not effective in initiating T cell activation or tumor elimination as they lacked co-stimulatory signaling required for full T cell activation. To improve the immune cell response to tumor cells, second-generation CARs were created by adding one of the T cell co-stimulatory molecules such as CD28, 4-1BB (CD137), ICOS, or OX40 (CD134) in addition to CD3 ζ . The third-generation CARs include three signaling domains (for example, CD3 ζ and two co-stimulatory domains) [34, 35]. There is no consistent evidence suggesting increased performance in the third—compared to the second—generation CARs. The type of co-stimulatory domain used can dramatically influence effector functions mediated by the CAR. For example, 4-1BB co-activation results in development of T memory cell characteristics, whereas CD28 co-activation results in enhanced T cell activation and expansion [36].

Generation of vectors for CARs

CARs can be generated against any of the surface antigens expressed on tumor cells. However, the target should have a specific antigen density to avoid “on-target, off-tumor” effects. Studies in T cells have shown that target density between 200 and 2000 copies per cell is optimal to avoid damage to the normal cells [37]. Generation of a CAR usually starts with an scFv derived from an antibody, composed of heavy and light chains [38]. Recently, nanobodies composed only of heavy chains are being utilized for CAR NK cell production [39]. These single-chain antibodies not only possess specificity comparable to scFvs but also are much smaller in size, facilitating transduction. However, their use in CAR NK cell production is currently at the experimental stage.


Usually, a signal peptide is introduced in the front of an scFv. A linker that connects heavy and light chains usually contains the (G₄S)₃ or (G₄S)₅ signature that

provides flexibility for accurate folding of connected protein domains [40, 41]. A spacer, also referred to as a hinge, usually derived from IgG, CD8 α , or CD28, plays a critical role in specificity of the CAR-bearing immune cells [41]. Adjusting the length of spacer may help optimize the distance between CAR-bearing cells and target cells for an optimal immunologic synapse (IS) formation and specificity. Use of CAR T cells created using spacers with suboptimal length results in off-target effects. The downstream CAR component(s) are co-stimulatory domain(s) that can be shared between CAR NK and CAR T cells [42, 43], followed by CD3 ζ . With the rapid development of DNA synthesis technology, many CAR sequences are now synthesized by a vendor if the DNA sequences encoding the scFv are known. All CAR components or a cassette are commonly cloned into a retrovirus- or lentivirus-based expression vector, which is then used to transduce primary NK cells or NK cell lines [44] (Fig. 1). In some instances, an inducible iCaspase can be incorporated as a safety switch for use in case of adverse reactions [10]. Different CAR designs and detailed engineering approaches have been reviewed elsewhere [45].

CAR T cells vs. CAR NK cells for cancer immunotherapy

CAR T cells targeting CD19 or other B cell markers such as CD20 have achieved remarkable responses in patients

with B cell malignancies including chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), and lymphoma [26, 46–48]. CTL019 (Kymriah, tisagenlecleucel) marketed by Novartis achieved 82.5% overall remission rate (ORR), which includes 40 complete remission (CR) and 12 CR with incomplete hematologic recovery (CRi) in 63 patients with ALL. A biologic license application for Kymriah was assessed based on B2202, a single-arm, open label, multi-center phase 2 study including 107 patients who signed consent. There were 19 patients who were not eligible due to death before the study began, unacceptable apheresis, incomplete screening before the cutoff date, or loss of eligibility for other reasons [49]. Additional twenty patients from the remaining 88 were further excluded due to adverse events, manufacturing failure, death, or pending infusion. The remaining 68 patients were included in the safety set, but five were further excluded from the efficacy set because the products were manufactured in Germany, and thus, 63 patients were included in the ORR calculation above. The ORR of 82.5% was an impressive outcome compared to less than 20% of the standard treatment, and thus, the product received an expedited approval by the US Food and Drug Administration (FDA) [36, 49, 50]. The intent-to-treatment (ITT) rate, as defined by the number of patients with ORR divided by the number of



patients consented, was 48.5% (52/107). In the Yescarta study, 111 patients were enrolled, but 10 withdrew [51]. Objective response rate was 72% (51% for advanced cases). In the treatment of DLBCL, CR rates were higher in the Yescarta study compared to the Kymriah study (72% vs. 50%). It is unknown whether this difference was influenced by differences in the number of cells infused (60–600 million cells were infused if the body weight was greater than 50 kg in the Kymriah study and 2 million cells per kg were infused in the Yescarta study).

CD19-CAR T cells are the first CAR T cell-based drug approved by the FDA. Studies to develop CAR T cells for treating other tumor types including solid tumors are increasing [52–55]. However, there have been reports of serious side effects in the clinic associated with CAR T cell therapy including cytokine release syndrome (CRS), “on-target, off-tumor” toxicity, neurologic symptoms, graft vs. host disease (GvHD) if used in an allogeneic setting, and tumor lysis syndrome. For example, 94% of the patients in the Yescarta study and 74% in the Kymriah study had CRS and at least four patients in each study died from CRS [51]. CRS is a rapid and massive release of inflammatory cytokines such as IFN- γ , TNF- α , and IL-6 into circulation [47]. It can present with symptoms such as high fever, malaise, fatigue, myalgia, nausea, anorexia, tachycardia/hypotension, capillary leak, or disseminated intravascular coagulation (reviewed by Bonifant et al. (2016) [56, 57]. Other studies have also shown that CRS by CAR T cells can be life-threatening [56–58]. In addition, CAR T cells may attack not only tumor but also normal cells, resulting in on-target, off-tumor toxicity. For example, HER2-specific CAR T treatment resulted in acute respiratory failure and death in a patient with metastatic colon cancer possibly due to low levels of HER2/neu expressed in the pulmonary parenchyma or vasculature [59]. Severe neurological toxicities such as acute cerebral edema resulting in brain death a few hours after initial appearance of neurological symptoms have also been reported in treatment with CD19-CAR T cells [60]. In the separate Kymriah and Yescarta studies, although, respectively, noted in 72% and 87% of the patients treated with CAR T cells, no deaths were attributed to neurotoxicity. CAR T cells obtained from allogeneic sources may also result in GvHD [61]. Disabling the endogenous TCR has the potential to avoid or lower GvHD risk. Endogenous TCR has been disabled using recent techniques such as CRISPR/Cas9, TALEN, zinc finger endonucleases, and receptor blockage using an array of proteins [61]. Some of these techniques are currently being evaluated for clinical use.

Compared to T cells, NK cells offer many advantages for cancer immunotherapy (Table 1). For example, it has been reported that allogeneic NK cells do not need

complete HLA matching and cause little or no GvHD compared to allogeneic T cells [61–63]. In fact, there is experimental data in animals showing that NK cells not only do not cause GvHD but also prevent it in some cases by inhibiting activated alloreactive T cells while retaining graft vs. tumor effects [63]. On the contrary, CAR T cells can cause GvHD even when HLA-matched as minor mismatches may still be provocative [64]. Additionally, the cytokine levels produced by NK cell infusions are moderate [65] compared to the levels seen in life-threatening cases of CRS observed in CAR T cell trials [61].

NK cells have a limited life span in circulation and do not produce memory cells except for some viruses. Therefore, a primary-NK cell-based CAR approach may not need to be modified to include a suicide gene as a safety switch [66]. A further potential advantage of using CAR NK cells is that, unlike CAR T cells, they can be derived from umbilical cord blood (UCB) or cell lines such as NK-92, both of which have the potential for use as an “off-the-shelf” therapeutic product [61].

The safety of CAR NK-92 cells was tested in acute myeloid leukemia (AML) patients in a recent “first-in-man” clinical trial targeting CD33 [67]. Although the CAR NK-92 cells did not seem significantly beneficial in that particular study, injection of approximately five billion irradiated cells per patient did not produce any significant adverse side effects, suggesting that injection with a large number of NK-92 cells may be safe. A recent clinical trial (NCT03056339) produced exciting evidence supporting the potential for production of “off-the-shelf” NK cells with significant benefits in relapsed or refractory CD19-positive lymphoma and leukemia [68]. HLA-mismatched NK cells isolated from UCB and transfected with a retrovirus-based vector expressing anti-CD19 CAR, iCaspase-9, and IL-15 were administered in 11 patients with CLL or non-Hodgkin’s lymphoma. Nine patients received haplo-mismatched CAR NK cells and 2 patients received completely mismatched CAR NK cells. Seven patients had a complete response and one had a partial response with remission of the Richter’s transformation component. These patients had relapsed/refractory tumors and were heavily pre-treated using conventional methods before starting CAR NK cell therapy. The patients had a median of four lines of therapies before beginning the CAR NK cell treatment. Five were CLL patients who failed to respond to ibrutinib and other therapies. Six were lymphoma patients, four of whom failed to respond to autologous hematopoietic stem cell transplantation (HSCT), and two had refractory disease. All but one patient had at least two prior relapses. An objective response of eight patients (73%) was observed during a follow-up period of 13.8 months. In all eight patients, a response was seen in the first month of the

Table 1 Comparison of clinical efficacy and future prospects in CART cell product during Kymriah and Yescarta clinical trials, complicating interpretation of the results [51]. This table has been partially adopted from Kurlander (2017), O’Leary (2017) [49, 51], and Klingemann (2014) [65]

Parameter	CART	Primary CAR NK	CAR NK-92
Source	PB	PB or UCB	Cell line
Need for in vitro expansion before infusion	Yes	Yes	No
Pre-activation and differentiation	Needs CD3/CD28 stimulation	Requires feeder cells and/or IL-2 or other cytokines for optimal growth	Requires IL-2 for optimal growth
Viral transduction efficiency	Variable	Variable	High transduction efficiency
Side effects			
Neurotoxicity	Yes (reported in 58–87% of the patients in the Kymriah and Yescarta clinical trials)	No	No
CPS	Yes (reported in 74–94% of the patients in the Kymriah and Yescarta clinical trials)	No	No
Long-term side effects	Yes, CART cells can persist in circulation for years	Limited in vivo survival without IL-15 and persistent growth for over a year with incorporated IL-15	Short persistence, the cell line needs to be irradiated prior to infusion
Killing activity	Kills only tumor cells carrying TAA in an MHC-independent manner; produces slow response	Kills tumor cells regardless of their MHC status with some preference in killing tumor cells lacking MHC-I; quick response without prior activation	Similar to primary CAR NK
ADCC activity	No	Yes, due to CD16 expression	Similar to primary CAR NK when NK-92 CAR NK cells are transduced with CD16
Prospects for clinical use	Impressive clinical outcomes have already been reported; CART treatment has been approved by the FDA (Kymriah for ALL and Kymriah and Yescarta for DLBCL [51])	Being developed and promising	Being developed and combined with other agents
"Serial" killing ability	Yes	Yes, one NK cell can kill up to 5 tumor cells [220]	Unknown because cells are irradiated before infusion
Production efficiency	Low; needs to be prepared individually for each patient and requires coordination between production facility and clinic ^a	High due to donor availability and the off-the-shelf potential	High; unlimited expansion potential of the cell line
Cost of treatment	Expensive; approximately \$373,000 USD [221]	Relatively less expensive due to off-the-shelf potential	Relatively less expensive than CAR T and primary CAR NK cells due to unlimited expansion potential
Potential for production of "off-the-shelf" anti-cancer products	Low to moderate; TCR knock-out studies are ongoing, but disabling TCR may lower T cell activity and/or survival and may be technically challenging	High; UCB NK cells may have better survival rates than PB NK cells after thawing or infusion	Very high; unlimited expansion potential

^a 10% of the patients died or were dropped and 80% needed bridge chemotherapy while waiting for the CART cell product during Kymriah and Yescarta clinical trials, complicating interpretation of the results [51]. This table has been partially adopted from Kurlander (2017), O’Leary (2017) [49, 51], and Klingemann (2014) [65]

^b DLBCL diffuse large B-cell lymphoma, PB peripheral blood, UCB umbilical cord blood

CAR NK treatment. An additional patient had a complete response, but his response was not attributed to CAR NK cell therapy. It was possible to bridge five of the eight patients who responded CAR NK therapy to HSCT, rituximab, venetoclax or lenalidomide treatment after CAR NK-cell-induced remission. None of the side effects seen with CAR T cells were noted. Excluding transient myelotoxicity, there was no neurotoxicity, elevation in IL-6 levels, CRS, or GvHD. The infused CAR NK cells expanded and persisted in vivo for at least 12 months [68].

Sources to obtain NK cells for immunotherapy

NK cell lines as well as primary NK cells from peripheral blood (PB), UCB, or induced pluripotent stem cells (iPSCs) have been used for CAR NK cell production. Several NK cell lines such as NK-92, NKG, YT, NK-YS, HANK-1, YTS, and NKL are commercially available [69]. NK-92 is the most widely used cell line and has already entered into clinical trials [70]. In comparison to PB NK cells, NK-92 cells are easier to expand and transduce and do not carry any contaminating T cells that may cause GvHD [71]. In addition, their production is inexpensive and easy to standardize, and they do not express fewer inhibitory KIRs that restrict NK effector functions [72, 73]. However, there are also several disadvantages associated with the use of NK-92 cells. For example, the cells must be irradiated before infusion into patients to prevent unrestricted cell growth, thereby limiting their capacity for in vivo expansion. Additionally, the NK-92 cell line carries Epstein–Barr virus and has an abnormal genome.

NK cells constitute only about 10% of the PB lymphocytes and thus need expansion in vitro to obtain a sufficient quantity for a therapeutic effect. Various methods to enhance expansion of NK cells isolated from PB have been detailed below in the "Challenges in expanding CAR NK cells" section. NK cells from PB, and consequently CAR NK cells produced from PB, are potent killers of tumor cells as they are fully matured [74, 75]. Another well-studied source of primary NK cells is UCB, which may offer several advantages. Approximately 15 to 30% of UCB lymphocytes are NK cells, which can result in more than NK cells found in PB [76, 77]. There are less contaminating T cells in UCB compared to PB, which reduces the risk of GvHD [78–80]. In addition, NK cells in UCB may be less likely to undergo rejection by the patient compared to PB NK cells. However, NK cells isolated from UCB are immunologically immature and initiate only a weak immune response against target cells after being activated by stimuli such as cytokines or alloantigens [81]. Therefore, CAR NK cells derived from UCB may be less cytotoxic than those derived from PB

[82]. UCB is easy to collect with no harm to the mother or child and almost never contaminated with Epstein–Barr virus or cytomegalovirus.

A recent study showed that CD19-CAR NK cells isolated from UCB transduced to express IL-15 and an iCaspase-9-based suicide gene had effective and persistent anti-tumor activity in NOD scid gamma (NSG) mice [83]. The incorporated IL-15 dramatically increased anti-tumor activity in the CD19-CAR NK cells. This study has been translated into a clinical trial with impressive outcomes [68]. Other less common sources for obtaining NK cells include pluripotent stem cells (human embryonic stem cells or iPSCs or adult stem cells (CD34+ hematopoietic stem cells) [84–88]. Recently, Li et al. (2018) utilized NK cells derived from human iPSC with a mesothelin-CAR modification as a potential source for "off-the-shelf" CAR NK cells for anti-cancer immunotherapy [89].

Targets for CAR NK cells

The choice of targets for CARs depends on the type of tumor studied. Several proteins expressed by tumor cells are being targeted in preclinical or clinical studies. In lymphoid malignancies, the most widely targeted cell surface proteins include CD19 and CD20 [90–97]. CAR NK cells for multiple myeloma (MM) have targeted CD138 and CS1 [44, 98]. CS1 is a cell surface glycoprotein highly and ubiquitously expressed on the surface of myeloma cells [99]. In our previous studies, we found that CS1-CAR NK cells were able to target MM cells and eradicate MM both in vitro and in vivo [44]. MM cancer stem cell-like cells (CSCs) expressed significantly higher levels of CS1 than any other cell type, making them especially vulnerable to being eradicated by CS1-CAR NK cells [100]. Thus, CS1-CAR NK cells may prevent MM relapse via targeting and eliminating MM CSCs. CD22, important in B cell trafficking, can be targeted for treatment of B-cell lymphoma [101]. CD7, important in lymphoid development and B/T cell cross talk, can be targeted in ALL, whereas CD33, a lectin inhibiting multiple cellular functions, can be targeted in AML [102].

As for solid tumors, a set of targets have been evaluated. A well-known target is HER2 that is up-regulated in several cancers including breast, colon, and ovarian cancers [103]. HER2-CAR cells using PB NK cells or NK-92 cells have been used to treat breast and ovarian cancers and neuroblastoma [104–108]. GD2 has been targeted for treating neuroblastoma using CAR NK-92 cells [109, 110]. GD2-CAR NK-92 cells effectively lyse multi-drug resistant neuroblastoma cell lines and show significant anti-tumor activity in xenograft mouse models [110]. Additional antigens being explored as CAR NK

cell targets in solid tumors include EGFR, GPA7, CD244, and EpCAM [44, 111–114].

EGFR is one of the most frequently amplified genes in glioblastoma, the most frequent and lethal type of brain tumor [115]. Approximately 20–40% of the EGFR-amplified tumors harbor the EGFR variant III (EGFRvIII). In our previous work, we found that EGFR-CAR NK cells targeted glioblastoma with both wild-type EGFR and EGFRvIII. EGFR-CAR NK cells were strongly cytotoxic, inhibited glioblastoma growth, and prolonged survival in tumor-bearing mice [112]. Use of CAR NK cells in combination with oncolytic viruses has been tested to eliminate solid tumors. Combination of EGFR-CAR NK-92 cells with herpes simplex virus 1-based oncolytic virus (oHSV-1) results in increased efficiency in killing a breast cancer cell line engrafted into the brain [116].

Additional targets being evaluated in recent clinical trials include a prostate-specific membrane antigen (PSMA) expressed over 100 times higher in prostate tumor tissues than normal prostate and the tumor differentiation antigen mesothelin in ovarian cancer [117, 118]. CAR NK cells and CAR NKT cells have been reported to target MUC1, a glycoprotein that creates a significant barrier preventing drugs from binding to their targets on cancer cells [119]. ROBO1, important in neuronal development, is being evaluated for treatment of pancreatic cancer [120].

Challenges in CAR NK production

Challenges in transducing CAR NK cells

Several methods including electroporation and viral transduction methods have not achieved full transduction efficiency in CAR NK cell production (Table 2). Major groups of viruses tested for transduction include retro-, lenti-, vaccinia-, and adenoviruses. Vaccinia- and adenoviruses are not well suited for NK cell transduction as they may change the phenotype of the cells or may not bind receptors on NK cells [121]. In addition, transduction with adenoviruses is transient as they rarely integrate into the host genome [122]. Although transduction with retro- and lentiviral vectors has the advantage of achieving stable transgene expression, they may induce oncogenic transformation due to their potential for insertional mutagenesis [66]. For example, leukemia due to insertional mutagenesis has been reported in patients undergoing gene therapy with CD34+ cells transduced with retroviruses to restore IL-2 receptor expression [123]. Several studies have reported that insertion of retroviruses is more common in transcriptionally active than silent genes and transcription factor binding sites and thus may not be a completely random event. Likewise, lentiviruses also show strong preference for integrating into the transcriptionally active genes [124, 125].

Retroviruses can only infect dividing cells. NK cells also have a short half-life in vivo, yet expressing membrane bound IL-15 increases viability and cytotoxicity in NK cells transduced with retroviruses [126]. Vectors based on lentiviruses, a subclass of retroviruses, are more popular likely because they are relatively safer [61].

Alternative methods developed to avoid risks associated with the viral delivery of genes have not been fully successful. A transposon-based gene delivery system has been reported, but its design still needs improvement [61]. mRNA delivery of CARs into NK cells has relatively high success rates, but the transfection is only transient. CD19-CAR NK generated by trogocytosis avoids side effects of viral transduction. However, this approach also has remained at the experimental stage [127]. Using nucleofection or electroporation for gene delivery results in 1–90% transient transfection efficiency with viability ranging from 45 to 97%, suggesting that the technology is not stable or very reliable.

Several approaches and compounds including dextran, PHA, and BX795 have been tested to improve transduction efficiency in NK cells (Table 2). Dextran, a cationic polymer, significantly increases transduction efficiency in human and mice primary NK cells transduced with lentiviruses [128]. Magnetofection improves lentivirus-mediated gene delivery [129]. A modified 3D nanochannel-electroporation (NEP) method that can transfect 100–200 cells at a single-cell resolution has been developed [130]. The method involves precisely positioning cells on nanochannels allowing precise transfection. The method has shown nearly 70% transfection efficiency using CAR NK cells. However, this technology also delivers CARs only transiently.

Challenges in expanding CAR NK cells

A second significant challenge in CAR NK research is the limited expansion of NK cells in vitro and especially in vivo. Expanding CAR NK cells, especially PB NK cells, to clinical scale in vitro is difficult because a large number of NK cells are needed for immunotherapy and NK cells comprise only about 10% of the PB lymphocytes and therefore need to be expanded in culture [131]. Several methods have been developed to improve in vitro expansion rates in purified PB NK cells [90, 132–135]. A common method involves co-culturing PB NK cells with K562 feeder cells genetically modified to express IL-15 and the adhesion molecule 4-1BBL [90]. Telomere shortening seen in NK cells expanded by this technology can be prevented by expressing membrane bound IL-21 [135]. An approximate 2000-fold expansion of the NK cells from fresh or cryopreserved UCB in two weeks was achieved using these artificial antigen-presenting cells [136]. Expanded cells retain high expression levels

Table 2 Studies aimed at improving transduction efficiency in NK cells

Vector	Cells used for transduction	Transduction efficiency	Main findings	References
Lentivirus	Primary NK cells	29% (range 16–41%) with excellent viability	Multiplicity of infection (MOI) of 25 is optimal. Higher MOI results in higher transduction but lower viability. Transgene expression peaked at 5 days post-transduction and was detectable for 2 weeks	[222]
Lentivirus	Primary NK cells and the YTS NK cell line	98% in the YTS NK cell line, up to 80% in primary NK cells	Transduction efficiency was 20% when fresh primary NK cells were transduced without cytokines. The efficiency increased up to 20–50% when IL-2 plus IL-12 were added and 80% when PHA, a lectin, was added to the culture. The transduction efficiency remained stable in the presence of PHA for 10 days in culture	[223]
Lentivirus	Primary NK cells	Depending on cell type, BX795 addition increased transduction efficiency up to tenfold	Inhibition of TBK1/IKKε complex by BX795 significantly improved transduction efficiency. Lentivirus RNA is probably recognized by pathways involving TBK1/IKKε complex involved in anti-viral response pathways, which may also include RIG-I, MDA-5, and TLR3	[224]
Lentivirus	Primary NK cells (mice)	40%	IL-2 is not required for transduction	[225]
mRNA transfection and lentivirus	Primary NK cells, NK-92, and NK cells from UCB	In NK-92 (56% with mRNA, 26% with lentivirus), in primary NK (approximately 10% with mRNA), and in UCB (12% to 73% with lentivirus)	mRNA delivery was more effective than lentiviral transduction in transducing the NK-92 cell line (56% vs. 26%)	[226]
Lentivirus	NK-92, LNK, YT, and DERL7 cell lines	15% in NK-92 and 30–40% in LNK, YT, and DERL7 cell lines	Lentiviruses are more efficient than retroviruses that require multiple rounds of transduction	[121]
Retrovirus	Primary NK cells	27–47% on days 5–6 of the culture and 52–75% 21 days after initial culture	MOI of 10 yields highest transfection efficiency	[227]
Retrovirus	UCB NK cells	49%	NK cells expressing IL-15 were detectable up to a year after infusion	[68]
Retrovirus	UCB NK cells	66.6%	NK cells purified from UCB and transduced with iCas-pase-9, IL-15, and a CD19-specific CAR showed significant toxicity toward CD19+ tumors	[83]

of NKp44, NKG2D, TRAIL, and FasL, and their killing activity does not decrease compared to the fresh NK cells [137]. Nevertheless, NK cell exhaustion and loss of expansion capacity remain as concerns in long-term culturing exceeding 4 weeks.

Limited *in vivo* expansion of NK cells after infusion due to unknown reasons is an especially significant challenge. Studies using iPSCs show that *in vivo* expansion rates in CAR NK and CAR T cells are correlated with their anti-tumor activity [89]. IL-2 diphtheria toxin fusion protein (IL-2DT) infusion may help *in vivo* expansion of haplo-identical NK cells in AML patients by inhibiting Tregs [138].

A recent phase 1/2 clinical trial has shown that incorporating IL-15 into anti-CD19 CAR NK cells results in significant improvement in *in vivo* expansion [68]. A UCB unit was thawed and cultured in the presence of K562 expressing IL-21 and 4-1BBL and IL-2. Cells were transduced with an anti-CD19 CAR NK vector expressing IL-15 and iCaspase-9 on day 6, expanded, and harvested for infusion on day 15. As noted above, eight out of the eleven (73%) patients had responses up to 13.8 months. Expansion, as measured by real-time PCR (RT-PCR) (copies of the viral genome per micrograms of genomic DNA), was detected three days after infusion and persisted for over a year. Expansion was seen only in patients that responded to the treatment.

Challenges in CAR NK cell therapy in solid tumors

The exciting results from Liu et al. observed in CAR NK cell therapy in liquid tumors [68] have not yet been seen in solid tumors possibly due to challenges such as

solid tumor heterogeneity and the hostile tumor micro-environment (TME). CD19 in leukemia and lymphoma is highly and homogeneously expressed on the surface of malignant B cells, and the CD19-CAR NK cells do not face substantial anatomical barriers in the blood compartment before establishing a contact with their targets. In solid tumors, however, the CAR NK cells must travel in bloodstream, migrate into the tissue, and establish a contact with the tumor cells while battling many inhibitory factors in the TME (Table 3).

CAR NK cell therapy seems more promising in breast, ovarian, and prostate cancers than other types of solid tumors because these tissues are easily and safely accessible and destruction of the normal tissue can be tolerated. In addition, specific antigen targets such as PSMA or prostate stem cell antigen (PSCA) are already available for targeting prostate cancer [139]. Indeed, transduction of YTS cell lines with DAP12-based anti-PSCA CAR induces strong phosphorylation of ZAP70 and fully eradicates tumors in a portion of the mice tested [140]. CAR NK-92 cells expressing EpCAM accumulate in prostate tumor and are highly cytotoxic against EpCAM+ prostate cancer cells even at low doses [141]. iPSCs-derived CAR-NK cells targeting mesothelin have shown to have similar efficacy similar to that of CAR-T cells but with fewer toxicities [89].

In breast cancer, NK-92 cells expressing a HER2-specific CAR enhance killing of human breast cancer cell lines and increase survival in mice breast and renal cell carcinoma (RCC) models [107]. CAR NK cells targeting EGFR derived from PB are effective in triple-negative breast cancer (TNBC) both *in vitro* and *in vivo* [142].

Table 3 Challenges in building more efficient CAR NK cells for treatment of solid tumors and possible solutions^a

Challenges	Possible solutions	References
Lack of available targets		
TAA expression not homogenous	Use of oncolytic viruses in combination therapy or bi-specific CARs	[116, 228]
Clonal evolution		
On-target, off-tumor effects		
Anatomical barriers		
Insufficient trafficking or infiltration	Intra-tumoral injection of CAR NK cells; focused ultrasound guided delivery of CAR NK cells into tumor; targeting tumor vasculature; CAR NK cells expressing a chemokine(s)	[149, 155]
	Locoregional injection, use of stronger co-stimulatory domains, lymph depletion, <i>in vivo</i> administration of chemokine or cytokine-expressing CARs	[146, 169]
Insufficient NK cell proliferation and/or activation <i>in vivo</i>		
Immunosuppressive TME	TGF- β inhibition through neutralizing antibodies or dominant negative receptors	[229]
	Use of cytokines such as IL-2, IL-15, IL-12, or IL-18 to simulate NK cell proliferation and activation	[68]
	Inhibition of checkpoint blockades using anti-PD1, anti-TIGIT, etc	[165]
Metabolic abnormalities	Metabolic stimulation through inhibition of CD73 or arginase	[162]
NKG2D inhibition by TME	NKG2D activation through histone deacetylase inhibitors	[160, 230]

^a TAA tumor associated antigens, TME tumor microenvironment

A recent study focused on tissue factor (TF) highly expressed in 50–85% of the TNBC patient tumors with a FVII as a ligand [143]. FVII-CAR NK-92MI cells expressing CD16 are effective in vivo and enhance tumor killing by TF-targeting antibody-like immunoconjugates (L-ICON), in which FVII was fused to the Fc fraction of IgG1. Major challenges in the use of CAR NK cells for treatment of solid tumors can be categorized into the following groups:

Lack of available targets and antigen heterogeneity The first step in designing CARs is to locate a target antigen highly and uniformly expressed on the tumor cell surface. In solid tumors, most TAAs are expressed by cells not only in tumor but also in vital organs, making it impossible to avoid “on-target, off-tumor” effects [144]. In addition, TAA expression may be dramatically different between single cell clones from the same tumor due to clonal evolution and downregulation of TAA expression by tumors to evade immune surveillance. Bispecific CARs, composed of two scFvs recognizing two different antigens on the same CAR and activated only when both antigens are engaged, have been successful in improving specificity in prostate cancer [145].

Anatomical barriers Local injection, intra-peritoneal injection, and focused ultrasound-guided delivery of CARs into tissues have been experimented to overcome the anatomical barriers faced by the immune cells carrying CARs in solid tumors. In orthotopic models mimicking human pleural malignancies, pleural injection of CAR T cells is more effective and results in longer functional persistence than the *i.v.* injection [146]. Intrapleurally administered CAR T cells were able to circulate and persist in the periphery, suggesting that locally accessible sites may serve as “distribution centers” for CAR T therapy. Regional administration of CAR T cells was additionally useful in reducing the therapeutic dose of the CAR T cells. Endothelial cells in tumor vasculature have been targeted and successfully lysed by anti-PSMA-CAR T cells [147], as PSMA is expressed in neovasculature [148]. Anti-HER2 CAR NK-92 cells were delivered into the brain of mice with metastatic breast cancer using focused ultrasound [149]. The authors applied ultrasound bursts through intact skull combined with *i.v.* injection of microbubbles that temporarily allowed passage of NK-92 cells from the blood brain barrier into the brain without significant tissue damage.

Limited CAR NK cell trafficking and infiltration into tumor sites The number of NK cells infiltrating into tumor has a prognostic value in solid tumors including gastric, lung, and colorectal cancers [150]. CARs that express

chemokine receptors (CCRs) matching CC chemokine ligands (CCLs) expressed by the tumor cells acquire increased infiltration capabilities. Wennerberg et al. (2015) reported that in vitro expansion of NK cells was associated with a tenfold increase in CXCR3 expression compared to resting NK cells. Expanded cells migrated strongly toward CXCL10-transfected melanoma, but not CXCL10-negative tumor cells [151]. NK cells expressing CXCR2 show increased migration toward tumors expressing CXCR2 ligands [152]. The cells additionally show increased calcium mobilization, cytotoxicity, and adhesion properties. Overexpression of CXCR4 in anti-EGFRvIII CAR NK cells results in increased chemotaxis toward CXCL12/SDF-1 α -secreting glioblastoma cells, complete tumor remission in a number of mice, and increased overall survival [153]. Oncolytic viruses are anti-tumorigenic and may also be anti-angiogenic through negative regulation of VEGF expression [154, 155]. In addition, oncolytic viruses induce an inflammatory immune response and enhance immune cell trafficking by modifying the TME [155]. Our group first explored the concept of combining anti-EGFR CAR NK cells and oncolytic virus therapy in a metastatic breast cancer animal model. The combination of EGFR-CAR NK-92 cells with oHSV-1 resulted in more efficient killing of breast cancer tumor cells engrafted into the brain compared to monotherapies and significantly improved survival of mice bearing breast cancer tumor cells in the brain [116].

Immunosuppression in the TME due to TGF- β Tumors express immunosuppressive factors including TGF- β , IL-10, PD-1, or arginase. In vitro expanded NK cells are highly cytotoxic against cancer cells. However, they lose this ability upon injection in vivo. TGF- β is produced by neutrophils, macrophages, and Tregs in the TME. Tregs and immunosuppressive myeloid-derived suppressor cells (MDSCs) are actively recruited to the TME to create a strong immunosuppressive environment promoting tumor growth [156].

Several strategies to minimize inhibitory effects of TGF- β have been reported. Administration of TGF- β kinase inhibitors along with NK cells preserves the cytotoxic potential and expression of activating NK receptors NKG2D and CD16 [157]. Use of fresolimumab, a TGF- β neutralizing antibody, and galunisertib, an inhibitor of TGF- β RI, have been reported as safe and tolerable in solid tumors in humans [158]. Hybrid CARs with an extracellular TGF- β receptor domain linked to an intracellular NKG2D domain enhance anti-tumor activity in NK-92 cells and knocking down SMAD3, the downstream mediator of TGF- β , in NK cells improves cytotoxicity in solid tumors [75, 159]. UCB NK cells expressing a dominant-negative TGF- β receptor retain their capacity

to secrete IFN- γ and kill glioblastoma cells in the presence of TGF- β . Entinostat, a narrow spectrum histone deacetylase inhibitor, increases MICA expression on tumor cells and NKG2D expression in primary NK cells even in hypoxic conditions, leading to enhancement of cytotoxicity of NK cells against tumor cells [160]. It enhances both tumor cell recognition and killing.

Immunosuppression in the TME due to metabolic disturbances The TME additionally suffers from hypoxia and nutrient depletion leading to acidosis, which further suppresses immune responses [156]. Hypoxia supports tumor growth by inducing metabolic disturbances and increasing angiogenesis and expression of tumor growth factors. In addition, it downregulates expression of NK cell activating receptors NKp30, NKp46, NKp44, and NKG2D for optimal tumor growth and metastasis [161]. CD73 induces expression of arginase, an immunosuppressive metabolite. Arginase is produced under hypoxic conditions and inhibits NK cell functions. CD73 inhibition increases homing of NKG2D-CAR NK cells, which target tumor cells expressing NKG2D ligands, to the tumor site and improves anti-tumor responses in animal models for lung cancer [162].

Immunosuppression in the TME due to checkpoints Tumor cells evade immune surveillance through expression of checkpoint proteins that prevent immune cell activation. TIGIT inhibits NK cell cytotoxicity by opposing CD226 [163]. PD-1 + NK cells show reduced proliferation and effector functions, whereas PD-L1 + NK cells possess better effector functions [164]. Inhibition of PD-1 or PD-L1 using checkpoint blocking agents reactivates exhausted immune cells and achieves durable clinical responses [165]. Combining CARs with blockers of checkpoint proteins such as PD-1, CTLA-4, LAG3, and TIGIT has the potential to achieve persistent therapeutic benefits [145]. NK-92 cells engineered to express high-affinity CD16, IL-2, and a PD-L1-specific CAR contain high levels of perforin and granzyme and lyse human cancer cell lines including breast, lung, and gastric cancers [166].

Efforts to build more efficient CARs

Although CAR NK cells and CAR T cells have opened a new era in immunotherapy, results from clinical studies suggest that improvement is still needed. Key areas that need further attention for improvement include modifying CAR structure for better efficiency and/or suitability to the TME, over-expressing activating and/or blocking inhibitory receptors on NK cells, improving cytotoxic mechanisms involving death receptors or granule-dependent pathways, and/or use of multiple

compounds to increase CAR efficacy. Efforts to optimize CAR NK cell efficacy are extensive but can be categorized into the groups below:

CARs targeting multiple proteins

Some of the recent CAR NK-92 constructs include TCR-CARs created by fusing the extracellular domain of TCR (both TCR α and TCR β) to a CAR-signaling tail composed of CD28 and CD3 ζ signaling domains to enable the cells to recognize an array of targets, possibly some of the proteins presented on the MHC molecules, rather than a single TAA [167]. Expression of TCR-CAR in NK-92 cells grafts the cells with target recognition specificity of TCRs. This design has the potential of allowing the CAR NK-92 cells to target the proteome derived from tumor cells [167]. It would be interesting to determine whether this approach would work in primary NK cells or NK cells from other sources.

CAR-modified immune cells secreting enzymes, interleukins, or chemokines

“Armored” CAR design involves expressing cytokines, chemokines, or their receptors in CAR-modified immune cells to respond to threats from the TME or protect the CAR-bearing cells from immune suppression. CAR T cells expressing enzymes such as catalase, which can convert H₂O₂ into harmless by-products, or cytokines such as IL-15 have been developed [168]. CAR NK cells secreting autocrine IL-2 and IL-15 and silencing inhibitory KIRs have improved in vivo persistence [169]. CARs specific for various CC chemokines and/or their receptors have been reviewed above in the section on “[Limited CAR trafficking and infiltration into tumor sites](#)”.

Converter CARs

While some CARs provide the immune cells with more “armors” against the hostile tumor cells or the TME, others have been designed to “convert” the hostile actions into desirable effector functions. This “converter” design includes combining ecto- and endodomains of different molecules that graft specificity of the former coupled with the signaling capacity of the latter onto the immune cell. For example, CARs that combine ectodomain of IL-4 with the endodomain of IL-7 convert the inhibitory effects of IL-4 in the TME into the proliferative actions of IL-7 [170]. Similarly, CARs converting immunosuppressive effects of TGF- β into activating functions of NKG2D by fusing the extracellular and transmembrane portions of TGF- β RII and the intracellular domains of NKG2D, an activating receptor, have been reported [171]. Other variants of converter CARs include double chimeric design that involves introducing two CARs with different scFvs but the same intracellular signaling domain

(dual-signaling CAR) [172]. A recent study sought to interfere with PD-1 signaling to reverse immunosuppression by PD-1 [173]. The authors fused the extracellular domain of PD-1, transmembrane and cytoplasmic domains of NKG2D, and the cytoplasmic domain of 4-1BB to create a chimeric co-stimulatory converting receptor-NK-92 cell line that reversed the inhibitory PD-1 signaling and increased anti-tumor activity against lung cancer cells [173].

Inducible CARs

A fourth-generation CAR, also called TRUCK, has been developed [172]. In this design, inducible IL-12 is expressed only after encounter with the tumor specific antigens. IL-12 release activates innate immune cells that kill tumor cells carrying the specific antigen targeted by CAR. A form of TRUCK activated by inducible HIF1 α in the TME has been reported [61]. Inducible expression of CAR can also be achieved using an injectable drug. A version of inducible CARs also termed “trans-signaling CARs” has been developed. These designs are especially useful in increasing safety and specificity of CARs when multiple proteins or isoforms of the same protein are targeted. Such CARs may engage antigens when either of multiple isoforms is available to increase range of efficacy or may require binding to all isoforms of the target for activation for increased specificity [172].

Modification of CARs using the CRISPR/Cas9 system

Recent advances in CRISPR/Cas9 technology provide additional opportunities by enabling researchers to insert a CAR permanently into a desired position in the genome rather than insertion into random locations employed by most of the current viral transduction methods [174]. Another advantage of the CRISPR/Cas9 system is that it can be used to screen the whole genome in loss-of-function studies to reveal genomic regions or pathways important in mediating CAR toxicity. This approach has identified FADD and TNFRSF10B (TRAIL-R2) in CAR T cells but has not yet been studied in CAR NK cells [175]. Simultaneous administration of compounds to regulate these pathways can improve CAR effector functions.

The CRISPR/Cas9 technology holds great promise in advancing the CAR NK cell research through boosting effector functions, especially for iPSC-derived CAR NK cells [89]. It can be used to knock out several genes simultaneously [176]. It is especially advantageous to use in modifying NK cells due to the fact that NK cell activation is achieved through a balance between inhibitory and activating signals. This technology can knock-in activating while simultaneously knocking-out inhibitory signals. It is known that viral integration rates are higher in genomic regions with active transcription or highly

accessible chromatin [177]. The CRISPR/Cas9 system can be used to insert a CAR cassette into these regions to generate stably transduced primary NK cells, improving both efficacy and safety of CAR NK cells [61]. CRISPR/Cas9 has already been used to “knock-in” a CAR while simultaneously “knocking-out” endogenous elements such as components of the TCR that may interfere with the CAR functions in CAR T cells [178, 179].

Only few CRISPR/Cas9 studies have been conducted using NK cells possibly because NK cells are more difficult to transduce [180]. Two studies that successfully transfected NK cells with Cas9 protein and target-site-specific gRNA or DNA with homology to sequences flanking DNA double strand breaks using electroporation have been reviewed elsewhere [180]. In one of those studies, 60% and 34% reduction at the mRNA and protein level, respectively, in TGF- β R II expression was achieved [181]. Another study improved post-transduction survival up to 80% while maintaining efficient reduction in *Ptprc* gene expression [182]. This study additionally revealed that both homo- and heterozygous deletion of the gene of interest are possible in the same reaction.

Despite the low number of studies reporting CAR NK cells created using the CRISPR/Cas9 technology, multiplex knockout of three genes (*TRAC*, *B2M*, and *PD-1*) in CD19-CAR T cells has already been achieved [183]. However, low knock-out rates for PD-1 were reported in that study. Another recent study was successful in efficiently knocking out PD-1 in CAR T cells targeting EGFRvIII [184]. The disruption did not change T cell phenotype or expression of other checkpoint receptors and enhanced activity of the CAR T cells. TGF- β R II has also been knocked out in CAR T cells, resulting in decreased T cell exhaustion [185].

Use of CARs along with monoclonal antibodies, nanoparticles, or chemical compounds for increased efficacy
NK cells can be used in tandem with antibodies, radiation, chemotherapy [186], romidepsin, a histone deacetylase inhibitor, nanoparticles [187, 188], and/or cabozantinib, a tyrosine kinase inhibitor [189]. Romidepsin improves efficacy of CD20-CAR NK cells [187], and cabozantinib enhances effects of CAR NK-92 directed against EGFR [189]. Use of CAR NK cells along with monoclonal antibodies widely used in cancer immunotherapy represents an interesting field of research. Bi- and tri-specific killer cell engagers (BiKEs and TriKEs) have been used to improve NK cell anti-tumor activity via the CD16 (Fc γ RIII) receptor on NK cells [190]. A majority of NK cells express CD16 that enables them to kill tumor cells opsonized by therapeutic monoclonal antibodies such as rituximab through ADCC [191]. An excellent

review on potential of NK cells in combination therapy with monoclonal antibodies is available elsewhere [192].

CAR-engineered hematopoietic stem cells

CAR-engineered stem cells represent a viable option to eradicate viral infections or cancers as they can differentiate into CAR T cells or CAR NK cells. Hematopoietic stem/progenitor cell (HSPC) populations transfected with a CAR against HIV provide long-term immunity against HIV infections [193]. The CAR-stem cells differentiated into functional NK and T cells and suppressed HIV replication in vivo and in vitro. Since stem cells are self-renewing, these CAR-modified immune cells may lead to permanent remission [194]. This method has implications for transplantation studies and immunity against viral infections in addition to cancer immunotherapy. Protocols for generating NK cells through differentiation of genetically modified hematopoietic stem cells have already been developed [195].

Optimization of CAR signaling domain

Incorporation of 4-1BB, a surface protein expressed by NK and activated T-cells, improves NK cell cytotoxicity and IFN γ production [93, 94]. Although not expressed in NK cells, CD28 is commonly incorporated into CAR design for NK cells to amplify extracellular signaling. NK-92 cells carrying CARs with CD28/CD3 ζ and 4-1BB/CD3 ζ signaling domains have stronger cytotoxicity than CARs with only CD3 ζ in killing HER2-expressing breast carcinoma cells [107]. Stimulation with 4-1BB is more effective than CD28 in expanding CD8 $^+$ T cells, and it has been shown that incorporation of both CD28 and 4-1BB co-stimulatory domains improves in vivo persistence in CAR T cells [196]. Additional potential co-stimulatory molecules being explored in CAR design for NK cells include 2B4 (CD224), DAP10, and DAP12 [94, 113, 140, 197].

Although 2B4 signaling alone only weakly induces NK cell activation, CARs including 2B4 along with CD3 ζ enhance all aspects of the NK cell activation against leukemia and neuroblastoma cells [113]. Additionally, CARs utilizing a signaling domain for DAP12, which is involved in signal transduction of activating receptors such as NKG2C, NKp44, and KIR3DS1 [198, 199], exhibit improved NK cell cytotoxicity compared to CARs with CD3 ζ as the sole signaling domain [140].

Manipulation of immunologic synapse formation and NK cell subsets

Immunologic synapse formation by CAR NK cells and possible differences in CAR production efficiency among different subtypes of NK cells have not been fully investigated in the literature. Synapse formation is

a step-wise process that involves distribution, rearrangement and clustering of signaling molecules, F-actin, and tumor antigens at the site of contact. Synapse formation between NK cells and various target cells has been extensively investigated [200]. The NK IS formation involves central and peripheral supramolecular activation cluster (SMAC). The central and peripheral SMACs involve actions by distinct sets of proteins. The central domain includes mostly perforin, whereas the peripheral domain consists of surface receptors and F-actin. Upon stimulation, surface receptors including CD2, CD11a, CD11b, and F-actin rapidly accumulate at peripheral SMAC. This accumulation is dependent on actin polymerization driven by Wiskott–Aldrich syndrome protein. Accumulation of perforin at central SMAC is, on the other hand, slow and appears only after peripheral SMAC activation. The strong need for studying IS formation in CAR NK cells as well as critical steps of IS formation during the killing of target cells by NK cells has been reviewed elsewhere [201].

Similarities and differences between the conventional and CAR NK cell signaling events occurring at IS have not been well understood. A better understanding and possible manipulation of the IS formation can significantly improve CAR efficacy [201] because IS is critical for successful elimination of the target [202]. During natural T cell receptor activation, CD4/CD8 T cells interact with MHC molecules on antigen presenting cells (APCs) resulting in activation of Lck which then phosphorylates ZAP70, resulting in signal 1 for T cell activation. Signal 2 required for activation is a result of CD28 on T cells interacting with CD80/CD86 on APCs creating an IS with a 15-nm gap [203]. The 15-nm distance physically excludes other inhibitory molecules such as CD45 with large domains from entering into IS. It is unknown whether these processes occur similarly at IS formed between the CAR NK cells and their targets.

Studying different NK cell subsets carrying the same CAR may reveal subtypes more efficient than others for specific purposes [204]. It is estimated that there are 6000–30,000 different NK cell phenotypes in a healthy donor due to the high number of activating receptors with different functions expressed at different stages of development [138]. However, only limited data are available on the characteristics of these subtypes of NK cells.

Enhancement of CAR NK cell activation and proliferation by cytokines

NK cells in patients with tumors are often functionally impaired and live shorter than those in healthy individuals. Cytokines, or their analogs with longer in vivo half-life, can be used to increase NK cell activation, proliferation, trafficking, and persistence. Effects of several

cytokines including IL-2, IL-12, IL-15, IL-18, and IL-21 on NK cell function have been reviewed in detail elsewhere [75]. The cytokines can be administered *in vivo* or used for *in vitro* stimulation before adoptive cell transfer. NK cells require IL-2 or IL-15 for optimal growth. NK-92MI, an NK-92 cell line constitutively expressing IL-2, has been generated [114, 143]. IL-2 and IL-15 are both included in the γ chain family of cytokines and share some common functions [114]. Co-expressing IL-15 and an EpCAM CAR in NK-92 cells resulted in a predominant intracellular accumulation of the cytokine and selection of CAR-bearing cells upon IL-2 withdrawal, which eliminated all but the transduced NK-92 cells. These CAR NK cells successfully lysed breast cancer cells in the absence of any other cytokine [114].

IL-2 is safe to administer in lung cancer patients and induces proliferation and cytotoxic capacity in NK cells, but it also attracts and stimulates immunosuppressive Tregs at high doses [205]. IL-15 may be more effective in improving NK cell function because it is anti-neoplastic, more potent than IL-2, and does not attract Tregs. CAR NK cells may be stimulated by IL-15 through co-culturing with feeder K562 cells expressing membrane-bound IL-15 and 4-1BB ligand [94]. Co-culturing PB NK cells with K562-based feeder cells expressing membrane-bound IL-21 significantly improves NK cell expansion *in vitro*. IL-12 at the dose of 500 ng/kg/day was initially found safe in a phase 1 clinical trial. However, the same dose resulted in serious side effects in 12 of 17 patients enrolled in the study and two patients died, leading to a ban by the FDA on all clinical trials involving IL-12 [206].

CAR NK activation and proliferation through manipulation of NK cell receptors

NK cell effector functions can be improved through inhibiting inhibitory KIRs such as KIR2DL-1, -2, or -3 and/or overexpressing activating KIRs. Inhibitory receptors (as listed by Pockley et al. [205] with their ligands on tumor indicated in parentheses) include PD-1 (PD-L1), KIRs (HLA-C), LAG3 (MHC II), NKG2A/CD94 complex (HLA-E), TIGIT (CD155), TIM3 (Galectin-9), and KLRG-1 (E-cadherin), whereas activating NK receptors include NKG2D (MICA/B), KIRs (HLA-C), CD226 or DNAM-1 (CD112 or Nectin-2, CD155 or PVR), and NKG2C/CD94 complex (HLA-E and Hsp70). In addition, receptors 2B4 (CD244), CS1 (CD319), and LLT1 (CLEC2D) have been identified on NK cells. 2B4 and CS1 regulate NK cell cytotoxicity, whereas LLT1 is additionally expressed in prostate and TNBC cells and plays a role in evading NK cell-mediated tumor cell killing [207].

One of the most studied NK cell receptors is NKG2D. Histone deacetylase inhibitors such as sodium valproate enhance NK-mediated cell killing by increasing

expression of NKG2D ligands MHC class I chain-related molecules MICA and MICB on tumor cells [208]. Parihar et al. (2019) created CAR NK cells expressing a chimeric receptor in which NKG2D is fused to CD3 ζ [209]. These CAR NK cells reduced tumor burden through killing intra-tumoral MDSCs that expressed high levels of NKG2D ligands and, unlike unmodified NK cells, released cytokines even in the TME. The CAR NK cells additionally improved CAR T cell trafficking to the tumor site through chemokine secretion. The authors concluded that these CAR NK cells should be used in combination with CAR T cells for optimum efficacy in the TME in solid tumors.

Use of anti-KIR antibodies such as lirilumab to enhance *in vivo* CAR NK effector functions represents a novel approach in the treatment of solid tumors. Lirilumab is a fully human IgG4 mAb [210] that improves NK cell anti-tumor functions by inhibiting KIR2D subgroup [211] receptors on NK cells and blocking the repressive KIR-MHC interactions [210]. Lirilumab efficacy has been tested in enhancing NK effector functions in both liquid and solid tumors [210]. Lirilumab, in combination with ipilimumab, is being tested for treatment of advanced solid tumors in a clinical trial (NCT01750580), but the trial is still recruiting patients [212]. *In vitro* evidence suggests that lirilumab enhances killing of head and neck squamous cell carcinoma (HNSCC) cells by NK cells [213]. Enhancement of anti-tumoral effects of lirilumab, in combination with rituximab, has been well demonstrated in hematologic malignancies both *in vivo* and *in vitro* [214]. In myelodysplastic syndrome patients, lirilumab is effective both as a monotherapy and in combination with azacitidine although the authors cautioned that further studies are needed to confirm their findings [215]. IPH2101, a prior version of lirilumab that recognizes the same epitope, has also been tested in the clinic [216–218]. It will be interesting to determine if abrogation of KIR inhibitory signaling by KIR antibodies will further improve anti-tumor activity of CAR NK cells.

Modifying cell metabolism to improve CAR NK effector functions

Enhancing CAR NK cell activity in solid tumors through modulation of tumor metabolism has received little attention in the literature. Extracellular adenosine generated from ATP present in the hypoxic tumor environment through actions of CD39 and CD73 plays critical roles in immune evasion, inhibition of NK cell maturation, and suppression of NK cell trafficking to the tumor site [162]. Adenosine inhibition using anti-CD39 and anti-CD73 antibodies improves targeted therapy in ovarian cancer [219]. CD73 is highly expressed in several solid tumors including glioblastoma, prostate, and lung cancer,

and its inhibition using neutralizing antibodies further improves therapeutic effects of NKG2D-engineered CAR NK cells in a mice model of lung cancer [162].

CAR NK cell clinical trials

Currently, there are 19 CAR NK clinical trials available on *ClinicalTrials.Gov* website (Table 4). Eleven and eight trials have targeted liquid and solid tumors, respectively. The most common targets (with number of clinical trials in parentheses) are CD19 (7), ROBO1 (3), and CD22 (2). One clinical trial has been completed, eight have been recruiting patients, one is suspended, and one is withdrawn. Five trials have not begun recruiting patients, and the status of the remaining three trials is unknown. Five trials are at early phase 1, four at phase 1, and ten are at phase 1 or 2 stages. Fourteen of the trials originated from China, three from the USA, one from Singapore, and one from Germany.

In one of those studies, a phase 1/2 trial (NCT02944162) targeting CD33 in AML patients had three patients who were administered nearly five billion irradiated CD33-CAR NK-92 cells. Although

therapeutic response was limited, the patients did not show any significant adverse reactions after administration of up to five billion NK-92 cells, suggesting that injection of a large number of NK-92 cells may be safe [67]. A recent promising phase 1/2 clinical trial in the USA (NCT03056339) is being conducted to investigate the safety and efficacy of primary NK cells obtained from UCB carrying a CD19-CD28-CD3 ζ -2A-iCasp9-IL15 construct in patients with relapsed/refractory B-cell ALL, CLL, and non-Hodgkin lymphoma. The CAR NK treatment was carried out in conjunction with lymphodepleting chemotherapy. Very encouraging results from this trial have recently been published [68]. As summarized above, eight of the 11 patients responded to the therapy. Importantly, NK cells persisted in patients for over 12 months possibly due to the incorporation of IL-15 in the CAR vector. A clinical trial being conducted in Germany has been investigating the safety and tolerability of NK-92 cells carrying anti-HER2 CAR in glioblastoma patients (NCT03383978).

Table 4 Current CAR NK clinical trials listed on *ClinicalTrials.Gov*

Tumor type	Tumor type	Target antigen	NCT number	Types of NK cells	Status	Phase	Country
ALL	Liquid	CD19	NCT00995137	Haploidentical donor NK cells	Completed	1	USA
Solid tumors	Solid	NKG2D ligands	NCT03415100	Autologous or allogeneic CAR NK cells targeting NKG2D ligands	Recruiting	1	China
B-lymphoid malignancy, ALL, CLL, lymphoma	Liquid	CD19	NCT03056339	UCB NK cells	Recruiting	1/2	USA
Multiple myeloma	Liquid	BCMA	NCT03940833	NK-92 cells	Recruiting	1/2	China
Solid tumors	Solid	ROBO1	NCT03940820	Unclear	Recruiting	1/2	China
Pancreatic cancer	Solid	ROBO1	NCT03941457	Unclear	Recruiting	1/2	China
Solid tumors	Solid	ROBO1	NCT03931720	Unclear	Recruiting	1/2	China
Glioblastoma	Solid	HER2	NCT03383978	NK-92 cells	Recruiting	1	Germany
B-cell acute lymphoblastic leukemia	Liquid	CD19	NCT01974479	Haploidentical donor NK cells	Suspended	1	Singapore
AML	Liquid	CD33	NCT02944162	NK-92 cells	Recruiting	1/2	China
Mantle cell, follicular, and B-cell lymphoma	Liquid	CD19	NCT03579927	UCB NK cells	Withdrawn	1/2	USA
Refractory B-cell lymphoma	Liquid	CD22	NCT03692767	Unclear	Not yet recruiting	Early 1	China
Refractory B-cell lymphoma	Liquid	CD19	NCT03690310	Unclear	Not yet recruiting	Early 1	China
Ovarian cancer	Solid	Mesothelin	NCT03692637	Unclear	Not yet recruiting	Early 1	China
Prostate cancer	Solid	PSMA	NCT03692663	Unclear	Not yet recruiting	Early 1	China
Refractory B-cell lymphoma	Liquid	CD19, CD22	NCT03824964	Unclear	Not yet recruiting	Early 1	China
ALL, CLL, mantle cell and follicular lymphoma	Liquid	CD19	NCT02892695	NK-92 cells	Unknown	1/2	China
ALL, T-cell leukemia and lymphoma	Liquid	CD7	NCT02742727	NK-92 cells	Unknown	1/2	China
Hepatic, lung, pancreas, breast, colon, and gastric cancer	Solid	MUC1	NCT02839954	Unclear	Unknown	1/2	China

Conclusions

CAR-engineered T cells have already achieved impressive responses in treating some hematologic malignancies. NK cells offer an additional advantage in that they may not require any immunological matching between donor and patients and thus have the potential for developing “off-the-shelf” products for cancer immunotherapy. Although some challenges still exist, the recent success involving CD19-CAR NK cells expressing IL-15 [68] suggests that a broader application and advancement in using “off-the-shelf” CAR NK cell products for the treatment of hematologic malignancies and solid tumors may occur in the near future.

Abbreviations

ADCC: Antibody-dependent cellular cytotoxicity; ALL: Acute lymphoblastic leukemia; AML: Acute myeloid leukemia; APC: Antigen presenting cell; CAR: Chimeric antigen receptor; CCL: CC chemokine ligand; CCR: CC chemokine receptor; CLL: Chronic lymphocytic leukemia; CR: Complete remission; CRi: Complete remission with incomplete hematologic recovery; CRS: Cytokine release syndrome; CSC: Cancer stem cell-like cell; DLBCL: Diffuse large B-cell lymphoma; FasL: Fas ligand; FDA: US Food and Drug Administration; GM-CSF: Granulocyte-macrophage colony stimulating factor; GvHD: Graft vs. host disease; HLA: Human leukocyte antigen; HNSCC: Head and neck squamous cell carcinoma; HSPC: Hematopoietic stem/progenitor cell; IFN- γ : Interferon- γ ; IL-2DT: IL-2 diphtheria toxin fusion protein; ILC: Innate lymphoid cell; iPSC: Induced pluripotent stem cell; IS: Immunologic synapse; ITAM: Immunoreceptor tyrosine-based activation motif; ITT: Intent-to-treatment; KIR: Killer immunoglobulin receptor; L-ICON: Antibody-like immunoconjugates; MDSC: Myeloid-derived suppressor cell; MM: Multiple myeloma; MOI: Multiplicity of infection; NCR: Natural cytotoxicity receptor; NEP: 3D nanochannel-electroporation; NK: Natural killer; NSG: NOD scid gamma; oHSV-1: Herpes simplex virus 1-based oncolytic virus; ORR: Overall remission rate; PB: Peripheral blood; PSMA: Prostate-specific membrane antigen; PSCA: Prostate stem cell antigen; RAG: Recombination activating gene; RCC: Renal cell carcinoma; RT-PCR: Real-time PCR; scFv: Single-chain antibody variable fragment; SMAC: Supramolecular activation cluster; TAA: Tumor-associated antigen; TCR: T cell receptor; TF: Tissue factor; TGF- β : Transforming growth factor-beta; TME: Tumor microenvironment; TNBC: Triple-negative breast cancer; TRAIL: TNF-related apoptosis-inducing ligand; UCB: Umbilical cord blood.

Acknowledgements

The authors regret that it was not possible to include many interesting studies in the field due to limited space.

Authors' contributions

All authors wrote and revised the initial draft, finalized the last version of the article, and approved the final version.

Funding

This work was supported by grants from the NIH (AI129582, CA247550, NS106170, CA223400, CA210087, CA068458, and CA163205), the Leukemia & Lymphoma Society (6503-17 and 1364-19), and the California Institute for Regenerative Medicine (DISC2COVID19-11947).

Availability of data and materials

Not applicable.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

M.A. Caligiuri and J. Yu are the co-founders of CytImmune Therapeutics, Inc. The other authors declare no competing interests.

Author details

¹The Ohio State University Comprehensive Cancer Center, Columbus, OH 43210, USA. ²Department of Hematology and Hematopoietic Cell Transplantation, City of Hope National Medical Center, 1500 E. Duarte Road, KCRB, Bldg. 158, 3rd Floor, Room 3017, Los Angeles, CA 91010, USA. ³Hematologic Malignancies and Stem Cell Transplantation Institute, City of Hope National Medical Center, Los Angeles, CA 91010, USA. ⁴Department of Immuno-Oncology, City of Hope Beckman Research Institute, Los Angeles, CA 91010, USA. ⁵City of Hope Comprehensive Cancer Center and Beckman Research Institute, Los Angeles, CA 91010, USA.

Received: 16 September 2020 Accepted: 12 November 2020

Published online: 07 December 2020

References

- Spits H, Artis D, Colonna M, Dieffenbach A, Di Santo JP, Eberl G, Koyasu S, Locksley RM, McKenzie AN, Mebius RE, et al. Innate lymphoid cells—a proposal for uniform nomenclature. *Nat Rev Immunol*. 2013;13(2):145–9.
- Ishizuka IE, Constantinides MG, Gudjonson H, Bendelac A. The innate lymphoid cell precursor. *Annu Rev Immunol*. 2016;34:299–316.
- Walker JA, Barlow JL, McKenzie AN. Innate lymphoid cells—how did we miss them? *Nat Rev Immunol*. 2013;13(2):75–87.
- Vivier E, Tomasello E, Baratin M, Walzer T, Ugolini S. Functions of natural killer cells. *Nat Immunol*. 2008;9(5):503–10.
- Yu J, Freud AG, Caligiuri MA. Location and cellular stages of natural killer cell development. *Trends Immunol*. 2013;34(12):573–82.
- Freud AG, Mundy-Bosse BL, Yu J, Caligiuri MA. The broad spectrum of human natural killer cell diversity. *Immunity*. 2017;47(5):820–33.
- Farag SS, Caligiuri MA. Human natural killer cell development and biology. *Blood Rev*. 2006;20(3):123–37.
- Hesslein DG, Lanier LL. Transcriptional control of natural killer cell development and function. *Adv Immunol*. 2011;109:45–85.
- Di Santo JP. Natural killer cell developmental pathways: a question of balance. *Annu Rev Immunol*. 2006;24:257–86.
- Mehta RS, Rezvani K. Chimeric antigen receptor expressing natural killer cells for the immunotherapy of cancer. *Front Immunol*. 2018;9:283.
- Lanier LL. NK cell recognition. *Annu Rev Immunol*. 2005;23:225–74.
- Moretta A, Marcenaro E, Parolini S, Ferlazzo G, Moretta L. NK cells at the interface between innate and adaptive immunity. *Cell Death Differ*. 2008;15(2):226–33.
- Morvan MG, Lanier LL. NK cells and cancer: you can teach innate cells new tricks. *Nat Rev Cancer*. 2016;16(1):7–19.
- Klingemann H, Boissel L, Toneguzzo F. Natural killer cells for immunotherapy—advantages of the NK-92 cell line over blood NK cells. *Front Immunol*. 2016;7:91.
- Waldhauer I, Steinle A. NK cells and cancer immunosurveillance. *Oncogene*. 2008;27(45):5932–43.
- Ljunggren HG, Malmberg KJ. Prospects for the use of NK cells in immunotherapy of human cancer. *Nat Rev Immunol*. 2007;7(5):329–39.
- Hervieu A, Rébé C, Végran F, Chalmin F, Bruchard M, Vabres P, Apetoh L, Ghiringhelli F, Mignot G. Dacarbazine-mediated upregulation of NKG2D ligands on tumor cells activates NK and CD8 T cells and restrains melanoma growth. *J Invest Dermatol*. 2013;133(2):499–508.
- Childs RW, Carlsten M. Therapeutic approaches to enhance natural killer cell cytotoxicity against cancer: the force awakens. *Nat Rev Drug Discov*. 2015;14(7):487–98.
- Mailliard RB, Son YI, Redlinger R, Coates PT, Giermasz A, Morel PA, Storkus WJ, Kalinski P. Dendritic cells mediate NK cell help for Th1 and CTL responses: two-signal requirement for the induction of NK cell helper function. *J Immunol*. 2003;171(5):2366–73.
- Delneste Y, Charbonnier P, Herbault N, Magistrelli G, Caron G, Bonnefoy JY, Jeannin P. Interferon-gamma switches monocyte differentiation from dendritic cells to macrophages. *Blood*. 2003;101(1):143–50.

21. Cheng M, Chen Y, Xiao W, Sun R, Tian Z. NK cell-based immunotherapy for malignant diseases. *Cell Mol Immunol*. 2013;10(3):230–52.
22. Han J, Alvarez-Breckenridge CA, Wang QE, Yu J. TGF- β signaling and its targeting for glioma treatment. *Am J Cancer Res*. 2015;5(3):945–55.
23. Salih HR, Rammensee HG, Steinle A. Cutting edge: down-regulation of MICA on human tumors by proteolytic shedding. *J Immunol*. 2002;169(8):4098–102.
24. Lichtenegger FS, Krupka C, Haubner S, Kohnke T, Subklewe M. Recent developments in immunotherapy of acute myeloid leukemia. *J Hematol Oncol*. 2017;10(1):142.
25. Hegde PS, Chen DS. Top 10 challenges in cancer immunotherapy. *Immunity*. 2020;52(1):17–35.
26. Grupp SA, Kalos M, Barrett D, Aplenc R, Porter DL, Rheingold SR, Teachey DT, Chew A, Hauck B, Wright JF, et al. Chimeric antigen receptor-modified T cells for acute lymphoid leukemia. *N Engl J Med*. 2013;368(16):1509–18.
27. Sadelain M, Riviere I, Brentjens R. Targeting tumours with genetically enhanced T lymphocytes. *Nat Rev Cancer*. 2003;3(1):35–45.
28. Brown CE, Alizadeh D, Starr R, Weng L, Wagner JR, Naranjo A, Ostberg JR, Blanchard MS, Kilpatrick J, Simpson J, et al. Regression of glioblastoma after chimeric antigen receptor T-cell therapy. *N Engl J Med*. 2016;375(26):2561–9.
29. Fernandez L, Metais JY, Escudero A, Vela M, Valentin J, Vallcorba I, Leivas A, Torres J, Valeri A, Patino-Garcia A, et al. Memory T cells expressing an NKG2D-CAR efficiently target osteosarcoma cells. *Clin Cancer Res*. 2017;23(19):5824–35.
30. Irving BA, Weiss A. The cytoplasmic domain of the T cell receptor zeta chain is sufficient to couple to receptor-associated signal transduction pathways. *Cell*. 1991;64(5):891–901.
31. Sadelain M, Brentjens R, Riviere I. The basic principles of chimeric antigen receptor design. *Cancer Discov*. 2013;3(4):388–98.
32. Dotti G, Gottschalk S, Savoldo B, Brenner MK. Design and development of therapies using chimeric antigen receptor-expressing T cells. *Immunol Rev*. 2014;257(1):107–26.
33. Jackson HJ, Rafiq S, Brentjens RJ. Driving CAR T-cells forward. *Nat Rev Clin Oncol*. 2016;13(6):370–83.
34. Gill S, Maus MV, Porter DL. Chimeric antigen receptor T cell therapy: 25 years in the making. *Blood Rev*. 2016;30(3):157–67.
35. Ramos CA, Heslop HE, Brenner MK. CAR-T cell therapy for lymphoma. *Annu Rev Med*. 2016;67:165–83.
36. Jain MD, Davila ML. Concise review: emerging principles from the clinical application of chimeric antigen receptor T cell therapies for B cell malignancies. *Stem Cells (Dayton, OH)*. 2018;36(1):36–44.
37. Watanabe K, Terakura S, Martens AC, van Meerden T, Uchiyama S, Imai M, Sakemura R, Goto T, Hanajiri R, Imahashi N, et al. Target antigen density governs the efficacy of anti-CD20-CD28-CD3 zeta chimeric antigen receptor-modified effector CD8+ T cells. *J Immunol*. 2015;194(3):911–20.
38. Zhang C, Liu J, Zhong JF, Zhang X. Engineering CAR-T cells. *Biomark Res*. 2017;5:22.
39. Hambach J, Riecken K, Cichutek S, Schütze K, Albrecht B, Petry K, Röckendorf JL, Baum N, Kröger N, Hansen T, et al. Targeting CD38-expressing multiple myeloma and burkitt lymphoma cells in vitro with nanobody-based chimeric antigen receptors (Nb-CARs). *Cells*. 2020;9(2):321.
40. Hoseini SS, Cheung NK. Acute myeloid leukemia targets for bispecific antibodies. *Blood Cancer J*. 2017;7(4):e552.
41. Stoiber S, Cadilha BL, Benmebarek MR, Lesch S, Endres S, Kobold S. Limitations in the design of chimeric antigen receptors for cancer therapy. *Cells*. 2019;8(5):472.
42. van der Stegen SJ, Hamieh M, Sadelain M. The pharmacology of second-generation chimeric antigen receptors. *Nat Rev Drug Discov*. 2015;14(7):499–509.
43. Fernandez L, Fernandez A, Mirones I, Escudero A, Cardoso L, Vela M, Lanzarot D, de Paz R, Leivas A, Gallardo M, et al. GMP-compliant manufacturing of NKG2D CAR memory T cells using CliniMACS prodigy. *Front Immunol*. 2019;10:2361.
44. Chu J, Deng Y, Benson DM, He S, Hughes T, Zhang J, Peng Y, Mao H, Yi L, Ghoshal K, et al. CS1-specific chimeric antigen receptor (CAR)-engineered natural killer cells enhance in vitro and in vivo antitumor activity against human multiple myeloma. *Leukemia*. 2014;28(4):917–27.
45. Guedan S, Calderon H, Posey AD Jr, Maus MV. Engineering and design of chimeric antigen receptors. *Mol Therapy Methods Clin Dev*. 2019;12:145–56.
46. Porter DL, Hwang WT, Frey NV, Lacey SF, Shaw PA, Loren AW, Bagg A, Marcucci KT, Shen A, Gonzalez V, et al. Chimeric antigen receptor T cells persist and induce sustained remissions in relapsed refractory chronic lymphocytic leukemia. *Sci Transl Med*. 2015;7(303):303ra139.
47. Lee DW, Kochenderfer JN, Stetler-Stevenson M, Cui YK, Delbrook C, Feldman SA, Fry TJ, Orentas R, Sabatino M, Shah NN, et al. T cells expressing CD19 chimeric antigen receptors for acute lymphoblastic leukaemia in children and young adults: a phase 1 dose-escalation trial. *Lancet*. 2015;385(9967):517–28.
48. Maude SL, Frey N, Shaw PA, Aplenc R, Barrett DM, Bunin NJ, Chew A, Gonzalez VE, Zheng Z, Lacey SF, et al. Chimeric antigen receptor T cells for sustained remissions in leukemia. *N Engl J Med*. 2014;371(16):1507–17.
49. O'Leary M. Biologic License Application Clinical Review Memorandum. Biologics license application 2017:12.
50. Gorovits B, Koren E. Immunogenicity of chimeric antigen receptor T-cell therapeutics. *BioDrugs*. 2019;33(3):275–84.
51. Kurlander R. Approval of CD19-directed CAR T cells: FDA licensure of new cell-based gene therapy products with a challenging safety profile. FDA clinical review memorandum 2017:9.
52. Lamers CH, Sleijfer S, van Steenbergen S, van Elzakker P, van Krimpen B, Groot C, Vulto A, den Bakker M, Oosterwijk E, Debets R, et al. Treatment of metastatic renal cell carcinoma with CAIX CAR-engineered T cells: clinical evaluation and management of on-target toxicity. *Mol Therapy*. 2013;21(4):904–12.
53. Kershaw MH, Westwood JA, Parker LL, Wang G, Eshhar Z, Mavroukakis SA, White DE, Wunderlich JR, Canevari S, Rogers-Freezer L, et al. A phase I study on adoptive immunotherapy using gene-modified T cells for ovarian cancer. *Clin Cancer Res*. 2006;12(20 Pt 1):6106–15.
54. Craddock JA, Lu A, Bear A, Pule M, Brenner MK, Rooney CM, Foster AE. Enhanced tumor trafficking of GD2 chimeric antigen receptor T cells by expression of the chemokine receptor CCR2b. *J Immunother*. 2010;33(8):780–8.
55. Stern LA, Jonsson VD, Priceman SJ. CAR T cell therapy progress and challenges for solid tumors. *Cancer Treat Res*. 2020;180:297–326.
56. Lee DW, Gardner R, Porter DL, Louis CU, Ahmed N, Jensen M, Grupp SA, Mackall CL. Current concepts in the diagnosis and management of cytokine release syndrome. *Blood*. 2014;124(2):188–95.
57. Bonifant CL, Jackson HJ, Brentjens RJ, Curran KJ. Toxicity and management in CAR T-cell therapy. *Mol Therapy Oncolyt*. 2016;3:16011.
58. Davila ML, Riviere I, Wang X, Bartido S, Park J, Curran K, Chung SS, Stefanski J, Borquez-Ojeda O, Olszewska M, et al. Efficacy and toxicity management of 19–28z CAR T cell therapy in B cell acute lymphoblastic leukemia. *Sci Transl Med*. 2014;6(224):224ra225.
59. Morgan RA, Yang JC, Kitano M, Dudley ME, Laurencot CM, Rosenberg SA. Case report of a serious adverse event following the administration of T cells transduced with a chimeric antigen receptor recognizing ERBB2. *Mol Therapy*. 2010;18(4):843–51.
60. Hirayama AV, Turtle CJ. Toxicities of CD19 CAR-T cell immunotherapy. *Am J Hematol*. 2019;94(S1):S42-s49.
61. Hu Y, Tian ZG, Zhang C. Chimeric antigen receptor (CAR)-transduced natural killer cells in tumor immunotherapy. *Acta Pharmacol Sin*. 2018;39(2):167–76.
62. Miller JS, Soignier Y, Panoskaltsis-Mortari A, McNearney SA, Yun GH, Fautsch SK, McKenna D, Le C, Defor TE, Burns LJ, et al. Successful adoptive transfer and in vivo expansion of human haploidentical NK cells in patients with cancer. *Blood*. 2005;105(8):3051–7.
63. Olson JA, Leveson-Gower DB, Gill S, Baker J, Beilhack A, Negrin RS. NK cells mediate reduction of GVHD by inhibiting activated, alloreactive T cells while retaining GVT effects. *Blood*. 2010;115(21):4293–301.
64. Graham C, Jozwik A, Pepper A, Benjamin R. Alloeneic CAR-T cells: more than ease of access? *Cells*. 2018;7(10):155.
65. Klingemann H. Are natural killer cells superior CAR drivers? *Oncoimmunology*. 2014;3:e28147.
66. Glienke W, Esser R, Priesner C, Suerth JD, Schambach A, Wels WS, Grez M, Kloess S, Arseniev L, Koehl U. Advantages and applications of CAR-expressing natural killer cells. *Front Pharmacol*. 2015;6:21.

67. Tang X, Yang L, Li Z, Nalin AP, Dai H, Xu T, Yin J, You F, Zhu M, Shen W, et al. First-in-man clinical trial of CAR NK-92 cells: safety test of CD33-CAR NK-92 cells in patients with relapsed and refractory acute myeloid leukemia. *Am J Cancer Res*. 2018;8(6):1083–9.
68. Liu E, Marin D, Banerjee P, Macapinlac HA, Thompson P, Basar R, Nassif Kerbauy L, Overman B, Thall P, Kaplan M, et al. Use of CAR-transduced natural killer cells in CD19-positive lymphoid tumors. *N Engl J Med*. 2020;382(6):545–53.
69. Cheng M, Zhang J, Jiang W, Chen Y, Tian Z. Natural killer cell lines in tumor immunotherapy. *Front Med*. 2012;6(1):56–66.
70. Tonn T, Schwabe D, Klingemann HG, Becker S, Esser R, Koehl U, Suttorp M, Seifried E, Ottmann OG, Bug G. Treatment of patients with advanced cancer with the natural killer cell line NK-92. *Cytotherapy*. 2013;15(12):1563–70.
71. Klingemann H. Challenges of cancer therapy with natural killer cells. *Cytotherapy*. 2015;17(3):245–9.
72. MacLeod RA, Nagel S, Kaufmann M, Greulich-Bode K, Drexler HG. Multicolor-FISH analysis of a natural killer cell line (NK-92). *Leuk Res*. 2002;26(11):1027–33.
73. Hermanson DL, Kaufman DS. Utilizing chimeric antigen receptors to direct natural killer cell activity. *Front Immunol*. 2015;6:195.
74. Messaoudene M, Frazao A, Gavlovsky PJ, Toubert A, Dulphy N, Caignard A. Patient's natural killer cells in the era of targeted therapies: role for tumor killers. *Front Immunol*. 2017;8:683.
75. Nayyar G, Chu Y, Cairo MS. Overcoming resistance to natural killer cell based immunotherapies for solid tumors. *Front Oncol*. 2019;9:51.
76. Luevano M, Daryouzeh M, Alnabhan R, Querol S, Khakoo S, Madrigal A, Saudemont A. The unique profile of cord blood natural killer cells balances incomplete maturation and effective killing function upon activation. *Hum Immunol*. 2012;73(3):248–57.
77. Kotylo PK, Baenziger JC, Yoder MC, Engle WA, Bolinger CD. Rapid analysis of lymphocyte subsets in cord blood. *Am J Clin Pathol*. 1990;93(2):263–6.
78. Rocha V, Labopin M, Sanz G, Arcese W, Schwerdtfeger R, Bosi A, Jacobsen N, Ruutu T, de Lima M, Finke J, et al. Transplants of umbilical-cord blood or bone marrow from unrelated donors in adults with acute leukemia. *N Engl J Med*. 2004;351(22):2276–85.
79. Eapen M, Rubinstein P, Zhang MJ, Stevens C, Kurtzberg J, Scaradavou A, Loberiza FR, Champlin RE, Klein JP, Horowitz MM, et al. Outcomes of transplantation of unrelated donor umbilical cord blood and bone marrow in children with acute leukaemia: a comparison study. *Lancet*. 2007;369(9577):1947–54.
80. Atsuta Y, Suzuki R, Nagamura-Inoue T, Taniguchi S, Takahashi S, Kai S, Sakamaki H, Kouzai Y, Kasai M, Fukuda T, et al. Disease-specific analyses of unrelated cord blood transplantation compared with unrelated bone marrow transplantation in adult patients with acute leukemia. *Blood*. 2009;113(8):1631–8.
81. Harris DT, Schumacher MJ, Locascio J, Besencon FJ, Olson GB, DeLuca D, Shenker L, Bard J, Boyse EA. Phenotypic and functional immaturity of human umbilical cord blood T lymphocytes. *Proc Natl Acad Sci USA*. 1992;89(21):10006–10.
82. Xing D, Ramsay AG, Gribben JG, Decker WK, Burks JK, Munsell M, Li S, Robinson SN, Yang H, Steiner D, et al. Cord blood natural killer cells exhibit impaired lytic immunological synapse formation that is reversed with IL-2 ex vivo expansion. *J Immunother*. 2010;33(7):684–96.
83. Liu E, Tong Y, Dotti G, Shaim H, Savoldo B, Mukherjee M, Orange J, Wan X, Lu X, Reynolds A, et al. Cord blood NK cells engineered to express IL-15 and a CD19-targeted CAR show long-term persistence and potent antitumor activity. *Leukemia*. 2018;32(2):520–31.
84. Woll PS, Grzywacz B, Tian X, Marcus RK, Knorr DA, Verneris MR, Kaufman DS. Human embryonic stem cells differentiate into a homogeneous population of natural killer cells with potent in vivo antitumor activity. *Blood*. 2009;113(24):6094–101.
85. Eguizabal C, Zenarruzabeitia O, Monge J, Santos S, Vesga MA, Maruri N, Arieta A, Rinon M, Tamayo-Orbegozo E, Amo L, et al. Natural killer cells for cancer immunotherapy: pluripotent stem cells-derived NK cells as an immunotherapeutic perspective. *Front Immunol*. 2014;5:439.
86. Knorr DA, Ni Z, Hermanson D, Hexum MK, Bendzick L, Cooper LJ, Lee DA, Kaufman DS. Clinical-scale derivation of natural killer cells from human pluripotent stem cells for cancer therapy. *Stem Cells Transl Med*. 2013;2(4):274–83.
87. Cany J, van der Waart AB, Tordoir M, Franssen GM, Hangalapura BN, de Vries J, Boerman O, Schaap N, van der Voort R, Spanholtz J, et al. Natural killer cells generated from cord blood hematopoietic progenitor cells efficiently target bone marrow-residing human leukemia cells in NOD/SCID/IL2Rg(null) mice. *PLoS ONE*. 2013;8(6):e64384.
88. Spanholtz J, Preijers F, Tordoir M, Trilsbeek C, Paardekooper J, de Witte T, Schaap N, Dolstra H. Clinical-grade generation of active NK cells from cord blood hematopoietic progenitor cells for immunotherapy using a closed-system culture process. *PLoS ONE*. 2011;6(6):e20740.
89. Li Y, Hermanson DL, Moriarity BS, Kaufman DS. Human iPSC-derived natural killer cells engineered with chimeric antigen receptors enhance anti-tumor activity. *Cell Stem Cell*. 2018;23(2):181–192.e185.
90. Shimasaki N, Fujisaki H, Cho D, Masselli M, Lockey T, Eldridge P, Leung W, Campana D. A clinically adaptable method to enhance the cytotoxicity of natural killer cells against B-cell malignancies. *Cytotherapy*. 2012;14(7):830–40.
91. Boissel L, Betancur M, Wels WS, Tuncer H, Klingemann H. Transfection with mRNA for CD19 specific chimeric antigen receptor restores NK cell mediated killing of CLL cells. *Leuk Res*. 2009;33(9):1255–9.
92. Li L, Liu LN, Feller S, Allen C, Shivakumar R, Fratantoni J, Wolfraim LA, Fujisaki H, Campana D, Chopas N, et al. Expression of chimeric antigen receptors in natural killer cells with a regulatory-compliant non-viral method. *Cancer Gene Ther*. 2010;17(3):147–54.
93. Imai C, Mihara K, Andreansky M, Nicholson IC, Pui CH, Geiger TL, Campana D. Chimeric receptors with 4-1BB signaling capacity provoke potent cytotoxicity against acute lymphoblastic leukemia. *Leukemia*. 2004;18(4):676–84.
94. Imai C, Iwamoto S, Campana D. Genetic modification of primary natural killer cells overcomes inhibitory signals and induces specific killing of leukemic cells. *Blood*. 2005;106(1):376–83.
95. Y C, A Y, J A, C V, M B, M C, M S C. Anti-CD20 chimeric antigen receptor (CAR) modified expanded natural killer (NK) cells significantly mediate rituximab sensitive and resistant burkitt lymphoma (BL) regression and improve survival in human BL xenografted NSG mice. *Cytotherapy* 2014, 16(4):S21.
96. Boissel L, Betancur-Boissel M, Lu W, Krause DS, Van Etten RA, Wels WS, Klingemann H. Retargeting NK-92 cells by means of CD19- and CD20-specific chimeric antigen receptors compares favorably with antibody-dependent cellular cytotoxicity. *Oncoimmunology*. 2013;2(10):e26527.
97. Muller T, Uherek C, Maki G, Chow KU, Schimpf A, Klingemann HG, Tonn T, Wels WS. Expression of a CD20-specific chimeric antigen receptor enhances cytotoxic activity of NK cells and overcomes NK-resistance of lymphoma and leukemia cells. *Cancer Immunol Immunother CII*. 2008;57(3):411–23.
98. Jiang H, Zhang W, Shang P, Zhang H, Fu W, Ye F, Zeng T, Huang H, Zhang X, Sun W, et al. Transfection of chimeric anti-CD138 gene enhances natural killer cell activation and killing of multiple myeloma cells. *Mol Oncol*. 2014;8(2):297–310.
99. Hsi ED, Steinle R, Balasa B, Szmania S, Draksharapu A, Shum BP, Huseni M, Powers D, Naniseti A, Zhang Y, et al. CS1, a potential new therapeutic antibody target for the treatment of multiple myeloma. *Clin Cancer Res*. 2008;14(9):2775–84.
100. Zhang Y, Chen L, Wang Y, Li X, Hughes T, Klingemann H, Benson DM, Yu J. Combination therapy with daratumumab and CAR-NK targeting CS1 for multiple myeloma. *Blood Rev*. 2016;128:1342.
101. Clark EA, Giltiay NV. CD22: A regulator of innate and adaptive B cell responses and autoimmunity. *Front Immunol*. 2018;9:2235.
102. De Propriis MS, Raponi S, Diverio D, Milani ML, Meloni G, Falini B, Foà R, Guarini A. High CD33 expression levels in acute myeloid leukemia cells carrying the nucleophosmin (NPM1) mutation. *Haematologica*. 2011;96(10):1548–51.
103. Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, Levin WJ, Stuart SG, Udove J, Ullrich A, et al. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science (NY)*. 1989;244(4905):707–12.
104. Zhang C, Burger MC, Jennewein L, Genssler S, Schonfeld K, Zeiner P, Hattingen E, Harter PN, Mittelbronn M, Tonn T, et al. ErbB2/HER2-specific NK cells for targeted therapy of glioblastoma. *J Natl Cancer Inst*. 2016. <https://doi.org/10.1093/jnci/djv375>.
105. Alsamh W, Romia Y. Modification of natural killer cells to target tumors. *Int J Pharm Clin Res*. 2014;6(1):97–100.

106. Uherek C, Tonn T, Uherek B, Becker S, Schnierle B, Klingemann HG, Wels W. Retargeting of natural killer-cell cytolytic activity to ErbB2-expressing cancer cells results in efficient and selective tumor cell destruction. *Blood*. 2002;100(4):1265–73.
107. Schonfeld K, Sahm C, Zhang C, Naundorf S, Brendel C, Odendahl M, Nowakowska P, Bonig H, Kohl U, Kloess S, et al. Selective inhibition of tumor growth by clonal NK cells expressing an ErbB2/HER2-specific chimeric antigen receptor. *Mol Ther*. 2015;23(2):330–8.
108. Kruschinski A, Moosmann A, Poschke I, Norell H, Chmielewski M, Seliger B, Kiessling R, Blankenstein T, Abken H, Charo J. Engineering antigen-specific primary human NK cells against HER-2 positive carcinomas. *Proc Natl Acad Sci USA*. 2008;105(45):17481–6.
109. Esser R, Muller T, Stefes D, Kloess S, Seidel D, Gillies SD, Aperlo-Iffland C, Huston JS, Uherek C, Schonfeld K, et al. NK cells engineered to express a GD2-specific antigen receptor display built-in ADCC-like activity against tumour cells of neuroectodermal origin. *J Cell Mol Med*. 2012;16(3):569–81.
110. Seidel D, Shibina A, Siebert N, Wels WS, Reynolds CP, Huebener N, Lode HN. Disialoganglioside-specific human natural killer cells are effective against drug-resistant neuroblastoma. *Cancer Immunol Immunother*. 2015;64(5):621–34.
111. Zhang G, Liu R, Zhu X, Wang L, Ma J, Han H, Wang X, Zhang G, He W, Wang W, et al. Retargeting NK-92 for anti-melanoma activity by a TCR-like single-domain antibody. *Immunol Cell Biol*. 2013;91(10):615–24.
112. Han J, Chu J, Keung Chan W, Zhang J, Wang Y, Cohen JB, Victor A, Meisen WH, Kim SH, Grandi P, et al. CAR-engineered NK cells targeting wild-type EGFR and EGFRvIII enhance killing of glioblastoma and patient-derived glioblastoma stem cells. *Sci Rep*. 2015;5:11483.
113. Altvater B, Landmeier S, Pscherer S, Temme J, Schweer K, Kailayangiri S, Campana D, Juergens H, Pule M, Rossig C. 2B4 (CD244) signaling by recombinant antigen-specific chimeric receptors costimulates natural killer cell activation to leukemia and neuroblastoma cells. *Clin Cancer Res*. 2009;15(15):4857–66.
114. Sahm C, Schonfeld K, Wels WS. Expression of IL-15 in NK cells results in rapid enrichment and selective cytotoxicity of gene-modified effectors that carry a tumor-specific antigen receptor. *Cancer Immunol Immunother*. 2012;61(9):1451–61.
115. Parsons DW, Jones S, Zhang X, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Siu LM, Gallia GL, et al. An integrated genomic analysis of human glioblastoma multiforme. *Science (N Y)*. 2008;321(5897):1807–12.
116. Chen X, Han J, Chu J, Zhang L, Zhang J, Chen C, Chen L, Wang Y, Wang H, Yi L, et al. A combinational therapy of EGFR-CAR NK cells and oncolytic herpes simplex virus 1 for breast cancer brain metastases. *Oncotarget*. 2016;7(19):27764–77.
117. Cao B, Liu M, Wang L, Liang B, Feng Y, Chen X, Shi Y, Zhang J, Ye X, Tian Y, et al. Use of chimeric antigen receptor NK-92 cells to target mesothelin in ovarian cancer. *Biochem Biophys Res Commun*. 2020;524(1):96–102.
118. Montagner IM, Penna A, Fracasso G, Carpanese D, Dalla Pietra A, Barbieri V, Zuccolotto G, Rosato A. Anti-PSMA CAR-engineered NK-92 cells: an off-the-shelf cell therapy for prostate cancer. *Cells*. 2020;9(6):1382.
119. Bollino D, Webb TJ. Chimeric antigen receptor-engineered natural killer and natural killer T cells for cancer immunotherapy. *Transl Res*. 2017;187:32–43.
120. Xia N, Haopeng P, Gong JU, Lu J, Chen Z, Zheng Y, Wang Z, Sun YU, Yang Z, Hoffman RM, et al. Robo1-specific CAR-NK immunotherapy enhances efficacy of (125I) seed brachytherapy in an orthotopic mouse model of human pancreatic carcinoma. *Anticancer Res*. 2019;39(11):5919–25.
121. Savan R, Chan T, Young HA. Lentiviral gene transduction in human and mouse NK cell lines. *Methods Mol Biol (Clifton, NJ)*. 2010;612:209–21.
122. Danthinne X, Imperiale MJ. Production of first generation adenovirus vectors: a review. *Gene Ther*. 2000;7(20):1707–14.
123. Hacein-Bey-Abina S, Garrigue A, Wang GP, Soulier J, Lim A, Morillon E, Clappier E, Caccavelli L, Delabesse E, Beldjord K, et al. Insertional oncogenesis in 4 patients after retrovirus-mediated gene therapy of SCID-X1. *J Clin Invest*. 2008;118(9):3132–42.
124. Ranzani M, Annunziato S, Adams DJ, Montini E. Cancer gene discovery: exploiting insertional mutagenesis. *Mol Cancer Res MCR*. 2013;11(10):1141–58.
125. Felice B, Cattoglio C, Cittaro D, Testa A, Miccio A, Ferrari G, Luzi L, Recchia A, Mavilio F. Transcription factor binding sites are genetic determinants of retroviral integration in the human genome. *PLoS ONE*. 2009;4(2):e4571.
126. Imamura M, Shook D, Kamiya T, Shimasaki N, Chai SM, Coustan-Smith E, Imai C, Campana D. Autonomous growth and increased cytotoxicity of natural killer cells expressing membrane-bound interleukin-15. *Blood*. 2014;124(7):1081–8.
127. Cho FN, Chang TH, Shu CW, Ko MC, Liao SK, Wu KH, Yu MS, Lin SJ, Hong YC, Chen CH, et al. Enhanced cytotoxicity of natural killer cells following the acquisition of chimeric antigen receptors through trogocytosis. *PLoS ONE*. 2014;9(10):e109352.
128. Nanbakhsh A, Best B, Riese M, Rao S, Wang L, Medin J, Thakar MS, Malarkannan S. Dextran enhances the lentiviral transduction efficiency of murine and human primary NK cells. *J Vis Exp JoVE*. 2018. <https://doi.org/10.3791/55063>.
129. Castellani S, Orlando C, Carbone A, Di Gioia S, Conese M. Magnetofection enhances lentiviral-mediated transduction of airway epithelial cells through extracellular and cellular barriers. *Genes*. 2016;7(11):103.
130. Chang L, Gallego-Perez D, Zhao X, Bertani P, Yang Z, Chiang CL, Malkoc V, Shi J, Sen CK, Odonnell L, et al. Dielectrophoresis-assisted 3D nanoelectroporation for non-viral cell transfection in adoptive immunotherapy. *Lab Chip*. 2015;15(15):3147–53.
131. Becker PS, Suck G, Nowakowska P, Ullrich E, Seifried E, Bader P, Tonn T, Seidl C. Selection and expansion of natural killer cells for NK cell-based immunotherapy. *Cancer Immunol Immunother*. 2016;65(4):477–84.
132. Fujisaki H, Kakuda H, Shimasaki N, Imai C, Ma J, Lockey T, Eldridge P, Leung WH, Campana D. Expansion of highly cytotoxic human natural killer cells for cancer cell therapy. *Can Res*. 2009;69(9):4010–7.
133. Lim SA, Kim TJ, Lee JE, Sonn CH, Kim K, Kim J, Choi JG, Choi IK, Yun CO, Kim JH, et al. Ex vivo expansion of highly cytotoxic human NK cells by cocultivation with irradiated tumor cells for adoptive immunotherapy. *Can Res*. 2013;73(8):2598–607.
134. Sutlu T, Stellan B, Gilljam M, Quezada HC, Nahi H, Gahrton G, Alici E. Clinical-grade, large-scale, feeder-free expansion of highly active human natural killer cells for adoptive immunotherapy using an automated bioreactor. *Cytotherapy*. 2010;12(8):1044–55.
135. Denman CJ, Senyukov VV, Somanchi SS, Phatarpekar PV, Kopp LM, Johnson JL, Singh H, Hurton L, Maiti SN, Huls MH, et al. Membrane-bound IL-21 promotes sustained ex vivo proliferation of human natural killer cells. *PLoS ONE*. 2012;7(1):e30264.
136. Shah N, Martin-Antonio B, Yang H, Ku S, Lee DA, Cooper LJ, Decker WK, Li S, Robinson SN, Sekine T, et al. Antigen presenting cell-mediated expansion of human umbilical cord blood yields log-scale expansion of natural killer cells with anti-myeloma activity. *PLoS ONE*. 2013;8(10):e76781.
137. Kloss S, Oberschmidt O, Morgan M, Dahlke J, Arseniev L, Huppert V, Granzin M, Gardlowski T, Matthies N, Soltenborn S, et al. Optimization of human NK cell manufacturing: fully automated separation, improved ex vivo expansion using IL-21 with autologous feeder cells, and generation of anti-CD123-CAR-expressing effector cells. *Hum Gene Ther*. 2017;28(10):897–913.
138. Martin-Antonio B, Sune G, Perez-Amill L, Castella M, Urbano-Ispizua A. Natural killer cells: angels and devils for immunotherapy. *Int J Mol Sci*. 2017;18(9):1868.
139. Pinthus JH, Waks T, Kaufman-Francis K, Schindler DG, Harmelin A, Kanety H, Ramon J, Eshhar Z. Immuno-gene therapy of established prostate tumors using chimeric receptor-redirected human lymphocytes. *Can Res*. 2003;63(10):2470–6.
140. Topfer K, Cartellieri M, Michen S, Wiedemuth R, Muller N, Lindemann D, Bachmann M, Fussel M, Schackert G, Temme A. DAP12-based activating chimeric antigen receptor for NK cell tumor immunotherapy. *J Immunol*. 2015;194(7):3201–12.
141. Meier R, Golovko D, Tavri S, Henning TD, Knopp C, Piontek G, Rudelius M, Heinrich P, Wels WS, Daldrop-Link H. Depicting adoptive immunotherapy for prostate cancer in an animal model with magnetic resonance imaging. *Magn Reson Med*. 2011;65(3):756–63.
142. Liu Y, Zhou Y, Huang KH, Fang X, Li Y, Wang F, An L, Chen Q, Zhang Y, Shi A, et al. Targeting epidermal growth factor-overexpressing triple-negative breast cancer by natural killer cells expressing a specific chimeric antigen receptor. *Cell Prolif*. 2020;53(8):e12858.
143. Hu Z. Tissue factor as a new target for CAR-NK cell immunotherapy of triple-negative breast cancer. *Sci Rep*. 2020;10(1):2815.

144. Wei J, Han X, Bo J, Han W. Target selection for CAR-T therapy. *J Hematol Oncol*. 2019;12(1):62.
145. Guo Y, Wang Y, Han W. Chimeric antigen receptor-modified T cells for solid tumors: challenges and prospects. *J Immunol Res*. 2016;2016:3850839.
146. Adusumilli PS, Cherkassky L, Villena-Vargas J, Colovos C, Servais E, Plotkin J, Jones DR, Sadelain M. Regional delivery of mesothelin-targeted CAR T cell therapy generates potent and long-lasting CD4-dependent tumor immunity. *Sci Transl Med*. 2014;6(261):261151.
147. Santoro SP, Kim S, Motz GT, Alatzoglou D, Li C, Irving M, Powell DJ Jr, Coukos G. T cells bearing a chimeric antigen receptor against prostate-specific membrane antigen mediate vascular disruption and result in tumor regression. *Cancer Immunol Res*. 2015;3(1):68–84.
148. Gordon IO, Tretiakova MS, Noffsinger AE, Hart J, Reuter VE, Al-Ahmadie HA. Prostate-specific membrane antigen expression in regeneration and repair. *Mod Pathol*. 2008;21(12):1421–7.
149. Alkins R, Burgess A, Kerbel R, Wels WS, Hynynen K. Early treatment of HER2-amplified brain tumors with targeted NK-92 cells and focused ultrasound improves survival. *Neuro-oncology*. 2016;18(7):974–81.
150. Carrega P, Morandi B, Costa R, Frumento G, Forte G, Altavilla G, Ratto GB, Mingari MC, Moretta L, Ferlazzo G. Natural killer cells infiltrating human nonsmall-cell lung cancer are enriched in CD56 bright CD16(–) cells and display an impaired capability to kill tumor cells. *Cancer*. 2008;112(4):863–75.
151. Wennerberg E, Kremer V, Childs R, Lundqvist A. CXCL10-induced migration of adoptively transferred human natural killer cells toward solid tumors causes regression of tumor growth in vivo. *Cancer Immunol Immunother* CII. 2015;64(2):225–35.
152. Kremer V, Ligtenberg MA, Zendejdel R, Seitz C, Duivenvoorden A, Wennerberg E, Colón E, Scherman-Plogell AH, Lundqvist A. Genetic engineering of human NK cells to express CXCR2 improves migration to renal cell carcinoma. *J Immunother Cancer*. 2017;5(1):73.
153. Muller N, Michen S, Tietze S, Topfer K, Schulte A, Lamszus K, Schmitz M, Schackert G, Pastan I, Temme A. Engineering NK cells modified with an EGFRvIII-specific chimeric antigen receptor to overexpress CXCR4 improves immunotherapy of CXCL12/SDF-1alpha-secreting glioblastoma. *J Immunother*. 2015;38(5):197–210.
154. Hou W, Chen H, Rojas J, Sampath P, Thorne SH. Oncolytic vaccinia virus demonstrates antiangiogenic effects mediated by targeting of VEGF. *Int J Cancer*. 2014;135(5):1238–46.
155. Sampath P, Li J, Hou W, Chen H, Bartlett DL, Thorne SH. Crosstalk between immune cell and oncolytic vaccinia therapy enhances tumor trafficking and antitumor effects. *Mol Ther*. 2013;21(3):620–8.
156. Heyman B, Yang Y. Chimeric antigen receptor T cell therapy for solid tumors: current status, obstacles and future strategies. *Cancers*. 2019;11(2):191.
157. Otegbeye F, Ojo E, Moreton S, Mackowski N, Lee DA, de Lima M, Wald DN. Inhibiting TGF-beta signaling preserves the function of highly activated, in vitro expanded natural killer cells in AML and colon cancer models. *PLoS ONE*. 2018;13(1):e0191358.
158. Zhang C, Hu Y, Shi C. Targeting natural killer cells for tumor immunotherapy. *Front Immunol*. 2020;11:60.
159. Wang QM, Tang PM, Lian GY, Li C, Li J, Huang XR, Lan HY. Enhanced Cancer Immunotherapy with Smad3-Silenced NK-92 Cells. *Cancer Immunol Res*. 2018;6(8):965–77.
160. Zhu S, Denman CJ, Cobanoglu ZS, Kiany S, Lau CC, Gottschalk SM, Hughes DP, Kleinerman ES, Lee DA. The narrow-spectrum HDAC inhibitor entinostat enhances NKG2D expression without NK cell toxicity, leading to enhanced recognition of cancer cells. *Pharm Res*. 2015;32(3):779–92.
161. Balsamo M, Manzini C, Pietra G, Raggi F, Blengio F, Mingari MC, Varesio L, Moretta L, Bosco MC, Vitale M. Hypoxia downregulates the expression of activating receptors involved in NK-cell-mediated target cell killing without affecting ADCC. *Eur J Immunol*. 2013;43(10):2756–64.
162. Wang J, Lupo KB, Chambers AM, Matosevic S. Purinergic targeting enhances immunotherapy of CD73(+) solid tumors with piggyBac-engineered chimeric antigen receptor natural killer cells. *J Immunother Cancer*. 2018;6(1):136.
163. Chan CJ, Martinet L, Gilfillan S, Souza-Fonseca-Guimaraes F, Chow MT, Town L, Ritchie DS, Colonna M, Andrews DM, Smyth MJ. The receptors CD96 and CD226 oppose each other in the regulation of natural killer cell functions. *Nat Immunol*. 2014;15(5):431–8.
164. Dong W, Wu X, Ma S, Wang Y, Nalin AP, Zhu Z, Zhang J, Benson DM, He K, Caligiuri MA, et al. The mechanism of anti-PD-L1 Antibody efficacy against PD-L1-negative tumors identifies NK cells expressing PD-L1 as a cytolytic effector. *Cancer Discov*. 2019;9(10):1422–37.
165. Grosser R, Cherkassky L, Chintala N, Adusumilli PS. Combination immunotherapy with CAR T cells and checkpoint blockade for the treatment of solid tumors. *Cancer Cell*. 2019;36(5):471–82.
166. Fabian KP, Padgett MR, Donahue RN, Solocinski K, Robbins Y, Allen CT, Lee JH, Rabizadeh S, Soon-Shiong P, Schlom J, et al. PD-L1 targeting high-affinity NK (t-haNK) cells induce direct antitumor effects and target suppressive MDSC populations. *J Immunother Cancer*. 2020. <https://doi.org/10.1136/jitc-2019-000450>.
167. Walseng E, Koksai H, Sektioglu IM, Fane A, Skorstad G, Kvalheim G, Gaudernack G, Inderberg EM, Walchli S. A TCR-based chimeric antigen receptor. *Sci Rep*. 2017;7(1):10713.
168. Ligtenberg MA, Mougiakakos D, Mukhopadhyay M, Witt K, Lladser A, Chmielewski M, Riet T, Abken H, Kiessling R. Coexpressed catalase protects chimeric antigen receptor-directed T cells as well as bystander cells from oxidative stress-induced loss of antitumor activity. *J Immunol*. 2016;196(2):759–66.
169. Carlsten M, Childs RW. Genetic manipulation of NK cells for cancer immunotherapy: techniques and clinical implications. *Front Immunol*. 2015;6:266.
170. Mohammed S, Sukumaran S, Bajgain P, Watanabe N, Heslop HE, Rooney CM, Brenner MK, Fisher WE, Leen AM, Vera JF. Improving chimeric antigen receptor-modified T cell function by reversing the immunosuppressive tumor microenvironment of pancreatic cancer. *Mol Ther*. 2017;25(1):249–58.
171. Wang Z, Guo L, Song Y, Zhang Y, Lin D, Hu B, Mei Y, Sandik D, Liu H. Augmented anti-tumor activity of NK-92 cells expressing chimeric receptors of TGF-betaR II and NKG2D. *Cancer Immunol Immunother* CII. 2017b;66(4):537–48.
172. Fan M, Li M, Gao L, Geng S, Wang J, Wang Y, Yan Z, Yu L. Chimeric antigen receptors for adoptive T cell therapy in acute myeloid leukemia. *J Hematol Oncol*. 2017;10(1):151.
173. Lu C, Guo C, Chen H, Zhang H, Zhi L, Lv T, Li M, Niu Z, Lu P, Zhu W. A novel chimeric PD1-NKG2D-41BB receptor enhances antitumor activity of NK92 cells against human lung cancer H1299 cells by triggering pyroptosis. *Mol Immunol*. 2020;122:200–6.
174. Schambach A, Morgan M. Retroviral vectors for cancer gene therapy. Recent results in cancer research. *Fortschritte der Krebsforschung Progres dans les recherches sur le cancer*. 2016;209:17–35.
175. Dufva O, Koski J, Maliniemi P, Ianevski A, Klievink J, Leitner J, Pölonen P, Hohtari H, Saeed K, Hannunen T, et al. Integrated drug profiling and CRISPR screening identify essential pathways for CAR T-cell cytotoxicity. *Blood*. 2020;135(9):597–609.
176. Ureña-Bailén G, Lamsfus-Calle A, Daniel-Moreno A, Raju J, Schlegel P, Seitz C, Atar D, Antony JS, Handgretinger R, Mezger M. CRISPR/Cas9 technology: towards a new generation of improved CAR-T cells for anticancer therapies. *Brief Funct Genom*. 2020;19(3):191–200.
177. Michieletto D, Lucic M, Marenduzzo D, Orlandini E. Physical principles of retroviral integration in the human genome. *Nat Commun*. 2019;10(1):575.
178. Ren J, Zhao Y. Advancing chimeric antigen receptor T cell therapy with CRISPR/Cas9. *Protein Cell*. 2017;8(9):634–43.
179. Ren J, Liu X, Fang C, Jiang S, June CH, Zhao Y. Multiplex genome editing to generate universal CAR T cells resistant to PD1 inhibition. *Clin Cancer Res*. 2017;23(9):2255–66.
180. Afolabi LO, Adeshakin AO, Sani MM, Bi J, Wan X. Genetic reprogramming for NK cell cancer immunotherapy with CRISPR/Cas9. *Immunology*. 2019;158(2):63–9.
181. Naeimi Kararoudi M, Dolatshad H, Trikha P, Hussain SA, Elmas E, Foltz JA, Moseman JE, Thakkar A, Nakkula RJ, Lamb M, et al. Generation of knock-out primary and expanded human NK cells using Cas9 ribonucleoproteins. *J Vis Exp JoVE*. 2018. <https://doi.org/10.3791/58237>.
182. Hautela J, Surgenor E, Huntington ND. Efficient genome editing of human natural killer cells by CRISPR RNP. *bioRxiv* 2018:406934.

183. Liu X, Zhang Y, Cheng C, Cheng AW, Zhang X, Li N, Xia C, Wei X, Liu X, Wang H. CRISPR-Cas9-mediated multiplex gene editing in CAR-T cells. *Cell Res*. 2017;27(1):154–7.
184. Nakazawa T, Natsume A, Nishimura F, Morimoto T, Matsuda R, Nakamura M, Yamada S, Nakagawa I, Motoyama Y, Park YS, et al. Effect of CRISPR/Cas9-mediated PD-1-disrupted primary human third-generation CAR-T cells targeting EGFRvIII on in vitro human glioblastoma cell growth. *Cells*. 2020;9(4):998.
185. Tang N, Cheng C, Zhang X, Qiao M, Li N, Mu W, Wei XF, Han W, Wang H. TGF- β inhibition via CRISPR promotes the long-term efficacy of CAR T cells against solid tumors. *JCI Insight*. 2020;5(4):e133977.
186. Michen S, Temme A. Genetically engineered natural killer cells as a means for adoptive tumor immunotherapy. *Crit Rev Immunol*. 2016;36(4):329–47.
187. Chu Y, Yahr A, Huang B, Ayello J, Barth M. M SC: Romidepsin alone or in combination with anti-CD20 chimeric antigen receptor expanded natural killer cells targeting Burkitt lymphoma in vitro and in immunodeficient mice. *Oncoimmunology*. 2017;6(9):e1341031.
188. Siegler EL, Kim YJ, Chen X, Siriwon N, Mac J, Rohrs JA, Bryson PD, Wang P. Combination cancer therapy using chimeric antigen receptor-engineered natural killer cells as drug carriers. *Mol Ther*. 2017;25(12):2607–19.
189. Zhang Q, Tian K, Xu J, Zhang H, Li L, Fu Q, Chai D, Li H, Zheng J. Synergistic effects of cabozantinib and EGFR-specific CAR-NK-92 cells in renal cell carcinoma. *J Immunol Res*. 2017;2017:6915912.
190. Felices M, Lenvik TR, Davis ZB, Miller JS, Vallera DA. Generation of BiKEs and TriKEs to improve NK cell-mediated targeting of tumor cells. *Methods Mol Biol*. 2016;1441:333–46.
191. Klingemann H, Boissel L. Targeted cellular therapy with natural killer cells. *Hormone and Metab Res*. 2008;40(2):122–5.
192. Battella S, Cox MC, Santoni A, Palmieri G. Natural killer (NK) cells and anti-tumor therapeutic mAb: unexplored interactions. *J Leukoc Biol*. 2016;99(1):87–96.
193. Zhen A, Kamata M, Rezek V, Rick J, Levin B, Kasparian S, Chen IS, Yang OO, Zack JA, Kitchen SG. HIV-specific immunity derived from chimeric antigen receptor-engineered stem cells. *Mol Ther*. 2015;23(8):1358–67.
194. Gschweng E, De Oliveira S, Kohn DB. Hematopoietic stem cells for cancer immunotherapy. *Immunol Rev*. 2014;257(1):237–49.
195. Lowe E, Truscott LC, De Oliveira SN. In vitro generation of Human NK cells expressing chimeric antigen receptor through differentiation of gene-modified hematopoietic stem cells. *Methods Mol Biol (Clifton, NJ)*. 2016;1441:241–51.
196. Tammana S, Huang X, Wong M, Milone MC, Ma L, Levine BL, June CH, Wagner JE, Blazar BR, Zhou X. 4-1BB and CD28 signaling plays a synergistic role in redirecting umbilical cord blood T cells against B-cell malignancies. *Hum Gene Ther*. 2010;21(1):75–86.
197. Chang YH, Connolly J, Shimasaki N, Mimura K, Kono K, Campana D. A chimeric receptor with NKG2D specificity enhances natural killer cell activation and killing of tumor cells. *Can Res*. 2013;73(6):1777–86.
198. Lanier LL, Corliss B, Wu J, Phillips JH. Association of DAP12 with activating CD94/NKG2C NK cell receptors. *Immunity*. 1998;8(6):693–701.
199. Carr WH, Rosen DB, Arase H, Nixon DF, Michaelsson J, Lanier LL. Cutting edge: KIR3DS1, a gene implicated in resistance to progression to AIDS, Encodes a DAP12-associated receptor expressed on NK cells that triggers NK cell activation. *J Immunol*. 2007;178(2):647–51.
200. Orange JS, Harris KE, Andzelm MM, Valter MM, Geha RS, Strominger JL. The mature activating natural killer cell immunologic synapse is formed in distinct stages. *Proc Natl Acad Sci USA*. 2003;100(24):14151–6.
201. Mukherjee M, Mace EM, Carisey AF, Ahmed N, Orange JS. Quantitative imaging approaches to study the CAR immunological synapse. *Mol Ther*. 2017;25(8):1757–68.
202. Xiong W, Chen Y, Kang X, Chen Z, Zheng P, Hsu YH, Jang JH, Qin L, Liu H, Dotti G, et al. Immunological synapse predicts effectiveness of chimeric antigen receptor cells. *Mol Ther*. 2018;26(4):963–75.
203. Srivastava S, Riddell SR. Engineering CAR-T cells: design concepts. *Trends Immunol*. 2015;36(8):494–502.
204. Shevtsov M, Multhoff G. Immunological and translational aspects of NK cell-based antitumor immunotherapies. *Front Immunol*. 2016;7:492.
205. Pockley AG, Vaupel P, Multhoff G. NK cell-based therapeutics for lung cancer. *Expert Opin Biol Ther*. 2020;20(1):23–33.
206. Lasek W, Zagożdżon R, Jakobisiak M. Interleukin 12: still a promising candidate for tumor immunotherapy? *Cancer Immunol Immunother*. 2014;63(5):419–35.
207. Buller CW, Mathew PA, Mathew SO. Roles of NK cell receptors 2B4 (CD244), CS1 (CD319), and LLT1 (CLEC2D) in cancer. *Cancers*. 2020;12(7):1755.
208. Armeanu S, Bitzer M, Lauer UM, Venturelli S, Pathil A, Krusch M, Kaiser S, Jobst J, Smirnow I, Wagner A, et al. Natural killer cell-mediated lysis of hepatoma cells via specific induction of NKG2D ligands by the histone deacetylase inhibitor sodium valproate. *Can Res*. 2005;65(14):6321–9.
209. Parihar R, Rivas C, Huynh M, Omer B, Lapteva N, Metelitsa LS, Gottschalk SM, Rooney CM. NK cells expressing a chimeric activating receptor eliminate MDSCs and rescue impaired CAR-T cell activity against solid tumors. *Cancer Immunol Res*. 2019;7(3):363–75.
210. Vey N, Karlin L, Sadot-Lebouvier S, Broussais F, Berton-Rigaud D, Rey J, Charbonnier A, Marie D, André P, Patuel C, et al. A phase 1 study of lirilumab (antibody against killer immunoglobulin-like receptor antibody KIR2D; IPH2102) in patients with solid tumors and hematologic malignancies. *Oncotarget*. 2018;9(25):17675–88.
211. Terszowski G, Klein C, Schmied L, Stern M. How to outsmart NK cell tolerance. *Oncoimmunology*. 2015;4(9):e1016708.
212. Aranda F, Vacchelli E, Eggermont A, Galon J, Fridman WH, Zitvogel L, Kroemer G, Galluzzi L. Trial watch: immunostimulatory monoclonal antibodies in cancer therapy. *Oncoimmunology*. 2014;3(1):e27297.
213. Faden DL, Concha-Benavente F, Chakka AB, McMichael EL, Chandran U, Ferris RL. Immunogenomic correlates of response to cetuximab monotherapy in head and neck squamous cell carcinoma. *Head Neck*. 2019;41(8):2591–601.
214. Kohrt HE, Thielens A, Marabelle A, Sagiv-Barfi I, Sola C, Chanuc F, Fuseri N, Bonnafous C, Czerwinski D, Rajapaksa A, et al. Anti-KIR antibody enhancement of anti-lymphoma activity of natural killer cells as monotherapy and in combination with anti-CD20 antibodies. *Blood*. 2014;123(5):678–86.
215. Yalniz FF, Daver N, Rezvani K, Kornblau S, Ohanian M, Borthakur G, DiNardo CD, Konopleva M, Burger J, Gasiot Y, et al. A pilot trial of lirilumab with or without azacitidine for patients with myelodysplastic syndrome. *Clin Lymphoma Myeloma Leukemia*. 2018;18(10):658–663. e652.
216. Carlsten M, Korde N, Kotecha R, Reger R, Bor S, Kazandjian D, Landgren O, Childs RW. Checkpoint inhibition of KIR2D with the monoclonal antibody IPH2101 induces contraction and hyporesponsiveness of NK cells in patients with myeloma. *Clin Cancer Res*. 2016;22(21):5211–22.
217. Benson DM Jr, Cohen AD, Jagannath S, Munshi NC, Spitzer G, Hofmeister CC, Efebera YA, Andre P, Zerbib R, Caligiuri MA. A phase I trial of the anti-kir antibody IPH2101 and lenalidomide in patients with relapsed/refractory multiple myeloma. *Clin Cancer Res*. 2015;21(18):4055–61.
218. Korde N, Carlsten M, Lee MJ, Minter A, Tan E, Kwok M, Manasanch E, Bhutani M, Tajeja N, Roschewski M, et al. A phase II trial of pan-KIR2D blockade with IPH2101 in smoldering multiple myeloma. *Haematologica*. 2014;99(6):e81–83.
219. Häusler SF, Del Barrio IM, Diessner J, Stein RG, Strohschein J, Höning A, Dietl J, Wischhusen J. Anti-CD39 and anti-CD73 antibodies A1 and 7G2 improve targeted therapy in ovarian cancer by blocking adenosine-dependent immune evasion. *Am J Transl Res*. 2014;6(2):129–39.
220. Bhat R, Watzl C. Serial killing of tumor cells by human natural killer cells—enhancement by therapeutic antibodies. *PLoS ONE*. 2007;2(3):e326.
221. Lin JK, Muffly LS, Spinner MA, Barnes JI, Owens DK, Goldhaber-Fiebert JD. Cost effectiveness of chimeric antigen receptor t-cell therapy in multiply relapsed or refractory adult large B-cell lymphoma. *J Clin Oncol*. 2019;37(24):2105–19.
222. Su S, Betterts DM, Ramanathan M, Keyvanfar K, Smith A, Feng X, Furutani E, Carlsten M, Lundqvist A, Childs R. Optimizing lentiviral transduction of human natural killer cells. *Blood*. 2011;118:4714.
223. Micucci F, Zingoni A, Piccoli M, Frati L, Santoni A, Galandrini R. High-efficient lentiviral vector-mediated gene transfer into primary human NK cells. *Exp Hematol*. 2006;34(10):1344–52.
224. Sutlu T, Nystrom S, Gilljam M, Stellan B, Applequist SE, Alici E. Inhibition of intracellular antiviral defense mechanisms augments lentiviral

- transduction of human natural killer cells: implications for gene therapy. *Hum Gene Ther.* 2012;23(10):1090–100.
225. Tran J, Kung SK. Lentiviral vectors mediate stable and efficient gene delivery into primary murine natural killer cells. *Mol Ther.* 2007;15(7):1331–9.
226. Boissel L, Betancur M, Lu W, Wels WS, Marino T, Van Etten RA, Klingemann H. Comparison of mRNA and lentiviral based transfection of natural killer cells with chimeric antigen receptors recognizing lymphoid antigens. *Leukemia Lymphoma.* 2012;53(5):958–65.
227. Guven H, Konstantinidis KV, Alici E, Aints A, Abedi-Valugerdi M, Christensson B, Ljunggren HG, Dilber MS. Efficient gene transfer into primary human natural killer cells by retroviral transduction. *Exp Hematol.* 2005;33(11):1320–8.
228. Dai H, Wu Z, Jia H, Tong C, Guo Y, Ti D, Han X, Liu Y, Zhang W, Wang C, et al. Bispecific CAR-T cells targeting both CD19 and CD22 for therapy of adults with relapsed or refractory B cell acute lymphoblastic leukemia. *J Hematol Oncol.* 2020;13(1):30.
229. Guillerey C, Huntington ND, Smyth MJ. Targeting natural killer cells in cancer immunotherapy. *Nat Immunol.* 2016;17(9):1025–36.
230. Rossi GR, Trindade ES, Souza-Fonseca-Guimaraes F. Tumor microenvironment-associated extracellular matrix components regulate NK cell function. *Front Immunol.* 2020;11:73.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

