



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

An Unusual Case of COVID-19: What Is This 3D Printed Model of the Biliary Tree?



Alexandre Cenatti,¹ Thomas R. McCarty,² and Diogo T. H. de Moura^{3,4}

¹Serviço de Endoscopia Do Hospital Saúde de Caxias do Sul, Rio Grande do Sul, Brazil; ²Division of Gastroenterology, Hepatology and Endoscopy, Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts;

³Gastroenterology Department, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil; and ⁴Instituto D'Or de Pesquisa e Ensino (IDOR), São Paulo, Brazil



Question: A 78-year-old woman presented with severe acute respiratory syndrome (SARS) owing to coronavirus disease 2019 (COVID-19). During the hospitalization, she developed abdominal pain, fever, and jaundice with leukocytosis (white blood count of $16 \times 10^9/L$), total bilirubin level of 21.1 mg/dL (reference range, 0.4–1.4 mg/dL), alkaline phosphatase of 870 U/L (reference range, 30–135 U/L), and serum alanine aminotransferase of 77 U/L (reference range, 10–55 U/L).

An abdominal ultrasound examination showed no dilation in the intrahepatic bile duct. Magnetic resonance cholangiopancreatography revealed a dilated distal common bile duct with filling defects in the proximal common bile duct and tapering of the intrahepatic bile ducts concerning for cholangitis. Endoscopic retrograde cholangiopancreatography (ERCP) was subsequently performed. During the biliary sweep with a retrieval basket, an uncommon structure—similar to a 3-dimensional-printed model of the biliary tract (Figure A)—was removed. Final cholangiography demonstrated progressive destruction and obliteration of the intrahepatic bile ducts (pruned tree). A plastic biliary stent was then placed.

The procedure was uncomplicated with improvement in clinical symptoms and laboratory derangements. Owing to the active COVID-19 infection, the patient was eventually discharged after 120 days of hospitalization. Despite the diagnosis of secondary sclerosing cholangitis in critically ill patients (SSC-CIP), the patient was doing well at one-month follow-up with plan for repeat ERCP in 2 months.

What is the diagnosis?

See the *Gastroenterology* web site (www.gastrojournal.org) for more information on submitting your favorite image to Clinical Challenges and Images in GI.

Correspondence

Address correspondence to: Diogo T. H. de Moura, MD, MSc, PhD, Hospital Vila Nova Star, Rua Dr. Alceu de Campos Rodrigues, 126, 04544-000, São Paulo, Brazil. e-mail: diogo.moura@vilanovastar.com.br.

Conflicts of interest

The authors disclose no conflicts.

© 2022 by the AGA Institute.

0016-5085/\$36.00

<https://doi.org/10.1053/j.gastro.2022.01.051>

Answer to: Image 5: Biliary Cast Syndrome

Biliary cast syndrome is characterized by the development of casts in the biliary tree. Although uncommon, it is usually associated with liver transplantation.¹ However, critical illness owing to COVID-19 may result in biliary complications such as bile duct casts and progressive destruction of the intrahepatic bile ducts.

The etiology of cast development is not fully understood but is likely to be multifactorial, including sludge as a prerequisite and related factors such as bile duct damage and ischemia, biliary infection, hemolysis, fasting, parenteral nutrition, abdominal surgery, hypovolemic shock, and polytrauma.^{1,2} Biliary cast syndrome is a hallmark of SSC-CIP. This is characterized by persistent cholestasis, biliary cast formation, and a stricturing fibrosing cholangiopathy.²

Different from our experience, endoscopic management has been shown to be effective in a minority of cases and surgical removal of casts is usually required.¹ However, SSC-CIP carries a high risk for hepatobiliary complications and, if progression toward cirrhosis occurs, the definitive treatment remains liver transplantation.²

Some cases of SSC-CIP among patients with SARS, especially owing to H1N1 influenza infection, have been reported; however, few reports have demonstrated a relationship to COVID-19.^{2,3} The combined rarity of SSC-CIP associated with SARS-CoV2 infection, successful management by endoscopy, as well as the removal of the entire biliary casts as 1 piece during ERCP, as described here, adds a valuable contribution to the literature, especially given the COVID-19 pandemic.

Keywords: Endoscopy; Gastrointestinal; Bleeding; COVID-19; ECMO.

References

1. Parry SD, Muiesan P. Cholangiopathy and the biliary cast syndrome. *Eur J Gastroenterol Hepatol* 2003;15:341–343.
2. Tunney R, Scott J, Rudralingam V, et al. Secondary sclerosing cholangitis following extracorporeal membrane oxygenation for acute respiratory distress in polytrauma. *Clin Case Rep* 2018;6:1849–1853.
3. Sanders D, Bomman S, Irani S. COVID-19-induced bile duct casts and cholangitis: a case report. *Cureus* 2021;13:e14560.