

Research Article

Amylenes Do Not Lead to Bacterial Mutagenicity in Contrast to Structurally Related Epoxides

Götz A. Westphal, Carolin Tüshaus, Christian Monsé, Nina Rosenkranz, Thomas Brüning, and Jürgen Büniger

IPA-Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Ruhr-Universität Bochum, 44789 Bochum, Germany

Correspondence should be addressed to Götz A. Westphal; westphal@ipa-dguv.de

Received 2 October 2013; Accepted 11 December 2013; Published 6 January 2014

Academic Editor: Masood Ahmad

Copyright © 2014 Götz A. Westphal et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Amylenes are unsaturated hydrocarbons (C_5H_{10}), such as 1-pentene, 2-pentene, 2-methyl-but-1-en (3-methyl-1-butene), 2-methyl-but-2-en (isopentene), and 3-methyl-but-1-en. We investigated bacterial mutagenicity of 1-pentene, 2-pentene, and 3-methyl-but-1-en in the Ames test. 2-Pentene was investigated as racemate and as pure diastereomers. We included the methyltransferase deficient *Salmonella* Typhimurium strain YG7108 and the application of a gas-tight preincubation to reduce the risk of false negative results. 1,2-Epoxy-pentane which may arise from 1-pentene was used as positive control. None of the investigated amylenes showed mutagenic effects, whereas 1,2-epoxy-pentane was mutagenic exceeding $100 \mu\text{g}$ per plate. An exceptional high reverse mutation in the negative control plates in the experiments with 1,2-epoxy-pentane was obviously caused by evaporation into the incubator which was shown by placing the control plates in a separate apparatus. No differences were seen upon use of YG7108 and its parent strain TA1535. In conclusion, 1,2-epoxy-pentane is most probably not a substrate of the deleted bacterial methyltransferases. The comparison of the bacterial mutagenicity of the investigated amylenes and 1,2-epoxy-pentane suggests that epoxidation of amylenes in the S9-mix does not proceed effectively or is counterbalanced by detoxifying reactions. The assessment of mutagenic effects of short chained aliphatic epoxides can be underestimated due to the evaporation of these compounds.

1. Introduction

Amylenes denote a group of unsaturated hydrocarbons (C_5H_{10}), including unbranched compounds such as 1- and 2-pentene (α - and β -amylyene) as well as branched compounds such as 2-methyl-but-2-en (isopentene) [CAS 513-35-9] and 3-methyl-1-butene [CAS 563-45-1] (Figure 1). Amylenes are used in the synthesis of amylyphenols, isoprene, and pentanols and can occur in gasoline or diesel fuel [1]. In addition amylenes are used as stabilizers in solvents; dichloromethane (DCM), for example, can contain up to 40 ppm.

Short chained unsaturated hydrocarbons react with biological molecules by electrophilic addition and can easily be epoxidized [2]. Both reactions can lead to mutagenic effects [3]. Theoretically, amylenes may react in a similar way. Short chained aliphatic epoxides such as ethylene oxide, 1,2-epoxypropane, and 1,2-epoxybutane are mutagenic in the

bacterial reverse mutation assay [3]. Epoxidation of 1-pentene is expected to yield 1,2-epoxy-pentane (propylloxirane). A study which is available at ECHA reports weak mutagenicity of 1,2-epoxy-pentane at and above $5000 \mu\text{g}/\text{plate}$ [4]. Therefore, mutagenic properties of amylenes are suspected. However, available toxicological data of most amylenes are fragmentary despite their frequent use. The available data show no bacterial mutagenicity for 1-pentene [5]. We were not able to identify studies which investigated bacterial mutagenicity of *cis*- and *trans*-2-pentene and 3-methyl-1-butene.

Amylenes and their possible oxidation products show close structural relationship to known mutagens and carcinogens. However, chain length and structure have a strong impact on the mutagenicity of such short chained epoxides: a methyl group in or near the oxirane ring prevents mutagenic




FIGURE 1

TABLE 1: Reverse mutations for positive and negative controls.

| Strain | | TA98 | TA100 | YG7108 |
|-----------------|-----|--------------------|--------------------|--------------------|
| Solvent control | -S9 | 18 ± 6 (N = 16) | 120 ± 17 (N = 17) | 20 ± 6 (N = 18) |
| | +S9 | 20 ± 7 (N = 20) | 145 ± 19 (N = 15) | 26 ± 9 (N = 19) |
| Gas-tight | -S9 | 19 ± 7 (N = 10) | 128 ± 18 (N = 9) | 24 ± 9 (N = 13) |
| | +S9 | 21 ± 4 (N = 20) | 141 ± 18 (N = 12) | 23 ± 6 (N = 17) |
| Not gas-tight: | | | | |
| 3-NBA, 5 ng | -S9 | 709 ± 328 (N = 12) | | |
| MMS, 200 µg | -S9 | | 1516 (N = 1) | 805 ± 283 (N = 10) |
| MMS, 500 µg | -S9 | | 968 ± 175 (N = 13) | |
| MMS, 1000 µg | -S9 | | 1543 ± 40 (N = 2) | |
| 2-AF | +S9 | 843 ± 228 (N = 19) | 646 ± 117 (N = 13) | |
| 2-AA, 5 µg | +S9 | | 781 ± 52 (N = 3) | |
| 2-AA, 10 µg | +S9 | 486 ± 175 (N = 3) | | |
| NDEA, 500 µg | +S9 | | | 225 ± 85 (N = 11) |
| NDEA, 1000 µg | +S9 | | | 414 ± 360 (N = 16) |

3-NBA: 3-nitrobenzanthrone; 2AF: 2-aminofluorene; 2-AA: 2-aminoanthracene; MMS: methyl methanesulfonate; NDEA: N-nitrosodimethylamine.

effects in case of the isoprene monoepoxide. Isoprene-dioxide (2-methyl-1,2,3,4-diepoxybutane) on the other hand is strong mutagenic [6]. The same is true for 2,3-epoxybutane which is a direct mutagen, whereas its 2-methyl derivative shows no mutagenicity [7].

The detection of mutagenic effects of low molecular weight compounds can be hampered by several factors such as evaporation [8, 9] and bacterial DNA repair [10]. The latter was first demonstrated using methyltransferase deficient tester strains such as YG7108 (*ogt⁻* and *ada⁻*) [11]. In addition, strong stereoselective effects regarding mutagenicity were reported in case of *cis*- and *trans*-2,3-epoxybutane [12] and for the diastereoisomers of 3-bromo-1,2-epoxycyclohexane [13]. We therefore compared 1-pentene, 2-pentene, *cis*- and *trans*-2-pentene, 3-methyl-but-1-en, and 1,2-epoxy-pentane in the bacterial reverse mutation test.

2. Methods

We investigated 1-pentene [CAS 109-67-1], 2-pentene [CAS 109-68-2], *cis*- [CAS 627-20-3] and *trans*-2-pentene [CAS

646-04-8], 3-methyl-1-butene [CAS 563-45-1] (Sigma-Aldrich, Steinheim, Germany), and 1,2-epoxy-pentane [CAS 1003-14-1] (CHEMOS GmbH, Regenstauf, Germany) using the bacterial reverse mutation assay, with the *S. Typhimurium* strains TA98, TA100, TA1535 [14] (TA1535 only in case of 1,2-epoxy-pentane), and the methyltransferase deficient tester strain YG7108 (*ogt⁻*, *ada⁻*) which is derived from TA1535 [11]. The test was carried out with and without metabolic activation by rat liver enzymes (S9) as well as gastight preincubation in gas-tight autosampler vials [8]. Since 3-methyl-1-butene is gaseous at room temperature it was withdrawn from a pressure cylinder and condensed at approximately -18°C using an ice/NaCl mixture. The condensate was filled in a headspace vial and transferred with a gas-tight syringe to the autosampler vials which contained the tester strains and the S9-mix or equivalent amounts of phosphate buffered saline.

We performed at least two independent experiments for every substance. The compounds were tested up to toxic concentrations but not above 5000 µg/plate. Acceptance criteria for a positive test were positive and negative controls within the range of the historical controls (see Table 1) and a

TABLE 2: Overview of all other results, not displayed as a figure.

| Compound | Strain | Condition | Result |
|-------------------------|--------|--|----------|
| 1-Pentene | TA98 | ±S9, gas-tight ($N = 3$) and not gas-tight ($N = 7$) | Negative |
| | TA100 | ±S9, gas-tight ($N = 2$) and not gas-tight ($N = 7$) | Negative |
| | YG7108 | ±S9, gas-tight ($N = 5$) and not gas-tight ($N = 7$) | Negative |
| 2-Pentene (racemate) | TA98 | ±S9, gas-tight ($N = 4$) and not gas-tight ($N = 4$) | Negative |
| | TA100 | ±S9, gas-tight ($N = 3$) and not gas-tight ($N = 4$) | Negative |
| | YG7108 | ±S9, gas-tight ($N = 3$) and not gas-tight ($N = 4$) | Negative |
| <i>cis</i> -2-pentene | TA98 | ±S9, not gas-tight ($N = 3$) | Negative |
| | TA100 | ±S9, not gas-tight ($N = 4$) | Negative |
| | YG7108 | ±S9, not gas-tight ($N = 3$) | Negative |
| <i>trans</i> -2-pentene | TA98 | ±S9, not gas-tight ($N = 3$) | Negative |
| | TA100 | ±S9, not gas-tight ($N = 4$) | Negative |
| | YG7108 | ±S9, not gas-tight ($N = 3$) | Negative |
| 3-Methyl-1-butene | TA98 | -S9, not gas-tight ($N = 2$) | Negative |
| | TA100 | -S9, not gas-tight ($N = 2$) | Negative |
| | YG7108 | -S9, not gas-tight ($N = 5$) | Negative |

reproducible, dose-dependent increase of reverse mutations across at least 2 consecutive concentrations, with a maximum exceeding the base rate by at least two times [15]. Counting was performed by an automatic colony counter (BioCount, Bio-Sys GmbH, Karben, Germany).

Chemicals and Test Compounds. Phenobarbital/ β -naphthoflavone induced S9 was purchased from MOLTOX, Trinova Biochem GmbH (Giessen, Germany). 1-Pentene, 2-pentene, *cis*- and *trans*-2-pentene, and 3-methyl-1-butene (95% purity, Sigma-Aldrich, Steinheim, Germany) were freshly dissolved in DMSO prior to each experiment. 1,2-Epoxy-pentane (98%), methyl methane sulfonate (99%), *N*-nitrosodiethylamine (99%), 2-aminofluorene (97%), and 2-aminoanthracene (95%) were purchased from Sigma-Aldrich, Steinheim, Germany. 3-Nitrobenzanthrone (99%) was purchased from Biochemical Institute for Environmental Carcinogens, Professor Dr. Gernot Grimmer-Foundation, Grosshansdorf, Germany.

Calculations of means and standard deviations were done using commercial software (GraphPad Prism version 4.00 for Windows, GraphPad Software, San Diego California USA).

3. Results

1,2-Epoxy-pentane was mutagenic in TA1535, YG7108 (Figures 2(a) and 2(b)), and TA100 (data not shown), exceeding the spontaneous mutation rate up to about 6-fold. No mutagenicity occurred in TA98 (data not shown). Notable differences of the response of 1,2-epoxy-pentane in TA1535 and YG7108 were not seen (Figures 2(a) and 2(b)). Remarkably the assay with 1,2-epoxy-pentane led to a strongly elevated spontaneous mutation rate (Figure 2(b), standard conditions). This was avoided by placing the control plates in a separate incubator (Figure 2(a)).

1-Pentene, 2-pentene, *cis*- and *trans*-2-pentene, and 3-methyl-1-butene were not mutagenic in any tester strain or

under any test condition, summarized in Table 2. All tests with the amylenes were done with and without metabolic activation.

Toxic effects occurred with and without addition of S9-mix, apparent by microcolony induction or reduced colony counts (latter is exemplarily shown in Figure 3). The racemate of 2-pentene revealed no toxicity in TA100 under standard conditions up to the highest tested amount of 5000 $\mu\text{g}/\text{plate}$ but according to the experiments using gas-tight preincubation the compound was toxic at and above 1000 $\mu\text{g}/\text{plate}$. Accordingly toxicity occurred in YG7108 at and above 1500 $\mu\text{g}/\text{plate}$ upon gas-tight preincubation, whereas no toxicity was seen up to 2500 $\mu\text{g}/\text{plate}$ under standard conditions.

In case of microcolony induction more closely spaced test doses were applied with the aim to exclude a dose dependency (microcolonies can be recognized as toxic artifacts by a missing dose dependency). Single experiments showed in fact a tendency towards a dose trend. Such indications of dose trends could however not be confirmed by use of more closely spaced test doses. Therefore, in some experiments we applied an unusual number of different test concentrations (see Figure 3). If no toxicity occurred the compounds were tested up to 5000 $\mu\text{g}/\text{plate}$.

4. Discussion

We observed no mutagenic effects of the short chained unsaturated hydrocarbons 1-pentene, 2-pentene, its stereo isomers *cis*-, *trans*-2-pentene, and 3-methyl-1-butene, even so we considered possible stereoselective effects in case of *cis*- and *trans*-2-pentene, possible evaporation of the compounds [8, 9], and bacterial repair [11]. YG7108 is sensitive for low molecular weight compounds which are false negative by use of the standard protocol and the parent strain TA1535. This was shown for compounds up to 4 C-atoms such as allyl chloride [10]. However, 1,2-epoxy-pentane exerted similar




FIGURE 2: Bacterial mutagenicity of 1,2-epoxy-pentane in TA1535 and YG7108 without addition of S9-mix. With the aim to reduce the elevated base rate (standard conditions (b)), the control plates (0 µg 1,2-epoxy-pentane) were placed in a separate incubator (a).




FIGURE 3: Results for 3-methyl-1-butene in TA98, TA100, and YG7108 with gas-tight preincubation and S9-mix. Three independent experiments were performed with differently spaced dose ranges.

bacterial mutagenicity in YG7108 and TA1535. Possibly C-5 compounds mark the cutoff for the substrates of the deleted bacterial methyltransferases. In addition these experiments show that the oxirane which can be formed in the metabolism of 2-pentene would exert mutagenic effects in the bacterial reverse mutation test.

Data on mutagenic effects of 1,2-epoxy-pentane are only available from the summary of a study report which is

cited by ECHA [4]. Mutagenicity in TA1535 was observed without S-9 mix from about 5000 µg/plate (factor 2.3–2.6) onward and with metabolic activation from about 2500 µg–5000 µg/plate (factor 1.7–2.0) onward [4]. In our experiments 1,2-epoxy-pentane was mutagenic at and above 100 µg per plate with a maximal increase of about 6-fold at 5000 µg per plate. This discrepancy can be explained by evaporation of 1,2-epoxy-pentane into the incubator. In our experiments this led to exceptional high reverse mutations in the negative control plates and quite weak mutagenic effects in the dosed samples. We attained a spontaneous reverse mutation rate within the historical range by placement of the control plates in a separate incubator (Figures 2(a) and 2(b)). Therefore, evaporation can lead to an underestimation of mutagenic effects of short chained oxiranes. This is in accordance with findings concerning ethylene oxide [3] and propylene oxide [3, 8, 16].

The question remains open why branched and unsaturated alylenes cannot be activated to mutagenic intermediates *in vitro*. Possibly the investigated alylenes are not efficiently epoxidized *in vitro* or oxidative metabolites which may occur are subjected to detoxification by the bacterial metabolism [17].

5. Conclusions

First, the investigated alylenes are not mutagenic, whereas the corresponding epoxide of 1-pentene (1,2-epoxy-pentane)

exerted clear bacterial mutagenicity. Epoxidation of the investigated amylenes using rat liver S9-mix may not be effective or is counterbalanced by detoxifying reactions. Second, weak bacterial mutagenicity of low molecular weight compounds such as 1,2-epoxipentane, ethylene-, or propylene oxide can be caused by evaporation of compounds.

Conflict of Interests

The authors declare that there is no conflict of interests.

Acknowledgment

The authors thank Dr. Takehiko Nohmi (National Institute of Health Sciences, Tokyo, Japan) for the donation of his tester strain YG7108.

References

- [1] HSDB, <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>.
- [2] D. Wistuba, H. P. Nowotny, O. Träger, and V. Schurig, "Cytochrome P-450-catalyzed asymmetric epoxidation of simple prochiral and chiral aliphatic alkenes: species dependence and effect of enzyme induction on enantioselective oxirane formation," *Chirality*, vol. 1, no. 2, pp. 127–136, 1989.
- [3] D. A. Canter, E. Zeiger, and S. Haworth, "Comparative mutagenicity of aliphatic epoxides in *Salmonella*," *Mutation Research*, vol. 172, no. 2, pp. 105–138, 1986.
- [4] ECHA, "Substance identification," <http://apps.echa.europa.eu/registered/data/dossiers/DISS-9eae805-c5d0-3646-044-00144f67d031/DISS-9eae805-c5d0-3646-e044-00144f67d031-DISS-9eae805-c5d0-3646-e044-00144f67d031.html#AGGR-a41a0eff-9de0-4379-bd0d-637cc33ac98d>.
- [5] ECHA, "Information on chemicals," <http://www.echa.europa.eu/web/guest/information-on-chemicals>.
- [6] P. G. Gervasi and V. Longo, "Metabolism and mutagenicity of isoprene," *Environmental Health Perspectives*, vol. 86, pp. 85–87, 1990.
- [7] C. de Meester, M. Mercier, and F. Poncelet, "Non-mutagenicity of 2-methyl-2,3-epoxybutane and factors influencing the mutagenicity of 2,3-epoxybutane," *Journal of Applied Toxicology*, vol. 2, no. 6, pp. 284–288, 1982.
- [8] G. A. Westphal, M. Blaszkewicz, M. Leutbecher, A. Muller, E. Hallier, and H. M. Bolt, "Bacterial mutagenicity of 2-chloro-1,3-butadiene (chloroprene) caused by decomposition products," *Archives of Toxicology*, vol. 68, no. 2, pp. 79–84, 1994.
- [9] G. A. Westphal, J. Bünger, T. G. Schulz, M. M. Müller, and E. Hallier, "Mutagenicity of N-nitrosodiethylamine in the Ames test with *S. typhimurium* TA1535 is due to volatile metabolites and is not dependent on cytochrome P4502E1 induction," *Archives of Toxicology*, vol. 74, no. 10, pp. 638–641, 2000.
- [10] B. Emmert, J. Bünger, K. Keuch et al., "Mutagenicity of cytochrome P450 2E1 substrates in the Ames test with the metabolic competent *S. typhimurium* strain YG7108pin3ERb₅," *Toxicology*, vol. 228, no. 1, pp. 66–76, 2006.
- [11] M. Yamada, B. Sedgwick, T. Sofuni, and T. Nohmi, "Construction and characterization of mutants of *Salmonella typhimurium* deficient in DNA repair of O⁶-methylguanine," *Journal of Bacteriology*, vol. 177, no. 6, pp. 1511–1519, 1995.
- [12] P. Castelain, B. Criado, M. Cornet, R. Laib, V. Rogiers, and M. Kirsch-Volders, "Comparative mutagenicity of structurally related aliphatic epoxides in a modified *Salmonella*/microsome assay," *Mutagenesis*, vol. 8, no. 5, pp. 387–393, 1993.
- [13] G. Bellucci, A. Lippi, and F. Marioni, "Structure activity relationship of epoxides: different mutagenicity of the two diastereoisomeric 3-bromo-1,2-epoxycyclohexanes," *Chemo-Biological Interactions*, vol. 51, no. 1, pp. 77–89, 1984.
- [14] D. M. Maron and B. N. Ames, "Revised methods for the *Salmonella* mutagenicity test," *Mutation Research*, vol. 113, no. 3-4, pp. 173–215, 1983.
- [15] K. Mortelmans and E. Zeiger, "The Ames *Salmonella*/microsome mutagenicity assay," *Mutation Research*, vol. 455, no. 1-2, pp. 29–60, 2000.
- [16] J. Bootman, D. C. Lodge, and H. E. Whalley, "Mutagenic activity of propylene oxide in bacterial and mammalian systems," *Mutation Research*, vol. 67, no. 2, pp. 101–112, 1979.
- [17] S. A. Ensign and J. R. Allen, "Aliphatic epoxide carboxylation," *Annual Review of Biochemistry*, vol. 72, pp. 55–76, 2003.