



# Peritoneal Dialysis After Liver Transplantation: A Systematic Review

Canadian Journal of Kidney Health and Disease  
Volume 8: 1–9  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/20543581211029722  
journals.sagepub.com/home/cjk



Jean Maxime Côté<sup>1,2,3</sup> , Isabelle Ethier<sup>1,2,4</sup>, Héroïse Cardinal<sup>1,2</sup>, and Marie-Noëlle Pépin<sup>1</sup>

## Abstract

**Background:** Chronic kidney disease following liver transplantation is a major long-term complication. Most liver transplant recipients with kidney failure will be treated with dialysis instead of kidney transplantation due to noneligibility and shortage in organ availability. In this population, the role of peritoneal dialysis (PD) as a modality of kidney replacement therapy (KRT) remains unclear.

**Objective:** To determine the feasibility regarding safety, technique survival, and dialysis efficiency of PD in liver transplant recipients requiring KRT for maintenance dialysis.

**Design:** Systematic review.

**Setting:** Interventional and observational studies reporting the use of PD after liver transplantation.

**Patients:** Adult liver transplant recipients with kidney failure treated with maintenance KRT.

**Measurements:** Extracted data included eligibility criteria, study design, demographics, and PD modality. The following outcomes of interest were extracted: rate of peritonitis and microorganisms involved, noninfectious peritoneal complications, technique survival, and kidney transplantation-censored technique survival. Non-PD complications included overall survival, liver graft dysfunction, and hospitalization rate.

**Methods:** The following databases were searched until July 2020: MedLine/PubMed, EMBASE, CINAHL, and Cochrane Library. Two reviewers independently screening all titles and abstracts of all identified articles. Due to the limited sample size, observational designs and study heterogeneity expected, no meta-analysis was pre-planned. Descriptive statistics were used to report all results.

**Results:** From the 5263 identified studies, 4 were included in the analysis as they reported at least 1 outcome of interest on a total of 21 liver transplant recipients, with an overall follow-up duration on PD of 19.0 (Interquartile range [IQR]: 9.5–29.5) months. Fifteen episodes of peritonitis occurred in a total cumulative PD follow-up of 514 patient-months, representing an incidence rate of 0.35 per year. These episodes did not result in PD technique failure, mortality, or impairment of liver graft function.

**Limitations:** Limitations include the paucity of studies in the field and the small number of patients included in each report, a risk of publication bias and the impossibility to directly compare hemodialysis to PD in this population. These results, therefore, must be interpreted with caution.

**Conclusions:** Based on limited data reporting the feasibility of PD in liver transplant recipients with kidney failure, no signal was associated with an increased risk of infectious complications. Long-term studies evaluating this modality need to be performed.

**Registration (PROSPERO):** CRD42020218374.

## Abrégé

**Contexte:** L'insuffisance rénale chronique est une complication majeure à long terme survenant après une transplantation hépatique. La plupart des transplantés du foie qui développent une insuffisance rénale seront traités par dialyse plutôt que par une greffe rénale; En raison de la pénurie d'organes et par non-éligibilité à la greffe rénale, la plupart des transplantés du foie qui développent une insuffisance rénale avancée seront traités par dialyse. L'importance de la dialyse péritonéale (DP) comme modalité de remplacement rénal demeure toutefois inconnue dans cette population de patients.

**Objectifs:** Étudier la faisabilité de la DP en matière d'innocuité, d'efficacité et de survie de la technique chez les receveurs d'une greffe hépatique qui nécessitent une thérapie de remplacement rénal comme dialyse d'entretien.

**Type d'étude:** Revue systématique

**Cadre:** Les études interventionnelles et observationnelles signalant l'utilisation de la DP après une transplantation hépatique.



**Sujets:** Des adultes ayant subi une transplantation hépatique et dont l'insuffisance rénale secondaire est traitée par thérapie de remplacement rénal.

**Mesures:** Les données extraites comprenaient les critères d'admissibilité, la méthodologie de l'étude, les caractéristiques démographiques des sujets et la modalité de DP. Les résultats d'intérêt suivants ont été extraits : le taux de péritonites et les microorganismes impliqués, les complications péritonéales non infectieuses, la survie de la technique et la survie de la technique censurée par la transplantation rénale. Les complications non liées à la DP comprenaient la survie globale, la défaillance du greffon hépatique et le taux d'hospitalisation.

**Méthodologie:** Les bases de données Medline/PubMed, Embase, CINAHL et Cochrane Library ont été consultées jusqu'en juillet 2020. Deux réviseurs indépendants ont examiné les titres et résumés de tous les articles recensés. Aucune méta-analyse n'a été planifiée en raison de la nature observationnelle et de l'hétérogénéité attendue des études retenues, et de la faible taille de l'échantillon. Des statistiques descriptives ont été utilisées pour présenter les données.

**Résultats:** Des 5263 études recensées, seules quatre ont été incluses dans l'analyse, rapportant un total de 21 transplantés hépatiques, dont la durée médiane sur DP s'établissait à 19.0 mois (IIQ: 95 à 29.5). Au cours d'un suivi cumulatif de 514 mois-patients sur DP, 15 épisodes de péritonite ont été observés, soit un taux d'incidence de 0,35 par année. Ces épisodes n'ont pas entraîné d'échec de la DP, ni de mortalité ou d'altération de la fonction du greffon hépatique.

**Limites:** Les limites comprennent le manque d'études sur le sujet, le faible nombre de patients inclus dans chaque rapport, un risque de biais de publication et l'impossibilité de comparer directement l'hémodialyse à la DP dans cette population. Les résultats doivent ainsi être interprétés avec prudence.

**Conclusion:** Selon les données limitées portant sur la faisabilité de la dialyse péritonéale chez les receveurs d'une greffe hépatique atteints d'insuffisance rénale, aucun signal notable n'est associé à un risque accru de complications infectieuses. Des études à long terme évaluant cette modalité sont nécessaires.

## Keywords

peritoneal dialysis, liver transplantation, systematic review, kidney failure, peritonitis, chronic kidney disease

Received March 7, 2021. Accepted for publication May 25, 2021.

## Introduction

In advanced liver disease, liver transplantation (LT) is the treatment of choice in eligible patients. Over time, graft and patient survival have improved significantly with the development of potent immunosuppression, including calcineurin inhibitors (CNIs).<sup>1</sup> The development of posttransplant chronic kidney disease (CKD) is a major long-term complication of LT, with stage-3 CKD occurring in 56.8%, stage 4 in 6.1%, and stage 5 in 2.3% of LT recipients 10 years post-transplant.<sup>2,3</sup> Since the introduction of the Model for End-stage Liver Disease (MELD) allocation system in 2002 that prioritizes LT candidates with pre-existing kidney dysfunction, the incidence of CKD after LT has progressed, where stage 4 or 5 occurred in up to 22% of all LT recipients 5 years after transplantation.<sup>4,5</sup> Among LT recipients who survived the first 3 months post-transplantation, 4% will eventually require maintenance dialysis.<sup>1</sup>

Post-transplant CKD has various causes, including pre-transplant insults such as infection, renal hypoperfusion, perioperative renal insults, and finally, postoperative offenses such as CNI toxicity.<sup>6</sup> The occurrence of CKD after nonkidney organ transplantation is not without consequence, with an increased risk of death when compared to solid-organ transplant recipients who have not developed CKD.<sup>1</sup>

Kidney transplantation confers a significant survival benefit in nonkidney solid-organ transplant recipients when compared to remaining wait-listed and on dialysis.<sup>5,7-9</sup> However, due to the shortage in organ availability as well as the noneligibility of a large proportion of these patients, most LT recipients who develop kidney failure need to initiate dialysis.<sup>8</sup> Currently, hemodialysis (HD) is the most common kidney replacement therapy (KRT) modality used in LT recipients suffering from kidney failure.<sup>8</sup> These patients represent 3% to 5% of the prevalent long-term HD population in the United States.<sup>1</sup>

<sup>1</sup>Division of Nephrology, Department of Medicine, Centre hospitalier de l'Université de Montréal, QC, Canada

<sup>2</sup>Research Center, Centre hospitalier de l'Université de Montréal, QC, Canada

<sup>3</sup>Clinical Research Centre, University College Dublin, Ireland

<sup>4</sup>Department of Nephrology, Princess Alexandra Hospital, Brisbane, Australia

### Corresponding Author:

Jean-Maxime Côté, Service of Nephrology, Centre hospitalier de l'Université de Montréal, 1000 St-Denis, Montréal, QC H2X 0C1, Canada.

Email: Jean-maxime.cote@umontreal.ca

In the general population of patient undergoing KRT, peritoneal dialysis (PD) and HD appear to be equivalent in terms of patient survival.<sup>10</sup> However, the role of PD in patients having received previous LT remains unclear. Possible concerns about technique success and infectious risks may explain the limited use of PD in this patient population.<sup>11</sup> Although a few publications have reported the success of the PD technique in LT recipients progressing to kidney failure, no study has systematically analyzed the feasibility of PD in this clinical context.

Our aim was to assess the feasibility regarding safety, technique survival, and dialysis efficiency of PD in patients who develop end-stage CKD after LT. To achieve this, we performed a systematic search of the literature to uncover the potential benefits and limitations of PD in this patient population.

## Method

This systematic review is reported following the Preferred Reporting Items for Systematic Reviews and Meta-analyses guidelines.<sup>12</sup> The protocol was registered into the PROSPERO international prospective register of systematic reviews (CRD42020218374).

### Eligibility Criteria

Eligible studies reported the use of PD for maintenance KRT in adult LT recipients. Studies addressing HD were included if data for at least one patient treated with PD were reported separately. Patients who received LT from living or deceased donors were included. Likewise, studies addressing maintenance KRT after various solid-organ transplantations were included if data for LT recipients were individually reported. No study design, date, or language limits were imposed on the search.

### Literature Search

According to the predetermined protocol, a systematic literature search of the following databases was performed from inception until July 2020 in cooperation with a trained medical librarian: MedLine/PubMed, EMBASE, Cochrane Library, and CINAHL. The literature search strategy was developed using medical subject headings and text words related to PD and LT (Supplementary File, Table S1). To ensure literature saturation, we also scanned the references list of all eligible articles and searched the gray literature for all abstracts listed into the annual meetings archives of the *American Society of Nephrology*, the *European Renal Association-European Dialysis & Transplant Association*, the *International Society of Peritoneal Dialysis*, and the *International Society of Nephrology*. Finally, we circulated a bibliography of the included articles to all authors, to add any other relevant publications.

### Study Selection

Eligible studies were clinical trials, cohort studies, case-control studies, case reports, and case series. Published abstracts were also eligible when at least one outcome of interest was described quantitatively. Literature search results were screened with *Rayyan QCR*. Two reviewers (J.M.C. and I.E.) independently screened the titles and abstracts of all identified articles. Disagreements were resolved by consensus-based discussions.

### Data Extraction and Quality Assessment

Extracted data included eligibility criteria, study type, demographics, PD modality, and results. The following prespecified outcomes of interest related to PD were extracted: rate of peritonitis and microorganisms involved, noninfectious peritoneal complications classified as abdominal hernia, exit-site or abdominal leak, or others; technique survival and kidney transplantation-censored technique survival. Non-PD complications included overall survival, liver graft dysfunction, and hospitalization rate.

Risk of bias was assessed using the *Newcastle-Ottawa Score* for cohort studies. Although not specified in the preplanned protocol, the *Joanna Briggs Institute Tool* was used to assess the risk of bias for case series.<sup>13</sup> When insufficient details were reported, the risk of bias was judged as unclear.


### Statistical Analysis and Synthesis of Results

As most studies were expected to be observational, no meta-analysis was preplanned. Descriptive statistics were reported as median with interquartile range (IQR) for continuous variables and proportion for dichotomic variables (SPSS 26; Armonk, NY; IBM Corp.). No statistical hypothesis test was performed due to the limited sample size and various follow-up definitions. Heterogeneity across all included studies was described narratively. When quantitative synthesis was not possible, a narrative synthesis provided additional relevant information.

## Results

### Study Selection

Study selection is depicted in Figure 1. After removal of duplicates, we identified 5,263 studies. Of these, 5,251 were excluded after an initial screening of titles and abstracts. A total of 12 unique studies were deemed eligible for full-text screening<sup>8,11,14-23</sup> from which 4 presented quantitative results for at least 1 prespecified outcome.<sup>15,16,20,21</sup> The remaining studies were excluded based on reasons specified in Figure 1. The studies by Shukla et al<sup>22</sup> and Cornelis et al<sup>17</sup> were excluded because cases reported were already described in the more recent publication by Saiprasertkit et al.<sup>20</sup>



**Figure 1.** Flow chart of included studies.

### Study and Patient Characteristics

An overview of study characteristics is presented in Table 1. All included studies, published between 2004 and 2018, were observational. One study had a comparator group of PD patients with no history of transplantation.<sup>16</sup> Two studies described cases from Canada, while the two others were from France and the United States, cumulating 21 LT recipients treated with PD as a modality of KRT. Patients' median age was 60 (IQR: 57-66) years and 43% of patients were female. The most common cause of liver failure leading to LT was viral hepatitis (63%), while CNI toxicity, based on clinical judgment, was involved in the cause of kidney failure for 81% of reported patients. Only one patient started KRT following acute kidney injury in the postoperative LT period and remained dialyzed for 7 months.<sup>15</sup> For the remaining, the time between LT and kidney failure requiring KRT initiation ranged from 2 to 24 years (median: 10.0; IQR: 7.1-15.8). At

the time of PD initiation, 38% of LT recipients had diabetes, and only 1 patient was anuric. The PD modality chosen was automated peritoneal dialysis (APD) for 11 patients and continuous ambulatory peritoneal dialysis (CAPD) for another 11 patients. For 19 patients (90%), the first modality of KRT received was PD, while 2 patients were previously treated with HD before transferring to PD. The overall follow-up duration on PD was 19.0 months (IQR: 9.5-29.5).

### Risk of Bias and Quality Assessment

The risks of bias for cohort studies and case series are presented in Supplementary Tables S2 and S3. As all studies included were either retrospective cohort studies, case series, or case reports, the overall quality of evidence resulting from this systematic review was low. Considering the limited sample size and the rare use of PD in LT recipients, there is a risk of

**Table 1.** Characteristics of Included Studies and Patients.

	Apiratpracha et al <sup>15</sup>	Buffet et al <sup>16</sup>	Saiprasertkit et al <sup>20</sup>	Shepherd and Holley <sup>21</sup>	Total
<b>Studies characteristics</b>					
Study type	Case report	Retrospective cohort	Case series	Case report	—
Sample size	1	5/13	14	1	21
Comparison group	—	Non-Tx pt on PD	—	—	—
Country	Canada	France	Canada	United States	—
Era (of follow-up)	2000-2005	2006-2016	1991-2016	2001-2004	—
Median follow-up duration on PD in months (IQR)	10.0	6.0 (3.5-15.5)	26.0 (14.5-37.8)	22.0	19.0 (9.5-29.5)
<b>Patients characteristics</b>					
Age at ESKD, median (IQR)	52	65 (43-67)	59 (54-64)	60	60 (57-66)
Male sex, no (%)	F	3 (60)	8 (57)	M	12 (57)
Diabetes, no (%)	0	4 (80)	4 (29)	0	8 (38)
Cause of liver failure, no					
Alcoholic	1	—	1 (7.1) <sup>a</sup>	1	3 (19) <sup>b</sup>
Hepatitis C virus or Hepatitis B virus	—	—	10 (71) <sup>a</sup>	—	10 (63) <sup>b</sup>
Nonalcoholic steatohepatitis	—	—	1 (7.1) <sup>a</sup>	—	1 (6.2) <sup>b</sup>
Other	—	—	4 (29) <sup>a</sup>	—	4 (25) <sup>b</sup>
Cause of renal failure, no (%)					
CNI toxicity	—	—	12 (86) <sup>a</sup>	1	13 (81) <sup>b</sup>
Glomerulonephritis	—	—	3 (21) <sup>a</sup>	—	3 (19) <sup>b</sup>
AKI	1	—	—	—	1 (6.2) <sup>b</sup>
Other	—	—	1 (7.1) <sup>a</sup>	—	1 (6.2) <sup>b</sup>
Anuric, %	No	0 (0)	1 (7.1)	No	1 (4.8) <sup>b</sup>
Immunosuppression, no (%)					
CNI only	1	—	5 (36)	—	—
CNI + Pred	—	—	4 (29)	—	—
CNI + Antimetabolites ± Pred	—	—	5 (36)	—	—
Median time from LT to ESKD in years (IQR)	0	13.8 (8.3-22.3)	9.5 (6.0-14.3)	10	10.0 (7.1-15.8)
<b>PD characteristics</b>					
<b>PD modality</b>					
Automated PD, no (%)	1 <sup>c</sup>	3 (60)	7 (50)	—	11 (52)
Continuous ambulatory PD, no (%)	1 <sup>c</sup>	2 (40)	7 (50)	1	11 (52)
hemodialysis prior to PD, no (%)	1	0 (0)	1 (7.1)	No	2 (9.5)
<b>Note</b>					
	Hemodialysis was initiated acutely post-LT, with transfer to PD at discharge. Dialysis was stopped 1 year after initiation due to renal recovery	Results were presented individually for all patients (n: 13). Only LT patients were included (n: 5)	Results were presented individually for all patients (n: 14)	Patient was primarily started on PD	

Note. PD = peritoneal dialysis; IQR= interquartile range; ESKD= end-stage kidney disease; CNI = calcineurin inhibitors; LT = liver transplantation.

<sup>a</sup>Patients may have more than 1 cause.

<sup>b</sup>Excluding Buffet et al cohort where data were not available.

<sup>c</sup>The case was treated with CAPD for 2 months, then switched to APD for 5 months.



**Table 2.** Outcomes Reported in Included Studies.

	Apiratpracha et al <sup>15</sup>	Buffet et al <sup>16</sup>	Saiprasertkit et al <sup>20</sup>	Shepherd and Holley <sup>21</sup>	Total
	n = 1	n = 5 (of 13) <sup>a</sup>	n = 14	n = 1	n = 21
<b>Infectious outcomes</b>					
Pt with peritonitis, no. (%)	0 (0)	3 (60)	10 (71)	0 (0)	<b>13 (62)</b>
Number of peritonitis, no.	0	4	11	0	<b>15</b>
Peritonitis event rate (1/patient-months)	—	1/11	1/40	—	<b>1/34</b>
<b>Microorganisms, no<sup>b</sup></b>					
<i>Corynebacterium</i>	—	1	1	—	—
<i>S. marcesens</i>		1			
<i>Coag.-neg Staphylococcus</i>		1	4		
<i>Staphylococcus aureus</i>			2		
<i>K. pneumoniae</i>		1	1		
<i>E. coli</i>			1		
<i>Acinetobacter baumannii</i>			1		
<i>Proteus mirabilis</i>			1		
<i>Candida</i> sp.			1		
No growth			1		
<b>Noninfectious PD-related outcomes</b>					
Abdominal hernia, no. (%)	1 (100)	0 (0)	2 (14)	0 (0)	<b>3 (14)</b>
Exit-site leak, no. (%)	0 (0)	0 (0)	1 (7.1)	0 (0)	<b>1 (4.8)</b>
Catheter dysfunction, no. (%)	0 (0)	2 (40)	0 (0)	0 (0)	<b>2 (9.5)</b>
<b>PD technique and Survival</b>					
Cumulated follow-up (patient-months)	10	44	438	22	<b>514</b>
<b>PD cessation, no. (%)</b>					
PD failure <sup>c</sup>	0 (0)	0 (0)	1 (7.1)	0 (0)	<b>1 (4.8)</b>
Kidney transplant	0 (0)	2 (40)	4 (29)	0 (0)	<b>6 (29)</b>
Death during follow-up, no. (%)	0 (0)	3 (60)	4 (29)	0 (0)	<b>7 (33)</b>
Graft dysfunction, no. (%)	0 (0)	0 (0)	0 (0)	0 (0)	<b>0 (0)</b>

Note. PD = peritoneal dialysis.

<sup>a</sup>Including only liver transplanted patients on PD from the cohort.

<sup>b</sup>Multiple organisms can be implicated for a single peritonitis episode.

<sup>c</sup>Defined as inadequate PD leading to transfer to hemodialysis.

publication bias, where only positive experiences regarding PD after LT might have been reported in the medical literature.

## Outcomes

**Peritonitis.** When combining all studies, 15 episodes of peritonitis occurred in a total cumulative PD follow-up of 514 patient-months, representing an incidence rate of peritonitis of 1 for every 34 patient-months or 0.35 episodes per patient-year. Clinical evolution and response to antimicrobial therapy were reported as similar to PD-related peritonitis in nontransplanted patients. Microorganism identification was specified in 2 studies<sup>16,20</sup> and is detailed in Table 2. *Staphylococcus* sp. were the most frequently reported. The comparison between the rate of peritonitis for these LT recipients with the complete local PD program for Saiprasertkit et al<sup>20</sup> is presented in Table 3. The study by Buffet et al<sup>16</sup> showed no difference in the peritonitis-free survival when comparing the whole cohort of solid-organ transplant recipients treated

with PD to nontransplant recipients (hazard ratio [HR]: 0.91; 95% CI: [0.37, 2.22]). One patient died after hospitalization for peritonitis-associated sepsis.<sup>20</sup> No study reported the antimicrobial regimen used for these peritonitis episodes.

**Noninfectious PD-related complications.** Abdominal hernias were reported for 3 patients (14%) of which 2 were corrected surgically with patients resuming PD after surgery<sup>15,20</sup> One was medically managed with conversion from CAPD to APD with dry day to minimize intra-abdominal pressure.<sup>15</sup> Exit-site leak was reported for 1 patient who required a temporary conversion to HD before resuming PD once the exit-site had healed.<sup>20</sup> Catheter dysfunction occurred in 2 patients, but no additional information regarding its impact was reported.<sup>16</sup>

**Dialysis adequacy and technique survival.** The overall dialysis adequacy was considered within current standards for all included studies. One patient (4.8%), from the Saiprasertkit et al study,<sup>20</sup> was transferred to HD after 16 months due to

**Table 3.** Risk of Peritonitis in Liver Transplant Recipients Compared to Nontransplanted Patients on Peritoneal Dialysis.

	Buffet et al <sup>16</sup>		Saiprasertkit et al <sup>20</sup>	
	PD in solid-organ transplanted pts (n: 13)	PD in nontransplanted pts (n: 370)	PD in liver transplant pts (n: 14)	Whole PD program (n: ?)
Peritonitis event rate, 1/patient-months	—	—	1/41	1/39
Survival without peritonitis event vs control, hazard ratio (95% CI)	0.91 (0.37-2.22)	—	—	—

Note. PD = peritoneal dialysis; ? = data not reported; CI = confidence interval.

poor ultrafiltration, reaching the endpoint of PD failure. The remaining LT recipients terminating PD either underwent kidney transplantation (n = 6, 29%) or died (n = 7, 33%) during follow-up. The median time from PD initiation until death was 10 months (IQR: 5-33, n = 7).

**Liver graft dysfunction.** No liver graft dysfunction or rejection was observed in any study. One group reported their home protocol for liver biopsies for PD patients.<sup>20</sup> The procedure was done in the absence of intraperitoneal PD solution, with antibiotic prophylaxis consisting of oral amoxicillin and intravenous tobramycin. Patients were allowed to resume PD exchange 48 hours after the procedure.

No data regarding the prespecified outcome of hospitalization rate were reported in the selected studies.

## Discussion

As LT recipient survival increases, CKD has become a major complication of LT, leading to increased morbidity and mortality. Most CKD risk factors in this population, such as the use of CNI, need for contrast procedures, or pre-existent CKD, cannot be easily mitigated.<sup>24</sup>

The impact of the dialysis modality on LT recipients who require KRT remains unknown. However, having the opportunity to offer a home-based KRT modality can be attractive for this patient population as LT recipients are on average younger than the general dialysis population.<sup>17</sup> Potential benefits of PD include reduction of the risk of hepatitis virus transmission, optimization of hemodynamic stability, decreasing risk of catheter-related bloodstream infection, and preservation of residual kidney function and vascular access.<sup>11,20,25-27</sup> However, infectious and anatomical concerns regarding the feasibility of PD in patients with an intra-abdominal transplant and remodeled abdomen may deter nephrologists from choosing PD when LT recipients develop end-stage CKD.

Here, we identified four studies reporting quantitative data for 21 patients concerning our prespecified outcomes of interest. The paucity of reports of PD use in LT recipients probably reflects the fear of an increased risk of peritonitis and associated complications in this immunocompromised population. Two studies allowed direct comparisons of the

rate of peritonitis in LT patients to that observed in cohorts of nontransplanted PD patients (see Table 3).<sup>16,17,20</sup> When pooling the reported experience from this systematic review, an overall peritonitis incidence rate of 1/34 patient-months was found, which represents 0.35 infections per patient-year. This rate is slightly higher than that recently reported for general PD population in Canada (0.27; 95% CI: [0.25, 0.30]) and Australia/New Zealand (0.29; 95% CI: [0.26, 0.32]) but similar to that observed in the United Kingdom (0.35; 95% CI: [0.30, 0.40]).<sup>28</sup>

Immunosuppression is a known risk factor for peritonitis and other intra-abdominal infections, and this risk cannot be easily minimized following solid-organ transplantations.<sup>29,30</sup> However, observational data from Australia and New Zealand showed that patients with ESKD who started PD after failed kidney transplants—most of whom were still receiving immunosuppressive therapy—had similar peritonitis-free survival (adj. HR: 0.92, 95% CI: [0.72, 1.16], *P* = .44) to those starting PD with native kidneys.<sup>31</sup> In addition, other studies have reported incidence rates of peritonitis of 0.36 episode per patient-year following heart and lung transplantation,<sup>32</sup> and 0.41 episode per patient-year following nonrenal solid-organ transplant in a second.<sup>17</sup> These figures are similar to the incidence rate we derived in this systematic review of LT recipients (0.35 episode per patient-year). However, when compared to nontransplanted PD populations receiving immunosuppressive therapy, the risk of peritonitis in LT patients was lower than for PD patients with lupus (0.68 episode per patient-year) but similar to PD patients with primary long-term glomerulonephritis (0.29 episode per patient-year).<sup>33</sup> Thus, in this review, no clinically significant trend of an increased risk of peritonitis episodes in LT patients receiving PD has been identified, when compared to nontransplanted patients receiving immunosuppression or to patients transplanted with solid organs other than the liver.

We found similar clinical evolution during treatment and no report of liver graft dysfunction following peritonitis episode. Some centers have already developed protocols to minimize the risk of infectious complications in patients requiring posttransplantation liver biopsy.<sup>20</sup> However, no international guideline has been published regarding the prophylactic management of such interventions in LT recipients on PD.

LT is a potential risk factor for impaired wound healing and postsurgery incisional hernia. The limited number of studies selected does not allow us to draw definite conclusions concerning the safety of PD with regards to noninfectious complications. Nevertheless, none of the patients who developed such complications had technique failure requiring permanent transfer to HD. To decrease intrabdominal pressure associated with PD exchange, LT recipients might be better candidates for APD than CAPD when possible. Careful planning of early PD catheter insertion in patients with progressive deterioration of kidney function may also reduce the risk of requiring urgent HD initiation prior to PD by permitting sufficient time for wound healing.<sup>11,20</sup>

Only 1 case report quantitatively mentioned dialysis adequacy using Kt/v.<sup>21</sup> However, all other studies reported that dialysis adequacy for these patients was within standard recommendations until the end of follow-up, death, or kidney transplantation, except for one. This patient, with ultrafiltration failure, was transferred to HD 16 months after PD initiation.<sup>20</sup> Overall, although LT is considered as major abdominal surgery, there is no evidence in this review showing any significant peritoneal membrane malfunction, lower PD efficiency, or limitation in the installation of a functioning catheter. However, due to the limited sample size, this review does not allow to definitely address this question.

On the contrary, the short median follow-up in these studies could have underestimated some late complications of PD such as sclerosing encapsulating peritonitis, for which both PD and LT patients are at increased risk.<sup>34,35</sup>

This review has several limitations. As the topic of PD after solid-organ transplantation is rarely described in the literature, a risk of publication bias exists, where it is possible that only positive experiences were published. In addition, the observational design of all included studies, as well as the small number of patients reported, limit the level of evidence resulting from this review. The level of details in the description of all cases reported varied across included studies. Finally, as no study directly compared patients treated with PD to similar patients treated with HD, we were unable to evaluate the impact of the dialysis modality itself in LT recipients requiring KRT.

## Conclusion

With the caveat of limited data availability, our systematic review of the literature shows that PD is feasible in LT recipients who require KRT, where the risk of peritonitis reported was similar than that reported in comparable PD populations receiving immunosuppression. These episodes of peritonitis had no significant impact on clinical evolution, technique survival, and liver graft function. Due to observational data mostly coming from limited case series, no definitive recommendation regarding this modality can be formulated. However, despite the possibility of publication bias, we found no signal suggesting that undertaking PD following

LT is putting the patient at an increased risk of infectious or technical complications. Large and long-term studies evaluating PD in this population need to be performed. Further collaborative multicenter studies may help to address this important question.

## Acknowledgments

The authors would like to thank Daniela Ziegler, biomedical librarian, Center hospitalier de l'Université de Montréal, for her assistance with the systematic literature search.

## Ethics Approval and Consent to Participate

Not applicable for this type of study.

## Consent for Publication

All authors consent to the publication of this study.

## Availability of Data and Materials

Data are available on request.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: No specific funding has been received for this study. J.M.C. was supported by the *Fonds de Recherche en Santé du Québec* and *Société Québécoise de Néphrologie*. J.M.C. and I.E. were supported by a Fellowship grant from the *Center Hospitalier de l'Université de Montréal* and the *Fondation du CHUM*. H.C. is a *Fonds de Recherche en Santé du Québec* Junior 2 Scholar.

## ORCID iD

Jean Maxime Côté  <https://orcid.org/0000-0002-3487-2670>

## Supplemental Material

Supplemental material for this article is available online.

## References

1. Ojo AO, Held PJ, Port FK, et al. Chronic renal failure after transplantation of a nonrenal organ. *N Engl J Med*. 2003;349:931-940.
2. Bloom RD, Reese PP. Chronic kidney disease after nonrenal solid-organ transplantation. *J Am Soc Nephrol*. 2007;18:3031-3041.
3. O'Riordan A, Wong V, McCormick PA, Hegarty JE, Watson AJ. Chronic kidney disease post-liver transplantation. *Nephrol Dial Transplant*. 2006;21:2630-2636.
4. Sharma P, Welch K, Eikstadt R, Marrero JA, Fontana RJ, Lok AS. Renal outcomes after liver transplantation in the model for end-stage liver disease era. *Liver Transpl*. 2009;15(9):1142-1148.
5. Yunhua T, Qiang Z, Lipeng J, et al. Liver transplant recipients with end-stage renal disease largely benefit from kidney transplantation. *Transplant Proc*. 2018;50(1):202-210.



6. Goldberg DS, Ruebner RL, Abt PL. The risk of end-stage renal disease among living donor liver transplant recipients in the United States. *Am J Transplant*. 2015;15(10):2732-2738.
7. Srinivas TR, Stephany BR, Budev M, et al. An emerging population: kidney transplant candidates who are placed on the waiting list after liver, heart, and lung transplantation. *Clin J Am Soc Nephrol*. 2010;5(10):1881-1886.
8. Al Riyami D, Alam A, Badovinac K, Ivis F, Trpeski L, Cantarovich M. Decreased survival in liver transplant patients requiring chronic dialysis: a Canadian experience. *Transplantation*. 2008;85:1277-1280.
9. Gonwa TA, Mai ML, Melton LB, Hays SR, Goldstein RM, Levy MF, Klintmalm GB. End-stage renal disease (ESRD) after orthotopic liver transplantation (OLT) using calcineurin-based immunotherapy: risk of development and treatment. *Transplantation*. 2001;72:1934-1939.
10. Wong B, Ravani P, Oliver MJ, et al. Comparison of patient survival between hemodialysis and peritoneal dialysis among patients eligible for both modalities. *Am J Kidney Dis*. 2018;71(3):344-351.
11. Perl J, Bargman JM, Jassal SV. Peritoneal dialysis after nonrenal solid organ transplantation: clinical outcomes and practical considerations. *Perit Dial Int*. 2010;30(1):7-12.
12. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
13. Munn Z, Barker TH, Moola S, et al. Methodological quality of case series studies: an introduction to the JBI critical appraisal tool. *JBI Evid Synth*. 2020;18(10):2127-2133.
14. Akhtar I, Bastani B. Role of albumin peritoneal dialysis for bilirubin removal after complicated liver transplant. *Journal of Applied Research*. 2006;6:153-156.
15. Apiratpracha W, Chung HV, Ali JA, Bashir K, Yoshida EM, Singh RS. Peritoneal dialysis after orthotopic liver transplant surgery. *Perit Dial Int*. 2007;27(1):99-100.
16. Buffet A, Guillouet S, Lobbedez T, Ficheux M., Lanot A., Béchade C. Safety of peritoneal dialysis after nonrenal solid-organ transplantation. *Perit Dial Int*. 2018;38(1):37-43.
17. Cornelis T, Rioux JP, Bargman JM, Chan CT. Home dialysis is a successful strategy in nonrenal solid organ transplant recipients with end-stage renal disease. *Nephrol Dial Transplant*. 2010;25(10):3425-3429.
18. Davis O, Wall BM. "Broom Straw Peritonitis" secondary to *Cedecia lapagei* in a liver transplant recipient. *Perit Dial Int*. 2006;26:512-513.
19. Faria B, Rodrigues A. Peritoneal dialysis in transplant recipient patients: Outcomes and management. *Scand J Urol Nephrol*. 2011;45(6):444-451.
20. Saiprasertkit N, Nihei CH, Bargman JM. Peritoneal dialysis in orthotopic liver transplantation recipients. *Perit Dial Int*. 2018;38(1):44-48.
21. Shepherd LA, Holley JL. Successful peritoneal dialysis after intra-abdominal transplantation. *Perit Dial Int*. 2004;24(1):90.
22. Shukla AM, Lobbedez T, Chu M, Izatt S, Bargman JM, Oreopoulos DG. Peritoneal dialysis in end-stage renal disease after liver transplantation. *Adv Perit Dial*. 2004;20:93-97.
23. Wilkie M. Peritoneal dialysis (PD) to support nonrenal solid-organ transplantation. *Perit Dial Int*. 2018;38(1):3.
24. Lee JP, Heo NJ, Joo KW, et al. Risk factors for consequent kidney impairment and differential impact of liver transplantation on renal function. *Nephrol Dial Transplant*. 2010;25(8):2772-2785.
25. Francois K, Bargman JM. Evaluating the benefits of home-based peritoneal dialysis. *Int J Nephrol Renovasc Dis*. 2014;7:447-455.
26. Blake PG. Integrated end-stage renal disease care: the role of peritoneal dialysis. *Nephrol Dial Transplant*. 2001;16(Suppl. 5):61-66.
27. Jansen MA, Hart AA, Korevaar JC, et al. Predictors of the rate of decline of residual renal function in incident dialysis patients. *Kidney Int*. 2002;62(3):1046-1053.
28. Perl J, Fuller DS, Bieber BA, et al. Peritoneal dialysis-related infection rates and outcomes: results from the Peritoneal Dialysis Outcomes and Practice Patterns Study (PDOPPS). *Am J Kidney Dis*. 2020;76(1):42-53.
29. Shaked A, DesMarais MR, Kopetskie H, et al. Outcomes of immunosuppression minimization and withdrawal early after liver transplantation. *Am J Transplant*. 2019;19(5):1397-1409.
30. Haidar G, Green M, American Society of Transplantation Infectious Diseases Community of Practice. Intra-abdominal infections in solid organ transplant recipients: Guidelines from the American Society of Transplantation Infectious Diseases Community of Practice. *Clin Transplant*. 2019;33(9):e13595.
31. Badve SV, Hawley CM, McDonald SP, et al. Effect of previously failed kidney transplantation on peritoneal dialysis outcomes in the Australian and New Zealand patient populations. *Nephrol Dial Transplant*. 2006;21(3):776-783.
32. Guru P, Prakash R, Sheth H, Bender F, Burr R, Piraino B. Comparison of survival of patients with heart and lung transplants on peritoneal dialysis and hemodialysis. *Perit Dial Int*. 2015;35(1):98-101.
33. Siu YP, Leung KT, Tong MK, Kwan TH, Mok CC. Clinical outcomes of systemic lupus erythematosus patients undergoing continuous ambulatory peritoneal dialysis. *Nephrol Dial Transplant*. 2005;20(12):2797-2802.
34. Merkle M, Wörnle M. Sclerosing peritonitis: a rare but fatal complication of peritoneal inflammation. *Mediators Inflamm*. 2012;2012:709673.
35. Mekeel K, Moss A, Reddy KS, Douglas D, Mulligan D. Sclerosing peritonitis and mortality after liver transplantation. *Liver Transpl*. 2009;15(4):435-439.