

# Rootstock–scion combination contributes to shape diversity and composition of microbial communities associated with grapevine root system

Ramona Marasco<sup>1</sup>,<sup>1\*</sup> Hend Alturkey,<sup>1</sup> Marco Fusi,<sup>1</sup>  
Michele Brandi,<sup>2</sup> Isabella Ghiglieno,<sup>3,4</sup>  
Leonardo Valenti<sup>3</sup> and Daniele Daffonchio<sup>1</sup><sup>1\*</sup>

<sup>1</sup>Biological and Environmental Sciences and Engineering Division (BESE), King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi Arabia.

<sup>2</sup>Marchesi Frescobaldi Società Agricola s.p.a. Fattoria Poggio a Remole, Sieci, Italy.

<sup>3</sup>Department of Agricultural and Environmental Sciences – Production, Landscape, Agroenergy, University of Milan, Milan, Italy.

<sup>4</sup>Department of Civil, Environmental, Architectural Engineering and Mathematics (DICATAM), University of Brescia, Agrofood Research Hub, Brescia, Italy.

## Summary

To alleviate biotic and abiotic stresses and enhance fruit yield, many crops are cultivated in the form of grafted plants, in which the shoot (scion) and root (rootstock) systems of different species are joined together. Because (i) the plant species determines the microbial recruitment from the soil to the root and (ii) both scion and rootstock impact the physiology, morphology and biochemistry of the grafted plant, it can be expected that their different combinations should affect the recruitment and assembly of plant microbiome. To test our hypothesis, we investigated at a field scale the bacterial and fungal communities associated with the root system of seven grapevine rootstock–scion combinations cultivated across 10 different vineyards. Following the soil type, which resulted in the main determinant of the grapevine root microbial community diversity, the rootstock–scion combination resulted more important than the two components taken alone. Notably, the microbiome differences among the rootstock–

scion combinations were mainly dictated by the changes in the relative abundance of microbiome members rather than by their presence/absence. These results reveal that the microbiome of grafted grapevine root systems is largely influenced by the combination of rootstock and scion, which affects the microbial diversity uptaken from soil.

## Introduction

The bacterial and fungal microbiota are essential component of the plant holobiont (Sánchez-Cañizares *et al.*, 2017; Hassani *et al.*, 2018; Ceja-Navarro *et al.*, 2021). These microorganisms add an external second metagenome to those of the plants, supplying key ecological functions and services (Zilber-Rosenberg and Rosenberg, 2008; Turner *et al.*, 2013; Vandenkoornhuysen *et al.*, 2015; Rosenberg *et al.*, 2016), including promotion of plant growth, fitness, health and adaptation to environmental challenges (Berendsen *et al.*, 2012; Rolli *et al.*, 2017; Mommer *et al.*, 2018; Pugnaire *et al.*, 2019). Understanding the dynamics and factors driving the recruitment and assembly of the plant microbiome is becoming more and more important to translate fundamental ecological knowledge into effective manipulative strategies aimed at the engineering of plant microbiomes (Lakshmanan *et al.*, 2014; Mueller and Sachs, 2015; Dini-Andreote and Raaijmakers, 2018; Toju *et al.*, 2018; Singh *et al.*, 2020).

Plants actively recruit and select microorganisms from the surrounding soil by means of the rhizosphere (*i.e.* the transition zone between root and soil influenced by root exudates) and a subset of such microorganisms could further cross the rhizoplane barrier to become endophytes (Van Der Heijden and Schlaeppi, 2015; Santoyo *et al.*, 2016; Zhainina *et al.*, 2018). Most studies have been conducted on conventional plant species (cultivated and wild), while little attention has been directed at ‘chimeric plants’; *i.e.* individual plants made up of two or multiple species. Examples of these chimaeras are given by grafting, a mechanism of plant–plant interaction that is widespread in nature, in which parts of two different

Received 15 February, 2022; accepted 6 May, 2022. \*For correspondence. E-mail [ramona.marasco@kaust.edu.sa](mailto:ramona.marasco@kaust.edu.sa); [daniele.daffonchio@kaust.edu.sa](mailto:daniele.daffonchio@kaust.edu.sa). Tel: +966 (012) 8082455.

plants are physically joined to create a new individual (Fuentes *et al.*, 2014; Warschefsky *et al.*, 2016; Gaion *et al.*, 2018). Typically, a grafted plant has two genetically different components: the root system of one plant species (rootstock) and the shoot (scion) from another plant cultivar or even species (Fuentes *et al.*, 2014). Grafting is widely used with trees and shrubs, but recently it was also applied in vegetables (Gaion *et al.*, 2018), to combine the rusticity and resistance of the rootstock, and the quality and productivity of the scion (Warschefsky *et al.*, 2016; Riaz *et al.*, 2019). Today, most flowering and fruit trees (e.g. apple, avocado, citrus fruit, grapevine, peach, and rose) and horticulture plants (e.g. up to 40% of tomatoes and eggplants, 75% of cucumbers, 5%–10% of peppers and 90% of watermelons) are grafted onto rootstocks (Koevoets *et al.*, 2016; Warschefsky *et al.*, 2016; Gaion *et al.*, 2018). Of these, grapevine is an iconic and the most widespread grafted crop, covering up to 7.5 million hectares worldwide (OIV The International Organisation of Vine and Wine, 2020). Since the 19th century, in Europe *Vitis vinifera* (the European grape) cultivars have been grown as scions grafted onto rootstocks of other species of the genus *Vitis* (*V. rupestris*, *V. berlandieri* and *V. riparia*, and their derived hybrids) to prevent vineyard failure due to the aphid-like hemipteran *Phylloxera* (Battey and Simmonds, 2005; Riaz *et al.*, 2019). To date, grafting on rootstocks is used as a general practice, because the rootstock can affect several physiological parameters of the plant and make the grapevine growing in otherwise unsuitable soils. Thus, rootstock selection and development are an important component of modern viticulture. Rootstocks have been initially developed for improving plant resistance to certain soil-borne pests and diseases (Wallis *et al.*, 2013; Warschefsky *et al.*, 2016), and for improving plant resistance to adverse climatic or soil conditions (Bert *et al.*, 2013; Henderson *et al.*, 2014; Berdeja *et al.*, 2015; Hamrouni *et al.*, 2015; Habran *et al.*, 2016). Grafting also influences the reproductive performance, scion vigour, biomass accumulation and distribution, grape yield and quality, and phenology (Whiting, 2003; Wallis *et al.*, 2013; Bonghi *et al.*, 2016; Warschefsky *et al.*, 2016). In grafting, plant vigour and productivity depend on the efficient integration mechanisms between the rootstock and the scion, including wound healing, tissue fusion, vascular reconnection and shoot–root signalling regulation (Pina *et al.*, 2017; Venema *et al.*, 2017; Melnyk *et al.*, 2018; Gautier *et al.*, 2019). The grafting process unequivocally results in a two-sided influence on the phenotype (physiological/morphological traits) of both components; *i.e.* the rootstock affects the scion and vice versa (Martínez-Ballesta *et al.*, 2010; Hamrouni *et al.*, 2015; Habran *et al.*, 2016; Warschefsky *et al.*, 2016; Ruan *et al.*, 2020). However, to date,

research has mainly been focused on how the rootstock affects the scion phenotype, with little attention paid to the reverse effect of scion on the rootstock phenotype and on the mechanisms of scion–rootstock interactions (Gautier *et al.*, 2019).

Based on the bilateral influence of rootstock and scion, it is plausible to suppose that both and their interaction may also affect the association of microorganisms with the plant, including their recruitment, selection and assembly. Studies of spatial and temporal variations in microbial communities associated with grapevine, including the above- and below-ground parts of the plant, have revealed a myriad of factors that can influence the dynamics of grapevine microbiome assembly (Bettenfeld *et al.*, 2021; Griggs *et al.*, 2021). Soil heterogeneity and vineyard agricultural practices represent the main factors that affect the diversity of taxa recruited by the grapevine in the root system (Marasco *et al.*, 2013; Campisano *et al.*, 2014; Vega-Avila *et al.*, 2015; Zarraonaindia *et al.*, 2015; Burns *et al.*, 2016; Manici *et al.*, 2017; Berlanas *et al.*, 2019; Vitulo *et al.*, 2019; Gamalero *et al.*, 2020; Swift *et al.*, 2021; Vink *et al.*, 2021a). Furthermore, a recent body of literature has demonstrated that the rootstock genotype strongly influences the diversity of bacterial and fungal communities recruited by the root from the surrounding soil (Wallis *et al.*, 2013; D'Amico *et al.*, 2018; Marasco *et al.*, 2018b; Berlanas *et al.*, 2019; Nerva *et al.*, 2021; Swift *et al.*, 2021; Dries *et al.*, 2021a; Darriaut *et al.*, 2022). However, the effect mediated by the interaction between different rootstocks and scions (*i.e.* grafting combinations) on the grapevine microbiome has been poorly studied. To date, only one study has explored this in grapevine, using grafted plants obtained from the interaction between four rootstocks and four cultivar scions (Vink *et al.*, 2021b). This study was limited to the bacterial community associated with the rhizosphere and was performed in a single soil in an experimental field. Given that the current literature has neglected the overall complexity of an extended-environmental context at ecosystem scale, in our study, we hypothesized that (i) the effect of genotype observed for the rootstock microbiome recruitment at the levels of root endosphere and rhizosphere for both bacteria and fungi is further tuned if the latest is combined with different scion cultivars, and (ii) it can be consistently detectable across heterogeneous environmental settings in a real agricultural ecosystem context. To test these hypotheses, we selected a large area devoted to grapevine cultivation, in which differential environmental and pedoclimatic conditions occur. We selected seven different rootstock–scion combinations cultivated in 10 vineyards located in two geographical areas in Tuscany, namely, Pomino and Nipozzano. These combinations are formed by four rootstocks (SO4, 420A, 110R and 3309C) and

three scions (Chardonnay, Sangiovese and Merlot). The bacterial 16S rRNA gene and fungal ITS from the root system (root and rhizosphere) and bulk soil sampled were amplified and sequenced to evaluate the effects of the rootstock, scion, and their combination, along with those of soil and geographical area, on the structure and diversity of the grapevine root system microbiome.

With this work we aim to demonstrate that the scion–rootstock combination factor is an important parameter to be considered in ecological studies of grafted plants. One requirement of the modern research in plant-associated microbiomes is to extend investigations from a confined range, e.g. microcosms or single fields (Vink *et al.*, 2021b), to wide ecosystem scales to unravel the selective forces that influence the microbiome associated with plant across the heterogeneous conditions of real-world ecosystems (Delgado-Baquerizo, 2022; Gobbi *et al.*, 2022). Ecosystem-scale studies can offer important contributions that can set the basis of future framework for targeted agriculture (Trivedi *et al.*, 2021) and microbial synthetic communities applications (Delgado-Baquerizo, 2022) to support productivity and sustainability. It is important to note that the use of soil microorganisms to promote plant productivity is a promising tool also in viticulture (Rolli *et al.*, 2017; Bettenfeld *et al.*, 2021; Dries *et al.*, 2021b; Darriaut *et al.*, 2022). This study can provide insights on the effect of different rootstock–scion combinations on the microbiome assembly and recruitment that should be taken into account during the development and use of targeted-agriculture applications (e.g. biofertilizers and bio-promoters), especially in the light of the predicted changes that viticulture will face soon (Morales-Castilla *et al.*, 2020).

## Experimental procedures

### Sample collection


The root system (root tissue and rhizosphere soil) of 420A, SO4, 110R and 3309C rootstocks, which are originated from the breeding of the most used *Vitis riparia*, *V. berlandieri* and *V. rupestris* (Supplementary Table S1), was collected at the end of October 2016 in 10 vineyards across the Nipozzano and Pomino areas owned by Marchesi Frescobaldi Società Agricola s.p.a. (north-eastern Tuscany, Italy; Supplementary Fig. S1; Supplementary Table S2). Onto the selected rootstocks were grafted the scions of *V. vinifera* cultivars Chardonnay, Merlot and Sangiovese, for a total of seven rootstock–scion combinations (Supplementary Table S2). Each of the vineyard studied here has a unique combination of environmental conditions, strongly dictated by the soil physicochemical characteristics and geoclimatic factors (Fig. 1; Supplementary Table S3), and for this reason different rootstocks were distributed across them (Ollat

*et al.*, 2016; Bianchi *et al.*, 2020). For instance, the SO4 rootstock (selection oppenheim, *V. berlandieri* × *V. riparia*) is one of the most used in Italy because it offers great adaptability to edaphic and pedo-climatic conditions, as well as good productivity. Rootstocks that push plant vigour, such as 110 Richter (110R, *V. berlandieri* × *V. rupestris*) are preferable in tough and dry hillside slopes, and to cope with warm climates and drought events (Bianchi *et al.*, 2020), often experience in Italy in the last decades (Ciais *et al.*, 2005). 420A rootstock (420A, *V. berlandieri* × *V. riparia*) confers low vigour to plants, limiting the yields, particularly during the first years of production, but it covers a wide range of soil type (*i.e.* deep, fine texture and alkaline soils), representing a possible compromise for vineyards with heterogeneous soils, such as the case of Pomino and Nipozzano estates. On the contrary, the 3309 Couderc rootstock (3309C, *V. riparia* × *V. rupestris*), with low to moderate tolerance to chlorosis and active limestone, and sensitivity to water stress, is not widespread in the Frescobaldi vineyards.

A total of six plants were selected in each vineyard; in the case of Rena vineyard, we have sampled two different combinations in the same field (Supplementary Table S2). The root samples ( $n = 66$ ) were collected at a depth of 20–40 cm; this range of depth allowed us to consistently collect the active and denser portion of root system across the different vineyards and for the different rootstocks. Six samples of bulk soil (*i.e.* soil not influenced by the root system) from each location were also collected. All samples were collected under sterile conditions using sterile tools. The recovered samples were stored at  $-20^{\circ}\text{C}$  for molecular analysis. The sampling was authorized by the vineyard owner at each location and no other specific permissions were required for this activity.

### Physico-chemical analysis of vineyard soils

The chemical and physical properties of the bulk soil from each vineyard were provided by the vineyard owner. Soil samples have been collected, pooled and homogenized before analysing soil texture and physicochemical parameters, including pH, cation exchange capacity (CEC), total and active limestone content ( $\text{CaCO}_3$ ), organic matter, total carbon, total nitrogen, C/N ratio and available elements/nutrients (calcium, magnesium, potassium and phosphorus), base saturation and overall acidity. The results of physicochemical analyses were analysed by performing principal coordinate analysis (PCoA) and permutational multivariate analysis of variance (PERMANOVA) on the Euclidean distance matrix with Primer v.6.1 (Anderson *et al.*, 2008); sample categories (two levels: Pomino and Nipozzano) were used as explanatory variables.



**Fig. 1.** A and B. Beta-diversity of microbial communities associated with grapevine rootstock root systems compartments and bulk soils in Tuscan vineyards. Principal coordinate analysis (PCoA) of (A) bacterial and (B) fungal communities based on Bray–Curtis similarity matrices; colours indicate the compartments (root tissues, rhizosphere and bulk soil).

C and D. Variation of dispersion (within-beta-diversity) as distance of each sample from the centroid of each compartment group is reported for bacteria and fungi respectively; ANOVAs are performed for the factor compartment; bacteria:  $F_{2,185} = 586.8$ ,  $p < 0.0001$  and fungi:  $F_{2,187} = 59.51$ ,  $p < 0.0001$ .

E–H. Alpha-diversity of the bacterial and fungal communities. E and F. OTUs richness (number of observed OTUs) and (G and H) Shannon index are reported for bacterial and fungal communities respectively, across root tissues (RT), rhizosphere (RH) and bulk soil (BS). ANOVAs are performed for the factor compartment; bacteria<sub>richness</sub>:  $F_{2,185} = 526.4$ ,  $p < 0.0001$  and bacteria<sub>Shannon</sub>:  $F_{2,185} = 237.0$ ,  $p < 0.0001$ ; fungi<sub>richness</sub>:  $F_{2,187} = 238.0$ ,  $p < 0.0001$  and fungi<sub>Shannon</sub>:  $F_{2,187} = 47.18$ ,  $p < 0.0001$ . C–H. Lowercase letters indicate the results of the Tukey's multiple comparison tests among the compartments (significance,  $p < 0.05$ ) for dispersion and alpha-diversity.

#### Total DNA extraction

The sampled roots with attached soil particles were placed in a sterile tube and 9 ml of physiological solution ( $9 \text{ g L}^{-1}$  NaCl) was added. The tubes were vortexed for 5 min to detach tightly attached soil particles, then centrifuged at 4000 rpm for 10 min. The roots were removed, and the supernatant was discarded to collect the remaining soil and use it as rhizosphere soil. The soil samples (rhizosphere and bulk soil) were homogenized with sterile pestles and mortars, and  $0.5 \pm 0.1 \text{ g}$  was further used to extract total DNA using the PowerSoil DNA Kit (Qiagen), following the manufacturer's instructions. In the case of the root tissues, the material obtained after the removal of the rhizospheric soil was further surface sterilized by soaking in 70% ethanol for 3 min, followed by sodium hypochlorite 2.5% for 5 min, 70% ethanol for 30 s, and finally by washing with sterile distilled water five times; the

efficacy of the sterilization method was verified by plating pieces of the sterilized root and the water from the last washing step on plates with tryptone soy agar. The plates were examined for bacterial growth after incubation at  $30^\circ\text{C}$  for 3 days. The root tissues were smashed with liquid nitrogen and further homogenized before performing genomic DNA extraction ( $1 \pm 0.5 \text{ g}$ ) using the DNeasy Plant Maxi Kit (Qiagen). Eluted DNA ( $5 \mu\text{l}$ ) was visualized on 0.8% agarose gel using electrophoresis to evaluate its quality, while quantification of DNA was performed using a Qubit 3.0 Fluorometer and dsDNA BR Assay kit. All extracted DNA samples were stored at  $-20^\circ\text{C}$ .

#### Amplification, sequencing and analysis

Illumina tag screening of the V3–V4 hypervariable regions of the bacterial 16S rRNA gene was applied on

DNA, using the primers 341f and 785r (Klindworth *et al.*, 2013) and following the protocol previously described (Mapelli *et al.*, 2018). Briefly, PCR reactions mixture of 30  $\mu$ l was performed for each sample using 1 U of Platinum<sup>®</sup> Taq DNA Polymerase, High Fidelity (Invitrogen) with 1 $\times$  High Fidelity Buffer, 1.5 mM of MgSO<sub>4</sub>, 0.3 mM dNTPs mix, 0.3  $\mu$ M each of forward and reverse primers, and ca. 10 ng of template DNA, along with the addition of 0.5  $\mu$ M each of pPNA and mPNA clamps to reduce the amplification of host chloroplasts and mitochondria (Fitzpatrick *et al.*, 2018; Deyett and Rolshausen, 2020). In the case of fungi, the fungal ITS2 gene was amplified using ITS3F and ITS4R primers (Marasco *et al.*, 2018a). The sequencing libraries were constructed using a 96 Nextera<sup>®</sup> XT Index kit (Illumina) and the products of amplification were cleaned using a SequalPrep<sup>™</sup> Normalization Plate kit. All libraries were sequenced on the Illumina MiSeq platform at the Bioscience Core Lab, King Abdullah University of Science and Technology. The raw sequencing data for bacterial and fungal communities were submitted to the National Center of Biotechnology Information (NCBI) Sequence Read Archive under accession number PRJNA807110.

Analysis of sequences was performed using a combination of the QIIME pipeline version 1.9 (Caporaso *et al.*, 2010) and UPARSE version 8 (Edgar, 2013). After paired-end merging and quality filtering, only high-quality reads were kept for further steps in the analysis. Reads were clustered as OTUs considering a 97% sequence distance similarity and taxonomy was assigned using UCLUST against SILVA 138 and UNITE databases for bacteria and fungi respectively. All OTUs non-assigned to bacteria [mitochondria (24 626 reads, 31 OTUs), chloroplasts (24 465 reads, 63 OTUs), archaea (740 reads, 11 OTUs) and unclassified (95 530 reads, 12 951 OTUs)] and fungi [non-fungi (90 833 reads, 326 OTUs) and unclassified (1 178 884 reads, 942 OTUs)] were removed. OTUs were filtered to keep only those with a relative abundance higher than 0.001% and the rarefaction curves were generated (Supplementary Fig. S2). By using *qiime feature-table rarefy* plug-in without replacement (Weiss *et al.*, 2017), the rarefaction threshold was set at 9590 and 13 400 reads per sample for bacteria and fungi respectively (Supplementary Table S3), after the removal of samples with low sampling depth (<5000 reads). In PRIMER v.6.1 (Anderson *et al.*, 2008), Bray–Curtis distance matrices were used to perform PCoA, permutational analysis of multivariate dispersion and canonical analysis of principal coordinates (CAP). We evaluated the most important factor accounting for the beta-diversity variability between microbial communities by using the function *best.rsq()* in the R package *mvabund* (Wang *et al.*, 2012). Since the effect of ‘compartment’ was

ranked as the most important factor (*i.e.* niche partitioning; Supplementary Table S4), we performed further analyses considering separately the root tissue, the rhizosphere and the bulk soil. For each compartment, we evaluated and quantified the effect of estate, field and rootstock–scion interaction factors by using a model-based approach for compositional data performed with the *anova.manyglm()* and the *best.rsq()* functions in the R package *mvabund* (Wang *et al.*, 2012). Our categorical explanatory variables were ‘estate’ (two levels), ‘field’ (10 levels), ‘rootstock’ (four), ‘scion’ (three) and their interaction ‘rootstock  $\times$  scion’ (seven levels). To disentangle the effect of the rootstock–scion interaction we conducted CAP, PERMANOVA and pairwise analysis using PRIMER v.6.1 (Anderson *et al.*, 2008). Shared (*i.e.* core microbiome) and specific microbial OTUs across the rootstock–scion combinations within each compartment (root tissue, rhizosphere and bulk soil) were defined using Venn diagram analysis with R. We further calculated the individual contribution of each OTUs to define the diversity of bacterial and fungal communities across rootstock–scion combinations by testing univariate *glm* for each OTU including the parameter *p.uni* = ‘adjusted’ in the function *anova.manyglm()* (Wang *et al.*, 2012). To pinpoint those bacterial and fungal OTUs that were enriched within each sample type, an indicator species analysis combining both the abundance and occurrence of a given OTU across all rootstock–scion combinations was used. Indicator species were calculated by comparing combinations’ communities within each compartment, using the *multipatt* function in the R statistical package *IndicSpecies* (De Cáceres *et al.*, 2020). This function creates combinations of the input clusters and compares each combination with the species in the input matrix. For each species it chooses the combination with a highest association value. Best matching patterns are tested for statistical significance of the associations. The association indices used was ‘*IndVal.g*’ that return the pattern that better matches the species observed pattern. The significance of each OTU–combination association was tested using a permutation test ( $n = 9999$ ) and non-significant associations ( $p > 0.05$ ) were discarded. Richness and Shannon diversity were calculated in R using the *Phyloseq* package (McMurdie and Holmes, 2013). We used ANOVA with Tukey’s multiple comparison tests to determine statistical significance across compartments for richness and Shannon diversity in bacteria and fungi and we used the machine learning algorithm Random Forest to rank the most important predictors of alpha diversity patterns by taking into account the mean decrease accuracy (%IncMSE) that indicates the increase of the mean squared error when a given variable is randomly permuted (Delgado-Baquerizo and Eldridge, 2019).

## Results

### *Geoclimatic and edaphic physicochemical characterization of vineyards*

Pomino and Nipozzano are estates belonging to the Frescobaldi company located in the north-eastern area of Tuscany (Italy), close to the Apennines (Supplementary Fig. S1). The two estates are characterized by different geology (PERMANOVA soil textures,  $F_{1,8} = 5.99$ ;  $p = 0.023$ ; Supplementary Fig. S3A; Supplementary Table S5); whereas in the Nipozzano estate the heterogeneity of the soil texture led to distinct vineyard clusters, in the Pomino estate the dominance of sand (range, 57%–75%) favoured the establishment of similar edaphic conditions across the vineyards (Supplementary Fig. S3B). The physicochemical characteristics of the vineyard soils (Supplementary Table S5) confirmed the differences between the two estates ( $F_{1,8} = 3.03$ ;  $p = 0.013$ ; Supplementary Fig. S3C). For instance, while all the vineyard soils were around neutral-slightly alkaline, with pH from 6.5 to 8 at Pomino and from 7.7 to 8.1 at Nipozzano, organic carbon, limestone, exchangeable ions and CEC determined the soil diversity (Supplementary Table S5; sequential test and AIC model in Supplementary Table S6). Additional details on the ecosystem and agronomic practices applied in the two estates are also provided in Supplementary Result S1. Both estates use the 110R, 3309C, SO4 and 420A rootstocks obtained by crossing of different *Vitis* species (*V. berlandieri*, *V. riparia* and *V. rupestris*). These rootstocks are grafted with Chardonnay, Merlot and Sangiovese, while ungrafted plants are not present in the Frescobaldi estates.

### *Compartmentalization of bacterial and fungal communities associated with grapevine root system*

The bacterial and fungal diversity associated with the root system (root tissue and rhizosphere) of seven rootstock–scion combinations obtained from four rootstocks and three scions, and related bulk soils not influenced by grapevine root systems have been measured (Supplementary Tables S1 and S2). PCoA revealed strong clustering of the microbial communities according to the different compartments sampled (root tissue, rhizosphere and bulk soil) for both bacteria and fungi (manyglm, bacteria:  $Dev_{2,185} = 206\,634$ ,  $p = 0.001$  and fungi:  $Dev_{2,187} = 29\,238$ ,  $p = 0.001$ ; Fig. 1A and B; Supplementary Table S4). The microbial communities of the compartments were mainly differentiated along the first axis of PCoA (Fig. 1A and B) with cross-validation of 100% for the samples within each compartment (CAP). A selection determined by the plant and a rhizosphere effect were consistently observed in bacteria and fungi:

we detected a horseshoe-shaped distribution in the ordination space with the three compartments succeeding each other (Fig. 1A and B). This pattern indicates the presence of a dual step process of recruitment and selection from the soil to the root tissues (Van Der Heijden and Schlaeppi, 2015). Notably, within each compartment two distinct sub-clusters based on the geographical/environmental origin of the samples were detected (Pomino vs. Nipozzano; Supplementary Fig. S4).

Different levels of dispersion (within-beta diversity) were observed along the root system compartments (Fig. 1C and D). The highest dispersion values were detected for root microbial communities, indicating that endophytes had the largest intrinsic variability, possibly due to their limited diversity (low richness, Fig. 1E and F) and uneven distribution (low Shannon index, Fig. 1G and H). In contrast, the rhizosphere and bulk soil, which harboured more diverse and more even microbial communities (Fig. 1E–H), had less dispersion (Fig. 1C and D). It is important to note that the compartment was also identified as the most important factor determining the differences observed in the alpha-diversity indices (Supplementary Table S7).

The niche partitioning described in terms of beta-diversity and alpha-diversity was confirmed by the differential composition of the main bacterial and fungal classes across the three compartments (Supplementary Fig. S5). Bacterial communities were dominated (on average) by *Proteobacteria* (37.5%: 21% *Alphaproteobacteria* and 16.5% *Gammaproteobacteria*), *Actinobacteria* (16%), *Bacilli* (8%), *Bacteroidia* (7%), *Blastocatellia* (6%) and *Clostridia* (3%), followed by *Vicinamibacteria*, *Thermoleophilia*, *Verrucomicrobiae*, *Polyangia*, *Gemmatimonadetes*, *Acidobacteriae*, *Acidimicrobiia*, *Anaerolineae* and the remaining classes with low relative abundance (<1%). Most bacterial classes were differentially distributed across the compartments, with *Actinobacteria* and *Clostridia* enriched in root tissues, *Gammaproteobacteria* in the rhizosphere, and *Blastocatellia*, *Gemmatimonadetes*, *Acidobacteriae*, *Verrucomicrobiae* and *Anaerolineae* in bulk soil (Supplementary Fig. S6). In the case of fungal communities, *Ascomycota* was the dominant phylum mainly represented by *Sordariomycetes* (40%), *Dothideomycetes* (17%), *Eurotiomycetes* (8%), *Leotiomycetes* (5%) and *Pezizomycetes* (3%), and it was followed by *Basidiomycota* (7% *Agaricomycetes* and 5% *Tremellomycetes*), *Mortierellomycetes* (8%) and *Mucoromycetes* (1%). The remaining part was constituted by rare fungal taxa (relative abundance <1%) and unclassified fungi (2% and 4% respectively). All of them were differentially distributed across compartments except for *Agaricomycetes*, *Leotiomycetes* and the dominant *Sordariomycetes*. Members of *Dothideomycetes* were enriched in the root tissues, while


*Eurotiomycetes* and *Mucoromycetes* were enriched in the rhizosphere, and *Pezizomycetes* in the bulk soil (Supplementary Fig. S7). The list of bacterial and fungal OTUs that most contributed to determine the differences across compartments is reported in Supplementary Table S8.

#### Effect of biotic and abiotic factors on beta-diversity and alpha-diversity of microbial communities associated with the root system of grafted grapevine and bulk soil

It is well known that the micro-ecosystem of each vineyard specifically influences the diversity and composition of edaphic microorganisms available for plant recruitment (Gobbi *et al.*, 2022). In our study the 'field' factor was confirmed to be the most important abiotic factor shaping the diversity of microbial communities associated with root system compartments (root and rhizosphere) and bulk soil (bacteria, Fig. 2A–C; fungi, Fig. 2D–F; manyglm in Table 1). The interaction between rootstock and scion was the second factor for importance in shaping the root system microbial diversity (Fig. 2; Table 1). It was consistently ranked as the first and most important plant-related factor, explaining a percentage of observed variability higher than that explained by the rootstock or the scion

considered alone (Fig. 2). Furthermore, when testing the best combination of factors for explaining microbial variability, the rootstock–scion interaction was always included, together with the field, as a meaningful explanatory factor in the best models for both bacterial and fungal communities in all compartments (see sequential test in Fig. 2). The scion, and in most cases the rootstock, were discarded because they did not provide the model with any additional explanatory power. The role of the interaction between rootstock and scion (*i.e.* rootstock–scion combination) in the assembly of microbial communities was corroborated by the low percentages of misclassification from cross-validation (CAP, Fig. 3A and D) for rhizospheric (8% and 22% for bacteria and fungi respectively) and endophytic (21% and 33% for bacteria and fungi respectively) microbial communities. Notably, while the pedo-climatic conditions of the Nipozzano and Pomino estates were more important than the scion and rootstock biotic components in shaping the edaphic microbial communities of the bulk soil (Fig. 2E and F), the formers gave the lowest contribution for the endophytic microbiomes (Fig. 2A and D).

In terms of microbial alpha-diversity, the field and the rootstock–scion interaction were confirmed as the most important factors explaining richness and Shannon




**Fig. 2.** Results of variation partitioning models used to identify the effects of abiotic (estate) and biotic (rootstock, scion and their interaction, *i.e.* rootstock–scion combination) factors analysed. The analyses are reported separately for (A–C) the bacterial and (D–F) the fungal community across (A and D) root tissues, (B and E) rhizosphere and (C and F) bulk soil compartments. Results of sequential test run considering all the factors are also reported as pie chart for bacterial and fungal communities at compartment level.

**Table 1.** Multivariate generalized linear model (*manyglm*) reporting bacterial and fungal communities' variability explained by each factor (estate, field and interaction between rootstock and scion) for the three compartments (RT, root tissue; RH, rhizosphere; BS, bulk soil).

Kingdom	Factor	RT	RH	BS
Bacteria	Estate	Dev <sub>1,55</sub> = 8199, <i>p</i> = 0.001	Dev <sub>1,64</sub> = 52 774, <i>p</i> = 0.001	Dev <sub>1,63</sub> = 60 243, <i>p</i> = 0.001
	Field	Dev <sub>9,46</sub> = 20 482, <i>p</i> = 0.001	Dev <sub>9,55</sub> = 77 717, <i>p</i> = 0.001	Dev <sub>9,54</sub> = 97 921, <i>p</i> = 0.001
	Rootstock × Scion	Dev <sub>6,46</sub> = 3861, <i>p</i> = 0.001	Dev <sub>6,55</sub> = 6339, <i>p</i> = 0.001	Dev <sub>6,54</sub> = 9318, <i>p</i> = 0.001
Fungi	Estate	Dev <sub>1,58</sub> = 4231, <i>p</i> = 0.001	Dev <sub>1,63</sub> = 13 432, <i>p</i> = 0.001	Dev <sub>1,62</sub> = 12 190, <i>p</i> = 0.001
	Field	Dev <sub>9,49</sub> = 17 778, <i>p</i> = 0.001	Dev <sub>9,54</sub> = 28 676, <i>p</i> = 0.001	Dev <sub>9,53</sub> = 29 202, <i>p</i> = 0.001
	Rootstock × Scion	Dev <sub>6,49</sub> = 1673, <i>p</i> = 0.001	Dev <sub>6,54</sub> = 2274, <i>p</i> = 0.001	Dev <sub>6,53</sub> = 2130, <i>p</i> = 0.001

Deviance and *p*-value are reported.



**Fig. 3.** Variability within endophytic and rhizospheric bacterial and fungal communities associated with the seven rootstock–scion combinations. A and B. Results of CAP cross-validation showed the percentage of samples confirmed in the different groups of rootstock–scion combinations are visualized by heat map for (A) bacterial and (B) fungal communities in root tissues (RT) and rhizosphere (RH). C–F. Statistical *p*-values of PERMANOVA pairwise analysis of beta-diversity in endophytic and rhizospheric bacterial and fungal communities associated with the seven rootstock–scion combinations. A: 110R–Chardonnay; B: 110R–Sangiovese; C: 3309–Chardonnay; D: 420A–Chardonnay; E: 420A–Merlot; F: 420A–Sangiovese; G: SO4–Chardonnay.

diversity (Table 2). However, in contrast to what was observed for beta-diversity, alpha-diversity was strongly influenced by the estates, which in certain circumstances represent the most important factor.

#### Identification of core and signature microbial discriminants across rootstock–scion combinations

For each compartment we evaluated the bacterial and fungal communities associated with different grapevine rootstock–scion combinations, independently of their field of origin. A complex picture emerged from the taxa

distribution in root tissue, rhizosphere and bulk soil across the seven rootstock–scion combinations (see summary of bacterial and fungal classes in Fig. 4). Even though the microbiomes associated with the different rootstock–scion combinations were significantly different, the shared OTUs – defining the core microbiome across the combinations – in the rhizosphere and bulk soil were 21% and 31.5% for bacteria respectively (contributing to 79.6% and 75.6% of total bacterial relative abundance), and 11.7% and 11% for fungi respectively (contributing to 69.5% and 67.5% of total fungal relative abundance; Fig. 4B and D). The edaphic core microbiomes included both abundant (>1%)



**Table 2.** Effect of abiotic (estate and field) and biotic (rootstock, scion and their interaction) factors on alpha-diversity (Shannon diversity and richness) in bacteria and fungi.

Kingdom	Factor		Shannon diversity (%IncMSE)			Richness (%IncMSE)		
			RT	RH	BS	RT	RH	BS
Bacteria	Abiotic	Estate	36.9	34.5	22.2	40.2	40.5	24.0
		Field	31.3	22.3	26.4	33.1	7.2	47.4
	Biotic	Rootstock	17.8	11.6	22.1	16.1	5.7	22.7
		Scion	23.1	8.7	22.9	25.7	4.9	22.5
		Rootstock × Scion	35.5	17.7	14.3	22.6	11.4	12.2
Fungi	Abiotic	Estate	28.8	1.2	12.4	21.3	38.1	31.2
		Filed	21.6	15.1	29.7	45.2	38.9	15.1
	Biotic	Rootstock	8.2	1.1	6.9	24.5	9.9	10.9
		Scion	8.4	3.2	3.6	22.9	20.7	14.6
		Rootstock × Scion	13.1	6.1	8.8	31.1	29.8	17.8

The most important factor that explains the pattern of alpha-diversity in root tissues (RT), rhizosphere (RH) and bulk soil (BS) was determined by Random Forest analysis. The mean decrease accuracy (%IncMSE) was reported for each tested factor.

and rare (<1%) OTUs, and they were dominated by bacteria belonging to *Alphaproteobacteria*, *Blastocatellia*, *Gammaproteobacteria*, *Actinobacteria* and *Bacteroidia*, and fungi belonging to *Sordariomycetes*, *Dothideomycetes*, *Mortierellomycetes* and *Tremellomycetes*. The total of OTUs specific for each rootstock–scion combination (*i.e.* present only in a single combination) were only 5% and 4.8% in the bacterial communities of the rhizosphere and bulk soil respectively, while they were up to 14.4% and 19.8% in those of fungi (Fig. 4B and D). However, these OTUs specific of each combination covered less than 3% of relative abundance, underlining the importance of rare species in community assembly processes and plant–microbes interactions (Zhang *et al.*, 2022). In contrast, the root microbiomes shared only 1% of bacterial OTUs and 3.4% of fungal OTUs, corresponding to 24% (13% *Alphaproteobacteria* and 6% *Bacilli*) and 46.4% (24% *Sordariomycetes* and 17.5% *Dothideomycetes*) of total abundance, along with a higher percentage of rootstock–scion combination-specific OTUs [34% (3% of relative abundance) and 48% (5.3% of relative abundance) respectively; Fig. 4B and D]. These combination-specific bacterial OTUs mainly belong to *Rhizobium* (*Alphaproteobacteria*) and *Alloprevotella* (*Bacteroidia*) in SO4–Chardonnay, *Nonlabens* (*Flavobacteria*) and *Arcticibacter* (*Sphingobacteriia*) in 420A–Sangiovese, *Candidatus* Phytoplasma (*Bacilli*) in 420A–Chardonnay, *Ezakiella* (*Clostridia*) and *Cutibacterium* (*Actinobacteria*) in 110R–Chardonnay, *Turicella* (*Actinobacteria*) in 110R–Sangiovese, *Flavobacterium* (*Flavobacteriia*) and *Chitinophaga* (*Chitinophagia*) in 3309C–Chardonnay, and *Corynebacterium* (*Actinobacteria*) and *Steroidobacter* (*Gammaproteobacteria*) in 420A–Merlot. In the case of fungi, we mainly detected members of *Psathyrella* (*Agaricomycetes*) and *Rhizophydium* (*Chytridiomycetes*) in 110R–Chardonnay, *Parasola* (*Agaricomycetes*) and the arbuscular mycorrhiza *Glomus* (*Glomeromycetes*) in

110R–Sangiovese, *Catapyrenium* (*Eurotiomycetes*) and *Hymenoscyphus* (*Ascomycetes*) in 3309C–Chardonnay, *Fusarium* (*Sordariomycetes*) and *Mortierella* (*Mucoromycota*) in 420A–Chardonnay, *Saccharomyces* (*Saccharomycetes*) and *Donadinia* (*Pezizomycetes*) in 420A–Merlot, the symbiotic *Claroideoglossum* (*Glomeromycetes*) in 420A–Sangiovese and *Thelebolus* (*Leotiomycetes*) in SO4–Chardonnay. The remaining OTUs were shared across the rootstock–scion combinations (minimum two, maximum six; see ‘Multi combo’ in Fig. 5B and D).

To distinguish the OTUs that could discriminate the microbial communities across the rootstock–scion combinations – not limited to their presence/absence – we analysed the deviance of OTUs abundance. Ranked by their importance values (*i.e.* highest deviance), the top 20 bacterial and fungal OTUs were reported (Fig. 5A–F; full list in Supplementary Table S9). In the rhizosphere, the discriminant bacterial OTUs mainly belonged to *Gammaproteobacteria* (11 OTUs), *Actinobacteria* (3) and *Alphaproteobacteria* (2), while in bulk soil they belonged to *Acidobacteria* (3), *Bacteroidia* (2), *Blastocatellia* (2), *Thermoleophilia* (2) and *Gammaproteobacteria* (2). Notably, in root tissue *Actinobacteria* (15) were the most important bacterial discriminants, along with *Candidatus* Phytoplasma OTUs (2) that were detected only in the 420A–Chardonnay bacterial communities (Fig. 5A). In the case of fungal communities, the rootstock–scion combination-discriminant OTUs were *Sordariomycetes* (five and six in the rhizosphere and bulk soil respectively), *Eurotiomycetes* (three and six) and *Dothideomycetes* (three and five); *Leotiomycetes* (1) *Mortierellomycetes* (1) *Mucoromycetes* (1) and *Tremellomycetes* (2) were rootstock–scion combination signatures only in the rhizosphere (Fig. 5D–F). Overall, the 20 OTUs distinguishing rootstock–scion combinations had a low level of abundance in the rhizosphere





Fig. 4. Legend on next page.



**Fig. 5.** Top 20 OTUs discriminating (A–C) bacterial and (D–F) fungal communities in the different rootstock–scion combinations; the importance (deviance) value of each OTU estimated by random forest model is reported in the bar chart for (A and D) root tissues (green), (B and E) rhizosphere (black) and (C and F) bulk soil (grey) compartments. The assigned taxonomy of each taxon is displayed at the class (or highest taxonomic rank if class is not available for an OTU). The heat maps on the right of each bar plot show the relative abundances of the selected OTUs across the rootstock–scion combinations; values are reported as mean relative abundance log-transformed. Star (\*) indicates that the OTU belong to the bacterial and fungal compartment core (*i.e.* bacterium/fungus shared across all the rootstock–scion combinations in a certain compartment).

and bulk soil bacterial communities (range, <1%–11%), while they reached up to 26% of relative abundance in those of fungi. The majority of these OTUs belong to the core microbiome (*i.e.* present across all rootstock–scion combinations; see stars in Fig. 5), suggesting that their distribution/relative abundance rather than their presence/absence make them microbial discriminants. In contrast, in root tissues the relative abundance of influential OTUs was consistently greater compared to that of edaphic compartments, reaching up to 50% and 53% in bacterial and fungal communities respectively. However,

in the heterogeneous, highly dispersed and uneven endophytic communities, only a few of these OTUs made up part of the core microbiome (see stars in Fig. 5).

*Microbial taxa as indicators of rootstock–scion combinations in grafted grapevine root system*

Indicator species analysis was performed to test the strength and statistical significance of the relationship between microbial OTUs and the different rootstock–scion combinations (Table 3), and thus to establish which

**Fig. 4.** Compositional diversity of (A and B) bacterial and (C and D) fungal communities associated with root tissues, rhizosphere and bulk soil compartments across the seven rootstock–scion combinations respectively; A: 110R–Chardonnay; B: 110R–Sangiovese; C: 3309–Chardonnay; D: 420A–Chardonnay; E: 420A–Merlot; F: 420A–Sangiovese; G: SO4–Chardonnay.

A and C. Relative abundance of bacterial and fungal classes is reported respectively; classes representing <1% of the total reads are grouped indicated as ‘Others’.

B and D. Distribution of bacterial and fungal operational taxonomic units (OTUs) respectively, across the seven different scion–rootstock combinations in the three compartments. Values are reported as percentage of OTUs; for the core and multiple-combination categories the total relative abundance of OTUs in the compartment is also reported.

**Table 3.** Microbial indicator species (OTUs) of rootstock–scion combinations in grapevine root system, namely, root tissues (RT) and rhizosphere (RH).

Indicator OTUs of rootstock–scion combination	Bacteria		Fungi	
	RT	RH	RT	RH
N. OTUs	2197	5150	941	1694
Percentage (%) of indicator OTUs of comb.	<b>12.7</b>	<b>35.1</b>	<b>44.4</b>	<b>27.3</b>
% indicator OTUs in 1 combination	10.3 (81)	17.7 (50)	38.1 (86)	18.5 (68)
% in 110R–Chardonnay	0.7	3.2	10.5	9.7
% in 110R–Sangiovese	6.4	14.6	6.9	9.9
% in 3309–Chardonnay	45.4	16.6	14.8	15.6
% in 420A–Chardonnay	3.9	8.1	22.5	10.2
% in 420A–Merlot	n.d.	1.3	6.5	2.2
% in 420A–Sangiovese	0.7	3.2	17.2	12.1
% in SO4–Chardonnay	23.6	3.5	7.2	8.2
Indicator OTUs in two combinations	2.3 (18)	10.8 (29)	4.8 (11)	6.3 (23)
Indicator OTUs in three combinations	0.1 (1)	5.1 (14)	1 (2)	1.8 (7)
Indicator OTUs in four combinations	n.d.	2.0 (6)	0.5 (1)	0.6 (2)
Indicator OTUs in five combinations	n.d.	0.2 (1)	0.1 (<1)	n.d.
Indicator OTUs in six combinations	n.d.	n.d.	n.d.	n.d.

Total number of OTUs and percentage of detected indicator OTUs (numbers in bold; significance,  $p < 0.05$ ) are reported. Within the indicator OTUs of rootstock–scion combinations, differentiation along the different groups of combinations (single and multiple) are specified; in the brackets is calculated the percentage of each group with respect to the percentage of the total indicator OTUs for each compartment. The percentage of indicator OTUs associated with specific combinations is reported in the table. The list of the indicator OTUs across the different groups is reported in Supplementary Table S10.

n.d., not determined by indicator species analysis (*IndicSpecies*).

OTUs are characteristic for individual or multiple (from two to six) combinations. The analysis detected 35% and 27.3% of bacterial and fungal OTUs respectively, as specifically associated with the rhizosphere of one or more rootstock–scion combinations, while in the case of endophytic communities we detected 12.7% and 44.4% of indicator bacterial and fungal OTUs respectively (Table 3). As a general trend, indicator species tended to be associated with only one rootstock–scion combination, with 81% and 86% bacterial and fungal indicator OTUs in the root tissues respectively, and 50% and 68% in the rhizosphere (see the group with one combination in Table 3). Considering the group containing two rootstock–scion combinations, we observed a decline in microbial indicator OTUs for root tissues (18% and 11% in bacteria and fungi respectively) and the rhizosphere (29% and 23% respectively). Bacterial endophytic indicators were dominated by members of *Rhizobiaceae* (*Alphaproteobacteria*), *Micromonosporaceae* and *Streptomyetaceae* (*Actinobacteria*), *Comamonadaceae* and *Steroidobacteraceae* (*Gammaproteobacteria*) and *Microscillaceae* (*Bacteroidia*), except for 420A–Chardonnay that presented *Candidatus* *Phytoplasma* (*Bacilli*) as indicator OTU (Supplementary Table S10A). The taxonomic diversity of indicator OTUs increased in the rhizosphere, with the presence of numerous families, including *Micromonosporaceae*, *Nocardioidaceae* and *Pseudonocardiaceae* within *Actinobacteria*, *Beijerinckiaceae*, *Rhizobiaceae*, *Sphingomonadaceae* and *Xanthobacteraceae* within *Alphaproteobacteria*, *Comamonadaceae*,

*Nitrosomonadaceae*, *Pseudomonadaceae* and *Steroidobacteraceae* within *Gammaproteobacteria*, and several members of *Bacteroidia*, *Gemmatimonadetes*, *Plantomycetes*, *Polyangia*, *Thermoleophilia*, *Verrucomicrobiae* and *Vicinamibacteria* (Supplementary Table S10A). In the case of fungi, the endophytic and rhizosphere indicator OTUs of rootstock–scion combinations were mainly composed of *Nectriaceae*, *Lasiosphaeriaceae*, *Chaetomiaceae* (*Sordariomycetes*) and unclassified of the same class, unclassified *Dothideomycetes*, unclassified *Agaricomycetes*, *Herpotrichiellaceae* and unclassified within the *Eurotiomycetes* class, *Rhytismataceae* and *Helotiaceae* (*Leotiomycetes*), *Glomeraceae* (*Glomeromycetes*), along with members of the *Mortierellomycetes*, *Peizizomycetes*, *Orbiliomycetes* and *Tremellomycetes* classes (Supplementary Table S10B).

## Discussion

Environmental factors, such as physicochemical characteristics of soil, along with plant-related factors (among others, plant genotype) have been reported as the main factors regulating grapevine performance, as well as recruitment and selection of its associated microbiome (Costa et al., 2006; Pancher et al., 2012; Marasco et al., 2013; Zarraonandia et al., 2015). However, in grafted plants the physical/morphological/chemical mechanisms modulating the plant–soil feedback (Hu et al., 2018; Olanrewaju et al., 2019) are influenced by the interaction

of, at least, two genotypes, *i.e.* the rootstock and the scion (Gautier *et al.*, 2019).

Belowground, the root metabolites, which act as growth substrates for suitable microbial partners, attract the surrounding edaphic microorganisms that form the rhizosphere microbiome (van Dam and Bouwmeester, 2016; Massalha *et al.*, 2017; Sasse *et al.*, 2018; Vives-Peris *et al.*, 2020). This 'selective pressure' imposed by the plant on its immediate surrounding results in a sharp decrease in microbial diversity with proximity to the plant (*i.e.* bulk soil > rhizosphere > root tissue; Fig. 1), consequently shaping the composition and relative abundance of microbial species associated with the grapevine compartments (Figs 3 and 4) (Bulgarelli *et al.*, 2013; Van Der Heijden and Schlaeppi, 2015; Zorraonaindia *et al.*, 2015; Saleem *et al.*, 2018). The root exudation pattern is genetically controlled by different factors innate to the plant, including the genotype, with inevitable consequences on the establishment of mutualistic relations with the surrounding edaphic microorganisms (Vives-Peris *et al.*, 2020; Chai *et al.*, 2022). Recent studies have shown that within a vineyard grapevine rootstock genotypes grafted with the same scion are associated with different rhizospheric and endophytic microbiomes (D'Amico *et al.*, 2018; Marasco *et al.*, 2018b; Berlanas *et al.*, 2019). However, here, we have shown that at ecosystem level across the heterogeneous conditions of different vineyards, when the interaction between rootstock and scion is considered, this is more important than the rootstock or scion considered alone (Fig. 2). This reinforces, at ecosystem scale, what was recently observed for young grapevines cultivated in an experimental field (Vink *et al.*, 2021b) and for apple trees from a single orchard (Chai *et al.*, 2022). In addition, our study provides evidence that this process is extended to the endophytic root communities, and to the fungal components of the grape microbiome.

The aboveground component of the plant (the scion) is involved in communication with the rootstock belowground for nutrient exchange, growth signalling and many other aspects of plant physiology (Tandonnet *et al.*, 2009; Aloni *et al.*, 2010; Martínez-Ballesta *et al.*, 2010; Bonghi *et al.*, 2016; Gautier *et al.*, 2019). Among them, we find the production and distribution of phytohormones. Auxins, which are produced in the shoot apices, are translocated to the roots where they regulate many aspects of plant vegetative and reproductive development, such as root system growth, architecture and functioning (Persello-Cartieaux *et al.*, 2001; Aloni *et al.*, 2010; Overvoorde *et al.*, 2010; Egamberdieva *et al.*, 2017). Although the mechanism was not fully elucidated, it has been shown that in young grapevines the scion genotype confers different root vigour possibly through signal molecules and hormones rather than by carbon supply from the shoot (Tandonnet *et al.*, 2009).

Since roots play a direct role in the establishment and maintenance of the interaction between host plants and soil microorganisms, any modification in their architecture (*i.e.* overall length, density, branching and biomass) and root morphology (*i.e.* diameter, surface area and root hairs) can affect the plant-associated microbial communities and the recruitment of beneficial microorganisms (Herms *et al.*, 2022). For instance, thinner root diameter recruits a more diverse rhizosphere community in different plants, including arboreal and herbaceous species, which can often result in a positive impact on plant health and productivity (Bardgett and van der Putten, 2014; Saleem *et al.*, 2019; Herms *et al.*, 2022).

Phytohormones, such as auxin and salicylic acid, are also key players in mediating the crosstalk between plants and beneficial bacteria, acting as reciprocal signalling molecules in plant–microbe interactions (Lebeis *et al.*, 2015; Khan *et al.*, 2020). Thus, modifications in the hormones' distribution mediated by the plant genotypes (and their interactions) could affect the composition of the plant microbiome. In our work, we found predominant amounts of *Proteobacteria* and *Actinobacteria* among bacteria, and *Ascomycota* and *Basidiomycota* among fungi in all the rhizosphere and root samples, but with differential distribution across the grafting combinations (Fig. 4). Microbial discriminants and indicator species were in part constituted by OTUs shared across all the combinations but with differential enrichment in terms of relative abundance (Fig. 5; Table 3). Such variability in the relative abundance possibly occurs because such OTUs are differentially supported by specific combination of scion/rootstock genotypes. As suggested by Semchenko *et al.* (2022), the identification of the factors that promote the plant association with microbial specialists and/or generalists will advance our understanding of the mechanisms regulating plant–soil feedback and of the ways plant–microbiome contributes to plant performances.

The compatibility of the plant genotype with the growth/association of/beneficial microorganisms is a *sine qua non* requisite for the plant to benefit from the services provided by such microorganisms and stochastic encounters with PGP microorganisms are not sufficient to explain the health benefits mediated by the plant microbiome (Haney *et al.*, 2015). For instance, during bacteria-based treatments conducted in the greenhouse with Barbera plantlets grafted onto 420A, SO4 and K5BB rootstocks, it has been observed that the water deficit-tolerance mediated by PGP bacteria was a rootstock dependent trait; the three rootstocks showed different physiological responses and phenotypes (*i.e.* biomass, photosynthetic activity, evapotranspiration and stomatal conductance) in response to the treatment with a given PGP bacterium (Rolli *et al.*, 2012, 2015). Similarly, by exploiting the availability of 196 naturally occurring

accessions of *Arabidopsis thaliana*, Haney *et al.* (2015) demonstrated that the genotype was an essential factor in determining whether a microbial community provides benefit or harm to its host. The authors showed that different plant genotypes within a single species (*A. thaliana*) have different rhizosphere microbiomes, which mainly differ in terms of the presence of beneficial *Pseudomonas* bacteria, with consequences for plant health. Extending this finding to our work, it is plausible to suppose that any change in the distribution of microorganisms driven by the interaction between the rootstock and the scion can have further influence on the ecological services carried to the host and to the ecosystem. One interesting example in our study is provided by *Candidatus* Phytoplasma, detected by different analyses exclusively as an indicator of 420A–Chardonnay. This suggests that this specific combination is more susceptible to colonization of the plant root by this intracellular parasite and the related insect vectors (Aryan *et al.*, 2014). Notably, in the root tissues of 420A grafted with different scions, including Merlot and Sangiovese (both from this work) and Barbera (Marasco *et al.*, 2018b), OTUs assigned to *Candidatus* Phytoplasma were not detected.

Since microbes interacting with plants are involved in sustaining plant fitness, growth promotion and disease control (Delgado-Baquerizo, 2022), the results reported here indicate that it is important to consider the rootstock–scion combination rather than only the rootstock to evaluate the process of recruitment and selection of edaphic microorganisms mediated by the plant. Such considerations are also valid when we consider the plant microbiome and its engineering for sustainable agriculture (Kaul *et al.*, 2021): the link between the agronomic features, grafting combinations and microbiome is instrumental to understand the performance reports, often contradictory, of grafted grapevines in different areas (Ibacache *et al.*, 2020). In the case of viticulture, the genotype-mediated selection carried out by the root system becomes a more important factor because it can also affect the microbiome associated with other plant organs, such as the fruit, potentially impacting the final product (Bokulich *et al.*, 2016; Griggs *et al.*, 2021).

## Conclusion

Considering the importance of the microbiome for several aspects of grapevine physiology, stress tolerance and pathogen protection, it is important to elucidate the role of plant components (*i.e.* rootstock, scion, and their combination) in ecological processes that shape plant–microbe interactions. Which genotype, which rootstock, which scion or which of their combinations, has the greatest effect on the selection of microorganisms? The data provided here underline the importance of considering the interaction between the genotypes that compose the overall host,

namely, rootstock and scion, as a factor that drives the recruitment and selection of the surrounding edaphic microbiome by the root system. In our study, the interaction of rootstock and scion resulted to be more important in shaping the root system microbiome than the rootstock and scion considered separately. Thus, selection of a given rootstock–scion combination is not only crucial for physiological adaptation of the grapevine to a certain vineyard/ecosystem but has also important repercussions on the overall metaorganism (plant and microbiome) and its fitness. Although the relative weight of each genotype in determining the composition of a microbial community has yet to be elucidated, the results obtained contribute towards creating a new perspective for research, targeted at understanding and explaining the role of rootstock–scion combination management in the ecology of the grapevine microbiome. This could also lay the foundations for including the microbiome in grapevine breeding programmes of scions and rootstocks and for improving grapevine physiology through microorganism/microbiome-based treatments.

## Acknowledgements

The authors thank Frescobaldi s.p.a. for allowing access to their vineyards and for their support in sampling. We thank the manager of the Pomino estate, Fabio Dini, the manager of the Nipozzano estate, Daniele Settesoldi and the manager of the production process, Francesca Pratesi for their invaluable logistical support during fieldwork and our visit to Frescobaldi s.p.a. We also thank Andrea Bellucci for coordinating the visit to the Frescobaldi vineyards, Taskeen Begun for her help in laboratory organization, and the BCL team for support during MiSeq sequencing.

## Availability of Data and Materials

The datasets analysed in this current study are available in the NCBI SRA repository under the BioProject PRJNA807110. The codes used to perform the analyses are available at [https://github.com/MarcoFusi1980/Tuscan\\_Vineyard\\_microbial\\_ecology](https://github.com/MarcoFusi1980/Tuscan_Vineyard_microbial_ecology).

## Authors' Contributions

R.M., M.F. and D.D.: Conceived and designed the study and experiments. R.M., M.F. and D.D.: Sample collection. R.M. and H.A.: Performed the experiments. R.M. and M.F.: Analysed the data. D.D.: Contributed reagents/materials/analysis tools. R.M., H.A. and M.F.: Wrote the paper. D.D., I.G. and L.V.: Critical revision of the manuscript. All the authors critically revised the manuscript.

## References

- Aloni, B., Cohen, R., Karni, L., Aktas, H., and Edelstein, M. (2010) Hormonal signaling in rootstock–scion interactions. *Sci Hortic* **127**: 119–126.
- Anderson, M.M.J.J., Gorley, R.N.R.N., and Clarke, K.R.R. (2008) *PERMANOVA + for PRIMER: guide to Software and Statistical Methods*. Plymouth, UK: PRIMER-E.
- Aryan, A., Brader, G., Mörtel, J., Pastar, M., and Riedle-Bauer, M. (2014) An abundant ‘*Candidatus* Phytoplasma solani’ tuf b strain is associated with grapevine, stinging nettle and *Hyalesthes obsoletus*. *Eur J Plant Pathol* **140**: 213–227.
- Bardgett, R.D., and van der Putten, W.H. (2014) Below-ground biodiversity and ecosystem functioning. *Nature* **515**: 505–511.
- Batley, N.H., and Simmonds, P.E. (2005) Phylloxera and the grapevine: a sense of common purpose? *J Exp Bot* **56**: 3029–3031.
- Berdeja, M., Nicolas, P., Kappel, C., Dai, Z.W., Hilbert, G., Peccoux, A., et al. (2015) Water limitation and rootstock genotype interact to alter grape berry metabolism through transcriptome reprogramming. *Hortic Res* **2**: 15012.
- Berendsen, R.L., Pieterse, C.M.J., and Bakker, P.A.H.M. (2012) The rhizosphere microbiome and plant health. *Trends Plant Sci* **17**: 478–486.
- Berlanas, C., Berbegal, M., Elena, G., Laidani, M., Cibriain, J.F., Sagües, A., and Gramaje, D. (2019) The fungal and bacterial rhizosphere microbiome associated with grapevine rootstock genotypes in mature and young vineyards. *Front Microbiol* **10**: 1–16.
- Bert, P.-F., Bordenave, L., Donnart, M., Hévin, C., Ollat, N., and Decroocq, S. (2013) Mapping genetic loci for tolerance to lime-induced iron deficiency chlorosis in grapevine rootstocks (*Vitis* sp.). *Theor Appl Genet* **126**: 451–473.
- Bettenfeld, P., Cadena I Canals, J., Jacquens, L., Fernandez, O., Fontaine, F., van Schaik, E., et al. (2021) The microbiota of the grapevine holobiont: a key component of plant health. *J Adv Res.* in press. doi:[10.1016/j.jare.2021.12.008](https://doi.org/10.1016/j.jare.2021.12.008)
- Bianchi, D., Caramanico, L., Grossi, D., Brancadoro, L., and De Lorenzis, G. (2020) How do novel M-rootstock (*Vitis* spp.) genotypes cope with drought? *Plan Theory* **9**: 1385.
- Bokulich, N.A., Collins, T.S., Masarweh, C., Allen, G., Heymann, H., Ebeler, S.E., and Mills, D.A. (2016) Associations among wine grape microbiome, metabolome, and fermentation behavior suggest microbial contribution to regional wine characteristics. *MBio* **7**: 1–12.
- Bonghi, C., Corso, M., Vannozzi, A., Ziliotto, F., Zouine, M., Maza, E., et al. (2016) Grapevine rootstocks differentially affect the rate of ripening and modulate auxin-related genes in Cabernet Sauvignon berries. *Front Plant Sci* **7**: 1–14.
- Bulgarelli, D., Schlaeppli, K., Spaepen, S., van Themaat, E. V.L., Schulze-Lefert, P., Loren, V., et al. (2013) Structure and functions of the bacterial microbiota of plants. *Annu Rev Plant Biol* **64**: 807–838.
- Burns, K.N., Bokulich, N.A., Cantu, D., Greenhut, R.F., Kluepfel, D.A., O’Geen, A.T., et al. (2016) Vineyard soil bacterial diversity and composition revealed by 16S rRNA genes: differentiation by vineyard management. *Soil Biol Biochem* **103**: 337–348.
- Campisano, A., Antonielli, L., Pancher, M., Yousaf, S., Pindo, M., and Pertot, I. (2014) Bacterial endophytic communities in the grapevine depend on pest management. *PLoS One* **9**: e112763.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**: 335–336.
- Ceja-Navarro, J.A., Wang, Y., Ning, D., Arellano, A., Ramanculova, L., Yuan, M.M., et al. (2021) Protist diversity and community complexity in the rhizosphere of switchgrass are dynamic as plants develop. *Microbiome* **9**: 1–18.
- Chai, X., Wang, X., Li, H., Xu, X., Wu, T., Zhang, X., et al. (2022) Apple scion cultivars regulate the rhizosphere microbiota of scion/rootstock combinations. *Appl Soil Ecol* **170**: 104305.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., et al. (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* **437**: 529–533.
- Costa, R., Götz, M., Mrotzek, N., Lottmann, J., Berg, G., and Smalla, K. (2006) Effects of site and plant species on rhizosphere community structure as revealed by molecular analysis of microbial guilds. *FEMS Microbiol Ecol* **56**: 236–249.
- D’Amico, F., Candela, M., Turrone, S., Biagi, E., Brigidi, P., Bega, A., et al. (2018) The rootstock regulates microbiome diversity in root and rhizosphere compartments of *Vitis vinifera* cultivar Lambrusco. *Front Microbiol* **9**: 1–11.
- Darriaut, R., Lailheugue, V., Masneuf-Pomarède, I., Marguerit, E., Martins, G., Compant, S., et al. (2022) Grapevine rootstock and soil microbiome interactions: keys for a resilient viticulture. *Hortic Res* **12**: 4683–4698.
- De Cáceres, M., Jansen, F., and Dell, N. (2020) Package ‘indicspecies.’ CRAN.
- Delgado-Baquerizo, M. (2022) Simplifying the complexity of the soil microbiome to guide the development of next-generation SynComs. *J Sustain Agric Environ* **1**: 1–7.
- Delgado-Baquerizo, M., and Eldridge, D.J. (2019) Cross-biome drivers of soil bacterial alpha diversity on a world-wide scale. *Ecosystems* **22**: 1220–1231.
- Deyett, E., and Rolshausen, P.E. (2020) Endophytic microbial assemblage in grapevine. *FEMS Microbiol Ecol* **96**: 1–11.
- Dini-Andreote, F., and Raaijmakers, J.M. (2018) Embracing community ecology in plant microbiome research. *Trends Plant Sci* **23**: 467–469.
- Dries, L., Bussotti, S., Pozzi, C., Kunz, R., Schnell, S., Löhnertz, O., and Vortkamp, A. (2021a) Rootstocks shape their microbiome-bacterial communities in the rhizosphere of different grapevine rootstocks. *Microorganisms* **9**: 822.
- Dries, L., Hendgen, M., Schnell, S., Löhnertz, O., and Vortkamp, A. (2021b) Rhizosphere engineering: leading towards a sustainable viticulture? *OENO One* **55**: 353–363.
- Edgar, R.C. (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* **10**: 996–998.
- Egamberdieva, D., Wirth, S.J., Alqarawi, A.A., Abd-Allah, E. F., and Hashem, A. (2017) Phytohormones and beneficial

- microbes: essential components for plants to balance stress and fitness. *Front Microbiol* **8**: 1–14.
- Fitzpatrick, C.R., Lu-Ingving, P., Copeland, J., Guttman, D.S., Wang, P.W., Baltrus, D.A., et al. (2018) Chloroplast sequence variation and the efficacy of peptide nucleic acids for blocking host amplification in plant microbiome studies. *Microbiome* **6**: 144.
- Fuentes, I., Stegemann, S., Golczyk, H., Karcher, D., and Bock, R. (2014) Horizontal genome transfer as an asexual path to the formation of new species. *Nature* **511**: 232–235.
- Gaion, L.A., Braz, L.T., and Carvalho, R.F. (2018) Grafting in vegetable crops: a great technique for agriculture. *Int J Veg Sci* **24**: 85–102.
- Gamalero, E., Bona, E., Novello, G., Boatti, L., Mignone, F., Massa, N., et al. (2020) Discovering the bacteriome of *Vitis vinifera* cv. Pinot Noir in a conventionally managed vineyard. *Sci Rep* **10**: 1–12.
- Gautier, A.T., Chambaud, C., Brocard, L., Ollat, N., Gambetta, G.A., Delrot, S., and Cookson, S.J. (2019) Merging genotypes: graft union formation and scion–rootstock interactions. *J Exp Bot* **70**: 747–755.
- Gobbi, A., Acedo, A., Imam, N., Santini, R.G., Ortiz-Álvarez, R., Ellegaard-Jensen, L., et al. (2022) A global microbiome survey of vineyard soils highlights the microbial dimension of viticultural terroirs. *Commun Biol* **5**: 1–9.
- Griggs, R.G., Steenwerth, K.L., Mills, D.A., Cantu, D., and Bokulich, N.A. (2021) Sources and assembly of microbial communities in vineyards as a functional component of winegrowing. *Front Microbiol* **12**: 673810.
- Habran, A., Commisso, M., Helwi, P., Hilbert, G., and Negri, S. (2016) Rootstocks/scion/nitrogen interactions affect secondary metabolism in the grape berry. *Front Plant Sci* **7**: 1134.
- Hamrouni, L., Hanana, M., Abdelly, C., and Hamed, K. (2015) Influence of the rootstock/scion combination on the grapevines behavior under salt stress. *J Plant Biochem Physiol* **03**: 154.
- Haney, C.H., Samuel, B.S., Bush, J., and Ausubel, F.M. (2015) Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nat Plants* **1**: 15051.
- Hassani, M.A., Durán, P., and Hacquard, S. (2018) Microbial interactions within the plant holobiont. *Microbiome* **6**: 58.
- Henderson, S.W., Baumann, U., Blackmore, D.H., Walker, A.R., Walker, R.R., and Gilliam, M. (2014) Shoot chloride exclusion and salt tolerance in grapevine is associated with differential ion transporter expression in roots. *BMC Plant Biol* **14**: 273.
- Herns, C.H., Hennessy, R.C., Bak, F., Dresbøll, D.B., and Nicolaisen, M.H. (2022) Back to our roots: exploring the role of root morphology as a mediator of beneficial plant–microbe interactions. *Environ Microbiol.* doi:10.1111/1462-2920.15926
- Hu, L., Robert, C.A.M., Cadot, S., Zhang, X., Ye, M., Li, B., et al. (2018) Root exudate metabolites drive plant–soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat Commun* **9**: 2738.
- Ibacache, A., Verdugo-Vásquez, N., and Zurita-Silva, A. (2020) Rootstock:scion combinations and nutrient uptake in grapevines. In *Fruit Crops*. Elsevier, pp. 297–316.
- Kaul, S., Choudhary, M., Gupta, S., and Dhar, M.K. (2021) Engineering host microbiome for crop improvement and sustainable agriculture. *Front Microbiol* **12**: 1–10.
- Khan, N., Bano, A., Ali, S., and Babar, M.A. (2020) Crosstalk amongst phytohormones from planta and PGPR under biotic and abiotic stresses. *Plant Growth Regul* **90**: 189–203.
- Klindworth, A., Priesse, E., Schweer, T., Peplies, J.J., Quast, C., Horn, M., et al. (2013) Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res* **41**: 1–11.
- Koevoets, I.T., Venema, J.H., Elzenga, J.T.M., and Testerink, C. (2016) Roots withstanding their environment: exploiting root system architecture responses to abiotic stress to improve crop tolerance. *Front Plant Sci* **07**: 1–19.
- Lakshmanan, V., Selvaraj, G., and Bais, H.P. (2014) Functional soil microbiome: belowground solutions to an aboveground problem. *Plant Physiol* **166**: 689–700.
- Lebeis, S.L., Paredes, S.H., Lundberg, D.S., Breakfield, N., Gehring, J., McDonald, M., et al. (2015) Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science* **349**: 860–864.
- Manici, L.M., Saccà, M.L., Caputo, F., Zanzotto, A., Gardiman, M., and Fila, G. (2017) Long-term grapevine cultivation and agro-environment affect rhizosphere microbiome rather than plant age. *Appl Soil Ecol* **119**: 214–225.
- Mapelli, F., Marasco, R., Fusi, M., Scaglia, B., Tsiamis, G., Rolli, E., et al. (2018) The stage of soil development modulates rhizosphere effect along a high Arctic desert chronosequence. *ISME J* **12**: 1188–1198.
- Marasco, R., Mosqueira, M.J., Fusi, M., Ramond, J., Merlino, G., Booth, J.M., et al. (2018a) Rhizosphere microbial community assembly of sympatric desert spargrasses is independent of the plant host. *Microbiome* **6**: 215.
- Marasco, R., Rolli, E., Fusi, M., Cherif, A., Abou-Hadid, A., El-Bahairy, U., et al. (2013) Plant growth promotion potential is equally represented in diverse grapevine root-associated bacterial communities from different biopedoclimatic environments. *Biomed Res Int* **2013**: 1–17.
- Marasco, R., Rolli, E., Fusi, M., Michoud, G., and Daffonchio, D. (2018b) Grapevine rootstocks shape underground bacterial microbiome and networking but not potential functionality. *Microbiome* **6**: 3.
- Martínez-Ballesta, M.C., Alcaraz-López, C., Muries, B., Mota-Cadenas, C., and Carvajal, M. (2010) Physiological aspects of rootstock–scion interactions. *Sci Hortic* **127**: 112–118.
- Massalha, H., Korenblum, E., Tholl, D., and Aharoni, A. (2017) Small molecules below-ground: the role of specialized metabolites in the rhizosphere. *Plant J* **90**: 788–807.
- McMurdie, P.J., and Holmes, S. (2013) phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* **8**: e61217.
- Melnyk, C.W., Gabel, A., Hardcastle, T.J., Robinson, S., Miyashima, S., Grosse, I., and Meyerowitz, E.M. (2018) Transcriptome dynamics at *Arabidopsis* graft junctions reveal an intertissue recognition mechanism that activates



- vascular regeneration. *Proc Natl Acad Sci U S A* **115**: E2447–E2456.
- Mommer, L., Cotton, T.E.A., Raaijmakers, J.M., Termorshuizen, A.J., van Ruijven, J., Hendriks, M., *et al.* (2018) Lost in diversity: the interactions between soil-borne fungi, biodiversity and plant productivity. *New Phytol* **218**: 542–553.
- Morales-Castilla, I., García de Cortázar-Atauri, I., Cook, B.I., Lacombe, T., Parker, A., van Leeuwen, C., *et al.* (2020) Diversity buffers winegrowing regions from climate change losses. *Proc Natl Acad Sci U S A* **117**: 2864–2869.
- Mueller, U.G., and Sachs, J.L. (2015) Engineering microbiomes to improve plant and animal health. *Trends Microbiol* **23**: 606–617.
- Nerva, L., Giudice, G., Quiroga, G., Belfiore, N., Lovat, L., Perria, R., *et al.* (2021) Mycorrhizal symbiosis balances rootstock-mediated growth-defence tradeoffs. *Biol Fertil Soils* **58**: 17–34.
- OIV The International Organisation of Vine and Wine. (2020) *The scientific and technical reference of the vine and wine world*. URL <https://www.oiv.int/en/the-international-organisation-of-vine-and-wine>.
- Olanrewaju, O.S., Ayangbenro, A.S., Glick, B.R., and Babalola, O.O. (2019) Plant health: feedback effect of root exudates-rhizobium interactions. *Appl Microbiol Biotechnol* **103**: 1155–1166.
- Ollat, N., Bordenave, L., Tandonnet, J.P., Boursiquot, J.M., and Marguerit, E. (2016) Grapevine rootstocks: origins and perspectives. *Acta Hort* **1136**: 11–22.
- Overvoorde, P., Fukaki, H., and Beeckman, T. (2010) Auxin control of root development. *Cold Spring Harb Perspect Biol* **2**: 1–16.
- Pancher, M., Ceol, M., Corneo, P.E., Longa, C.M.O., Yousaf, S., Pertot, I., and Campisano, A. (2012) Fungal endophytic communities in grapevines (*Vitis vinifera* L.) respond to crop management. *Appl Environ Microbiol* **78**: 4308–4317.
- Persello-Cartieaux, F., David, P., Sarrobert, C., Thibaud, M. C., Achouak, W., Robaglia, C., and Nussaume, L. (2001) Utilization of mutants to analyze the interaction between *Arabidopsis thaliana* and its naturally root-associated *Pseudomonas*. *Planta* **212**: 190–198.
- Pina, A., Cookson, S.J., Calatayud, A., Trinchera, A., and Errea, P. (2017) Physiological and molecular mechanisms underlying graft compatibility. In *Vegetable Grafting: Principles and Practices*. Wallingford: CAB, pp. 132–154.
- Pugnaire, F.I., Morillo, J.A., Peñuelas, J., Reich, P.B., Bardgett, R.D., Gaxiola, A., *et al.* (2019) Climate change effects on plant-soil feedbacks and consequences for biodiversity and functioning of terrestrial ecosystems. *Sci Adv* **5**: eaaz1834.
- Riaz, S., Pap, D., Uretsky, J., Laucou, V., Boursiquot, J.-M., Kocsis, L., and Andrew Walker, M. (2019) Genetic diversity and parentage analysis of grape rootstocks. *Theor Appl Genet* **132**: 1847–1860.
- Rolli, E., Marasco, R., Saderi, S., Corretto, E., Mapelli, F., Cherif, A., *et al.* (2017) Root-associated bacteria promote grapevine growth: from the laboratory to the field. *Plant Soil* **410**: 369–382.
- Rolli, E., Marasco, R., Vigani, G., Ettoumi, B., de Angelis, M., Casati, E., *et al.* (2012) The plant growth promoting microbiome increases grapevine resistance to drought stress: a collaborative study between Fondazione Bussolera Bianca, Le Fracce Farm and Milan Universities. *Bull l'OIV* **85**: 189–199.
- Rolli, E., Marasco, R., Vigani, G., Ettoumi, B., Mapelli, F., Deangelis, M.L., *et al.* (2015) Improved plant resistance to drought is promoted by the root-associated microbiome as a water stress-dependent trait. *Environ Microbiol* **17**: 316–331.
- Rosenberg, E., Zilber-Rosenberg, I., Manwani, D., Mortha, A., Xu, C., Faith, J., *et al.* (2016) Microbes drive evolution of animals and plants: the hologenome concept. *MBio* **7**: e01395-15.
- Ruan, Y., Wang, T., Guo, S., Ling, N., and Shen, Q. (2020) Plant grafting shapes complexity and co-occurrence of rhizobacterial assemblages. *Microb Ecol* **80**: 643–655.
- Saleem, M., Hu, J., and Jousset, A. (2019) More than the sum of its parts: microbiome biodiversity as a driver of plant growth and soil health. *Annu Rev Ecol Syst* **50**: 145–168.
- Saleem, M., Law, A.D., Sahib, M.R., Pervaiz, Z.H., and Zhang, Q. (2018) Impact of root system architecture on rhizosphere and root microbiome. *Rhizosphere* **6**: 47–51.
- Sánchez-Cañizares, C., Jorrin, B., Poole, P.S., and Tkacz, A. (2017) Understanding the holobiont: the interdependence of plants and their microbiome. *Curr Opin Microbiol* **38**: 188–196.
- Santoyo, G., Moreno-Hagelsieb, G., del Carmen Orozco-Mosqueda, M., and Glick, B.R. (2016) Plant growth-promoting bacterial endophytes. *Microbiol Res* **183**: 92–99.
- Sasse, J., Martinoia, E., and Northen, T. (2018) Feed your friends: do plant exudates shape the root microbiome? *Trends Plant Sci* **23**: 25–41.
- Semchenko, M., Barry, K.E., Vries, F.T., Mommer, L., Moora, M., and Maciá-Vicente, J.G. (2022) Deciphering the role of specialist and generalist plant-microbial interactions as drivers of plant-soil feedback. *New Phytol*. doi: [10.1111/nph.18118](https://doi.org/10.1111/nph.18118)
- Singh, B.K., Trivedi, P., Egidi, E., Macdonald, C.A., and Delgado-Baquerizo, M. (2020) Crop microbiome and sustainable agriculture. *Nat Rev Microbiol* **18**: 601–602.
- Swift, J.F., Hall, M.E., Harris, Z.N., Kwasniewski, M.T., and Miller, A.J. (2021) Grapevine microbiota reflect diversity among compartments and complex interactions within and among root and shoot systems. *Microorganisms* **9**: 92.
- Tandonnet, J.-P., Cookson, S.J., Vivin, P., and Ollat, N. (2009) Scion genotype controls biomass allocation and root development in grafted grapevine. *Aust J Grape Wine Res* **16**: 290–300.
- Toju, H., Peay, K.G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., *et al.* (2018) Core microbiomes for sustainable agroecosystems. *Nat Plants* **4**: 247–257.
- Trivedi, P., Mattupalli, C., Eversole, K., and Leach, J.E. (2021) Enabling sustainable agriculture through understanding and enhancement of microbiomes. *New Phytol* **230**: 2129–2147.
- Turner, T.R., James, E.K., and Poole, P.S. (2013) The plant microbiome. *Genome Biol* **14**: 1–10.
- van Dam, N.M., and Bouwmeester, H.J. (2016) Metabolomics in the rhizosphere: tapping into belowground chemical communication. *Trends Plant Sci* **21**: 256–265.

- Van Der Heijden, M.G.A.A., and Schlaeppi, K. (2015) Root surface as a frontier for plant microbiome research. *Proc Natl Acad Sci U S A* **112**: 2299–2300.
- Vandenkoornhuysen, P., Quaiser, A., Duhamel, M., Le Van, A., and Dufresne, A. (2015) The importance of the microbiome of the plant holobiont. *New Phytol* **206**: 1196–1206.
- Vega-Avila, A.D., Gumiere, T., Andrade, P.A.M., Lima-Perim, J.E., Durrer, A., Baigori, M., et al. (2015) Bacterial communities in the rhizosphere of *Vitis vinifera* L. cultivated under distinct agricultural practices in Argentina. *Antonie Van Leeuwenhoek* **107**: 575–588.
- Venema, J.H., Giuffrida, F., Paponov, I., Albacete, A., Pérez-Alfocea, F., and Dodd, I.C. (2017) Rootstock–scion signaling: key factors mediating scion performance. In *Vegetable Grafting: Principles and Practices*. Wallingford: CAB, pp. 94–131.
- Vink, S., Chrysargyris, A., Tzortzakis, N., and Salles, J.F. (2021a) Bacterial community dynamics varies with soil management and irrigation practices in grapevines (*Vitis vinifera* L.). *Appl Soil Ecol* **158**: 103807.
- Vink, S., Dini-Andreote, F., Höfle, R., Kicherer, A., and Salles, J. (2021b) Interactive effects of scion and rootstock genotypes on the root microbiome of grapevines (*Vitis* spp. L.). *Appl Sci* **11**: 1615.
- Vitolo, N., Lemos, W.J.F., Calgaro, M., Confalone, M., Felis, G.E., Zapparoli, G., and Nardi, T. (2019) Bark and grape microbiome of *Vitis vinifera*: influence of geographic patterns and agronomic management on bacterial diversity. *Front Microbiol* **10**: 1–12.
- Vives-Peris, V., de Ollas, C., Gómez-Cadenas, A., and Pérez-Clemente, R.M. (2020) Root exudates: from plant to rhizosphere and beyond. *Plant Cell Rep* **39**: 3–17.
- Wallis, C.M., Wallingford, A.K., and Chen, J. (2013) Grapevine rootstock effects on scion sap phenolic levels, resistance to *Xylella fastidiosa* infection, and progression of Pierce's disease. *Front Plant Sci* **4**: 1–9.
- Wang, Y., Naumann, U., Wright, S.T., and Warton, D.I. (2012) mvabund – an R package for model-based analysis of multivariate abundance data. *Methods Ecol Evol* **3**: 471–474.
- Warschewsky, E.J., Klein, L.L., Frank, M.H., Chitwood, D.H., Londo, J.P., Von Wettberg, E.J.B., and Miller, A.J. (2016) Rootstocks: diversity, domestication, and impacts on shoot phenotypes. *Trends Plant Sci* **21**: 418–437.
- Weiss, S., Xu, Z.Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., et al. (2017) Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome* **5**: 27.
- Whiting, J. (2003) *Selection of Grapevine Rootstocks and Clones*. Department of Primary Industries. College of Agricultural Sciences Cooperative Extension, PennState.
- Zarraonandia, I., Owens, S.S.M., Weisenhorn, P., West, K., Hampton-Marcell, J., Lax, S., et al. (2015) The soil microbiome influences grapevine-associated microbiota. *MBio* **6**: e02527-14.
- Zhalnina, K., Louie, K.B., Hao, Z., Mansoori, N., Da Rocha, U.N., Shi, S., et al. (2018) Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly. *Nat Microbiol* **3**: 470–480.
- Zhang, Z., Lu, Y., Wei, G., and Jiao, S. (2022) Rare species-driven diversity–ecosystem multifunctionality relationships are promoted by stochastic community assembly. *MBio*: e0044922. doi:10.1128/mbio.00449-22
- Zilber-Rosenberg, I., and Rosenberg, E. (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* **32**: 723–735.

### Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

**Appendix S1.** Supporting Information.

**Supplementary Table S3.** Number of sequences and OTUs detected after quality trimming, exclusion of non-target co-amplicons and unclassified reads for bacterial and fungal communities associated with bulk soil, rhizosphere and root tissue. Refer to Marasco et al 2021\_Supplementary Table S3.

**Supplementary Table S8.** List of OTUs and related variance detected by multivariate generalized linear model for discriminating bacterial and fungal communities across compartments. The assigned taxonomy of each OTU is displayed at the phylum and class level. Refer to Marasco et al 2021\_Supplementary Table S8.

**Supplementary Table S9.** List of OTUs and related variance detected by multivariate generalized linear model for discriminating bacterial and fungal communities across rootstock–scion combinations in each compartment. The assigned taxonomy of each OTU is displayed at the class level. Refer to Marasco et al 2021\_Supplementary Table S9.

**Supplementary Table S10.** List of bacterial and fungal indicator OTUs of rootstock–scion combinations in root tissues and rhizosphere compartments. Refer to Marasco et al 2021\_Supplementary Table S10.