

RESEARCH ARTICLE

Diagnostic performance of GeneXpert in tuberculosis–HIV co-infected patients at Asella Teaching and Referral Hospital, Southeastern Ethiopia: A cross sectional study

Abebe Sorsa ^{*}, Muhammedawel Kaso

Arsi University College of Health Science and Referral Hospital, Asella, Ethiopia

^{*} nathanabebe08@gmail.com

Abstract

Background

GeneXpert is a new introduction in the diagnostic modality to fight tuberculosis (TB) among people living with HIV (PLHIV) under the program of intensified TB case finding. This study aimed to evaluate the diagnostic performance of GeneXpert under the program of intensified TB cases finding among PLHIV.

Methods

Cross-sectional study was conducted by recruiting individuals attending an HIV clinic from February 2018 to January 2019. Data on clinical parameters were collected using a standardized tool. Two-morning sputum samples were collected and processed for smear microscopy and GeneXpert. SPSS 21 used for data analysis. Proportion, percentage, and mean with SD were used to describe variables. Univariate and multivariable logistic regressions were used to assess factors associated with the GeneXpert. Values for which the 95% CI interval not includes 1 and for which $P < 0.05$ were considered significant.

Result

A total of 384 presumptive TB-HIV co-infection cases were included, of which 166 (43%) were diagnosed to have TB. Fifty-four (32.5%) TB cases were smear AFB positive while 79 (47.7%) TB cases were GeneXpert positive. The GeneXpert detection rate was almost two-fold of that of smear microscopy and all smear positive TB cases were detected by GeneXpert. Moreover, GeneXpert was able to detect an additional third of TB confirmed cases among smear AFB negative cases. Advanced stage of the disease, high viral load and presence of anemia were significantly associated with TB. The WHO TB screening tool remained least sensitive with the lowest positive predictive value.

OPEN ACCESS

Citation: Sorsa A, Kaso M (2021) Diagnostic performance of GeneXpert in tuberculosis–HIV co-infected patients at Asella Teaching and Referral Hospital, Southeastern Ethiopia: A cross sectional study. PLoS ONE 16(1): e0242205. <https://doi.org/10.1371/journal.pone.0242205>

Editor: Frederick Quinn, The University of Georgia, UNITED STATES

Received: February 28, 2020

Accepted: October 28, 2020

Published: January 27, 2021

Copyright: © 2021 Sorsa, Kaso. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data underlying the study are within the manuscript and its [Supporting Information](#) files.

Funding: The author(s) received no specific funding for this work

Competing interests: No authors have competing interests.

Conclusion

GeneXpert demonstrated two-fold case detection rate compared to the sputum smear microscopy and additional third TB case detection rate among smear AFB negative cases. Clinical screening tool for evaluation of TB-HIV co-infection showed poor performance in TB case notification.

Background

Tuberculosis (TB) is among the most widespread and serious of all human infectious diseases. Owing to widespread poverty, inequity and conflict, suboptimal health services in many countries and the impact of HIV/AIDS pandemic, there are more cases of TB today than at any time previously in human history. Close to 95% of all cases, and 98% of deaths due to TB, occur in tropical countries [1]. TB is the most frequent life-threatening opportunistic infection and a leading cause of death among PLHIV. HIV-positive people with latent TB infection have a 10% annual and 50% lifetime risk of developing active TB disease [2]. Similarly, about 79% of HIV-positive clients were screened for TB; of these 11% were found to have active TB. World Health Organization (WHO) recently introduced the algorithm of intensified TB case finding among PLHIV [3–5]. WHO recommends that TB case findings should be intensified in all human immunodeficiency virus (HIV) testing and counseling services for HIV-positive clients by using a set of simple questions for early identification of TB suspects [6–8]. Diagnosis of TB is challenging in HIV-positive individuals, especially when the stage of the disease is advanced. Standard TB diagnostic approaches and clinical algorithms should be followed to guide the diagnosis of TB in PLHIV. Sputum smear Acid fast bacilli (AFB)- microscopy has been in clinical use for decades for the diagnosis of TB, and sensitivity varies between 30–70%. In individuals co-infected with HIV, it varies between 20% and 50% [9–11]. The development of GeneXpert represents a paradigm shift in the diagnosis of TB and drug-resistant. It simplifies molecular testing by fully integrating and automating the three processes required for real-time PCR-based molecular testing (that is, specimen preparation, amplification, and detection) and it detects both live and dead bacteria [9]. GeneXpert detects *M. tuberculosis* as well as mutation that confer rifampicin resistance using three specific primers and five unique molecular probes to ensure a high degree of specificity [10]. The assay provides results directly from sputum in less than 2 hours [9, 10]. The sensitivity of the GeneXpert test is much better than smear microscopy test and comparable to solid culture. Previous studies reported the sensitivity and specificity of GeneXpert to be 72–77% and 99% in smear-negative adults and 98–99% and 99–100% in smear-positive adults, respectively. The negative predictive value (NPV) was 90.6%, and the positive predictive value (PPV) was 100%. Limit of detection is 5 genome copies of purified DNA per reaction or 131 colony forming units/ mL in *M. tuberculosis* spiked sputum in comparison of identification of TB bacilli by microscopic examination requires at least 10,000 bacilli per mL of sputum [9–16]. In spite of early initiation of highly active antiretroviral therapy (HAART), TB remains a significant cause of morbidity and mortality among people living with HIV infection in Sub-Saharan Africa [10]. In Ethiopia, even though more than 400,000 HIV-positive people were enrolled in chronic HIV care with an annual enrolment of around 30,000 newly diagnosed HIV positives in 2016, only 0.4% of them were diagnosed to have active TB while only 40% of the newly enrolled HIV positives were reported to access preventive therapy as per national recommendation [15–17]. This implies a large segment of HIV-positive TB cases did not reach care. Prompt and accurate diagnosis of TB and timely

initiation of appropriate treatment play a pivotal role in reducing the suffering of the patients and contributes a remarkable role in halting TB transmission [5, 6].

The national TB-HIV collaborative work emphasized on three I's which include Intensive case findings, INH Preventive Therapy, and Infection control to mitigate the morbidity and mortality associated with TB-HIV co-infection. The Ethiopia national TB prevention and control program stressed the intensified TB case-finding approach by devising simple screening symptoms that include any one of the following (current cough, weight loss, and fever). Limited information exists in Ethiopia regarding the performance of GeneXpert under the program for intensified TB case finding among PLHIV [5]. Therefore, this study was aimed to evaluate the diagnostic performance of GeneXpert in TB- HIV co-infected patients at Asella Teaching and Referral Hospital in Southeastern Ethiopia.

Methods and material

Ethical consideration

Research proposal was initially submitted to Arsi University College of Health Science Research and Ethical Review Board (IRB) and ethical approval was obtained to pursue the study with reference number of AU_TB_10/2017. After briefing about the purpose and benefit of the study, a consent form was signed by the patient before commencing data collection. Confidentiality of information about the patients' medical problem was secured with maximum effort and was not exposed to any other third party.

Study setting

The study was conducted at Arsi University Asella Teaching and Referral Hospital one of the referral hospitals in the South east part of the country, delivering service for close to 3–4 million people. The region is one of the highest TB prevalent area, and Asella Teaching and Referral Hospital has been reporting more than 1000 annual TB cases [7, 18]. Hence, the Hospital was given priority during the initial phase of GeneXpert service roll out program and equipped with the machine in 2016 [7, 18]. During the first three years, the diagnostic service of this new technology was indicated for special group patients which include TB-HIV co-infection, childhood TB and suspected MDR-TB but since mid of 2019, GeneXpert became the diagnostic of choice for all suspected TB cases despite frequent interruption of the service because of shortage of cartridge [7]. The Hospital has been also one of the treatment centers for PLHIV for more than a decade and at the study time there were about 3560 PLHIV on follow-up at the infectious disease clinic of the hospital as reported from HIV care, treatment and control registry book.

Study design

We conducted a Hospital based cross-sectional study. The source populations were individuals attending an HIV clinic from February 2018 to January 2019 at Asella Teaching and Referral Hospital.

Inclusion criteria. Patients fulfilling the revised national/WHO screening tool for TB assessment among PLHIV were included [5, 7]. As per the National/WHO TB screening tool among PLHIV, we included all patients who fulfilled one of these clinical features (current cough, fever or weight loss) during their regular follow up visit.

Exclusion criteria. Those patients already diagnosed to have TB and receiving anti-TB medications at the time of the survey and patients fulfill clinical definition but who couldn't give sputum for examination were excluded.

Data collection tools and procedures

Patients who fulfilled inclusion criteria were interviewed for socio-demographic variables and evaluated for clinical parameters. Data were annexed to the structured check list tool. Then patients were asked to provide two morning spot sputum specimens (>2mL each) for AFB microscopy and GeneXpert with 1 hour apart. Sputum samples were not induced, but collection was directly observed by one investigator. Smear microscopy was performed by two experienced laboratory technologists throughout the study period. Direct smear microscopy for AFB was performed using Ziehl-Neelsen staining using LED microscopy as per SOP [7, 18]. The test results were reported in accordance with WHO/International Union Against TB and Lung Diseases recommendations [2]. Whenever there were discordant smear AFB reports, the findings were correlated with other clinical findings and GeneXpert reports results to make a final decision.

After sputum sample smear microscopy was used, the remainder of the sputum specimen was tested for GeneXpert v 4.3 (Cepheid, Sunnyvale, CA) according to manufacturer's instructions and national implementation Guideline for GeneXpert MTB/RIF Assay in Ethiopia [5]. Interpretation, Case definitions, Registration, and Management of TB and RR cases based on Xpert MTB/RIF test results were per the WHO (2013) update on "definitions and reporting framework for Tuberculosis" [5]. The laboratory technologists who performed the two tests (smear AFB microscopy and GeneXpert) were independent and were blinded for the results of each test.

Data processing and analysis

Data were coded and entered to EPI-info version 7.1.1 for clean-up and then exported to SPSS 21 for further analysis. Descriptive statistics like frequency, percentage, proportion, and mean with standard deviation (SD), median with interquartile range (IQR) were used. Univariate logistic regression was used to assess factors associated with the yield of sputum microscopy and GeneXpert. Variables with p-value of <0.2 were further analyzed using multivariable regression. Frequencies were calculated for categorical variables and mean with SD and median with IQR were calculated for continuous variables. The factors were considered to be associated when p value was <0.05 and or when 95% confidence interval did not include 1

Operational definition [6–8, 18]

Compatible chest X-ray with pulmonary TB. Chest X-ray report with any of the following finding; bronchiectasis, consolidation, patchy, cavitory, and miliary TB.

TB cases. All forms of TB which include both pulmonary and disseminated TB.

Pulmonary TB. A form of TB contained only to the lung.

Disseminated TB. A clinical spectrum of TB involving lung and one or more of other organ systems.

Presumptive TB case. Any patient who fulfills the national screening for clinical symptoms of TB.

Probable TB cases. If patient met two or more of the following criteria; history of tuberculous contact, clinical features suggestive of TB, reactive tuberculin, skin test >10 mm, or radiographic findings compatible with TB (miliary disease, cavitory lesions, hilar lymphadenopathy, or primary complex), no response to broad spectrum antibiotic trial.

Results

A total of 384 presumptive TB-HIV co-infection cases were enrolled. Males accounted for 199 (51.9%) and the mean age of patient was 37±12.4 years. Most patients were WHO HIV/AIDS

clinical stage II and stage III, each accounting for 119 (30.9%) and 130 (33.9%), respectively. The median CD₄ count at the time of TB consideration was 256 cell/ml with IQR of (129–452), and about 104 (38.2%) had CD₄ cell count of less than 200 cell/ml. Of those patients who had HIV viral load determination during the last six months at the time of TB screening, 134 (78.8%) had viral suppression less than 1000copies/ml. Cough as positive screening symptoms was reported by 372 (97.1%) patients while fever was reported by 298 (77.8%) patients (Table 1). All patients have been on antiretroviral therapy (ART) with duration of ART treatment ranging from 0–132 months with a median duration of 52 months. A total of 166 (43.2%) TB cases were diagnosed during the study period and 79 (49.0%) TB cases were bacteriologically confirmed either with GeneXpert or sputum AFB.

Concordance of GeneXpert assay and sputum acid-fast smear (AFB)

The case notification rates of both sputum AFB and GeneXpert was evaluated among presumptive TB cases and the case detection rate of sputum smear AFB was 51 (13.3%) while that of GeneXpert was almost two-fold higher, 79(20.6%). GeneXpert increased the TB detection rate by 25 cases (31.6%) compared with sputum smear AFB. The agreement between the two tests was found to be substantial with *kappa* value of 0.74.

Sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of national/WHO symptom screening tool of TB among PLHIV

The sensitivity, specificity, PPV, and NPV of national/WHO screening tool of TB among PLHIV patients varied based on whether the tools were used singly, or in combination; if cough used as sole screening tool the sensitivity, specificity, PPV and NPV were found to be

Table 1. Demographic and some clinical data of presumptive TB-HIV co-infection patients among people living with HIV, Asella Teaching and Referral Hospital 2019. (n = 384).

Characteristics	N (%)		
Sex	Male	199(51.8)	
	Female	185(48.2)	
Age category(years)	= <18 years	45(11.7)	
	18–60 years	324(84.4)	
Clinical characteristics at base line	Cough	372(97.1)	
	Fever	298(77.8)	
	Weight loss	257(67.1)	
	Previous TB treatment	84(21.9)	
	INH prophylaxis	161(42.0)	
	CPT prophylaxis	302(78.9)	
	WHO clinical stages	III	53(13.8)
		I	119(30.9)
II		130(33.9)	
Laboratory	Hematocrit	Normal	224(59.4)
		Anemia	153(40.6)
	CD4 count	<200cell/ml	104(38.2)
		> = 200cell/ml	168(61.8)
	HIV viral load	<1000copy/ml	39(48.1)
		> = 1000copy/ml	42(51.9)

<https://doi.org/10.1371/journal.pone.0242205.t001>

3.6%, 100%, 21.2% and 100% respectively. When cough and fever were combined as screening tool, the sensitivity, specificity, PPV, and NPV became 25.5%, 84.5%, 22.6%, and 86.8% respectively, and if the three screening symptoms were positive, the sensitivity, specificity, PPV and NPV showed 44.4%, 60.0%, 21.8% and 80.4% respectively (Table 2).

Clinical factors associated with results of GeneXpert and smear AFB

Using univariate analysis, body mass index (BMI), duration of cough, chronic medical illness, previous TB treatment, isoniazid preventive therapy (IPT), WHO clinical staging of HIV/AIDS were positively associated with the result of both sputum AFB microscopy and GeneXpert (Table 3). However, using multivariable logistic regression, longer duration of cough, advanced stage of the disease, presence of anemia, and high HIV viral load were significantly associated with positive sputum AFB microscopy and GeneXpert results (Table 4). Lower BMI was associated with positive sputum AFB microscopy but not statistically associated with positive GeneXpert results.

Patients who have never taken IPT during their course of HIV illness were at significant risk of developing TB, inferring the preventive efficacy of IPT to be 78%.

Laboratory finding associated with yield of GeneXpert and smear AFB

Using univariate logistic regression: anemia, increases in erythrocyte sedimentation rate (ESR), CD4 cell count, HIV Viral load were associated with positive results of GeneXpert and smear AFB. However, using multiple logistic regression, anemia and high HIV viral load were significantly associated with positive sputum AFB microscopy and GeneXpert (Table 4).

Discussion

Our study focused on the role of GeneXpert in TB case notification under the program of WHO intensified TB case finding among PLHIV. GeneXpert is a relatively a new diagnostic modality in the battle to combat TB as a global public health problem, and is believed that it is to improve bacteriology confirmed TB cases with shorter turnaround time as shown in previous studies [1, 4]. In this study, the case detection rate of GeneXpert was compared with the conventional smear microscopy and revealed that it showed almost two-fold case detection rate. All patients who turned positive for smear AFB were also positive for GeneXpert, implying that it can replace the conventional smear AFB microscopy in the clinical care of TB [9, 12, 13]. GeneXpert detected additional 31.6% among smear AFB negative cases which is congruent with the finding of the study from north-western part of Ethiopia and other multicenter studies [1, 4, 7, 13]. However, the study finding reported by Habte et al indicated additional TB case detection of GeneXpert among smear AFB negative cases was 64.3% [19], which was higher than our finding while the report from Cochrane review

Table 2. Sensitivity, specificity, PPV and NPV TB screening tool among PLHIV, Asella Teaching and Referral Hospital 2019.

Screening tools	GeneXpert confirmed TB cases, N = 79			
	Sensitivity	Specificity	PPV	NPV
Cough +fever+ weight loss	44.4%	60.0%	21.8%	80.4%
Cough + fever	25.5%	84.5%	22.6%	86.8%
Cough +weight loss	34.5%	66.7%	20.7%	80.2%
Cough only	3.6%	100%	21.2%	100%

NPV-negative predictive value, PLHIV-people living with HIV, PPV-positive predictive value

<https://doi.org/10.1371/journal.pone.0242205.t002>

Table 3. Clinical and laboratory findings among patients with sputum AFB microscopy and GeneXpert positive results, Asella Teaching and Referral Hospital 2019.

Variables	Smear AFB positive TB cases, N = 54		GeneXpert positive = 79	
	Number (%)	p-value	Number (%)	P value
Duration of cough (in weeks)	<2	16(26.0)	14(17.7)	0.03
	≥2	38(74.0)	65(82.3)	
Previous TB treatment	Yes	17(31.5%)	22(27.8)	0.17
	No	37(68.5%)	57(72.2)	
Chronic medical illness	Yes	31(57.4%)	31(39.2)	0.001
	No	23(42.6)	48(60.8)	
Weight loss	Yes	29(53.7)	43(54.4)	0.58
	No	26(46.3)	36(45.6)	
IPT*	Yes	11(18.7)	17(22.7)	<0.001
	No	43(81.3)	62(77.2)	
WHO* clinical staging of HIV/AIDS	III	40(74.1)	43(62.3)	0.18
	IV	14(25.9)	36(37.7)	
Anemia	Yes	45(83.3)	62(78.5)	0.001
	No	9(16.7)	17(21.5)	
CD ₄ count(cell/ml)	<200	26(48.4)	67(85)	<0.001
	≥200	28(51.6)	12(15)	
HIV Viral load(copy/ml)	<1000	13(25)	60(75.7)	0.013
	≥1000	41(75.0)	19(24.3)	
BMI (kg/m ²)	<18.5	33(61.0)	47(60)	0.17
	≥18.5	21(39.0)	32(40)	

ESR = erythrocyte sedimentation rate, IPT = Isoniazid preventive therapy, BMI = body mass index, WHO = world health organization, hr = hours, AFB = Acid fast bacilli, AIDS = acquired immunodeficiency syndrome, HIV-human immunodeficiency virus, TB = tuberculosis

<https://doi.org/10.1371/journal.pone.0242205.t003>

and meta-analysis by Steingart and his colleagues, and studies from other areas were in the range of quarter which were lower than our finding [5, 6, 9, 14, 17, 20–25]. The difference could be related with enrollment criteria of study participants in which our study was used the revised National/WHO screening tool protocol for TB-HIV co-infection among PLHIV which tended to indicate very low positive predictive value.

The sensitivity, specificity, PPV, and NPV of the revised national/WHO screening tool were evaluated in our study [6–8, 18]. Accordingly, if a cough is used as a sole screening tool of TB, the sensitivity is very poor implying that patients who were coughing were not actually having the disease. However, the specificity of these screening tools was very good, indicating that most patients who were not coughing or having fever were less likely to have TB. The good specificity of the screening tool for TB is within the recommended minimum proposed specificity requirement of 70% proposed by the WHO for a TB triage test [4, 8]. This strategy has been limited in part by a low sensitivity of symptom screening leading to a large number of patients were in need of tests even though the results in most cases turned to be negative resulting in resource exploitation and manpower burden. The argument here is that it is more important not to miss even a single case of TB at an expense of resource and manpower exploitation [8].

In our study, patients who were having anemia were found to be highly likely of having TB which could be explained by the actual presence of chronic illness which actually causes anemia of chronic illness. Similar findings were reported by Temesgen et al and other reports [3, 4].

Table 4. Univariate and multivariate logistic regression assessing factors associated with the result of GeneXpert and smear AFB, Asella Teaching and Referral Hospital 2019.

		Smear positive TB cases, N = 54		GeneXpert positive, N = 79	
		COR, 95%CI	AOR, 95%CI	COR (95%CI)	AOR (95%CI)
BMI* (kg/m ²)	<18.5	1	1	1	1
	≥18.5	1.97(1.27–4.44)	2.00(1.07–3.70)	1.74(1.14–2.65)	1.051(0.67–3.70)
Duration of cough (in weeks)	<2	1	1	1	1
	≥2	1.28(1.16–2.47)	2.55(1.96–5.53)	1.31(0.67–2.57)	2.39(1.28–4.48)
Previous TB treatment	Yes	2.08(1.21–3.58)	1.22(0.76–1.95)	1.78(1.05–3.02)	1.44(0.89–2.31)
	No	1		1	1
Chronic medical illness	Yes	1.95(1.13–4.61)	2.42(1.24–5.73)	3.63(1.33–8.08)	2.688(1.41–6.92)
	No	1	1	1	1
IPT*	Yes	1	1	1	1
	No	5.7(3.2–10.2)	7.5(4.5–12.6)	3.34(1.62–6.90)	3.21 (1.79–5.75)
WHO* clinical staging of AIDS	III	1	1	1	1
	IV	10.21(1.35–77.4)	6.32(1.35–35.43)	10(1.35–100.00)	25(3.33–90.91)
Anemia	Yes	4.33(2.05–9.18)	4.71(2.10–10.58)	4.88(2.22–10.69)	3.528(2.10–6.31)
	No	1	1	1	1
CD ₄ count (cell/ml)	<200	1.48(0.69–3.17)	1.05(0.45–2.48)	2.25(0.69–7.34)	1.73(0.63–4.80)
	>= 200	1	1	1	1
HIV Viral load(copy/ml)	<1000	1	1	1	1
	≥1000	0.67(0.24–1.87)	1.13(0.31–4.15)	2.67(1.08–6.583)	1.87(1.21–4.085)

ESR = erythrocyte sedimentation rate, IPT = Isoniazid preventive therapy, BMI = body mass index, WHO = world health organization, hr = hours, AFB = Acid fast bacilli, AIDS = acquired immunodeficiency syndrome, HIV-human immunodeficiency virus, TB = tuberculosis

<https://doi.org/10.1371/journal.pone.0242205.t004>

Our study also concluded that HIV viral load above 1000copy/ml was significantly associated with the occurrence of TB while CD4 has no association with TB diagnosis, implying that HIV viral load was more reliable indicator of predicting the degree of HIV/AIDS disease status and opportunistic infection [19, 20, 26–28].

Other clinical factors like advanced WHO clinical stage of HIV/AIDS was significantly associated with the development of TB, implicating that the higher the stage of the disease, the weaker the immune system to contain the bacilli leading to flare-up of the disease. Consistent findings were reported from other studies [15, 29–31].

One of the strategies to prevent the development of TB among PLHIV is the provision of isoniazid preventive therapy (IPT). In our report, among the presumptive TB cases, a total of 161(42%) PLHIV took IPT. The risk of developing TB among patients who didn't take IPT was five-fold which is comparable with reports from other studies [15, 32].

In conclusion, the introduction of GeneXpert significantly improved the case detection of TB compared to the conventional smear microscopy and it additionally detected third TB cases among smear AFB negative presumptive HIV-TB co-infection. Advanced stage of the disease and poorly suppressed viral load were the two independent factors predicting the results of GeneXpert. Using symptom based screening of TB-HIV co-infection showed poor validity in TB case notification.

Supporting information

S1 File.

(BIN)

S1 Data.
(XLS)

Acknowledgments

We are very much grateful for patients attending follow-up at Asella Teaching and Referral Hospital ART clinic for their unreserved cooperation during the study. Next, we are equally grateful to the ART health professional team for their continuous support during data collection. Lastly but not least, we would like to extend heartfelt gratitude to Arsi University for allowing us to carry out the research.

Author Contributions

Conceptualization: Abebe Sorsa.

Data curation: Abebe Sorsa.

Formal analysis: Abebe Sorsa, Muhammedawel Kaso.

Funding acquisition: Abebe Sorsa, Muhammedawel Kaso.

Investigation: Abebe Sorsa.

Methodology: Abebe Sorsa, Muhammedawel Kaso.

Project administration: Abebe Sorsa.

Resources: Abebe Sorsa, Muhammedawel Kaso.

Software: Abebe Sorsa, Muhammedawel Kaso.

Supervision: Abebe Sorsa.

Validation: Abebe Sorsa.

Visualization: Abebe Sorsa, Muhammedawel Kaso.

Writing – original draft: Abebe Sorsa, Muhammedawel Kaso.

Writing – review & editing: Abebe Sorsa.

References

1. Maynard-smith L, Larke N, Peters JA, Lawn SD. Diagnostic accuracy of the Xpert MTB / RIF assay for extra pulmonary and pulmonary tuberculosis when testing non-respiratory samples: a systematic review. *BMC Infect Dis.* 2014; 14:709. <https://doi.org/10.1186/s12879-014-0709-7> PMID: 25599808
2. Habte D, Melese M, Hiruy N, Gashu Z, Jerene D, Moges F, et al. The additional yield of Gene Xpert MTB / RIF test in the diagnosis of pulmonary tuberculosis among household contacts of smear positive TB cases. *Int J Infect Dis.* 2016; 49: 179–84. <https://doi.org/10.1016/j.ijid.2016.07.002> PMID: 27401584
3. Temesgen B, Kibret GD, Alamirew NM, Melkamu MW, Hibstie YT, Petrucka P, et al. Incidence and predictors of tuberculosis among HIV-positive adults on antiretroviral therapy at Debre Markos referral hospital, Northwest Ethiopia: a retrospective record review. *BMC Public Health.* 2019; 19(1):1566. <https://doi.org/10.1186/s12889-019-7912-9> PMID: 31771552
4. Rasool G, Khan AM, Mohy-ud-din R, Riaz M. Detection of Mycobacterium tuberculosis in AFB smear-negative sputum specimens through MTB culture and GeneXpert MTB / RIF assay. *Int J Immunopathol Pharmacol.* 2019; 33. Available from: <https://doi.org/10.1177/20587384198271>
5. Dravid A, Natarajan K, Medisetty M, Gawali R, Mahajan U, Kulkarni M, et al. Incidence of tuberculosis among HIV infected individuals on long term antiretroviral therapy in private healthcare sector in Pune, Western India. *BMC Infect Dis.* 2019; 19(1):1–12. <https://doi.org/10.1186/s12879-018-3567-x> PMID: 30606108
6. Ethiopia federal minister of health: National comprehensive HIV prevention, care and treatment in Ethiopia 5th ed, 2018.

7. Ethiopia federal minister of health: Guidelines for clinical and programmatic management of TB, leprosy and TB/HIV in Ethiopia 6th ed, 2018.
8. WHO: Guidelines for intensified tuberculosis case-finding and isoniazid preventive therapy for people living with HIV in resource constrained settings, 2011.
9. Pinyopornpanish K, Chaiwarith R, Pantip C, Keawwichit R, Wongworapat K, Khamnoi P, et al. Comparison of Xpert MTB / RIF assay and the conventional sputum microscopy in detecting *Mycobacterium tuberculosis* in Northern Thailand. *Tuberculosis Research and Treatment*. 2015; 2015. <https://doi.org/10.1155/2015/571782> PMID: 26064681
10. Ejeta E, Beyene G, Bonga Z, Abebe G. Xpert MTB/RIF assay for the diagnosis of *Mycobacterium tuberculosis* and Rifampicin resistance in high Human Immunodeficiency Virus setting in Gambella regional state, Southwest Ethiopia. *J Clin Tuberc Other Mycobact Dis* [Internet]. 2018; 12(June):14–20. Available from: <https://doi.org/10.1016/j.jctube.2018.06.002>.
11. Steingart KR, Schiller I, Horne DJ, Pai M, Boehme CC. Xpert® MTB/RIF assay for pulmonary tuberculosis and Rifampicin resistance in adults. *Cochrane Database Syst Rev*. 2014. Available from: <https://doi.org/10.1002/14651858.CD009593.pub3> PMID: 24448973
12. Lange B, Khan P, Kalmambetova G, Al-Darraj HA, Alland D, Antonenka U, et al. Diagnostic accuracy of the Xpert® MTB/RIF cycle threshold level to predict smear positivity: A meta-analysis. *Int J Tuberc Lung Dis*. 2017; 21(5): 493–502. <https://doi.org/10.5588/ijtld.16.0702> PMID: 28399963
13. Rasheed W, Rao NA, Adel H, Baig MS, Adil SO. Diagnostic accuracy of Xpert MTB/RIF in sputum smear-negative pulmonary tuberculosis. *Cureus*. 2019; 11(8). <https://doi.org/10.7759/cureus.5391> PMID: 31620317
14. Davis JL, Kawamura LM, Chaisson LH, Grinsdale J, Benhammou J, Ho C, et al. Impact of GeneXpert MTB / RIF on patients and tuberculosis programs in a low-burden setting a hypothetical trial. *Am J Respir Crit Care Med*. 2014; 189: 1551–9. <https://doi.org/10.1164/rccm.201311-1974OC> PMID: 24869625
15. Zeka AN, Tasbakan S, Cavusoglu C. Evaluation of the GeneXpert MTB / RIF assay for rapid diagnosis of tuberculosis and detection of Rifampin resistance in pulmonary and extrapulmonary specimens. *J Clin Microbiol*. 2011; 49(12): 4138–41. <https://doi.org/10.1128/JCM.05434-11> PMID: 21956978
16. Gunda DW, Maganga SC, Nkandala I, Kilonzo SB, Mpondo BC, Shao ER, et al. Prevalence and risk factors of active TB among adult HIV patients receiving ART in Northwestern Tanzania: a retrospective cohort study. *Can J Infect Dis Med Microbiol*. 2018; 2018. <https://doi.org/10.1155/2018/1346104> PMID: 30073038
17. Tadesse M, Aragaw D, Rigouts L, Abebe G. Increased detection of smear-negative pulmonary tuberculosis by GeneXpert MTB/RIF® assay after bleach concentration. *Int J Mycobact* [Internet]. 2016; 5(2):211–8. Available from: <https://doi.org/10.1016/j.ijmyco.2016.03.005> PMID: 27242234
18. Ethiopia federal minister of health: Implementation guideline for GeneXpert MTB/RIF assay in Ethiopia 2014.
19. Akanbi MO, Achenbach C, Taiwo B, Idoko J, Ani A, Isa Y, et al. Evaluation of gene Xpert for routine diagnosis of HIV-associated tuberculosis in Nigeria: a prospective cohort study. *BMC Pulm Med*. 2017; 17(1): 1–10. <https://doi.org/10.1186/s12890-016-0353-7> PMID: 28049457
20. Mollel EW, Maokola W, Todd J, Msuya SE, Mahande MJ. Incidence rates for tuberculosis among HIV infected patients in Northern Tanzania. *Front Public Heal*. 2019; 7(October): 19. <https://doi.org/10.3389/fpubh.2019.00306> PMID: 31709218
21. Kempker RR, Chkhartishvili N, Kinkladze I, Schechter MC, Harrington K, Rukhadze N, et al. High yield of active tuberculosis case finding among HIV-infected patients using Xpert MTB/RIF testing. *Open Forum Infect Dis*. 2019; 6(6): 1–6. <https://doi.org/10.1093/ofid/ofz233> PMID: 31211163
22. Tesfaye B, Alebel A, Gebrie A, Zegeye A, Tesema C, Kassie B. The twin epidemics: prevalence of TB/HIV co-infection and its associated factors in Ethiopia; a systematic review and meta-analysis. *PLoS One*. 2018; 13(10): 1–18. <https://doi.org/10.1371/journal.pone.0203986> PMID: 30281631
23. Glaziou P, Sismanidis C, Floyd K, Raviglione M. Global epidemiology of tuberculosis. *Cold Spring Harb Perspect Med*. 2015; 5: 1–18.
24. Evans CA. GeneXpert—a game-changer for tuberculosis control? *PLoS Medicine*. 2011; 8(7): 8–11. <https://doi.org/10.1371/journal.pmed.1001064> PMID: 21814497
25. Boehme CC, Nicol MP, Nabeta P, Michael JS, Gotuzzo E, Tahirli R, et al. Feasibility, diagnostic accuracy, and effectiveness of decentralized use of the Xpert MTB / RIF test for diagnosis of tuberculosis and multidrug resistance: a multicenter implementation study. *Lancet* [Internet]. 2011; 377(9776):1495–505. Available from: [https://doi.org/10.1016/S0140-6736\(11\)60438-8](https://doi.org/10.1016/S0140-6736(11)60438-8) PMID: 21507477
26. Lawn SD, Kerkhoff AD, Vogt M, Ghebrekristos Y, Whitelaw A, Wood R. Characteristics and early outcomes of patients with Xpert MTB/RIF-negative pulmonary tuberculosis diagnosed during screening

- before antiretroviral therapy. *Clin Infect Dis*. 2012; 54(8): 1071–9. <https://doi.org/10.1093/cid/cir1039> PMID: 22318975
27. Al-darraj HAA, Razak HA, Ng KP, Altice FL. The diagnostic performance of a single GeneXpert MTB / RIF assay in an intensified tuberculosis case finding survey among HIV-infected prisoners in Malaysia. *PLoS One*. 2015; 8(9): 1–10.
 28. Blakemore R, Story E, Helb D, Kop JA, Banada P, Owens MR, et al. Evaluation of the analytical performance of the Xpert MTB/RIF assay. *J Clin Microbiol*. 2010; 48(7): 2495–501. <https://doi.org/10.1128/JCM.00128-10> PMID: 20504986
 29. Balcha TT, Sturegård E, Winqvist N, Skogmar S, Reepalu A, Jemal ZH, et al. Intensified tuberculosis case-finding in HIV-positive adults managed at Ethiopian health centers: diagnostic yield of xpert MTB/RIF compared with smear microscopy and liquid culture. *PLoS One*. 2014; 9(1). <https://doi.org/10.1371/journal.pone.0085478> PMID: 24465572
 30. Albert H, Nathavitharana RR, Isaacs C, Pai M, Denkinger CM, Boehme CC. Development, roll-out and impact of Xpert have we learnt and how can we do better? *Eur Respir J*. 2016; 1–10. Available from: <https://doi.org/10.1183/13993003.00543-2016>.
 31. Pandey P, Pant ND, Rijal KR, Shrestha B, Kattel S, Banjara MR, et al. Diagnostic accuracy of GeneXpert MTB / RIF assay in comparison to conventional drug susceptibility testing method for the diagnosis of multidrug-resistant tuberculosis. *PLoS One*. 2017; 12(1).
 32. Bucher HC, Griffith LE, Guyatt GH, Sudre P, Naef M, Sendi P, et al. Isoniazid prophylaxis for tuberculosis in HIV infection: a meta-analysis of randomized controlled trials *AIDS*. 1999; 13: 501–507. <https://doi.org/10.1097/00002030-199903110-00009> PMID: 10197379