

Review

The Impact of the Ca²⁺-Independent Phospholipase A₂β (iPLA₂β) on Immune Cells

Tayleur D. White †, Abdulaziz Almutairi †, Ying Gai Tusing, Xiaoyong Lei and Sasanka Ramanadham *

Department of Cell, Developmental and Integrative Biology, Comprehensive Diabetes Center, University of Alabama at Birmingham, Birmingham, AL 35294, USA; twhite1@uab.edu (T.D.W.); aziiz@uab.edu (A.A.); yingg@uab.edu (Y.G.T.); xlei@uab.edu (X.L.)

* Correspondence: sramvem@uab.edu; Tel.: +1-205-996-5973

† These authors contributed equally.

Abstract: The Ca²⁺-independent phospholipase A₂β (iPLA₂β) is a member of the PLA₂ family that has been proposed to have roles in multiple biological processes including membrane remodeling, cell proliferation, bone formation, male fertility, cell death, and signaling. Such involvement has led to the identification of iPLA₂β activation in several diseases such as cancer, cardiovascular abnormalities, glaucoma, periodontitis, neurological disorders, diabetes, and other metabolic disorders. More recently, there has been heightened interest in the role that iPLA₂β plays in promoting inflammation. Recognizing the potential contribution of iPLA₂β in the development of autoimmune diseases, we review this issue in the context of an iPLA₂β link with macrophages and T-cells.

Keywords: iPLA₂β; macrophages; T-cells; inflammation; eicosanoids; resolvins



Citation: White, T.D.; Almutairi, A.; Tusing, Y.G.; Lei, X.; Ramanadham, S. The Impact of the Ca²⁺-Independent Phospholipase A₂β (iPLA₂β) on Immune Cells. *Biomolecules* **2021**, *11*, 577. <https://doi.org/10.3390/biom11040577>

Academic Editor: Jesús Balsinde

Received: 5 March 2021

Accepted: 12 April 2021

Published: 15 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Phospholipases A₂ (PLA₂s) hydrolyze the *sn*-2 substituent of glycerophospholipids to release a lysophospholipid and a free fatty acid [1]. Among the family of PLA₂s are the group VI Ca²⁺-independent PLA₂s (iPLA₂s), which include iPLA₂β (VIA), iPLA₂γ (VIB), iPLA₂δ (VIC), iPLA₂ε (VID), iPLA₂ζ (VIE), and iPLA₂η (VIF) [2]. The cytosolic iPLA₂β enzyme, along with the membrane-associated iPLA₂γ, are the most studied of the group VI PLA₂s. Because of its emerging link with inflammation, the focus of this review will be on iPLA₂β.

Encoded by *PLA2G6*, the iPLA₂β protein (84–88 kDa) has a serine lipase consensus sequence (GTSGT), contains eight N-terminal ankyrin repeats, an ATP binding cassette (GGGVKG), a caspase-3 cleavage site (DVDT), and two calmodulin (CAM) binding domains [2,3]. Crystal structure studies suggest that iPLA₂β forms a dimer through the interaction of the catalytic domains and that CAM binds to the dimer to cause a closed state, denying the access of substrates to the active site [4]. Relief from this inhibitory state is achieved through activation of calmodulin kinase IIβ, which forms a signaling complex with iPLA₂β [5]. Lipidomics-based LC/MS approaches have revealed that among glycerophospholipids, iPLA₂β exhibited the greatest activity towards 1-palmitoyl-2-arachidonoyl-*sn*-glycero-3-phosphoethanolamine (PAPE) and, among *sn*-2 substituents, a greater selectivity for linoleate or myristate [6,7]. The currently available selective inhibitors of iPLA₂β include the irreversible *S*-BEL (*S*-bromo-enol lactone) and the reversible 1,1,1-trifluoro-6-(naphthalen-2-yl)hexan-2-one (FKGK18) [8,9]. iPLA₂β is widely expressed and has many proposed roles, including signal transduction, and is recognized to contribute to neurodegenerative disorders, cancers, myocardial complications, and metabolic dysfunction (reviewed extensively elsewhere [2,3]).

While localized in the cytosol under basal conditions [10], iPLA₂β mobilizes to subcellular organelles (e.g., the endoplasmic reticulum (ER), mitochondria, nucleus) upon stimulation. Its activation leads to the hydrolysis of the *sn*-2 fatty acid substituent from membrane glycerophospholipids [11–16] to yield a free fatty acid (e.g., arachidonic acid,

eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA)) and a 2-lysophospholipid [17]. Subsequent metabolism of arachidonic acid by cyclooxygenases (COX), lipoxygenase (LOX), and cytochrome P450 (CYP) pathways leads to the generation of bioactive oxidized eicosanoids, several of which are proinflammatory [18] and recognized contributors to inflammatory diseases [19–27]. Some of the most potent inflammatory eicosanoids [21] are prostaglandin E₂ (PGE₂), leukotrienes (LTs), hydroxyeicosatetraenoic acids (HETEs), and dihydroxyeicosatetraenoic acids (DHETEs), and they contribute to inflammation and autoimmune diseases [22]. It is not unexpected that iPLA₂β activation can play critical roles in the initiation and progression of inflammatory pathways, which if not curbed can lead to the evolution of a variety of disorders. In contrast, other unsaturated fatty acids such as EPA and DHA can be metabolized to generate pro-resolving lipids, designated as specialized resolving mediators (SPMs) [28–30]. Among the proinflammatory lipids generated, PGs and LTs are the first to be produced [28]. When there is injury to the tissue, the generated PGs cause pain, swelling, and edema. To resolve the inflammation and clear antigenic debris, SPMs are generated in attempt to restore homeostasis, a process referred to as “lipid mediator class switching” [31,32].

Immune cells express iPLA₂β [33–36] and iPLA₂β-derived lipids (iDLs) contribute to cell proliferation [37], cell cycle progression [38], cell division [39], monocyte migration [40], and superoxide generation [41]. Inhibition of iPLA₂β reduces reactive oxygen species (ROS) generation [42] and is reported to be effective against autoimmune- [43] and inflammation-based [44–47] diseases.

Although extensive literature exists linking a number of secretory PLA₂s (sPLA₂s) or cytosolic PLA₂α (cPLA₂α) with inflammatory responses, only a few studies have described a link between iPLA₂β and macrophages, and even fewer have considered a link between iPLA₂β and T-cells or B-cells. To provide context, in this paper, the basic functionality of these cells is discussed first, followed by a review of the specific impact of iPLA₂β in these immune cells, in the context of different pathophysiologies.

2. Macrophages

Macrophages are recognized to play significant roles in the development of autoimmune diseases, including rheumatoid arthritis (RA), multiple sclerosis (MS), and type-1 diabetes (T1D) [48]. As fundamental components of the innate immune system, they contribute to tissue homeostasis, dead cell and antigen phagocytosis, and crucial induction of adaptive immunity [49]. They do so by recognizing pathogen-associated molecular patterns (PAMPs) of foreign microorganism or damage-associated molecular patterns (DAMPs) from damaged host cells through pattern recognition receptors (PRR). PRRs are found in all innate immune cells including macrophages. Following their recognition, macrophages internalize the PAMPs or DAMPs via phagocytosis and break them down for antigen presentation through major histocompatibility complex II (MHC-II) [50], which is another fundamental role of macrophages [51]. T lymphocytes recognize self-peptides that are presented on the MHC Class-II molecules and expressed mainly by antigen presenting cells (APC) like dendritic cells and macrophages [52]. All tissues have resident macrophages, which emerge from three different developmental stages—the yolk sac, fetal liver, and bone marrow [53,54]. Although fetal-liver and bone marrow-derived monocytes are the main source of macrophages after birth, yolk sac-derived macrophages give rise to microglia and a small fraction of tissue-derived macrophages [55].

Macrophages are essential in both inflammation and healing processes. However, the functionality of macrophages is subject to macrophage polarization, defined as the “M1-M2 paradigm” [56]. Tissue-resident macrophages are mainly of the M2 (anti-inflammatory) phenotype [57] and affect tissue homeostasis and the resolution of inflammation, which also involves debris clearance [58]. Polarization of macrophages to the M2 phenotype is induced by Th2 cytokines, including interleukins 4, 13, and 10 (IL-4, IL-13, and IL-10). This leads to upregulation of arginase-1 (Arg1), chemokine (C-C motif) ligands 17 and 24 (CCL17 and CCL24), and secretion of IL-10 and transforming growth factor beta (TGFβ) [59]. In contrast,

the M1 macrophages express a proinflammatory phenotype and are induced by Th1 cytokines that include interferon gamma (IFN γ) and TNF α . This promotes upregulation of Arg2, iNOS, cyclooxygenase-2 (COX-2), and other proinflammatory factors, leading to increased production of nitric oxide (NO), TNF α , IL-1 α , IL-1 β , IL-6, and IL-12 [60]. Whereas M2-like macrophages represent the majority of tissue-resident macrophages [57,61], M1-like macrophages are found significantly in inflamed tissues [60].

Emerging studies highlight the importance of lipids in macrophage functionality [62,63]. Although macrophage signaling has long been studied in the context of cytokine and chemokine production, recent studies suggest a profound impact of eicosanoids and SPMs on macrophage functions [31,64]. Eicosanoids have been linked to modifying macrophage inflammatory response in multiple diseases [23,65,66]. In humans, PGE₂ induces LOX-class switching from leukotriene B₄ (LTB₄) to lipoxins, which represents a stop signal for polymorphonuclear (PMN) recruitment and initiation of a resolution phase that promotes an anti-inflammatory macrophage phenotype and function (i.e., phagocytosis) [30]. Facilitating the resolution process are SPMs, which counter-regulate the early initiators (PGs and LTs) of acute inflammation, leading to inhibition of proinflammatory cytokines and upregulation of anti-inflammatory cytokines (e.g., IL-10) [29].

3. T-Cells and B-Cells

Lymphocyte development in mammals occurs in the central lymphoid organs such as the bone marrow (B-cell development) and the thymus (T-cell development) [67]. From these organs, T and B lymphocytes continue to migrate to other peripheral lymphoid tissues. Differences in both lymphocyte populations occurs in mature stages, where T-cell development slows down in the thymus, and they divide outside of the central lymphoid organs to maintain the number of mature T-cells [67].

The development of a cluster of either differentiation 4 or differentiation 8 (CD4⁺ or CD8⁺) T-cells depends on the thymocyte maturation process. This involves hematopoietic stem cells in the bone marrow migrating to the thymus, hence the “T” in T-cells, to mature into functional T-cells [68]. The earliest thymocytes are double-negative—they do not express either CD4 or CD8. In the double negative (DN) stage, these cells are negative for T-cell receptor (TCR) expression, are not able to bind to MHC, and do not carry out effector functions. The DN stage occurs in the cortex and has four subsets that are important for the full development of T-cells, designated as DN1–4. DN3 is a critical step in T-cell development and encompasses: (1) thymocytes becoming restricted to a T-lineage and generation of mature $\alpha\beta$ and $\gamma\delta$ T-cells, (2) T-cell receptor β -chain (TCR β) *V(D)J* recombination, (3) expression of functional pre-TCR complex, and (4) thymocytes becoming early pre-T-cells that are committed to the $\alpha\beta$ T-cell lineage. The DN4 cells start to express low levels of CD4 and CD8 and present themselves as double-positive (DP) [68]. During the DP stages, CD4 and CD8 expression levels are upregulated and the cells migrate to the medulla, where a negative selection process confers single positivity for CD4 or CD8.

Naïve CD4 T-cells are destined for four distinct fates that are determined by different cytokine patterns after interaction with antigens. These cytokine signals upregulate distinct transcription factors to generate specific cell populations, which include the T helper cells 1, 2, and 17 (Th1, Th2, Th17), and regulatory T-cells (Tregs) [69].

B lymphocytes also go through different stages, including (1) activation, (2) selection, and (3) maturation. Like T lymphocytes, B-cells are derived from hematopoietic stem cells (HSCs) that go through stages in the bone marrow. These stages include the pro-B-cell (progenitor B-cell), the pre-B-cell (precursor B-cell), the immature naïve B-cell, and lastly the mature naïve B-cell stage [70]. After maturation, the naïve B-cells migrate to a secondary lymphoid tissue (e.g., lymph nodes), where they can be activated by antigens to generate memory B cells and plasma antibody-secreting cells.

T and B lymphocytes are major cellular components for autoimmunity. B-cells are responsible for antigen presentation, antibody production, and cytokine production. However, T-cells take part in both “helper” and cytotoxic activity. T-helper cells, also known as

CD4⁺ T-cells, are a type of T-cells that play an essential role in adaptive immune responses. Cytotoxic T-cells, CD8⁺ T-cells, are identified as T-cells that kill cancer cells or infected cells.

Both CD4 and CD8 T lymphocyte types contribute to the adaptive immunity component of the immune system. There are two distinct types of immunity based on B- and T-cell function: humoral and cell-mediated. Humoral immunity is based on the antibodies produced by plasma cells, which are derived from mature naïve B-cells. This allows for an individual exposed to a disease to be administered antibodies already generated by another individual previously exposed to the same disease. Cell-mediated immunity, however, involves T-cells, specifically the cytotoxic (CD8) T-cells. Similar to humoral immunity, T-helper and cytotoxic cells from an individual who was exposed to and survived a disease are administered to an individual who has been freshly exposed to the same disease. The helper T-cells are able to activate other immune cells and cytotoxic T-cells, initiating the removal process of those pathogens [71]. Hence, both humoral and cell-mediated immunity result in the binding of the antibodies to the antigen and affect their elimination.

4. Immune Cells and Lipids

Proinflammatory lipids promote CD4 Th1/Th17 differentiation [72], inhibit Tr1 differentiation [72], induce NF- κ B [73], inhibit Treg cell differentiation [73], modulate local activation of T-cells [74], induce NO [75], participate in oxidative stress pathways [76], increase the expression of cytokine genes and CCL2, also known as monocyte chemoattractant protein-1 (MCP-1) [77,78], amplify ER stress [79–81], and reduce inflammation-resolving processes [82–85]. Furthermore, reduced cellular debris clearance [86] and apoptotic clearance defects in macrophages and dendritic cells of non-obese diabetic (NOD) mice, an autoimmune spontaneous T1D-prone model, have been attributed to PGE₂ [87], which is significantly higher in infiltrated versus insulinitis-free NOD islets [74]. Other lipids, in particular lysophosphatidic acid (LPA), are powerful chemoattractants [40,42,88,89]. LOX products induce migration of leukocytes, p38 mitogen-activated protein kinase pathway, ROS-generating NADPH oxidase (NOX), proinflammatory cytokine production, and β -cell apoptosis [90–95]. 12-LOX-deficiency protects mice from streptozotocin (STZ)-induced T1D [96] and reduces insulinitis and T1D incidence in the NOD [97]. 12-LOX is detectable in islet cells in recent-onset T1D but absent in non-diabetic subjects [98].

Several publications over the years have introduced the concept that lipid mediators and fatty acids influence immune responses [99,100]. One of the most relevant T-cell discoveries has been the characterization of signaling platforms enriched in cholesterol and glycosphingolipids, which are named lipid rafts. These rafts are important for downstream signaling pathways for enzymes, scaffold proteins, and adaptors [101].

5. Deleterious Impact of iPLA₂ β Activation

5.1. Cellular Compass

Immune responses are propagated through the recruitment of peripheral blood leukocytes to inflamed areas. The chemokine CCL2 plays a critical role in this process by signaling through receptor CC chemokine receptor 2 (CCR2) [102]. The Cathcart group, examining the role of CCL2 in promoting directed migration of monocytes, identified distinct roles for iPLA₂ β and cPLA₂ α [40,103]. Upon CCL2 stimulation, cPLA₂ α translocated from the cytosol to the endoplasmic reticulum (ER) and regulated monocyte migration via arachidonic acid and its metabolites. In contrast, iPLA₂ β was recruited to the membrane-enriched pseudopod. Here, iPLA₂ β was proposed to catalyze hydrolysis of the *sn*-2 fatty acid from PLD-derived phosphatidic acid (PA) [104]. This phospholipid is among the negatively charged lipid substrates preferred by iPLA₂ β [7,105], and leads to the accumulation of LPA. Utilizing the iPLA₂ β -selective inhibitor (*S*-BEL) [106] and antisense oligodeoxyribonucleotide (ODN), the authors demonstrated that iPLA₂ β -generated LPA regulates actin polymerization to facilitate directionality to the migrating monocytes. They go on to suggest that iPLA₂ β may manifest a cellular compass role or be an integral component of the cellular

compass. Consistently, lipids derived from iPLA₂β activation, but not other PLA₂s, have been demonstrated to promote monocyte chemotaxis [42,88,107,108] (Figure 1).




Figure 1. Proposed mechanisms by which iPLA₂β activation contributes to monocyte chemotaxis, foam cell formation, vascular and bladder inflammation, tumor cell survival, and macrophage polarization. LPA, lysophosphatidic acid; iDLs, iPLA₂β-derived lipids; ROS, reactive oxygen species; EC, endothelial cell (↑, increased expression/activity of iPLA₂β; ↑, indicates increase in).

5.2. Foam Cell Formation

Macrophages contribute to atherosclerosis development, and it requires their conversion to lipid-laden foam cells via a toll-like receptor (TLR)-mediated process [109]. Lipopolysaccharide (LPS) plays a critical role in this process through the generation of ROS [110,111]. Lee et al. [35] demonstrated that LPS binding to TLR4 induces NOX1 expression and, as a consequence, ROS production. Both S-BEL and small interfering RNA (siRNA) directed against iPLA₂β, but not R-BEL or siRNA directed against membrane-associated iPLA₂γ (iPLA₂γ), were found to decrease NOX1 expression, and consequently, ROS production. This was associated with mitigation of foam cell formation. They further reported that iPLA₂β effects are signaled through the Akt pathway, and although the specific lipid signal involved was not identified, they hypothesized that iDLs promote Akt phosphorylation. However, this remains to be elucidated.

5.3. Vascular Injury

Neointima formation leads to several vascular-related pathologies and macrophages are critical contributors to vascular inflammation [112]. Liu et al. [107] using S-BEL and siRNA directed against iPLA₂β, iPLA₂β-deficient mice, and mice that selectively overexpress iPLA₂β in smooth muscle cells, demonstrated that iPLA₂β participates in ligation-induced neointima formation. They further demonstrated that this was associated with increased production of proinflammatory cytokines and vascular infiltration by macrophages. By comparing the effects of inhibiting the arachidonic acid metabolism pathways COX2 with indomethacin, 5-LOX with NDGA, 12-LOX with baicalein, CYP with 17-octadecynoic acid, and 12/15-LOX with luteolin, they deduced that the products of 12-LOX and 15-LOX are complicit in the formation of neointima. In view of reports suggesting that sterol regulatory element-binding protein 1 (SREBP-1) expression is induced in the injured vascular wall [95] and SREBP-1 induces iPLA₂β [113,114], they speculated that SREBP-1-mediated induction of iPLA₂β occurs in their model of ligation-induced neointima formation. Interestingly, they found that the overexpression of iPLA₂β in the smooth muscle cells does not promote neointima formation and that it is only amplified upon injury, leading to increased TNFα production. As basally iPLA₂β is inactivated through an interaction with calmodulin [5,115,116], they postulate that TNFα causes disassociation of the complex to unmask iPLA₂β activity. Analogously, under basal conditions, β-cells overexpressing

iPLA₂β do not exhibit a difference in death rate; however, upon stimulation with ER stressors or pro-inflammatory cytokines, they exhibit exacerbated apoptosis [117,118].

5.4. Cigarette Smoke

An interesting link between iPLA₂β and platelet activating factor (PAF) was recognized by McHowat's group in their study of smoke-induced bladder inflammation [119]. This appears to involve a series of events initiated by the tethering of inflammatory cells to the apical endothelial cell surface, rolling of the inflammatory cells across the endothelium and tight adherence, and subsequent PAF-mediated transmigration between neighboring endothelial cells [120,121]. One route of PAF production, predominant during inflammation, is via remodeling initiated by the hydrolysis of the *sn*-2 substituent from phosphatidylcholine by PLA₂ activation to generate alkyl-lysophosphatidyl choline (LPC). Subsequently, an acetyl group is added at the *sn*-2 position to generate PAF. McHowat's group observed that upon exposure to cigarette smoke, human and rodent bladder endothelial cells (ECs) exhibited higher PAF accumulation, decreased activity of platelet-activating factor acetylhydrolase (PAF-AH), which degrades PAF, and increased inflammatory cell adherence. These outcomes were reversed by chemical inhibition or genetic knockout of iPLA₂β, and by PAF receptor (PAFR) antagonism. Cellular adhesion is facilitated by adhesion molecules such as platelets/endothelial cells (P/E)-selectin and intercellular adhesion molecule 1 (ICAM-1) and these are produced by mast cells [120], which are key contributors to chronic inflammation. Mast cells express PAFR and also release tryptase, which can induce iPLA₂β activation in bladder epithelial cells [122]. The authors proposed that mast-cell-induced PAF generation by endothelial cells via iPLA₂β activation can prolong the inflammatory state, thus leading to chronic inflammation. Of interest, Ueno et al. [123] reported that cPLA₂α, but not iPLA₂β, contributes to arachidonic acid production by bone marrow-derived mast cells, facilitating their maturation. Curiously, although mast cells have been reported not to be involved in the development of T1D, which is a consequence of autoimmune destruction of β-cells, in the NOD rodent model [124], they have been suggested to be potentially important in promoting β-cell dysfunction and death in human T1D [125].

5.5. Early-Stage Disease

A common blood cancer is chronic lymphocytic leukemia (CLL), which is characterized by immune-incompetent B-CLL lymphocytes [126,127]. These cells express COX2 and have increased production of PGE₂, PGF₂, and LTB₄. The latter signals through the BLT1 receptor to activate CD40-dependent chronic B lymphocytic leukemia cells to prolong B-CLL survival [128,129]. Greater expression of PLA₂ enzymes has been noted in tumor cells, relative to healthy cells, supporting their role in tumor cell proliferation and metastasis [38,130,131]. The work of Guriec et al. suggests that both cPLA₂α and iPLA₂β are expressed in the tumor cells, with cPLA₂α expression increasing higher in patients with advanced-stage disease and with iPLA₂β expression being higher in the early disease stages [132]. The disease progression was observed to be associated with dysregulation of arachidonic acid metabolizing pathways (COX, 5-LOX) and of LTA₄ hydrolase (LTA₄H), which generates LTB₄. The progressive increases in PGs and LTB₄, the latter of which activates its receptors BLT1 and BLT2, are proposed to support tumor cell proliferation and survival.

5.6. Metabolic Stress

It is well recognized that type 2 diabetes (T2D) is associated with increased coronary artery disease and atherosclerosis [133] and that these involve recruitment of macrophages to inflamed vascular sites [134]. Additionally, cellular oxidative stress appears to be a compounding factor. ROS are generated as byproducts of oxygenases or primary products of NOX. Among the NOX variants [135], although both Nox4 and Nox2 are expressed in macrophages [136], Nox4 promotes monocyte chemotaxis and macrophage recruitment during diabetic metabolic stress (DMS) [137]. Tan et al. [42] reported that under conditions mimicking DMS (i.e., high glucose and LDL), overexpression of iPLA₂β amplified

macrophage NOX4 expression, ROS production, and CCL2-induced migration. Conversely, these outcomes were mitigated by S-BEL and siRNA directed against iPLA₂β. The same interventions had no effect on NOX2. Interestingly, NOX2 is localized primarily in the plasma membrane, whereas NOX4 is localized in the mitochondria, ER and nuclear membranes [138,139], which are also the subcellular organelles that iPLA₂β mobilizes to upon the induction of stress [11,12,14,15,140]. Tan et al. further suggested that iPLA₂β-derived LPA manifested these effects, as LPA receptor antagonism prevented NOX4 induction and LPA addition was able to rescue outcomes inhibited by S-BEL.

5.7. Macrophage Polarization

In view of the reported impact of iPLA₂β on macrophage function, we sought to identify the role of iDLs in this process [39,141]. We found that iPLA₂β activation induces macrophage polarization, favoring the M1 inflammatory phenotype. Utilizing selective inhibitors, we identified that both COX and LOX products contribute to iPLA₂β-modulated macrophage polarization. Such an outcome was mitigated in macrophages from iPLA₂β-deficient mice upon classical activation (LPS+IFNγ). This was reflected by reductions in M1 markers (*Arg2* and *Nos2*) and increases in several M2 markers, including *Arg1* and *Ccl2*. Interestingly, *Ptgs2* and *Alox12*, which encode COX2 and 12-LOX, also decreased, suggesting feedback regulation by products of these enzymes. Among the iDLs identified as inducers of the M1 phenotype were 6-keto PGF₁α, PGE₂, and LTB₄. The inhibition of secretory PLA₂s (sPLA₂s) GIIA, GV, and GX or cPLA₂α did not alter the classically activated macrophage production of eicosanoids, suggesting a select role for iDLs on macrophage polarization. However, other groups have reported a role for sPLA₂ or cPLA₂α in eicosanoid production from macrophages [142–147]. Those studies were performed with macrophage-like cells (RAW264.7 and P388D1) or peritoneal macrophages from rats and mice, with only LPS (10–1000 ng/mL) or zymosan stimulation for 6–48 h, and the involvement of various PLA₂s was discerned with chemical inhibitors, some of which effected multiple PLA₂s. In contrast, our studies were performed with primary mouse peritoneal macrophages, which have been demonstrated to behave differently from the macrophage-like cells [36], from wild-type and genetically- modified iPLA₂β-deficient mice, and importantly, treated with LPS+IFNγ, conditions that induce a M1 macrophage phenotype. Interestingly, while iPLA₂β-deficiency skewed macrophage polarization towards an M2 anti-inflammatory phenotype, inhibition of downstream oxygenases did not necessarily promote an M2 phenotype. We posit that a reduction in macrophage-iPLA₂β activity, as a consequence of an attenuated inflammatory landscape, promotes a shift to a recovery/repair milieu.

5.8. T1D Development

Immune cells play a significant role in promoting β-cell death, which leads to T1D development [148,149]. In T1D-prone individuals, macrophages are among the first immune cells that migrate to pancreatic islets to initiate inflammatory responses and secrete proinflammatory cytokines and ROS, which lead to β-cell death [150,151]. Two different activation states of macrophages have been described—M1 proinflammatory macrophages [152], which are classically activated (e.g., by IFNγ, LPS, TNFα); and M2 anti-inflammatory macrophages, which are alternatively-activated (e.g., by IL-4 or IL-10) [153]. Although M1 macrophages are recognized to be causative factors in T1D development [154], M2 macrophages protect against T1D development [155].

Macrophages are the only resident myeloid cells found in pancreatic islets under normal conditions in all mice strains [156]. However, the cellular composition of pancreata from diabetic individuals and NOD mice revealed a considerable presence of macrophage infiltration into pancreatic islets, with a predominant M1-like phenotype [33,156,157]. Macrophages found in close contact with β-cells present diabetogenic antigens, predominantly insulin peptides [158–160]. The proinflammatory M1 macrophages produce ROS, which, along with proinflammatory cytokines such as IL-1β, IFNγ, and TNFα, cause β-cell destruction [49,161]. It is well-recognized that macrophages work in concert with other

immune cells to promote β -cell destruction in T1D [149,162,163]. Furthermore, proinflammatory eicosanoids are linked to macrophage phagocytosis, adhesion, apoptosis, and amplifying macrophage-derived eicosanoid release [24,164–166]. In the context of T1D, 12-LOX-null NOD has a lowered T1D incidence, which is associated with reduced macrophage production of proinflammatory cytokines [97].

It is also well-established that in human T1D patients and NOD mice, both CD4 and CD8 T-cells are the major components of the islet infiltrate [162]. Different stages of disease onset show varying compositions of CD4 and CD8 T-cells, depending on the timeframe of progression. First, autoreactive T-cells are activated by β -cell antigens presented by antigen-presenting cells (APCs). The activated T-helper cells are required to activate CD8 T-cells. Second, the activated CD4 T-cells infiltrate the pancreas and are thought to contribute to β -cell destruction via the activation of macrophages [154].

CD4 T-cells recognize MHC class II peptides presented by APCs in order to carry out β -cell destruction. CD4 T-cell-mediated β -cell destruction can be caused by the production of proinflammatory cytokines such as $\text{IFN}\gamma$, which are toxic to the β -cell, and indirectly, by activating local innate cells such as macrophages and dendritic cells (DCs) to enhance infiltration [154]. CD4 T-cells are also thought to interact with CD8 T-cells by facilitating their activation [154]. CD8 T-cells can directly kill β -cells by interacting with MHC class I molecules and by means of perforin and granzyme secretion [149]. It has been suggested that the MHC class I/CD8 T-cell interaction is required for T1D in the early stages of development [162] and antigen presentation to CD4 T-cells within pancreatic islets is essential for β -cell destruction [154]. Other reports have suggested that the MHC class I/CD8⁺ T-cell interaction is required for T1D in the early stages of development [167] and that antigen presentation to CD4 T-cells within pancreatic islets are essential for β -cell destruction [154]. Taken together, the literature establishes that immune cells work in concert to promote β -cell destruction in T1D [21].

Evidence linking iDLs with T1D development (Figure 2). Several reports have provided evidence of an association between deleterious outcomes in experimental and clinical diabetes and $\text{iPLA}_2\beta$ activation [41,168].

In view of these observations, we explored the possibility that inhibition of $\text{iPLA}_2\beta$ can ameliorate T1D [33,169]. We found that administration of FKGG18 to female NOD mice promoted several positive outcomes. There was a significant reduction in insulinitis, as reflected by reductions in islet abundances of CD4⁺ T-cells and B-cells. Glucose homeostasis was also improved, as reflected by β -cell preservation and higher circulating insulin. Consequentially, a significant reduction in T1D was achieved. Inhibition of $\text{iPLA}_2\beta$ resulted in decreased production of $\text{TNF}\alpha$ from CD4⁺ T-cells and antibodies from B-cells, suggesting that iDLs modulate immune cell responses. This was supported by the recapitulation of the mitigated $\text{TNF}\alpha$ production by select $\text{iPLA}_2\beta$ inhibitors, with COX and 12-LOX inhibition. $\text{TNF}\alpha$ acts as a powerful chemoattractant [89] and is produced by CD4 T-cells within inflamed islets during T1D development [170]. $\text{TNF}\alpha$ overexpression exacerbates insulinitis, whereas the opposite occurs in $\text{TNF}\beta$ -receptor-null mice [171]. To date, ours are the first and only reports of the modulation of T- and B-cell functions by iDLs. Similar findings in a genetically modified NOD model with reduced $\text{iPLA}_2\beta$ expression [169] further support this possibility.

These findings prompted us to further examine iDL production by macrophages. Not surprisingly, a dramatically more profound proinflammatory landscape was evident in macrophages from the NOD, relative to the spontaneous diabetes-resistant C57BL/6J, mouse [169]. Lipidomic assessments in the NOD model identified select iDLs (PGE_2 , PGD_2 , hydroxyeicosatetraenoic acids 5 and 15 (5-HETE and 15-HETE), LTC_4) that correlated with T1D development [169]. Importantly, a similar lipid signature was revealed in the plasma of human subjects at high risk of developing T1D [169]. We therefore posit that select iDLs contribute to T1D onset and that these could be targeted for therapeutics and, in conjunction with autoantibodies, serve as early biomarkers of pre-T1D.




Figure 2. Proposed contribution of iDLs generated by immune cells to β -cell destruction and type-1 diabetes (T1D) development. \uparrow , increased expression/activity of $iPLA_2\beta$; \uparrow , indicates increase in; and \downarrow , indicates decrease in).

6. Protective Consequences of $iPLA_2\beta$ Activation

6.1. Cancer Development

Inflammation is a key contributor to cancer development [172] and cytokines released by macrophages and T-cells are integral to this process [173]. In view of their earlier observations that $iPLA_2\beta$ -null mice are more susceptible to various inflammatory-based disorders [174–176], Inhoffen et al. [177] assessed the ability of immune cells from $iPLA_2\beta$ -null mice to produce cytokines following exposure to CD95/FasL, a trigger of proinflammatory cytokine production [178]. They found that $iPLA_2\beta$ -deficiency increased apoptosis in the liver, spleen, and mesenteric lymph nodes (MLN). Although Kupffer cells (i.e., satellite macrophages in the liver) generated a lower production of proinflammatory cytokines $TNF\alpha$ and IL-6, splenocytes became primed to release proinflammatory Th1-/Th17-related cytokines ($IFN\gamma$ /IL-17 α). These findings led to the suggestion that $iPLA_2\beta$ -deficiency can reduce age-related MLN lymphoma development. Mechanistically, they attributed this to the decreased availability of “find-me” and “eat-me” signals derived via $iPLA_2\beta$ activation. These include LPC, a find-me signal [88], which promotes the clearance of apoptotic debris, the accumulation of which triggers and amplifies subsequent immune responses [179,180] (Figure 3).




Figure 3. Proposed mechanisms by which iPLA₂β inactivation or activation can have protective outcomes in tumor development and promoting host defenses. (↓, decreased expression/activity of iPLA₂β; ↑, increased expression/activity of iPLA₂β; ↑, indicates increase in; ↓, indicates decrease in).

6.2. Inflammatory Bowel Disease (IBD)

An inflammatory disorder such as IBD is a consequence of the dysfunction of the intestinal epithelial barrier and the mucosal immune system [181]. Jiao et al. [175] further explored the role of “find-me” signals derived via iPLA₂β activation in the context of dextran sodium sulfate-induced IBD. They found that iPLA₂β-deficiency promotes the accumulation of infiltrating macrophages and dendritic cells in the colon lamina. These were associated with increased production of inflammatory cytokines (TNFα, IL-1β, and IL-6), macrophage inflammatory proteins (MIP-1α and MIP-1β), and CCL2, leading to intestinal epithelial cell apoptosis and mucus barrier damage. With a concurrent decrease in LPC levels, they speculated that iPLA₂β-deficiency mitigated the availability of a “find-me” signal, which limited the clearance of apoptotic debris and the amplification of immune responses. Further, they predicted that the iPLA₂β-deficiency also reduced LPA levels, impacting the cellular compass, and preventing the optimal phagocytic function of macrophages. Subsequently, Murase et al. [182], comparing the involvement of various PLA₂s, suggested that iPLA₂β-deficiency did not worsen the clinical scoring associated with IBD, relative to controls. As they did not reconcile the two studies, it is likely that differences between the source of the iPLA₂β^{-/-} mice and DSS concentrations used may be partly responsible. Furthermore, while a 7-day DSS regimen was employed by both groups, Jiao’s group maintained the mice for an additional 3 days without DSS, prior to analyses. It may also be noted that Murase’s study did demonstrate similar lower clinical scores in the wild-type (WT) and iPLA₂β-deficient groups between days 1 and 6, relative to mice with deficiencies in cPLA₂α or in a variety of sPLA₂s; however, at day 7, the scores in the iPLA₂β-deficient group appeared to be as high as in the other PLA₂-deficient groups. As such, the role of iPLA₂β in this model remains to be clarified.

6.3. Chagas Disease

A further link between iPLA₂β and PAF was reported by McHowat’s group [183], in the context of Chagas disease. This disease is caused through infection by the protozoan parasite [184] *Trypanosoma cruzi* (*T. cruzi*) and can lead to various cardiac abnormalities [185]. The sequela of infection begins with induction of an inflammatory response and upregulation of endothelial adhesion molecules [186], followed by attempts at resolution through the generation of proinflammatory cytokines and the induction of signaling pathways to promote the chemotaxis of immune cells to mitigate the invasion. McHowat’s group re-

ported that infection of iPLA₂β-deficient mice resulted in lowered PAF and NO production by cardiac endothelial cells, but that neither the expression of adhesion molecules nor the development of myocardial inflammation was affected. However, significant increases in parasite pseudocysts were noted in the myocardium of iPLA₂β-deficient mice. The authors suggested that this was due to an impairment in parasite clearance as a consequence of decreased iPLA₂β-mediated LPA production and, as a result, Nox4 expression and NO production. Thus, they surmised that the absence of iPLA₂β mitigates parasite clearance due to the reduced recruitment of inflammatory cells to the infected myocardial areas.

6.4. Negative Modulation of Inflammation by AAT and SLP1

In sequential reports, Grau's group constructed events that participated in regulating IL-1β activation and release. IL-1β is critical to host defense against infections, but the generation of excessive active IL-1β can lead to deleterious inflammatory consequences [187]. The release of IL-1β occurs via two signals. The first is an external stimulus that induces the synthesis of pro-IL-1β. The second signal has been suggested to be ATP, which when released from damaged cells activates the purinergic receptor P2 × 7R, promoting the loss of K⁺ current and triggering the assembly of NLRP3 (NLR family pyrin domain containing 3)-containing inflammasome [188]. This leads to caspase-1 activation, cleavage of pro-IL-1β, and release of IL-1β from the immune cell. Present in inflammatory cells, alpha-1 antitrypsin (AAT) is a strong inducer of anti-inflammatory processes and its upregulation during systemic inflammation has been associated with the decreased production of proinflammatory cytokines, including IL-1β [189]. In examining the potential role of AAT in ATP-dependent regulation of IL-1β in their first study [190], Siebers et al. found that AAT signaling through the CD36 receptor activates iPLA₂β, which leads to the release of a low molecular weight factor (LMWF). The LMWF is released from the cell and binds to nicotinic acetylcholine receptor (nAChR), leading to inhibition of P2X7R function, prevention of inflammasome assembly, and the processing of pro-IL-1β to active IL-1β, and its release from the cell. These outcomes were significantly mitigated with selective inhibition or knockdown of iPLA₂β and were also not evident in PBMCs from iPLA₂β-deficient mice. In the second study [191], Zakrzewicz et al. demonstrated that the secretory leukocyte protease inhibitor (SLP1) also interferes with ATP-dependent regulation of IL-1β via iPLA₂β activation. SLP1 is also present in inflammatory cells [192] and its actions promote anti-inflammatory outcomes. Their results suggested that the SLP1 signals released through annexin 2, a membrane binding protein for SLP1 [193], activate iPLA₂β, leading to the production of the LMWF and subsequent inhibition of IL-1β maturation and release. Although both pathways were found to mitigate inflammation, unfortunately neither study explored the identity of the LMWF. Interestingly, AAT administration to recent-onset T1D subjects improved β-cell function, which was correlated with reduced IL-1β production from monocytes and myeloid dendritic cells [194].

7. Summary

iPLA₂β is a member of the family of PLA₂s that hydrolyzes the *sn*-2 substituent from membrane glycerophospholipids. Thus, activation of iPLA₂β can lead to the production of a variety of bioactive lipid mediators. As eicosanoids generated subsequently to iPLA₂β-mediated hydrolysis of *sn*-2 arachidonic acid can exhibit profound proinflammatory effects and SPMs are produced from *sn*-2 substituents EPA and DHA, the impact of iPLA₂β on the inflammatory sequelae is profound. In recent years, the impact of iPLA₂β at the immune cell level is being recognized and those studies, as well as those from our laboratory, suggest that iPLA₂β activation modulates the maturation, polarization, activation, and functionality of macrophages, T-cells, and B-cells. The continuation of studies addressing these actions of iPLA₂β is important and warranted in order to gain a better understanding of the events that lead to the onset, maintenance, and amplification of inflammation. Identifying the relevant selective iDLs with effects on these processes could lead to the development of new strategies to treat autoimmune- and other inflammatory-based diseases.

Author Contributions: T.D.W., A.A., and S.R. contributed equally to the writing and editing, X.L. contributed to the editing, and Y.G.T. collected the reference materials. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by NIH/NIDDK R01DK110292, DK079626, and NIH/NIAID 1 R21 AI 146743.

Informed Consent Statement: Not Applicable.

Data Availability Statement: Not Applicable.

Acknowledgments: The authors would like to thank the Department of CDIB and the Comprehensive Diabetes Center for their administrative support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gijon, M.A.; Leslie, C.C. Phospholipases A₂. *Semin. Cell Dev. Biol.* **1997**, *8*, 297–303. [[CrossRef](#)] [[PubMed](#)]
2. Turk, J.; White, T.D.; Nelson, A.J.; Lei, X.; Ramanadham, S. iPLA₂b and its role in male fertility, neurological disorders, metabolic disorders, and inflammation. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **2019**, *1864*, 846–860. [[CrossRef](#)] [[PubMed](#)]
3. Ramanadham, S.; Ali, T.; Ashley, J.W.; Bone, R.N.; Hancock, W.D.; Lei, X. Calcium-independent phospholipases A₂ and their roles in biological processes and diseases. *J. Lipid Res.* **2015**, *56*, 1643–1668. [[CrossRef](#)] [[PubMed](#)]
4. Malley, K.R.; Koroleva, O.; Miller, I.; Sanishvili, R.; Jenkins, C.M.; Gross, R.W.; Korolev, S. The structure of iPLA₂beta reveals dimeric active sites and suggests mechanisms of regulation and localization. *Nat. Commun.* **2018**, *9*, 765. [[CrossRef](#)]
5. Wang, Z.; Ramanadham, S.; Ma, Z.A.; Bao, S.; Mancuso, D.J.; Gross, R.W.; Turk, J. Group VIA phospholipase A₂ forms a signaling complex with the calcium/calmodulin-dependent protein kinase IIbeta expressed in pancreatic islet beta-cells. *J. Biol. Chem.* **2005**, *280*, 6840–6849. [[CrossRef](#)] [[PubMed](#)]
6. Mouchlis, V.D.; Armando, A.; Dennis, E.A. Substrate-specific inhibition constants for phospholipase A₂ acting on unique phospholipid substrates in mixed micelles and membranes using lipidomics. *J. Med. Chem.* **2019**, *62*, 1999–2007. [[CrossRef](#)]
7. Mouchlis, V.D.; Chen, Y.; Mccammon, J.A.; Dennis, E.A. Membrane allostery and unique hydrophobic sites promote enzyme substrate specificity. *J. Am. Chem. Soc.* **2018**, *140*, 3285–3291. [[CrossRef](#)] [[PubMed](#)]
8. Ali, T.; Kokotos, G.; Magrioti, V.; Bone, R.N.; Mobley, J.A.; Hancock, W.; Ramanadham, S. Characterization of FKGK18 as inhibitor of group VIA Ca²⁺-independent phospholipase A₂ (iPLA₂b): Candidate drug for preventing beta-cell apoptosis and diabetes. *PLoS ONE* **2013**, *8*, e71748. [[CrossRef](#)] [[PubMed](#)]
9. Kokotos, G.; Hsu, Y.H.; Burke, J.E.; Baskakis, C.; Kokotos, C.G.; Magrioti, V.; Dennis, E.A. Potent and selective fluoroketone inhibitors of group VIA calcium-independent phospholipase A₂. *J. Med. Chem.* **2010**, *53*, 3602–3610. [[CrossRef](#)]
10. Gross, R.W.; Ramanadham, S.; Kruszka, K.K.; Han, X.; Turk, J. Rat and human pancreatic islet cells contain a calcium ion independent phospholipase A₂ activity selective for hydrolysis of arachidonate which is stimulated by adenosine triphosphate and is specifically localized to islet beta-cells. *Biochemistry* **1993**, *32*, 327–336. [[CrossRef](#)]
11. Bao, S.; Jin, C.; Zhang, S.; Turk, J.; Ma, Z.; Ramanadham, S. Beta-cell calcium-independent group VIA phospholipase A₂ (iPLA₂b): Tracking iPLA₂b movements in response to stimulation with insulin secretagogues in INS-1 cells. *Diabetes* **2004**, *53*, S186–S189. [[CrossRef](#)]
12. Lei, X.; Zhang, S.; Bohrer, A.; Ramanadham, S. Calcium-independent phospholipase A₂ (iPLA₂b)-mediated ceramide generation plays a key role in the cross-talk between the endoplasmic reticulum (ER) and mitochondria during ER stress-induced insulin-secreting cell apoptosis. *J. Biol. Chem.* **2008**, *283*, 34819–34832. [[CrossRef](#)]
13. Ma, Z.; Ramanadham, S.; Wohltmann, M.; Bohrer, A.; Hsu, F.F.; Turk, J. Studies of insulin secretory responses and of arachidonic acid incorporation into phospholipids of stably transfected insulinoma cells that overexpress group VIA phospholipase A₂ (iPLA₂b) indicate a signaling rather than a housekeeping role for iPLA₂b. *J. Biol. Chem.* **2001**, *276*, 13198–13208. [[CrossRef](#)]
14. Ma, Z.; Zhang, S.; Turk, J.; Ramanadham, S. Stimulation of insulin secretion and associated nuclear accumulation of iPLA₂b in INS-1 insulinoma cells. *Am. J. Physiol. Endocrinol. Metab.* **2002**, *282*, E820–E833. [[CrossRef](#)] [[PubMed](#)]
15. Ramanadham, S.; Hsu, F.F.; Zhang, S.; Jin, C.; Bohrer, A.; Song, H.; Bao, S.; Ma, Z.; Turk, J. Apoptosis of insulin-secreting cells induced by endoplasmic reticulum stress is amplified by overexpression of group VIA calcium-independent phospholipase A₂ (iPLA₂b) and suppressed by inhibition of iPLA₂b. *Biochemistry* **2004**, *43*, 918–930. [[CrossRef](#)] [[PubMed](#)]
16. Song, H.; Bao, S.; Lei, X.; Jin, C.; Zhang, S.; Turk, J.; Ramanadham, S. Evidence for proteolytic processing and stimulated organelle redistribution of iPLA₂b. *Biochim. Biophys. Acta* **2010**, *1801*, 547–558. [[CrossRef](#)] [[PubMed](#)]
17. Ramanadham, S.; Hsu, F.F.; Bohrer, A.; Zhongmin, M.; Turk, J. Studies of the role of group VI phospholipase A₂ in fatty acid incorporation, phospholipid remodeling, lysophosphatidylcholine generation, and secretagogue-induced arachidonic acid release in pancreatic islets and insulinoma cells. *J. Biol. Chem.* **1999**, *275*, 13915–13927. [[CrossRef](#)]
18. Hanna, V.S.; Hafez, E.A.A. Synopsis of arachidonic acid metabolism: A review. *J. Adv. Res.* **2018**, *11*, 23–32. [[CrossRef](#)]
19. Abdulkhaleq, L.A.; Assi, M.A.; Abdullah, R.; Zamri-Saad, M.; Taufiq-Yap, Y.H.; Hezmee, M.N.M. The crucial roles of inflammatory mediators in inflammation: A review. *Vet. World* **2018**, *11*, 627–635. [[CrossRef](#)] [[PubMed](#)]

20. Khanapure, S.P.; Garvey, D.S.; Janero, D.R.; Letts, L.G. Eicosanoids in inflammation: Biosynthesis, pharmacology, and therapeutic frontiers. *Curr. Top. Med. Chem.* **2007**, *7*, 311–340. [[CrossRef](#)]
21. Luo, P.; Wang, M.H. Eicosanoids, beta-cell function, and diabetes. *Prostaglandins Other Lipid Mediat.* **2011**, *95*, 1–10. [[CrossRef](#)]
22. Tessaro, F.H.; Ayala, T.S.; Martins, J.O. Lipid mediators are critical in resolving inflammation: A review of the emerging roles of eicosanoids in diabetes mellitus. *Biomed. Res. Int.* **2015**, *2015*, 568408. [[CrossRef](#)]
23. Umamaheswaran, S.; Dasari, S.K.; Yang, P.; Lutgendorf, S.K.; Sood, A.K. Stress, inflammation, and eicosanoids: An emerging perspective. *Cancer Metastasis Rev.* **2018**, *37*, 203–211. [[CrossRef](#)]
24. Zaitseva, L.; Vaisburd, M.; Shaposhnikova, G.; Mysyakin, E. Role of eicosanoids in regulation of macrophage phagocytic functions by platelet-activating factor during endotoxic shock. *Bull. Exp. Biol. Med.* **2000**, *130*, 879–881. [[CrossRef](#)] [[PubMed](#)]
25. Kuhn, H.; O'donnell, V.B. Inflammation and immune regulation by 12/15-lipoxygenases. *Prog. Lipid Res.* **2006**, *45*, 334–356. [[CrossRef](#)] [[PubMed](#)]
26. Issan, Y.; Hochhauser, E.; Guo, A.; Gotlinger, K.H.; Kornowski, R.; Leshem-Lev, D.; Lev, E.; Porat, E.; Snir, E.; Thompson, C.I.; et al. Elevated level of pro-inflammatory eicosanoids and EPC dysfunction in diabetic patients with cardiac ischemia. *Prostaglandins Other Lipid Mediat.* **2013**, *100–101*, 15–21. [[CrossRef](#)]
27. Gilroy, D.W.; Newson, J.; Sawmynaden, P.; Willoughby, D.A.; Croxtall, J.D. A novel role for phospholipase A₂ isoforms in the checkpoint control of acute inflammation. *FASEB J.* **2004**, *18*, 489–498. [[CrossRef](#)]
28. De Jong, A.J.; Kloppenburg, M.; Toes, R.E.; Ioan-Facsinay, A. Fatty acids, lipid mediators, and T-cell function. *Front. Immunol.* **2014**, *5*, 483. [[CrossRef](#)]
29. Serhan, C.N. Treating inflammation and infection in the 21st century: New hints from decoding resolution mediators and mechanisms. *FASEB J.* **2017**, *31*, 1273–1288. [[CrossRef](#)] [[PubMed](#)]
30. Serhan, C.N.; Levy, B.D. Resolvins in inflammation: Emergence of the pro-resolving superfamily of mediators. *J. Clin. Investig.* **2018**, *128*, 2657–2669. [[CrossRef](#)] [[PubMed](#)]
31. Serhan, C.N. Pro-resolving lipid mediators are leads for resolution physiology. *Nature* **2014**, *510*, 92–101. [[CrossRef](#)]
32. Serhan, C.N.; Chiang, N.; Dalli, J.; Levy, B.D. Lipid mediators in the resolution of inflammation. *Cold Spring Harb. Perspect. Biol.* **2014**, *7*, a016311. [[CrossRef](#)] [[PubMed](#)]
33. Bone, R.N.; Gai, Y.; Magrioti, V.; Kokotou, M.G.; Ali, T.; Lei, X.; Tse, H.M.; Kokotos, G.; Ramanadham, S. Inhibition of Ca²⁺-independent phospholipase A₂β (iPLA₂β) ameliorates islet infiltration and incidence of diabetes in NOD mice. *Diabetes* **2015**, *64*, 541–554. [[CrossRef](#)] [[PubMed](#)]
34. Gil-De-Gomez, L.; Astudillo, A.M.; Guijas, C.; Magrioti, V.; Kokotos, G.; Balboa, M.A.; Balsinde, J. Cytosolic group IVA and calcium-independent group VIA phospholipase A₂s act on distinct phospholipid pools in zymosan-stimulated mouse peritoneal macrophages. *J. Immunol.* **2014**, *192*, 752–762. [[CrossRef](#)] [[PubMed](#)]
35. Lee, S.H.; Park, D.W.; Park, S.C.; Park, Y.K.; Hong, S.Y.; Kim, J.R.; Lee, C.H.; Baek, S.H. Calcium-independent phospholipase A₂β-Akt signaling is involved in lipopolysaccharide-induced NADPH oxidase 1 expression and foam cell formation. *J. Immunol.* **2009**, *183*, 7497–7504. [[CrossRef](#)] [[PubMed](#)]
36. Bao, S.; Li, Y.; Lei, X.; Wohltmann, M.; Jin, W.; Bohrer, A.; Semenkovich, C.F.; Ramanadham, S.; Tabas, I.; Turk, J. Attenuated free cholesterol loading-induced apoptosis but preserved phospholipid composition of peritoneal macrophages from mice that do not express group VIA phospholipase A₂. *J. Biol. Chem.* **2007**, *282*, 27100–27114. [[CrossRef](#)] [[PubMed](#)]
37. Ma, Z.; Bohrer, A.; Wohltmann, M.; Ramanadham, S.; Hsu, F.F.; Turk, J. Studies of phospholipid metabolism, proliferation, and secretion of stably transfected insulinoma cells that overexpress group VIA phospholipase A₂. *Lipids* **2001**, *36*, 689–700. [[CrossRef](#)]
38. Song, Y.; Wilkins, P.; Hu, W.; Murthy, K.S.; Chen, J.; Lee, Z.; Oyesanya, R.; Wu, J.; Barbour, S.E.; Fang, X. Inhibition of calcium-independent phospholipase A₂ suppresses proliferation and tumorigenicity of ovarian carcinoma cells. *Biochem. J.* **2007**, *406*, 427–436. [[CrossRef](#)]
39. Ashley, J.W.; Hancock, W.D.; Nelson, A.J.; Bone, R.N.; Tse, H.M.; Wohltmann, M.; Turk, J.; Ramanadham, S. Polarization of macrophages toward M2 phenotype is favored by reduction in iPLA₂β (Group VIA Phospholipase A₂). *J. Biol. Chem.* **2016**, *291*, 23268–23281. [[CrossRef](#)]
40. Mishra, R.S.; Carnevale, K.A.; Cathcart, M.K. iPLA₂β: Front and center in human monocyte chemotaxis to MCP-1. *J. Exp. Med.* **2008**, *205*, 347–359. [[CrossRef](#)]
41. Ayilavarapu, S.; Kantarci, A.; Fredman, G.; Turkoglu, O.; Omori, K.; Liu, H.; Iwata, T.; Yagi, M.; Hasturk, H.; Van Dyke, T.E. Diabetes-induced oxidative stress is mediated by Ca²⁺-independent phospholipase A₂ in neutrophils. *J. Immunol.* **2010**, *184*, 1507–1515. [[CrossRef](#)]
42. Tan, C.; Day, R.; Bao, S.; Turk, J.; Zhao, Q.D. Group VIA phospholipase A₂ mediates enhanced macrophage migration in diabetes mellitus by increasing expression of nicotinamide adenine dinucleotide phosphate oxidase 4. *Arter. Thromb. Vasc. Biol.* **2014**, *34*, 768–778. [[CrossRef](#)] [[PubMed](#)]
43. Kalyvas, A.; Baskakis, C.; Magrioti, V.; Constantinou-Kokotou, V.; Stephens, D.; Lopez-Vales, R.; Lu, J.Q.; Yong, V.W.; Dennis, E.A.; Kokotos, G.; et al. Differing roles for members of the phospholipase A₂ superfamily in experimental autoimmune encephalomyelitis. *Brain J. Neurol.* **2009**, *132*, 1221–1235. [[CrossRef](#)] [[PubMed](#)]
44. Mchowat, J.; Gullickson, G.; Hoover, R.G.; Sharma, J.; Turk, J.; Kornbluth, J. Platelet-activating factor and metastasis: Calcium-independent phospholipase A₂β deficiency protects against breast cancer metastasis to the lung. *Am. J. Physiol. Cell Physiol.* **2011**, *300*, C825–C832. [[CrossRef](#)] [[PubMed](#)]

45. Nicotera, T.M.; Schuster, D.P.; Bourhim, M.; Chadha, K.; Klaich, G.; Corral, D.A. Regulation of PSA secretion and survival signaling by calcium-independent phospholipase A₂β in prostate cancer cells. *Prostate* **2009**, *69*, 1270–1280. [[CrossRef](#)] [[PubMed](#)]
46. Scuderi, M.R.; Anfuso, C.D.; Lupo, G.; Motta, C.; Romeo, L.; Guerra, L.; Cappellani, A.; Ragusa, N.; Cantarella, G.; Alberghina, M. Expression of Ca²⁺-independent and Ca²⁺-dependent phospholipases A₂ and cyclooxygenases in human melanocytes and malignant melanoma cell lines. *Biochim. Biophys. Acta* **2008**, *1781*, 635–642. [[CrossRef](#)] [[PubMed](#)]
47. Thayer, T.C.; Delano, M.; Liu, C.; Chen, J.; Padgett, L.E.; Tse, H.M.; Annamali, M.; Piganelli, J.D.; Moldawer, L.L.; Mathews, C.E. Superoxide production by macrophages and T cells is critical for the induction of autoreactivity and type 1 diabetes. *Diabetes* **2011**, *60*, 2144–2151. [[CrossRef](#)]
48. Jiang, Z.; Jiang, J.X.; Zhang, G.X. Macrophages: A double-edged sword in experimental autoimmune encephalomyelitis. *Immunol. Lett.* **2014**, *160*, 17–22. [[CrossRef](#)]
49. Burg, A.R.; Tse, H.M. Redox-sensitive innate immune pathways during macrophage activation in type 1 diabetes. *Antioxid. Redox Signal.* **2018**, *29*, 1373–1398. [[CrossRef](#)]
50. Hirayama, D.; Iida, T.; Nakase, H. The phagocytic function of macrophage-enforcing innate immunity and tissue homeostasis. *Int. J. Mol. Sci.* **2017**, *19*, 92. [[CrossRef](#)]
51. Schenten, D.; Medzhitov, R. The control of adaptive immune responses by the innate immune system. *Adv. Immunol.* **2011**, *109*, 87–124. [[PubMed](#)]
52. Colbert, J.D.; Cruz, F.M.; Rock, K.L. Cross-presentation of exogenous antigens on MHC I molecules. *Curr. Opin. Immunol.* **2020**, *64*, 1–8. [[CrossRef](#)] [[PubMed](#)]
53. Epelman, S.; Lavine, K.J.; Randolph, G.J. Origin and functions of tissue macrophages. *Immunity* **2014**, *41*, 21–35. [[CrossRef](#)]
54. Ginhoux, F.; Guilliams, M. Tissue-resident macrophage ontogeny and homeostasis. *Immunity* **2016**, *44*, 439–449. [[CrossRef](#)] [[PubMed](#)]
55. Theret, M.; Mounier, R.; Rossi, F. The origins and non-canonical functions of macrophages in development and regeneration. *Development* **2019**, 146. [[CrossRef](#)]
56. Mantovani, A.; Sica, A.; Locati, M. Macrophage polarization comes of age. *Immunity* **2005**, *23*, 344–346. [[CrossRef](#)]
57. Davies, L.C.; Jenkins, S.J.; Allen, J.E.; Taylor, P.R. Tissue-resident macrophages. *Nat. Immunol.* **2013**, *14*, 986–995. [[CrossRef](#)]
58. Mantovani, A.; Biswas, S.K.; Galdiero, M.R.; Sica, A.; Locati, M. Macrophage plasticity and polarization in tissue repair and remodelling. *J. Pathol.* **2013**, *229*, 176–185. [[CrossRef](#)]
59. Kurowska-Stolarska, M.; Stolarski, B.; Kewin, P.; Murphy, G.; Corrigan, C.J.; Ying, S.; Pitman, N.; Mirchandani, A.; Rana, B.; Van Rooijen, N.; et al. IL-33 amplifies the polarization of alternatively activated macrophages that contribute to airway inflammation. *J. Immunol.* **2009**, *183*, 6469–6477. [[CrossRef](#)]
60. Shapouri-Moghaddam, A.; Mohammadian, S.; Vazini, H.; Taghadosi, M.; Esmaeili, S.A.; Mardani, F.; Seifi, B.; Mohammadi, A.; Afshari, J.T.; Sahebkar, A. Macrophage plasticity, polarization, and function in health and disease. *J. Cell Physiol.* **2018**, *233*, 6425–6440. [[CrossRef](#)]
61. Davies, L.C.; Taylor, P.R. Tissue-resident macrophages: Then and now. *Immunology* **2015**, *144*, 541–548. [[CrossRef](#)] [[PubMed](#)]
62. Weigert, A.; Strack, E.; Snodgrass, R.G.; Brüne, B. mPGES-1 and ALOX5/-15 in tumor-associated macrophages. *Cancer Metastasis Rev.* **2018**, *37*, 317–334. [[CrossRef](#)] [[PubMed](#)]
63. Dalli, J.; Serhan, C.N. Pro-resolving mediators in regulating and conferring macrophage function. *Front. Immunol.* **2017**, *8*, 1400. [[CrossRef](#)]
64. Norris, P.C.; Dennis, E.A. A lipidomic perspective on inflammatory macrophage eicosanoid signaling. *Adv. Biol. Regul.* **2014**, *54*, 99–110. [[CrossRef](#)] [[PubMed](#)]
65. Li, X.; Bi, X.; Wang, S.; Zhang, Z.; Li, F.; Zhao, A.Z. Therapeutic Potential of ω-3 polyunsaturated fatty acids in human autoimmune diseases. *Front. Immunol.* **2019**, *10*, 2241. [[CrossRef](#)]
66. Wei, J.; Gronert, K. Eicosanoid and specialized proresolving mediator regulation of lymphoid cells. *Trends Biochem. Sci.* **2019**, *44*, 214–225. [[CrossRef](#)]
67. Janeway, C.A., Jr.; Travers, P.; Walport, M. *Generation of Lymphocytes in Bone Marrow and Thymus*; Garland Science: New York, NY, USA, 2001.
68. Li, M.O.; Rudensky, A.Y. T cell receptor signalling in the control of regulatory T cell differentiation and function. *Nat. Rev. Immunol.* **2016**, *16*, 220–233. [[CrossRef](#)] [[PubMed](#)]
69. Zhu, J.; Paul, W.E. CD4 T cells: Fates, functions, and faults. *Blood* **2008**, *5*, 1557–1569. [[CrossRef](#)]
70. Mak, T.W.; Saunders, M.E.; Jett, B.D. B cell development, activation and effector functions. *Primer Immune Response* **2014**, *2*, 111–142. [[CrossRef](#)]
71. Bonilla, F.A.; Oettgen, H.C. Adaptive immunity. *J. Allergy Clin. Immunol.* **2010**, *125*, S33–S40. [[CrossRef](#)]
72. Hooper, K.M.; Kong, W.; Ganea, D. Prostaglandin E₂ inhibits Tr1 cell differentiation through suppression of c-Maf. *PLoS ONE* **2017**, *12*, e0179184. [[CrossRef](#)]
73. Aoki, T.; Frosen, J.; Fukuda, M.; Bando, K.; Shioi, G.; Tsuji, K.; Ollikainen, E.; Nozaki, K.; Laakkonen, J.; Narumiya, S. Prostaglandin E₂-EP2-NF-κB signaling in macrophages as a potential therapeutic target for intracranial aneurysms. *Sci. Signal.* **2017**, *10*. [[CrossRef](#)]
74. Ganapathy, V.; Gurlo, T.; Jarstadmarken, H.O.; Von Grafenstein, H. Regulation of TCR-induced IFN-γ release from islet-reactive non-obese diabetic CD8⁺ T cells by prostaglandin E₂ receptor signaling. *Int. Immunol.* **2000**, *12*, 851–860. [[CrossRef](#)]

75. Ling, J.J.; Sun, Y.J.; Zhu, D.Y.; Chen, Q.; Han, X. Potential role of NO in modulation of COX-2 expression and PGE2 production in pancreatic beta-cells. *Acta Biochim. Biophys. Sin.* **2005**, *37*, 139–146. [[CrossRef](#)] [[PubMed](#)]
76. Boizel, R.; Bruttman, G.; Benhamou, P.Y.; Halimi, S.; Stanke-Labesque, F. Regulation of oxidative stress and inflammation by glycaemic control: Evidence for reversible activation of the 5-lipoxygenase pathway in type 1, but not in type 2 diabetes. *Diabetologia* **2010**, *53*, 2068–2070. [[CrossRef](#)]
77. Chakrabarti, S.K.; Cole, B.K.; Wen, Y.; Keller, S.R.; Nadler, J.L. 12/15-lipoxygenase products induce inflammation and impair insulin signaling in 3T3-L1 adipocytes. *Obesity* **2009**, *17*, 1657–1663. [[CrossRef](#)]
78. Imai, Y.; Dobrian, A.D.; Morris, M.A.; Taylor-Fishwick, D.A.; Nadler, J.L. Lipids and immunoinflammatory pathways of beta cell destruction. *Diabetologia* **2016**, *59*, 673–678. [[CrossRef](#)] [[PubMed](#)]
79. Bauman, J.A.; Li, S.D.; Yang, A.; Huang, L.; Kole, R. Anti-tumor activity of splice-switching oligonucleotides. *Nucleic Acids Res.* **2010**, *38*, 8348–8356. [[CrossRef](#)] [[PubMed](#)]
80. Inceoglu, B.; Bettaieb, A.; Haj, F.G.; Gomes, A.V.; Hammock, B.D. Modulation of mitochondrial dysfunction and endoplasmic reticulum stress are key mechanisms for the wide-ranging actions of epoxy fatty acids and soluble epoxide hydrolase inhibitors. *Prostaglandins Other Lipid Mediat.* **2017**, *133*, 68–78. [[CrossRef](#)] [[PubMed](#)]
81. Jung, T.W.; Hwang, H.J.; Hong, H.C.; Choi, H.Y.; Yoo, H.J.; Baik, S.H.; Choi, K.M. Resolvin D1 reduces ER stress-induced apoptosis and triglyceride accumulation through JNK pathway in HepG2 cells. *Mol. Cell. Endocrinol.* **2014**, *391*, 30–40. [[CrossRef](#)] [[PubMed](#)]
82. Huang, H.; Weng, J.; Wang, M.H. EETs/sEH in diabetes and obesity-induced cardiovascular diseases. *Prostaglandins Other Lipid Mediat.* **2016**, *125*, 80–89. [[CrossRef](#)]
83. Jouihan, S.A.; Zuloaga, K.L.; Zhang, W.; Shangraw, R.E.; Krasnow, S.M.; Marks, D.L.; Alkayed, N.J. Role of soluble epoxide hydrolase in exacerbation of stroke by streptozotocin-induced type 1 diabetes mellitus. *J. Cereb. Blood Flow Metab.* **2013**, *33*, 1650–1656. [[CrossRef](#)]
84. Luther, J.M.; Brown, N.J. Epoxyeicosatrienoic acids and glucose homeostasis in mice and men. *Prostaglandins Other Lipid Mediat.* **2016**, *125*, 2–7. [[CrossRef](#)] [[PubMed](#)]
85. Rodriguez, M.; Clare-Salzler, M. Eicosanoid imbalance in the NOD mouse is related to a dysregulation in soluble epoxide hydrolase and 15-PGDH expression. *Ann. N. Y. Acad. Sci.* **2006**, *1079*, 130–134. [[CrossRef](#)] [[PubMed](#)]
86. Kim, R.; Emi, M.; Tanabe, K. Cancer cell immune escape and tumor progression by exploitation of anti-inflammatory and pro-inflammatory responses. *Cancer Biol. Ther.* **2005**, *4*, 924–933. [[CrossRef](#)] [[PubMed](#)]
87. Atkinson, M.A. Mechanisms underlying the loss of self tolerance in NOD mice. *Res. Immunol.* **1997**, *148*, 301–306. [[CrossRef](#)]
88. Lauber, K.; Bohn, E.; Krober, S.M.; Xiao, Y.J.; Blumenthal, S.G.; Lindemann, R.K.; Marini, P.; Wiedig, C.; Zobywalski, A.; Baksh, S.; et al. Apoptotic cells induce migration of phagocytes via caspase-3-mediated release of a lipid attraction signal. *Cell* **2003**, *113*, 717–730. [[CrossRef](#)]
89. Ming, W.J.; Bersani, L.; Mantovani, A. Tumor necrosis factor is chemotactic for monocytes and polymorphonuclear leukocytes. *J. Immunol.* **1987**, *138*, 1469–1474.
90. Ma, K.; Nunemaker, C.S.; Wu, R.; Chakrabarti, S.K.; Taylor-Fishwick, D.A.; Nadler, J.L. 12-Lipoxygenase products reduce insulin secretion and {beta}-cell viability in human islets. *J. Clin. Endocrinol. Metab.* **2010**, *95*, 887–893. [[CrossRef](#)]
91. McDuffie, M.; Maybee, N.A.; Keller, S.R.; Stevens, B.K.; Garmey, J.C.; Morris, M.A.; Kropf, E.; Rival, C.; Ma, K.; Carter, J.D.; et al. Nonobese diabetic (NOD) mice congenic for a targeted deletion of 12/15-lipoxygenase are protected from autoimmune diabetes. *Diabetes* **2008**, *57*, 199–208. [[CrossRef](#)]
92. Prasad, K.M.; Thimmalapura, P.R.; Woode, E.A.; Nadler, J.L. Evidence that increased 12-lipoxygenase expression impairs pancreatic beta cell function and viability. *Biochem. Biophys. Res. Commun.* **2003**, *308*, 427–432. [[CrossRef](#)]
93. Weaver, J.R.; Holman, T.R.; Imai, Y.; Jadhav, A.; Kenyon, V.; Maloney, D.J.; Nadler, J.L.; Rai, G.; Simeonov, A.; Taylor-Fishwick, D.A. Integration of pro-inflammatory cytokines, 12-lipoxygenase and NOX-1 in pancreatic islet beta cell dysfunction. *Mol. Cell. Endocrinol.* **2012**, *358*, 88–95. [[CrossRef](#)] [[PubMed](#)]
94. Wen, Y.; Gu, J.; Vandenhoff, G.E.; Liu, X.; Nadler, J.L. Role of 12/15-lipoxygenase in the expression of MCP-1 in mouse macrophages. *Am. J. Physiol. Heart Circ. Physiol.* **2008**, *294*, H1933–H1938. [[CrossRef](#)] [[PubMed](#)]
95. Zhou, R.H.; Pesant, S.; Cohn, H.I.; Eckhart, A.D. Enhanced sterol response element-binding protein in postintervention restenotic blood vessels plays an important role in vascular smooth muscle proliferation. *Life Sci.* **2008**, *82*, 174–181. [[CrossRef](#)] [[PubMed](#)]
96. Bleich, D.; Chen, S.; Zipser, B.; Sun, D.; Funk, C.D.; Nadler, J.L. Resistance to type 1 diabetes induction in 12-lipoxygenase knockout mice. *J. Clin. Invest.* **1999**, *103*, 1431–1436. [[CrossRef](#)]
97. Green-Mitchell, S.M.; Tersey, S.A.; Cole, B.K.; Ma, K.; Kuhn, N.S.; Cunningham, T.D.; Maybee, N.A.; Chakrabarti, S.K.; McDuffie, M.; Taylor-Fishwick, D.A.; et al. Deletion of 12/15-lipoxygenase alters macrophage and islet function in NOD-Alox15(null) mice, leading to protection against type 1 diabetes development. *PLoS ONE* **2013**, *8*, e56763. [[CrossRef](#)] [[PubMed](#)]
98. Grzesik, W.J.; Nadler, J.L.; Machida, Y.; Nadler, J.L.; Imai, Y.; Morris, M.A. Expression pattern of 12-lipoxygenase in human islets with type 1 diabetes and type 2 diabetes. *J. Clin. Endocrinol. Metab.* **2015**, *100*, E387–E395. [[CrossRef](#)]
99. Bennett, M.; Gilroy, D.W. Lipid mediators in inflammation. *Microbiol. Spectr.* **2016**, *4*. [[CrossRef](#)]
100. Serhan, C.N.; Chiang, N.; Van Dyke, T.E. Resolving inflammation: Dual anti-inflammatory and pro-resolution lipid mediators. *Nat. Rev. Immunol.* **2008**, *8*, 349–361. [[CrossRef](#)]
101. Tuosto, L.; Xu, C. Editorial: Membrane lipids in T cell functions. *Front. Immunol.* **2018**, *9*, 1608. [[CrossRef](#)]
102. Gu, L.; Tseng, S.C.; Rollins, B.J. Monocyte chemoattractant protein-1. *Chem. Immunol.* **1999**, *72*, 7–29. [[PubMed](#)]

103. Cathcart, M.K. Signal-activated phospholipase regulation of leukocyte chemotaxis. *J. Lipid Res.* **2009**, *50*, S231–S236. [[CrossRef](#)]
104. Kam, Y.; Exton, J.H. Phospholipase D activity is required for actin stress fiber formation in fibroblasts. *Mol. Cell. Biol.* **2001**, *21*, 4055–4066. [[CrossRef](#)]
105. Tang, J.; Kriz, R.W.; Wolfman, N.; Shaffer, M.; Seehra, J.; Jones, S.S. A novel cytosolic calcium-independent phospholipase A₂ contains eight ankyrin motifs. *J. Biol. Chem.* **1997**, *272*, 8567–8575. [[CrossRef](#)] [[PubMed](#)]
106. Jenkins, C.M.; Han, X.; Mancuso, D.J.; Gross, R.W. Identification of calcium-independent phospholipase A₂ (iPLA₂) beta, and not iPLA₂gamma, as the mediator of arginine vasopressin-induced arachidonic acid release in A-10 smooth muscle cells. Enantioselective mechanism-based discrimination of mammalian iPLA₂s. *J. Biol. Chem.* **2002**, *277*, 32807–32814. [[PubMed](#)]
107. Liu, S.; Xie, Z.; Zhao, Q.; Pang, H.; Turk, J.; Calderon, L.; Su, W.; Zhao, G.; Xu, H.; Gong, M.C.; et al. Smooth muscle-specific expression of calcium-independent phospholipase A₂beta (iPLA₂b) participates in the initiation and early progression of vascular inflammation and neointima formation. *J. Biol. Chem.* **2012**, *287*, 24739–24753. [[CrossRef](#)] [[PubMed](#)]
108. Zhao, X.; Wang, D.; Zhao, Z.; Xiao, Y.; Sengupta, S.; Xiao, Y.; Zhang, R.; Lauber, K.; Wesselborg, S.; Feng, L.; et al. Caspase-3-dependent activation of calcium-independent phospholipase A₂ enhances cell migration in non-apoptotic ovarian cancer cells. *J. Biol. Chem.* **2006**, *281*, 29357–29368. [[CrossRef](#)] [[PubMed](#)]
109. Lee, J.G.; Lim, E.J.; Park, D.W.; Lee, S.H.; Kim, J.R.; Baek, S.H. A combination of Lox-1 and Nox1 regulates TLR9-mediated foam cell formation. *Cell Signal.* **2008**, *20*, 2266–2275. [[CrossRef](#)] [[PubMed](#)]
110. Funk, J.L.; Feingold, K.R.; Moser, A.H.; Grunfeld, C. Lipopolysaccharide stimulation of RAW 264.7 macrophages induces lipid accumulation and foam cell formation. *Atherosclerosis* **1993**, *98*, 67–82. [[CrossRef](#)]
111. Taniyama, Y.; Griendling, K.K. Reactive oxygen species in the vasculature: Molecular and cellular mechanisms. *Hypertension* **2003**, *42*, 1075–1081. [[CrossRef](#)]
112. Shirai, T.; Hilhorst, M.; Harrison, D.G.; Goronzy, J.J.; Weyand, C.M. Macrophages in vascular inflammation. From atherosclerosis to vasculitis. *Autoimmunity* **2015**, *48*, 139–151. [[CrossRef](#)]
113. Lei, X.; Zhang, S.; Barbour, S.E.; Bohrer, A.; Ford, E.L.; Koizumi, A.; Papa, F.R.; Ramanadham, S. Spontaneous development of endoplasmic reticulum stress that can lead to diabetes mellitus is associated with higher calcium-independent phospholipase A₂ expression: A role for regulation by SREBP-1. *J. Biol. Chem.* **2010**, *285*, 6693–6705. [[CrossRef](#)]
114. Seashols, S.J.; Del Castillo Olivares, A.; Gil, G.; Barbour, S.E. Regulation of group VIA phospholipase A₂ expression by sterol availability. *Biochim. Biophys. Acta* **2004**, *1684*, 29–37. [[CrossRef](#)]
115. Smani, T.; Zakharov, S.I.; Csutora, P.; Leno, E.; Trepakova, E.S.; Bolotina, V.M. A novel mechanism for the store-operated calcium influx pathway. *Nat. Cell. Biol.* **2004**, *6*, 113–120. [[CrossRef](#)] [[PubMed](#)]
116. Wolf, M.J.; Wang, J.; Turk, J.; Gross, R.W. Depletion of intracellular calcium stores activates smooth muscle cell calcium-independent phospholipase A₂. A novel mechanism underlying arachidonic acid mobilization. *J. Biol. Chem.* **1997**, *272*, 1522–1526. [[CrossRef](#)]
117. Lei, X.; Bone, R.N.; Ali, T.; Wohltmann, M.; Gai, Y.; Goodwin, K.J.; Bohrer, A.E.; Turk, J.; Ramanadham, S. Genetic modulation of islet beta-cell iPLA₂b expression provides evidence for its impact on beta-cell apoptosis and autophagy. *Islets* **2013**, *5*, 29–44. [[CrossRef](#)] [[PubMed](#)]
118. Lei, X.; Bone, R.N.; Ali, T.; Zhang, S.; Bohrer, A.; Tse, H.M.; Bidasee, K.R.; Ramanadham, S. Evidence of contribution of iPLA₂b-mediated events during islet beta-cell apoptosis due to proinflammatory cytokines suggests a role for iPLA₂b in T1D development. *Endocrinology* **2014**, *155*, 3352–3364. [[CrossRef](#)] [[PubMed](#)]
119. Marentette, J.; Kolar, G.; Mchowat, J. Increased susceptibility to bladder inflammation in smokers: Targeting the PAF-PAF receptor interaction to manage inflammatory cell recruitment. *Physiol. Rep.* **2015**, *3*. [[CrossRef](#)] [[PubMed](#)]
120. Green, M.; Filippou, A.; Sant, G.; Theoharides, T.C. Expression of intercellular adhesion molecules in the bladder of patients with interstitial cystitis. *Urology* **2004**, *63*, 688–693. [[CrossRef](#)]
121. Rastogi, P.; White, M.C.; Rickard, A.; Mchowat, J. Potential mechanism for recruitment and migration of CD133 positive cells to areas of vascular inflammation. *Thromb. Res.* **2008**, *123*, 258–266. [[CrossRef](#)]
122. Rickard, A.; Portell, C.; Kell, P.J.; Vinson, S.M.; Mchowat, J. Protease-activated receptor stimulation activates a Ca²⁺-independent phospholipase A₂ in bladder microvascular endothelial cells. *Am. J. Physiol. Renal Physiol.* **2005**, *288*, F714–F721. [[CrossRef](#)]
123. Ueno, N.; Taketomi, Y.; Yamamoto, K.; Hirabayashi, T.; Kamei, D.; Kita, Y.; Shimizu, T.; Shinzawa, K.; Tsujimoto, Y.; Ikeda, K.; et al. Analysis of two major intracellular phospholipases A₂ (PLA₂) in mast cells reveals crucial contribution of cytosolic PLA₂alpha, not Ca²⁺-independent PLA₂beta, to lipid mobilization in proximal mast cells and distal fibroblasts. *J. Biol. Chem.* **2011**, *286*, 37249–37263. [[CrossRef](#)] [[PubMed](#)]
124. Gutierrez, D.A.; Fu, W.; Schonefeldt, S.; Feyereabend, T.B.; Ortiz-Lopez, A.; Lampi, Y.; Liston, A.; Mathis, D.; Rodewald, H.R. Type 1 diabetes in NOD mice unaffected by mast cell deficiency. *Diabetes* **2014**, *63*, 3827–3834. [[CrossRef](#)] [[PubMed](#)]
125. Martino, L.; Masini, M.; Bugliani, M.; Marselli, L.; Suleiman, M.; Boggi, U.; Nogueira, T.C.; Filipponi, F.; Occhipinti, M.; Campani, D.; et al. Mast cells infiltrate pancreatic islets in human type 1 diabetes. *Diabetologia* **2015**, *58*, 2554–2562. [[CrossRef](#)]
126. Ryan, E.P.; Pollock, S.J.; Kaur, K.; Felgar, R.E.; Bernstein, S.H.; Chiorazzi, N.; Phipps, R.P. Constitutive and activation-inducible cyclooxygenase-2 expression enhances survival of chronic lymphocytic leukemia B cells. *Clin. Immunol.* **2006**, *120*, 76–90. [[CrossRef](#)] [[PubMed](#)]
127. Secchiero, P.; Barbarotto, E.; Gonelli, A.; Tiribelli, M.; Zerbinati, C.; Celeghini, C.; Agostinelli, C.; Pileri, S.A.; Zauli, G. Potential pathogenetic implications of cyclooxygenase-2 overexpression in B chronic lymphoid leukemia cells. *Am. J. Pathol.* **2005**, *167*, 1599–1607. [[CrossRef](#)]

128. Ray, D.M.; Akbiyik, F.; Bernstein, S.H.; Phipps, R.P. CD40 engagement prevents peroxisome proliferator-activated receptor gamma agonist-induced apoptosis of B lymphocytes and B lymphoma cells by an NF-kappaB-dependent mechanism. *J. Immunol.* **2005**, *174*, 4060–4069. [[CrossRef](#)]
129. Ricciotti, E.; Fitzgerald, G.A. Prostaglandins and inflammation. *Arterioscler. Thromb. Vasc. Biol.* **2011**, *31*, 986–1000. [[CrossRef](#)]
130. Roshak, A.K.; Capper, E.A.; Stevenson, C.; Eichman, C.; Marshall, L.A. Human calcium-independent phospholipase A₂ mediates lymphocyte proliferation. *J. Biol. Chem.* **2000**, *275*, 35692–35698. [[CrossRef](#)]
131. Patel, M.L.; Singh, J.; Niknami, M.; Kurek, C.; Yao, M.; Lu, S.; Maclean, F.; King, N.J.; Gelb, M.H.; Scott, K.F.; et al. Cytosolic phospholipase A₂-alpha: A potential therapeutic target for prostate cancer. *Clin. Cancer Res.* **2008**, *14*, 8070–8079. [[CrossRef](#)]
132. Guriec, N.; Le Jossic-Corcoss, C.; Simon, B.; Ianotto, J.C.; Tempescul, A.; Dreano, Y.; Salaun, J.P.; Berthou, C.; Corcos, L. The arachidonic acid-LTB₄-BLT2 pathway enhances human B-CLL aggressiveness. *Biochim. Biophys. Acta* **2014**, *1842*, 2096–2105. [[CrossRef](#)] [[PubMed](#)]
133. Haffner, S.M.; Lehto, S.; Ronnema, T.; Pyorala, K.; Laakso, M. Mortality from coronary heart disease in subjects with type 2 diabetes and in nondiabetic subjects with and without prior myocardial infarction. *N. Engl. J. Med.* **1998**, *339*, 229–234. [[CrossRef](#)] [[PubMed](#)]
134. Brownlee, M. Biochemistry and molecular cell biology of diabetic complications. *Nature* **2001**, *414*, 813–820. [[CrossRef](#)] [[PubMed](#)]
135. Panday, A.; Sahoo, M.K.; Osorio, D.; Batra, S. NADPH oxidases: An overview from structure to innate immunity-associated pathologies. *Cell Mol. Immunol.* **2015**, *12*, 5–23. [[CrossRef](#)] [[PubMed](#)]
136. Lee, C.F.; Qiao, M.; Schroder, K.; Zhao, Q.; Asmis, R. Nox4 is a novel inducible source of reactive oxygen species in monocytes and macrophages and mediates oxidized low density lipoprotein-induced macrophage death. *Circ. Res.* **2010**, *106*, 1489–1497. [[CrossRef](#)] [[PubMed](#)]
137. Ullevig, S.; Zhao, Q.; Lee, C.F.; Seok Kim, H.; Zamora, D.; Asmis, R. NADPH oxidase 4 mediates monocyte priming and accelerated chemotaxis induced by metabolic stress. *Arterioscler. Thromb. Vasc. Biol.* **2012**, *32*, 415–426. [[CrossRef](#)]
138. Ago, T.; Kuroda, J.; Pain, J.; Fu, C.; Li, H.; Sadoshima, J. Upregulation of Nox4 by hypertrophic stimuli promotes apoptosis and mitochondrial dysfunction in cardiac myocytes. *Circ. Res.* **2010**, *106*, 1253–1264. [[CrossRef](#)]
139. Block, K.; Gorin, Y.; Abboud, H.E. Subcellular localization of Nox4 and regulation in diabetes. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 14385–14390. [[CrossRef](#)]
140. Lei, X.; Zhang, S.; Bohrer, A.; Bao, S.; Song, H.; Ramanadham, S. The group VIA calcium-independent phospholipase A₂ participates in ER stress-induced INS-1 insulinoma cell apoptosis by promoting ceramide generation via hydrolysis of sphingomyelins by neutral sphingomyelinase. *Biochemistry* **2007**, *46*, 10170–10185. [[CrossRef](#)]
141. Nelson, A.J.; Stephenson, D.J.; Cardona, C.L.; Lei, X.; Almutairi, A.; White, T.D.; Tusing, Y.G.; Park, M.A.; Barbour, S.E.; Chalfant, C.E.; et al. Macrophage polarization is linked to Ca²⁺-independent phospholipase A₂beta-derived lipids and cross-cell signaling in mice. *J. Lipid Res.* **2020**, *61*, 143–158. [[CrossRef](#)]
142. Gil-De-Gomez, L.; Monge, P.; Rodriguez, J.P.; Astudillo, A.M.; Balboa, M.A.; Balsinde, J. Phospholipid arachidonic acid remodeling during phagocytosis in mouse peritoneal macrophages. *Biomedicines* **2020**, *8*, 274. [[CrossRef](#)] [[PubMed](#)]
143. Goldsmith, M.; Daka, A.; Lamour, N.F.; Mashlach, R.; Glucksam, Y.; Meijler, M.M.; Chalfant, C.E.; Zor, T. A ceramide analog inhibits cPLA₂ activity and consequent PGE₂ formation in LPS-stimulated macrophages. *Immunol. Lett.* **2011**, *135*, 136–143. [[CrossRef](#)]
144. Naraba, H.; Murakami, M.; Matsumoto, H.; Shimbara, S.; Ueno, A.; Kudo, I.; Oh-Ishi, S. Segregated coupling of phospholipases A₂, cyclooxygenases, and terminal prostanoid synthases in different phases of prostanoid biosynthesis in rat peritoneal macrophages. *J. Immunol.* **1998**, *160*, 2974–2982.
145. Qi, H.Y.; Shelhamer, J.H. Toll-like receptor 4 signaling regulates cytosolic phospholipase A₂ activation and lipid generation in lipopolysaccharide-stimulated macrophages. *J. Biol. Chem.* **2005**, *280*, 38969–38975. [[CrossRef](#)]
146. Ruiperez, V.; Casas, J.; Balboa, M.A.; Balsinde, J. Group V phospholipase A₂-derived lysophosphatidylcholine mediates cyclooxygenase-2 induction in lipopolysaccharide-stimulated macrophages. *J. Immunol.* **2007**, *1790*, 631–638. [[CrossRef](#)] [[PubMed](#)]
147. Zhu, C.; Zhou, J.; Li, T.; Mu, J.; Jin, L.; Li, S. Urocortin participates in LPS-induced apoptosis of THP-1 macrophages via S1P-cPLA₂ signaling pathway. *Eur. J. Pharmacol.* **2020**, *887*, 173559. [[CrossRef](#)] [[PubMed](#)]
148. Szablewski, L. Role of immune system in type 1 diabetes mellitus pathogenesis. *Int. Immunopharmacol.* **2014**, *22*, 182–191. [[CrossRef](#)]
149. Mathis, D.; Vence, L.; Benoist, C. Beta-cell death during progression to diabetes. *Nature* **2001**, *414*, 792–798. [[CrossRef](#)]
150. Pugliese, A. Autoreactive T cells in type 1 diabetes. *J. Clin. Investig.* **2017**, *127*, 2881–2891. [[CrossRef](#)]
151. Padgett, L.E.; Broniowska, K.A.; Hansen, P.A.; Corbett, J.A.; Tse, H.M. The role of reactive oxygen species and proinflammatory cytokines in type 1 diabetes pathogenesis. *Ann. N. Y. Acad. Sci.* **2013**, *1281*, 16–35. [[CrossRef](#)]
152. Mantovani, A.; Sica, A.; Sozzani, S.; Allavena, P.; Vecchi, A.; Locati, M. The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol.* **2004**, *25*, 677–686. [[CrossRef](#)] [[PubMed](#)]
153. Martinez, F.O.; Gordon, S. The M1 and M2 paradigm of macrophage activation: Time for reassessment. *F1000Prime Rep.* **2014**, *6*, 13. [[CrossRef](#)] [[PubMed](#)]
154. Calderon, B.; Suri, A.; Unanue, E.R. In CD4+ T-cell-induced diabetes, macrophages are the final effector cells that mediate islet beta-cell killing: Studies from an acute model. *Am. J. Pathol.* **2006**, *169*, 2137–2147. [[CrossRef](#)] [[PubMed](#)]

155. Parsa, R.; Andresen, P.; Gillett, A.; Mia, S.; Zhang, X.M.; Mayans, S.; Holmberg, D.; Harris, R.A. Adoptive transfer of immunomodulatory M2 macrophages prevents type 1 diabetes in NOD mice. *Diabetes* **2012**, *61*, 2881–2892. [[CrossRef](#)] [[PubMed](#)]
156. Zakharov, P.N.; Hu, H.; Wan, X.; Unanue, E.R. Single-cell RNA sequencing of murine islets shows high cellular complexity at all stages of autoimmune diabetes. *J. Exp. Med.* **2020**, *217*. [[CrossRef](#)]
157. Chen, C.; Cohrs, C.M.; Stertmann, J.; Bozsak, R.; Speier, S. Human beta cell mass and function in diabetes: Recent advances in knowledge and technologies to understand disease pathogenesis. *Mol. Metab.* **2017**, *6*, 943–957. [[CrossRef](#)]
158. Unanue, E.R. Macrophages in endocrine glands, with emphasis on pancreatic islets. *Microbiol. Spectr.* **2016**, *4*. [[CrossRef](#)]
159. Calderon, B.; Suri, A.; Miller, M.J.; Unanue, E.R. Dendritic cells in islets of Langerhans constitutively present beta cell-derived peptides bound to their class II MHC molecules. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 6121–6126. [[CrossRef](#)] [[PubMed](#)]
160. Mohan, J.F.; Levisetti, M.G.; Calderon, B.; Herzog, J.W.; Petzold, S.J.; Unanue, E.R. Unique autoreactive T cells recognize insulin peptides generated within the islets of Langerhans in autoimmune diabetes. *Nat. Immunol.* **2010**, *11*, 350–354. [[CrossRef](#)]
161. Feduska, J.M.; Tse, H.M. The proinflammatory effects of macrophage-derived NADPH oxidase function in autoimmune diabetes. *Free Radic. Biol. Med.* **2018**, *125*, 81–89. [[CrossRef](#)]
162. Burrack, A.L.; Martinov, T.; Fife, B.T. T Cell-mediated beta cell destruction: Autoimmunity and alloimmunity in the context of type 1 diabetes. *Front. Endocrinol.* **2017**, *8*, 343. [[CrossRef](#)]
163. Dimeglio, L.A.; Evans-Molina, C.; Oram, R.A. Type 1 diabetes. *Lancet* **2018**, *391*, 2449–2462. [[CrossRef](#)]
164. Sun, Y.; Wang, Y.; Li, J.-H.; Zhu, S.-H.; Tang, H.-T.; Xia, Z.-F. Macrophage migration inhibitory factor counter-regulates dexamethasone-induced annexin 1 expression and influences the release of eicosanoids in murine macrophages. *Immunology* **2013**, *140*, 250–258. [[CrossRef](#)]
165. Duan, L.; Gan, H.; Arm, J.; Remold, H.G. Cytosolic phospholipase A₂ participates with TNF- α in the induction of apoptosis of human macrophages infected with Mycobacterium tuberculosis H37Ra. *J. Immunol.* **2001**, *166*, 7469–7476. [[CrossRef](#)]
166. Nikolic, D.M.; Gong, M.C.; Turk, J.; Post, S.R. Class A scavenger receptor-mediated macrophage adhesion requires coupling of calcium-independent phospholipase A₂ and 12/15-lipoxygenase to Rac and Cdc42 activation. *J. Biol. Chem.* **2007**, *282*, 33405–33411. [[CrossRef](#)] [[PubMed](#)]
167. Dilorenzo, T.; Graser, R.; Ono, T.; Christianson, G.; Chapman, H.; Roopenian, D.; Nathenson, S.; Serreze, D. Major histocompatibility complex class I-restricted T cells are required for all but the end stages of diabetes development in nonobese diabetic mice and use a prevalent T cell receptor alpha chain gene rearrangement. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 12438–12543. [[CrossRef](#)] [[PubMed](#)]
168. Rahnema, P.; Shimoni, Y.; Nygren, A. Reduced conduction reserve in the diabetic rat heart: Role of iPLA₂ activation in the response to ischemia. *Am. J. Physiol. Heart Circ. Physiol.* **2011**, *300*, H326–H334. [[CrossRef](#)] [[PubMed](#)]
169. Nelson, A.J.; Stephenson, D.J.; Bone, R.N.; Cardona, C.L.; Park, M.A.; Tusing, Y.G.; Lei, X.; Kokotos, G.; Graves, C.L.; Mathews, C.E.; et al. Lipid mediators and biomarkers associated with type 1 diabetes development. *JCI Insight* **2020**, *5*. [[CrossRef](#)] [[PubMed](#)]
170. Held, W.; Macdonald, H.R.; Weissman, I.L.; Hess, M.W.; Mueller, C. Genes encoding tumor necrosis factor alpha and granzyme A are expressed during development of autoimmune diabetes. *Proc. Natl. Acad. Sci. USA* **1990**, *87*, 2239–2243. [[CrossRef](#)]
171. Herrera, P.L.; Harlan, D.M.; Vassalli, P. A mouse CD8 T cell-mediated acute autoimmune diabetes independent of the perforin and Fas cytotoxic pathways: Possible role of membrane TNF. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 279–284. [[CrossRef](#)]
172. Aggarwal, B.B.; Vijayalakshmi, R.V.; Sung, B. Targeting inflammatory pathways for prevention and therapy of cancer: Short-term friend, long-term foe. *Clin. Cancer Res.* **2009**, *15*, 425–430. [[CrossRef](#)]
173. Lin, W.W.; Karin, M. A cytokine-mediated link between innate immunity, inflammation, and cancer. *J. Clin. Investig.* **2007**, *117*, 1175–1183. [[CrossRef](#)]
174. Jiao, L.; Gan-Schreier, H.; Tuma-Kellner, S.; Stremmel, W.; Chamultrat, W. Sensitization to autoimmune hepatitis in group VIA calcium-independent phospholipase A₂-null mice led to duodenal villous atrophy with apoptosis, goblet cell hyperplasia and leaked bile acids. *Biochim. Biophys. Acta* **2015**, *1852*, 1646–1657. [[CrossRef](#)] [[PubMed](#)]
175. Jiao, L.; Inhoffen, J.; Gan-Schreier, H.; Tuma-Kellner, S.; Stremmel, W.; Sun, Z.; Chamultrat, W. Deficiency of group VIA phospholipase A₂ (iPLA₂b) renders susceptibility for chemical-induced colitis. *Dig. Dis. Sci.* **2015**, *60*, 3590–3602. [[CrossRef](#)] [[PubMed](#)]
176. Xu, W.; Tuma, S.; Katava, N.; Pathil-Warth, A.; Stremmel, W.; Chamultrat, W. Deficiencies of calcium-independent phospholipase A₂ B in vivo causes reduced systemic lipids and lipoproteins concomitant with increased hepatic apoptosis and inflammation. In *Journal of Hepatology, Proceeding of The International Liver Congress™ (EASL), Vienna, Austria, 10–14 April 2012*; EASL: Barcelona, Spain, 2012.
177. Inhoffen, J.; Tuma-Kellner, S.; Straub, B.; Stremmel, W.; Chamultrat, W. Deficiency of iPLA₂beta primes immune cells for proinflammation: Potential involvement in age-related mesenteric lymph node lymphoma. *Cancers* **2015**, *7*, 2427–2442. [[CrossRef](#)] [[PubMed](#)]
178. Park, D.R.; Thomsen, A.R.; Frevert, C.W.; Pham, U.; Skerrett, S.J.; Kiener, P.A.; Liles, W.C. Fas (CD95) induces proinflammatory cytokine responses by human monocytes and monocyte-derived macrophages. *J. Immunol.* **2003**, *170*, 6209–6216. [[CrossRef](#)] [[PubMed](#)]
179. Albert, M.L. Death-defying immunity: Do apoptotic cells influence antigen processing and presentation? *Nat. Rev. Immunol.* **2004**, *4*, 223–231. [[CrossRef](#)] [[PubMed](#)]
180. Scaffidi, P.; Misteli, T.; Bianchi, M.E. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* **2002**, *418*, 191–195. [[CrossRef](#)]
181. Xavier, R.J.; Podolsky, D.K. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* **2007**, *448*, 427–434. [[CrossRef](#)]

182. Murase, R.; Sato, H.; Yamamoto, K.; Ushida, A.; Nishito, Y.; Ikeda, K.; Kobayashi, T.; Yamamoto, T.; Taketomi, Y.; Murakami, M. Group X secreted phospholipase A₂ releases omega3 polyunsaturated fatty acids, suppresses colitis, and promotes sperm fertility. *J. Biol. Chem.* **2016**, *291*, 6895–6911. [[CrossRef](#)]
183. Sharma, J.; Blase, J.R.; Hoft, D.F.; Marentette, J.O.; Turk, J.; Mchowat, J. Mice with genetic deletion of group via phospholipase a2beta exhibit impaired macrophage function and increased parasite load in *Trypanosoma cruzi*-induced myocarditis. *Infect. Immun.* **2016**, *84*, 1137–1142. [[CrossRef](#)]
184. Bern, C.; Verastegui, M.; Gilman, R.H.; Lafuente, C.; Galdos-Cardenas, G.; Calderon, M.; Pacori, J.; Del Carmen Abastoflor, M.; Aparicio, H.; Brady, M.F.; et al. Congenital *Trypanosoma cruzi* transmission in Santa Cruz, Bolivia. *Clin. Infect. Dis.* **2009**, *49*, 1667–1674. [[CrossRef](#)]
185. Rassi, A., Jr.; Rassi, A.; Little, W.C. Chagas' heart disease. *Clin. Cardiol.* **2000**, *23*, 883–889.
186. Lannes-Vieira, J.; Silverio, J.C.; Pereira, I.R.; Vinagre, N.F.; Carvalho, C.M.; Paiva, C.N.; Silva Da, A.A. Chronic *Trypanosoma cruzi*-elicited cardiomyopathy: From the discovery to the proposal of rational therapeutic interventions targeting cell adhesion molecules and chemokine receptors—How to make a dream come true. *Mem. Inst. Oswaldo Cruz* **2009**, *104*, 226–235. [[CrossRef](#)] [[PubMed](#)]
187. Dinarello, C.A.; Simon, A.; Van Der Meer, J.W. Treating inflammation by blocking interleukin-1 in a broad spectrum of diseases. *Nat. Rev. Drug Discov.* **2012**, *11*, 633–652. [[CrossRef](#)]
188. Lamkanfi, M.; Dixit, V.M. Mechanisms and functions of inflammasomes. *Cell* **2014**, *157*, 1013–1022. [[CrossRef](#)]
189. Ehlers, M.R. Immune-modulating effects of alpha-1 antitrypsin. *Biol. Chem.* **2014**, *395*, 1187–1193. [[CrossRef](#)] [[PubMed](#)]
190. Siebers, K.; Fink, B.; Zakrzewicz, A.; Agne, A.; Richter, K.; Konzok, S.; Hecker, A.; Zukunft, S.; Kullmar, M.; Klein, J.; et al. Alpha-1 Antitrypsin Inhibits ATP-Mediated Release of Interleukin-1beta via CD36 and Teceptors. *Front. Immunol.* **2018**, *9*, 877. [[CrossRef](#)] [[PubMed](#)]
191. Zakrzewicz, A.; Richter, K.; Zakrzewicz, D.; Siebers, K.; Damm, J.; Agne, A.; Hecker, A.; Mcintosh, J.M.; Chamulitrat, W.; Krasteva-Christ, G.; et al. SLPI Inhibits ATP-mediated maturation of IL-1beta in human monocytic leukocytes: A Tyer. *Front. Immunol.* **2019**, *10*, 664. [[CrossRef](#)]
192. Moreau, T.; Baranger, K.; Dade, S.; Dallet-Choisy, S.; Guyot, N.; Zani, M.L. Multifaceted roles of human elafin and secretory leukocyte proteinase inhibitor (SLPI), two serine protease inhibitors of the chelonianin family. *Biochimie* **2008**, *90*, 284–295. [[CrossRef](#)]
193. Ma, G.; Greenwell-Wild, T.; Lei, K.; Jin, W.; Swisher, J.; Hardegen, N.; Wild, C.T.; Wahl, S.M. Secretory leukocyte protease inhibitor binds to annexin II, a cofactor for macrophage HIV-1 infection. *J. Exp. Med.* **2004**, *200*, 1337–1346. [[CrossRef](#)] [[PubMed](#)]
194. Gottlieb, P.A.; Alkanani, A.K.; Michels, A.W.; Lewis, E.C.; Shapiro, L.; Dinarello, C.A.; Zipris, D. alpha1-Antitrypsin therapy downregulates toll-like receptor-induced IL-1beta responses in monocytes and myeloid dendritic cells and may improve islet function in recently diagnosed patients with type 1 diabetes. *J. Clin. Endocrinol. Metab.* **2014**, *99*, E1418–E1426. [[CrossRef](#)] [[PubMed](#)]