



Canis aureus (Carnivore: Canidae)

PATRICIA D. MOEHLMAN AND VIRGINIA HAYSEN

www.mammalogy.org

Box 2013, Arusha, Tanzania; EcoHealth Alliance, New York, NY 1001-2320, USA; pdmhbf@aol.com (PDM)
Department of Biological Sciences, Smith College, Northampton, MA 01063, USA; vhayssen@smith.edu (VH)

Abstract: *Canis aureus* (Linnaeus, 1758), the golden jackal, is a medium-sized, wide spread, terrestrial carnivore. It is 1 of 7 species found in the genus *Canis*. It ranges from Africa to Europe, the Middle East, Central Asia, and Southeast Asia. Due to its tolerance of dry habitats and its omnivorous diet, *C. aureus* can live in a wide variety of habitats. It normally lives in open grassland habitat but also occurs in deserts, woodlands, mangroves, and agricultural and rural habitats in India and Bangladesh. It ranges from sea level in Eritrea to 3,500 m in the Bale Mountains of Ethiopia and 2,000 m in India. *C. aureus* is listed as “Least Concern” by the International Union for Conservation of Nature and Natural Resources Red List of Threatened Species version 2016.1.

Key words: canid, carnivore, cooperative breeder, golden jackal, opportunistic omnivore

Synonymies completed 1 June 2016

DOI: 10.1093/mspecies/sey002

Version of Record, first published online April 17, 2018, with fixed content and layout in compliance with Art. 8.1.3.2 ICZN.

Nomenclatural statement.—A life science identifier (LSID) number was obtained for this publication: urn:lsid:zoobank.org:pub:urn:lsid:zoobank.org:pub:9FC747F8-BAA4-4E2A-BC5B-53A20A0A03FA

Canis aureus Linnaeus, 1758

Golden Jackal

Canis aureus Linnaeus, 1758:40. Type locality “oriente;” restricted to “Benná Mts., Laristan, S. Persia” by Thomas (1911:135).

Lupus aureus Linnaeus, 1758:40. No type locality. From a pre-1758 name.

Vulpes indiae orientalis Linnaeus, 1758:41. No type locality. From a pre-1758 name.

[*Canis*] *anthus* F. Cuvier, 1820:plate. Type locality “Senegal.”

Canis variegatus Cretzschmar, 1826:31, plate 10. Type locality “Nubien. Oberes Egypten;” preoccupied.

Canis syriacus Hemprich and Ehrenberg, 1830:sig z, plate 16. Type locality “e monte Lebano.”

Canis lupaster Hemprich and Ehrenberg, 1833:sig ff. Type locality “Fayum, Egypt” (not seen cited in Allen 1939:194).

Canis sacer Hemprich and Ehrenberg, 1833:sig ff. Type locality “Fayum, Egypt” (not seen cited in Allen 1939:194).

Canis riparius Hemprich and Ehrenberg, 1833:sig ff. Type locality “Coast of Abyssinia, near Arkiko” (not seen cited in Allen 1939:195).

Canis aureus indicus Hodgson, 1833:237. Type locality “Himalaya.”

Canis aureus var. *moreotica* I. Geoffroy Saint-Hilaire, 1835:plate 1. Type locality “de Morée.”

Thous anthus: Hamilton Smith, 1839:195. Name combination.

Thous variegatus: Hamilton Smith, 1839:198. Name combination.

Thous sengalensi Hamilton Smith, 1839:201, plate 13. Type locality “Gambia and Senegal.”

Sacalius aureus: Hamilton Smith, 1839:214. Name combination.



Fig. 1.—Adult *Canis aureus* in the Serengeti, Tanzania, © Patricia D. Moehlman.

© The Author(s) 2018. Published by Oxford University Press on behalf of American Society of Mammalogists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

- Sacalius barbarus* [Hamilton Smith, 1839:218](#). Type locality “Tunis;” preoccupied by *Canis barbarus* [Shaw, 1800:311](#) synonym for *Vulpes vulpes*.
- Sacalius indicus*: [Hamilton Smith, 1839:219](#). Name combination.
- C[anis]. dalmatinus* [Wagner, 1841:383](#). No type locality; Dalmatia implied.
- Canis graecus* [Wagner, 1841:383](#). Type locality “Europaei.” Renaming of *C. aureus moreotica* I. Geoffroy Saint-Hilaire, 1835.
- C[anis]. aureus vulgaris* [Wagner, 1841:383](#). No type locality.
- Canis aureus algirensis* [Wagner, 1841:384](#). Type locality “d’Alger” from ‘chacal d’Alger’ I. Geoffroy Saint-Hilaire, 1835:22.
- C[anis]. aureus tripolitanus* [Wagner, 1841:384](#). No type locality; Tripoli, Tunis implied.
- Canis aureus typicus* [Kolenati, 1858:96](#). No type locality; Persia, Georgia, Turkey implied.
- Canis aureus typicus* var *caucasica* [Kolenati, 1858:96](#). No type locality; Caucasus implied.
- Canis aureus* var *syriaco* [Kolenati, 1858:96](#). Unauthorized emendation of *syriacus* [Hemprich and Ehrenberg, 1830](#).
- Canis aureus* var *indica* [Kolenati, 1858:96](#). Unauthorized emendation of *indicus* [Hodgson, 1833](#).
- Canis aureus* var *nubica* [Kolenati, 1858:96](#). Renaming of *Canis variegatus* [Cretzschmar, 1826](#) but Kolenati attributes the name to Rüppell without giving a date.
- Canis aureus* var *algira* [Kolenati, 1858:96](#). Unauthorized emendation of *algirensis* [Wagner, 1841](#).
- Canis aureus* var *senegalensis*: [Kolenati, 1858:96](#). Name combination.
- Canis aureus* var *barbarus*: [Kolenati, 1858:96](#). Name combination.
- Canis aureus balcanicus* [Brusina, 1892:317](#). Type locality “Nardu — nadomak Valpova — na obali Drave, Slavonija.”
- Canis hadramaticus* [Noack, 1896:356](#). Type locality “Hadramaut in Arabien.” A jackal-wolf composite. The jackal was chosen as the lectotype ([Morrison-Scott 1939](#)).
- Canis mensesi* [Noack, 1897:518](#). Type locality “Ostafrikanische.”
- Canis lupus minor* [Mojsisovico, 1897:202, 241, 244](#). Preoccupied by *Canis spelaeus minor* [Wagner, 1829](#) according to [Kretzoi, 1947](#).
- Canis cruesemanni* [Matschie, 1900:145](#). Type locality “Siam.”
- Canis anthus soudanicus* [Thomas, 1903:295](#). Type locality “El Obeid,” Kordofan, Sudan.
- Canis somalicus* [Lorenz, 1906:306](#). Type locality “Ireso bei Agada.”
- Canis gallaënsis* [Lorenz, 1906:307](#). Type locality “Ginea (Arussi).”
- Canis doederleini* [Hilzheimer, 1906a:116](#). Type locality “Oberägypten.”
- Canis thooides* [Hilzheimer, 1906b:364](#). Type locality “Sennaar.”
- Canis lupaster grayi* [Hilzheimer, 1906b:367](#). Type locality “Nordafrika.”
- Canis studei* [Hilzheimer, 1906b:368](#). Type locality “Tunis.”
- C[anis]. mensesi lamperti* [Hilzheimer, 1906b:371](#). No type locality.
- Thos aureus bea* [Heller, 1914:5](#). Type locality “Loita Plains, British East Africa,” Kenya.
- Canis indicus kola* [Wroughton, 1916:651](#). Type locality “Palampur,” India.
- Canis naria* [Wroughton, 1916:651](#). Type locality “Coorg,” west coast of India.
- Canis lanka* [Wroughton, 1916:652](#). Type locality “Mankeni,” Sri Lanka.
- Thos lupaster maroccanus* [Cabrera, 1921:263](#). Type locality “Mogador (Marruecos).”
- Th[os]. [lupaster]. Algirensis*: [Cabrera, 1921:263](#). Name combination.
- Th[os]. a[ureus]. Riparius*: [Cabrera, 1921:264](#). Name combination.
- Th[os]. aureus nubianus* [Cabrera, 1921:264](#). Renaming of *variegatus* [Cretzschmar, 1826](#).
- Canis aureus hungaricus* [Éhik, 1938:11](#). Type locality “Tyukod (Marshy-land Ecsed), Comitat Szatmár, Hungary;” Preoccupied according to [Kretzoi, 1947](#).
- Canis aureus minor*: [Éhik, 1938:13](#). Name combination; preoccupied.
- Thos aureus algirensis*: [Allen, 1939:194](#). Name combination.
- Thos aureus anthus*: [Allen, 1939:194](#). Name combination.
- Thos aureus lupaster*: [Allen, 1939:194](#). Name combination.
- Thos aureus maroccanus*: [Allen, 1939:195](#). Name combination.
- Thos aureus soudanicus*: [Allen, 1939:195](#). Name combination.
- Thos gallaensis*: [Allen, 1939:195](#). Name combination.
- Thos lamperti*: [Allen, 1939:195](#). Name combination.
- [*Canis aureus*] *moreoticus* [Kretzoi, 1947:287](#). Unjustified emendation of *moreotica* I. Geoffroy Saint-Hilaire, 1835.
- Thos aureus ecsedensis* [Kretzoi, 1947:287](#). Renaming of *C. a. hungaricus* [Éhik, 1938](#).
- Canis lupaster doederleini*: [Saleh and Basuony, 2014:49](#). Name combination.
- Canis aureus qattarensis* [Saleh and Basuony, 2014:49](#). Type locality “northern region of the Egyptian Western Desert.”

CONTENT AND CONTEXT. Order Carnivora, family Canidae, subfamily Caninae, genus *Canis*. As many as 13 subspecies of *C. aureus* are distinguished through Europe and Africa ([Allen 1939](#); [Ellerman and Morrison-Scott 1951](#); [Coetsee 1977](#)). However, these subspecies may be part of a 3-species complex (see “Nomenclatural Notes”). Unfortunately, much of the literature does not specify geographic location or subspecies. Here, *C. aureus* is used for the global range from Africa into Eurasia ([Wozencraft 2005](#)). Therefore, we list the historical 13 subspecies of *C. aureus* ([Heller 1914](#); [Allen 1939](#); [Ellerman and Morrison-Scott 1951](#); [Coetsee 1977](#); [Wozencraft 2005](#)). The newly named subspecies, *qattarensis* [Saleh and Basuony, 2014](#), is tentatively treated as a synonym of *syriacus*. Subspecies designations for the *orientalis* [Linnaeus, 1758](#) and *studei* [Hilzheimer, 1906b](#) are not known.

- C. a. algirensis* Wagner, 1841. See above; synonyms are *algira* Kolenati, 1858; *barbarus* (Hamilton Smith, 1839); *grayi* Hilzheimer, 1906b; *tripolitanus* Wagner, 1841.
- C. a. anthus* F. Cuvier, 1820. See above; synonym is *senegalen-sis* (Hamilton Smith, 1839).
- C. a. aureus* Linnaeus, 1758. See above; synonyms are *balcani-cus* Brusina, 1892; *caucasica* Kolenati, 1858; *hadramauti-cus* Noack, 1896; *maroccanus* (Cabrera, 1921); *orientalis* (Linnaeus, 1758); *typicus* Kolenati, 1858; *vulgaris* Wagner, 1841.
- C. a. bea* (Heller, 1914). See above.
- C. a. cruesemanni* Matschie, 1900. See above.
- C. a. ecsedensis* (Kretzoi, 1947). See above; synonyms are *hun-garicus* Ehi, 1938; *minor* Mojsisovico, 1897 (preoccupied).
- C. a. indicus* Hodgson, 1833. See above; synonyms are *indica* Kolenati, 1858; *kola* Wroughton, 1916.
- C. a. lupaster* Hemprich and Ehrenberg, 1833. See above; syn-onym is *sacer* Hemprich and Ehrenberg, 1833.
- C. a. moreotica* I. Geoffroy Saint-Hilaire, 1835. See above; synonyms are *dalmatinus* Wagner, 1841; *graecus* Wagner, 1841.
- C. a. naria* Wroughton, 1916. See above; synonym is *lanka* Wroughton, 1916.
- C. a. riparius* Hemprich and Ehrenberg, 1833. See above; syn-onyms are *gallaënsis* Lorenz, 1906; *hagenbecki* Noack, 1897; *lamperti* Hilzheimer, 1906b; *mengesi* Noack, 1897; *somalicus* Lorenz, 1906.
- C. a. soudanicus* Thomas, 1903. See above, synonyms are *doederleini* Hilzheimer, 1906a; *nubianus* (Cabrera, 1921); *nubica* Kolenati, 1858; *thooides* Hilzheimer, 1906b; *varie-gatus* Cretzschmar, 1826 (preoccupied).
- C. a. syriacus* Hemprich and Ehrenberg, 1830. See above, syn-onym are *qattarensis* Saleh and Basuony, 2014; *syriaco* Kolenati, 1858.

NOMENCLATURAL NOTES. The above-named list may represent 3 taxa of canids: *Canis anthus* (African wolf), *C. aureus*, and *C. lupus* (gray wolf). Genetic analyses of *C. aureus* and *C. lupus* suggest that *C. aureus* from Africa and Eurasia are distinct clades and should be considered to be separate species, that is *C. anthus* (Africa) and *C. aureus* (Eurasia—Koepfli et al. 2015). As noted above, the literature does not distinguish these species and we include both as *C. aureus*.

In addition, large *C. aureus* in Egypt (*lupaster*) may be a cryptic subspecies of *C. anthus* (Rueness et al. 2011; Gaubert et al. 2012). *Canis aureus lupaster* (Hemprich and Ehrenberg 1833) is classified as a golden jackal; however Ferguson (1981) suggested that the taxon *C. aureus lupaster*, which is present in arid areas of Egypt and Libya (Osborn and Helmy 1980), may actually represent a small *C. lupus* rather than a large jackal. Mitochondrial DNA analyses suggest that *C. a. lupaster* represents an ancient wolf lineage that may have colonized Africa prior to the northern hemisphere radiation (Rueness et al. 2011). Mitochondrial DNA analyses assigned individuals from Algeria, Mali, and Senegal to *C. lupus lupaster* (Gaubert et al. 2012) and

identified *C. lupus lupaster* mtDNA haplotypes in *C. aureus* from Senegal questioning the genetic differentiation between the proposed *C. anthus* and *C. aureus*. We retain *C. aureus lupaster* as a form of the golden jackal not as a subspecies of *C. lupus* following Wozencraft (2005—Qumsiyeh 1996; Ferguson 2002).

The specific name is Latin for golden. Common names in-clude Asiatic jackal, common jackal, ibn awa or ibn awee (clas-sic Arab), wa wie (spoken Arab), cakalli (Albanian), čagalj (Croatian), šakal obecný (Czech), sjakal (Danish, Swedish), jakhal (Dutch), šakal (Estonian), sakaali (Finnish), Sjakalur (Faeroese), chacal doré (French), Goldschakal (German), τσάκαλ (Greek), aranyakál (Hungarian), sciacallo dorato (Italian), chagal, turg (Kurdish), zeltainais šakālis (Latvian), xakali (Maltese), gullsjakal (Nowegian), shoghal (Persian), szakal zło-cisty (Polish), chacal-dourado (Portuguese), șakal (Rumanian), šakal obyčajný (Slovakian, Slovenian), chacal (Spanish), çakal (Turkish—Hatt 1959; P. D. Moehlman, in litt.). Indigenous names include Amharic: tera kebero (Ethiopia); Fulani: sundu; Hausa: dila; Hindi: Giddhad; Kanada: nuree; Kiswahili: bweha wa mbugani, bweha dhahabu (Tanzania); Marathi (India): kolha; Nepali (Nepal), Bengali, Gujarati and Kutchi (India): shiyal; Singhelese: nariya; Songhai: nzongo; Tamil (India): peria naree; Wolof: Tili (P. D. Moehlman, in litt.).

DIAGNOSIS

Canis aureus occurs sympatrically with the silver-backed jackal (*Canis mesomelas*) and the side-striped jackal (*Canis adustus*) in parts of East Africa. All have skull lengths of 141–147 mm with minimal divergence in size (van Valkenburgh and Wayne 1994). *C. mesomelas* is distinguished by reddish flanks and limbs, and the skull may have a less developed parietal crest (Clutton-Brock et al. 1976), the angle of its ears which are more upright, and, usually, the prominent dark “silver” saddle com-posed of black and white hairs (Walton and Joly 2003). *C. aureus* may have a dark saddle seasonally, which is not prominent. *C. adustus* characteristically has shorter ears set more to the side of the skull, a pale side-stripe, and a white-tipped tail (Moehlman and Jhala 2013). *C. lupus* is much larger with a heavier body and variable pelage coloring. The skull is much larger with a well-developed interparietal crest, strong jaw, and large canine and carnassial teeth (Clutton-Brock et al. 1976).

GENERAL CHARACTERS

Canis aureus (Fig. 1) is a medium-sized canid, considered the most typical representative of the genus *Canis* (Clutton-Brock et al. 1976). Basic coat color is golden but varies from pale creamy yellow to a dark tawny hue on a seasonal basis. The pelage on the back is often a mixture of black, brown, and white hairs, such that they can appear to have a dark saddle sim-ilar to the *C. mesomelas* (Walton and Joly 2003; Moehlman and Jhala 2013). The long contour hairs have an extensive black tip

and a wider penultimate white band (Pocock 1938). In winter, a copious under wool is present (Pocock 1938). *C. aureus* inhabiting rocky, mountainous terrain may have a grayer coat shade (Sheldon 1992). The belly and underparts are a lighter pale-tinger to cream. The tail is bushy with a tan to black tip. Females have 7–8 mammae (Poché et al. 1987), comprising 4 pairs (Sheldon 1992). Eye color varies from pale yellow to amber (Lewis et al. 1968).

Means and parenthetical ranges of external measurements (mm) from 6 males and 3 females, respectively, in Gujarat, India (Y. V. Jhala, in litt.) were: length of head and body, 793 (760–840), 760 (740–800); length of tail, 220 (200–240), 205 (200–210); length of ear, 76 (68–90), 80 (75–85); body mass (kg) of 6 male and 4 females was 8.8 (7.6–9.8) and 7.3 (6.5–7.8), respectively. In Tanzania, the external measurements (mm or kg) from 2 males in the Serengeti (Moehlman and Jhala 2013) were: length of head and body, 740, 785; length of tail, 270, 280; length of ear, 110, 110; mass, 6.3, 7.7. In Egypt, means and parenthetical ranges (mm) for 9 adult *C. a. lupaster* of unknown sex were: length of head and body, 872 (822–893); length of tail, 312 (290–347); length of hind foot, 200 (190–212); length of ear, 112 (104–121); and body mass (kg) of 4 adults was 13 (10–15—Osborn and Helmy 1980). In Lebanon, external measurements (mm) of 3 adult *C. a. syriacus* were: total length, 988.3 (950–1015); length of tail, 275 (265–290); length of head and body, 713.3 (680–750); length of hind foot, 156 (140–170); length of ear from notch, 74.7 (70–80—Lewis et al. 1968). In Bangladesh, mean measurements (mm, *SD*, *n*) of females and males, respectively, were: total length, 1009.3 (33.67, 6), 1058.6 (20.30, 7); length of tail, 220.1 (17.16, 7), 245.0 (21.41, 6); length of hind paw, 58.0 (4.93, 7), 56.3 (4.83, 7); length of hind foot, 146.2 (24.66, 5), 162.6 (8.52, 7); length of ear, 69.4 (4.80, 5), 74.1 (5.49, 7); mass (kg), 8.5 (0.73, 7), 10.3 (0.95, 7—Poché et al. 1987). Mean adult mass (kg) of 5 females and 3 males, respectively, was 10.0 and 12.0 (Palmqvist et al. 2002). On average, females are 12% lighter than males: 5.8 kg versus 6.6 kg (Moehlman and Hofer 1997).

Mean cranial measurements (mm, *SD*; Fig. 2) on 10 mandibles and upper jaws of adult *C. aureus* in Iran were: length of skull, 169, 31.9; length of cranium, 101, 23.7; nasal length, 67, 16.6; cranial width, 72, 4.4; length of mandible, 112, 34.6; lateral alveolar root to mental foramen, 25, 1.7; mental foramen to caudal mandibular border, 87, 3.8; mandibular foramen to base of mandible, 12, 0.4; caudal border of mandible to below mandibular foramen, 11, 0.8; maximum mandibular height, 60, 5.8 (Mofared 2013). Mean measurements (mm, range) for 3 Lebanese adults were: greatest length of skull, 148.6 (135.2–157.4); condylobasal length, 142.9 (131.9–149.1); zygomatic breadth, 83.9 (73.3–90.2); breadth of braincase, 48.8 (46.0–50.3); interorbital constriction, 29.4 (25.5–32.7); maxillary tooththrow, 62.9 (60.4–65.0); mandibular tooththrow, 72.2 (58.2–74.9); mandibular length, 114.1 (104.1–121.0—Lewis et al. 1968). Mean cranial measurements (mm; with range and *n*) of Egyptian adult *C. a. lupaster* were: condyloincisive length, 185.2 (173.5–196.0, 13); zygomatic width, 101.4 (93.5–111.3, 14); rostral width, 33.8



Fig. 2.—Dorsal, ventral, and lateral views of skull and lateral view of mandible of an adult male *Canis aureus aureus* (Field Museum of Natural History, Chicago, Illinois, [FMNH] #103909) from Maimana, Fariab, Afghanistan, 2,900 feet. Greatest length of skull is 177 mm. Photograph by Bruce Patterson used with permission.

(31.2–37.8, 14); postorbital width, 34.8 (31.8–38.9, 13); basioccipital width at tempoparietal suture, 54.4 (51.8–59.1, 13); nasal length, 72.0 (65.9–84.8, 14); width across upper molars, 58.2 (54.2–63.0, 14); distance from anterior most surface of canine to posterior of 2nd molar, 80.3 (75.6–86.8, 14); skull height, 66.8 (62.0–74.0, 9—Osborn and Helmy 1980). In Bulgaria, mean cranial and tooth measurements (mm) for 92 males and 56 females, respectively, were: total skull length, 166, 161; neurocranium length, 89, 86.5; greatest length of nasals, 59.7, 57.9; greatest neurocranium breadth, 52.5, 52.6; total length of mandible, 121.9, 118.6; height of vertebral ramus, 48.6, 46.7; length of P1 to P4, 33.9, 33.4; length of M1 to M3, 31.8, 31.0 (Stoyanov 2012). Mean mandible length and depth at 2nd molar (mm)

for 5 females and 3 males, respectively, was f: 102.8, 14.3; m: 108.9, 15.3 (Palmqvist 2002). Condylobasal length (mm) from 10 males was 154, whereas it was 140 from 7+ female skulls (Pocock 1938).

DISTRIBUTION

Canis aureus is widespread in North and East Africa, Europe, the Middle East, Central Asia, and Southeast Asia (Fig. 3). They range from sea level in Eritrea to 3,500 m in the Bale Mountains of Ethiopia and 2,000 m in India (Prater 1971; Admasu et al. 2004).

In Greece and Dalmatia, *C. aureus* is documented from the Holocene (Sommer and Benecke 2005; Malez 1984 in Rutkowski et al. 2015) and the ancient Mediterranean populations have persisted and merged with jackals coming from Asia (Fabbri et al. 2014; Rutkowski et al. 2015). The expanding population of *C. aureus* in Europe is due to natural long-distance dispersal (Arnold et al. 2012; Rutkowski et al. 2015). *C. aureus* expanded from southeastern Europe to Hungary in the late 1800s (Tóth et al. 2009). In the late 20th century, the expansion was rapid and is still ongoing (Kryštufek et al. 1997; Arnold et al. 2012).

FOSSIL RECORD

The fossil record of *Canis aureus* is limited with many fossils that are controversial as to their specific attribution. A right lower carnassial from a Paleolithic Lebanon cave may be *C. aureus*

(Hooijer 1961). Fossil remains attributed to *C. aureus* dating from the Upper Pleistocene were found in North Africa (Savage 1977). Palestinian fossils attributed to this species have been reattributed to *C. lupaster* (Kurten 1974) and hence might be a separate species (Spassov 1989). Reliable fossils of *C. aureus* have not been found in the Pleistocene of Europe (Spassov 1989). *C. aureus* appears to have arrived in sub-Saharan Africa after *C. mesomelas* and *C. adustus*, which have a fossil record in eastern and southern Africa from 500,000 to 2.5 million years (Ewer 1956; Savage 1977).

FORM AND FUNCTION

The dental formula is i 3/3, c 1/1, p 4/4, m 2/3, total 42 (Fig. 2). Of 31 *Canis aureus* skulls, 13% had broken teeth (van Valkenburgh 2009). Baculum length is 62.7 mm (Dixson 1995), but 2 bacula of *C. a. lupaster* were 70 and 73 mm (Osborn and Helmy 1980).

Male and female skulls can be differentiated but cranial measures overlap (Stoyanov 2012). Differentiation was not dependent on age or collection site. Only juveniles below the age of 6 months could be differentiated from adults. *C. aureus* from Bulgaria did not differ from European or Asian skulls and encompassed both African and Dalmatian jackals, but skulls from *Canis aureus lupaster* were larger, longer, and broader (Stoyanov 2012). Croatian skulls were of a shorter length and broader (Stoyanov 2012). In Bulgarian skulls, age could be assessed in 223 of 228 from craniometrics differences (Stoyanov 2012). Across 7 populations, sexual dimorphism in skull length




Fig. 3.—Geographic distribution of *Canis aureus* includes subspecies that may belong to *C. anthus* or *C. lupus*. The map shows the likely ranges of the following subspecies, 1, *C. a. algirensis*; 2, *C. a. anthus*; 3, *C. a. aureus*; 4, *C. a. bea*; 5, *C. a. cruesmanni*; 6, *C. a. ecsedensis*; 7, *C. a. indicus*; 8, *C. a. lupaster* (dispersed stars on map); 9, *C. a. moreotica* (isolated records presented as blue dots on map); 10, *C. a. naria*; 11, *C. a. riparius*; 12, *C. a. soudanicus*; 13, *C. a. syriacus*; 14, *C. a.* subspecies undetermined.

and canine diameter was significant for East African and northern Indian *C. aureus* but not north African, north central African, Middle Eastern, Pakistani, or southern Indian animals (Van Valkenburgh and Wayne 1994). In Bangladesh, mean organ weights (g, *SD*, *n*) for adult females and males, respectively, were: heart, 71.0 (7.35, 3), 79.2 (14.02, 4); liver, 297.7 (14.52, 3), 365.5 (72.06, 4); paired kidneys, 45.7 (8.73, 3), 64.0 (4.37, 4); spleen, 75.3 (21.1, 3), 43.0 (10.61, 3); lungs, 64.7 (11.47, 3), 85.7 (18.87, 3—Poché et al. 1987).

Adult *C. aureus* can be immobilized for handling up to 55 min with 130 mg ketamine hydrochloride and 15 mg promazine hydrochloride (Fuller et al. 1989). For 3 female and 4 male *C. aureus*, doses of 15–80 mg/kg of ketamine hydrochloride generated recovery times of 32–300 min with no apparent side effects (Poché et al. 1987). Time to immobilization was 2–3 min (Poché et al. 1987). Trapped jackals were anesthetized with an intramuscular injection of a mixture of medetomidine 0.09 mg/kg and ketamine 2.8 mg/kg and time to immobilization was an average of 5.0 min (Admasu et al. 2004).

ONTOGENY AND REPRODUCTION

Ontogeny.—The estimated weight from a captive born litter of 2 female and 2 male *Canis aureus* neonates was 189 g (Moehlman and Hofer 1997). A male pup weight at day 8 was 249 g, day 13: 500 g, and at day 15: 590 g. A female pup at day 5 weighed 295 g, and at day 21 weighed 670 g (Seitz 1959). Pups are blind at birth with eye opening at 8–11 days and tooth eruption at 11 days (Seitz 1959; Wandry 1975). In captivity, canine teeth erupted at 28 days (Seitz 1959) and pups emerged from the den box at approximately 4 weeks (Wandry 1975). At 13 months of age a female had her 1st litter and hence was reproductively mature at 11 months of age (Seitz 1959).

In the Serengeti, Tanzania, the pups are born and remain in underground dens until they are 3 weeks old during which time their mother spends most of her time with them (Moehlman 1983). Pups emerge at approximately 3 weeks of age and in addition to nursing, eat regurgitated food (Moehlman 1986). Some families have helpers which are offspring from the previous year's litter. During pregnancy, the female is provisioned by her mate and while she is lactating she is provisioned by her mate and the "helpers" which are her older offspring (Moehlman and Hofer 1997). One of us (PDM) has observed that helpers also provision the pups which are their full siblings. They are weaned at 8–10 weeks of age and stay within several hundred meters of the den until they are 10 weeks old. At 14 weeks of age they are well coordinated and start to forage with the adults. They will continue to receive some food from their parents until they are 8 months old. Even when they are helpers (11–18 months) they may infrequently receive food from their parents (Moehlman 1986).

Adults can be distinguished from juveniles by frontoparietal ridges that are fused posteriorly to the frontoparietal suture

creating a high sagittal ridge (Osborn and Helmy 1980). Tooth wear can be used to age *C. aureus* (Raichev 2011).

Reproduction.—In the Serengeti mating typically occurs from October to December (Moehlman 1983, 1986) and birth from December to March (Moehlman 1983). In Egypt, litters have been observed in March, April, and May (Flower 1932). In India reproductive behavior begins in February–March (Jhala and Moehlman 2004) and in Israel between October and March (Golani and Keller 1975; Ginsberg and Macdonald 1990). In Bangladesh, pups are present year-round (Poché et al. 1987). Gestation is about 63 days (Sheldon 1992). Litter size ranges from 1 to 9, average 5.7 (Moehlman and Hofer 1997). In Tanzania, an average of 2 pups emerged from the natal den at 3 weeks (Wyman 1967). Lactation lasts for 8–10 weeks (Moehlman 1986). Six females captured in Kenya in June to August were not lactating (Fuller et al. 1989).

ECOLOGY

Population characteristics.—In Tanzania, densities up to 2 adults per km² may occur (Moehlman 1983, 1986, 1989). In northern Dalmatia, Croatia, territory density was 0.61–1.15 groups per 10 km² (Krofel 2008), whereas in southern Dalmatia density was 6–7 groups per 10 km² (Krofel 2007). In Kenya, juveniles comprised 47% of captured *Canis aureus* (Fuller et al. 1989). In the Serengeti, maximum longevity is about 14 years (Moehlman and Hofer 1997). In captivity, individuals have lived up to 18 years 10 months (Weigl 2005). Sex ratio at approximately 3 weeks when pups emerge from the den is 1:0.8 (*n* = 33), surviving at age 14 weeks (1:0.8, *n* = 44), helping as juveniles at approximately 52 weeks (1:1, *n* = 16), and reproductive adults (1:1, *n* = 46—Moehlman 1986). The sex ratio is equal for all age groups and sex ratio is equal for helpers.

Space use.—Tolerance of dry habitats and an omnivorous diet allow *Canis aureus* access to a wide variety of habitats (Jhala and Moehlman 2008). They normally live in open grassland habitat but also occur in deserts, woodlands, mangroves, and agricultural and rural habitats in India and Bangladesh (Jhala and Moehlman 2008).

In the Serengeti, *C. aureus* maintains year-round exclusive territories of 0.5–7.0 km² (Moehlman 1983) and 2–5 km² in Ngorongoro (van Lawick and van Lawick-Goodall 1970), but will make excursions beyond these territorial boundaries to gain access to fresh carcasses. In Algeria, seasonal territories as may be as small as 0.39 km² (Khidas 1990). Territories are within larger home ranges. Home range size of radiocollared individuals in some locales can vary according to habitat and age. An adult pair in *Acacia* woodland in Kenya had a home range of 2.4 km² and 2 juvenile females had home ranges of 5.6 and 21.7 km² (Fuller et al. 1989), although range size over a 16-month period in the Bale Mountains of Ethiopia varied from 7.9 to 48.2 km² for adults and from 24.2 to 64.8 km² for subadults (Admasu et al. 2004). Subadults may have larger home ranges as they

explore and search for a mate and a territory. Home ranges of individuals within a social group tend to overlap (van Lawick and van Lawick-Goodall 1970; Khidas 1990; Admasu et al. 2004). In Bangladesh, breeding pairs defend an average of 37.3 ha of cover but not the peripheral foraging areas (Jaeger et al. 1996, 2007) and a male and female had home ranges of 1.1 and 0.6 km², respectively (Poché et al. 1987). In Israel, home and core ranges (km²) of *C. aureus* near villages were about one-third of those in more natural areas (mean, *SD*): home, 6.6 (4.5) versus 21.2 (9.3); core, 1.2 (0.9) versus 3.5 (1.6—Rotem et al. 2011).

In the Serengeti, territories are maintained for up to 8 years albeit with changes in boundaries and size through time. Both members of the pair mark and defend the territory (Moehlman 1986).

Diet.—In Bangladesh, rodents are the most common food (Jaeger et al. 2007). Similarly, in Bulgaria, rodents comprise 59.3% of the diet with brown hares (*Lepus europaeus*, 20.1%) and plants (primarily fruit, 19.7%) as the next most common components (Markov and Lanszki 2012). In Rajasthan, India, rodents make up 45% of the diet (Mukherjee et al. 2004). In Algeria, mammals comprise about a one-third of the diet with 17–24% poultry and 12–27% insects depending on location (Amroun et al. 2006). In Israel, domestic carrion may form a significant component of the diet along with fruit, birds, small mammals, and invertebrates (Borkowski et al. 2011). In Bangladesh, 12 stomachs contained primarily carrion, poultry, and vegetation but no rodents (Poché et al. 1987).

Single *C. aureus* typically hunt smaller prey like rodents and birds, using their hearing to locate rodents in the grass and then pouncing on them by leaping through the air, or digging out rodents from their burrows. However, single animals will also hunt Thomson's gazelle (*Eudorcas tomsonii*) fawns. They have been observed to hunt young, old, and infirm ungulates that are sometimes 4–5 times their body weight (van Lawick and van Lawick-Goodall 1970; Eisenberg and Lockhart 1972). *C. aureus* in Bale, Ethiopia, may be more solitary than elsewhere in the range, with individuals having been observed foraging alone on 87% of occasions (Admasu et al. 2004). This is likely because food resources were widely dispersed and rarely concentrated enough for jackals to forage in groups (Admasu et al. 2004).

In the Serengeti, mated pairs will hunt cooperatively and regularly kill Thomson Gazelle fawns and occasionally adults; pairs have a higher kill rate than individuals (Wyman 1967; Kruuk 1972; Rosevear 1974). Cooperative hunting permits them to harvest much larger prey in areas where they are available Temu et al. 2017. In the Ngorongoro Crater, *C. aureus* pairs cooperatively killed Thomson's gazelle fawns and occasionally killed adult Thomson's gazelle. Individuals successfully hunted Abdim's storks (*Ciconia abdimii*) and during Wildebeest calving season consumed the abundantly available placentas. There was one observation of *C. aureus* cooperatively killing and eating a bat-eared fox (*Otocyon megalotis*) (Temu et al. 2017). Golden jackals consumed significantly more grams of food per hour of foraging when compared with sympatric Silver-backed jackals

(Temu et al. 2016). In some areas, particularly where food resources are clumped, aggregations of *C. aureus* may occur, such as 14 jackals on a carcass in Ngorongoro (van Lawick and van Lawick-Goodall 1970) and aggregations of 5–18 jackals scavenging on carcasses of large ungulates in India (Y. V. Jhala, in litt.). *C. aureus* will cache excess food (van Lawick and van Lawick-Goodall 1970). In India, they will kill poultry as well as young and weak sheep and goats (Jhala and Moehlman 2004), and will venture into human habitation at night to feed on garbage (Poché et al. 1987). In the Golan Heights, 1.5–1.9% of domestic calves are killed by predators, primarily *C. aureus* (Yom-Tov et al. 1995).

In the Serengeti, *C. aureus* scavenges the kills of spotted hyenas (*Crocuta crocuta*) and lions (*Panthera leo*) and leopards (*Panthera pardus*—van Lawick and van Lawick-Goodall 1970; Moehlman 1986). In the Serengeti during the dry season, *C. aureus* does not have access to free standing water and can tolerate dry habitats (Moehlman and Jhala 2013).

Diseases and parasites.—In the Serengeti, blood of 2 *Canis aureus* males was negative for leptospirosis, canine distemper, canine brucellosis, rabies, rinderpest, African horse sickness, and rift valley fever. Both individuals had titers for canine adenovirus and canine coronavirus. The older male was positive for canine parvovirus (Karesh 1990 in litt.). In 1994 during an outbreak of distemper, one *C. aureus* tested positive for canine distemper virus (Roelke-Parker et al. 1996). In 2011, samples collected from *C. aureus* pups were positive for canine distemper virus, but only 1 death was observed (Olarte-Castillo et al. 2015). In Kenya (1987–1988) testing for canine parvovirus (CPV-2), canine distemper virus, *Ehrlichia canis*, and rabies indicated that 9 of 16 individuals were positive for CPV-2; all were negative for canine distemper virus, *E. canis*, and rabies (Alexander et al. 1994).

In Israel, a serological survey indicated the presence of canine distemper virus, *E. canis*, and *Leshmania infantum*. Blood smears in one individual revealed *Hepatozoon canis* (Shamir et al. 2001). *Leshmania tropica* is also known from Israeli jackals (Talmi-Frank et al. 2010). Six of 8 *C. aureus* were positive for *Toxoplasmosis gondii* in the United Arab Emirates (Dubey et al. 2010).

Canis aureus in Serbia had an infection rate of 46 out of 447 jackals (10.3%) of at least one helminth species and in total 12 species were identified: 2 trematodes, *Alaria alata* and *Pseudamphistomum truncatum*, 3 nematodes, *Toxocara canis*, *Ancylostoma caninum*, and *Gongylonema*, and 7 cestodes, *Taenia pisiformis*, *Taenia hydatigena*, *Multiceps multiceps*, *Multiceps serialis*, *Mesocostoides lineatus*, *Macrocanthorhynchus hirudinaceus*, and *Onicola canis* (Ćirović et al. 2015). In Israel, fecal samples had *A. caninum* and *Dipylidium caninum* (Shamir et al. 2001). In Tanzania, fecal samples from *C. aureus* had *Coccidia*, *Ancylostoma*, *Psoroptes*, and *T. canis* (W. B. Karesh, in litt.). In Rumania, the roundworm, *Trichinella britovi* occurs in *C. aureus* (Blaga et al. 2008). In Uzbekistan and Turkmenistan,

the Ministries of Health (Berdyeva 1998; L. M. Isaev Institute of Medical Parasitology 1998) reported that *C. aureus* may be infected with the guinea worm, *Dracunculiasis medinensis*; thus, *C. aureus* potentially can infect water sources. In Italy, a subadult male had 2 ectoparasitic species, *Dermacentor reticulatus*, *Ixodes ricinus*, and 1 endoparasitic species, *Metagonimus yokogawai* (Lapini et al. 2009). One *C. aureus* from Tunisia had 72 tapeworms, *Echinococcus granulosus* while 4 others had fecal evidence of the species (Lahmar et al. 2009). The protozoan, *Babesia gibsoni*, may use *C. aureus* as a host (Patton 1910).

Interspecific interactions.—In East Africa, *Canis aureus* is sympatric with *C. mesomelas* and *C. adustus*. Skull characters indicate that *C. aureus* are more carnivorous than *C. adustus* and less carnivorous than *C. mesomelas* (van Valkenburgh and Wayne 1994). In Hungary, dietary overlap between sympatric *C. aureus* and red foxes (*Vulpes vulpes*) is high at 73% as both primarily eat small mammals < 50 g (Lanszki et al. 2006).

In East Africa, spotted hyenas will kill and feed on *C. aureus* (Kruuk 1972; Kingdon 1977). In the Serengeti, *C. aureus* give a “warning yowl” when spotted hyenas approach their dens. The adult jackals then chase the hyenas and bite them on the rump or genitals. Honey badgers (*Mellivora capensis*) have been observed near *C. aureus* dens, but have always been chased away by the parents and helpers. In India, striped hyenas (*Hyaena hyaena*) may prey on *C. aureus* (Y. V. Jhala, in litt.) and pythons (*Python molurus*) can be major predators (Singh 1983). In India, when *C. aureus* approaches human habitation it may be killed by feral domestic dogs and death due to road kills increased during its breeding season (Y. V. Jhala, in litt.). *C. aureus* may be killed as a side effect of predator control programs when it scavenges poisoned carcasses (Moehlman and Jhala 2013).

Miscellaneous.—Capture methods include rubber-padded, steel, foot-hold traps, cage-type live traps, large box traps, and Victor steel traps with off-set jaws (Poché et al. 1987; Fuller et al. 1989; Admasu et al. 2004; Rotem et al. 2011). Differences in white markings on throat and chest may allow individual identification (Macdonald 1979). Unique facial scars and naturally notched ears can be used to identify individuals (Moehlman 1983, 1986).

HUSBANDRY

As of March 2000, approximately 193 *Canis aureus* were in Indian zoos (Jhala and Moehlman 2004). *C. aureus* have been raised in captivity for behavioral studies (Seitz 1959; Wandry 1975). Individuals were housed in a kennel of 225 m² which included 5 separate and interconnected areas/cages. Each cage had a large box with a tunnel-like entrance (Wandry 1975).

BEHAVIOR

Grouping behavior.—*Canis aureus* are typically monogamous but can potentially be socially flexible depending on demography and food resources (Macdonald 1979; Moehlman

1983, 1986, 1989; Fuller et al. 1989; Moehlman and Hofer 1997; Admasu et al. 2004). The basic social unit is the breeding pair and some offspring from the previous year’s litter may remain with the parents and help to raise the current litter of pups (Moehlman 1983, 1986, 1989). In Tanzania, *C. aureus* usually form long-term pair bonds which can last for up to 6 years. The bonded pair is territorial and both male and female mark within and on the boundaries of their territory and defend their territories against intruders. Same sex aggression is the norm with the territorial male attacking intruding males and females attacking intruding females. The pair hunt together, share food, and cooperatively rear their young (Moehlman 1983, 1986, 1989). Moehlman and Hofer (1997) give average group size as 2.5 (Serengeti, Tanzania), similar to average group size (3; *n* = 7) in Velavadar N. P., India (Y. V. Jhala, in litt.).

Reproductive behavior.—*Canis aureus* has a strong pair bond which normally lasts for its lifetime (Moehlman 1983, 1986, 1989). Mating involves a copulatory tie that lasts for several minutes (Golani and Mendelsohn 1971; Golani and Keller 1975). Pups are born in an underground den and normally emerge at about 3 weeks of age. Once pups emerge from the den they are fed solid food by adults regurgitating to them in addition to nursing. The origin of the dens can vary. Some authors (e.g., Rosevear 1974) report the use of existing aardvark *Oryzomys* or warthog *Phacochoerus* burrows, but in the Serengeti this is unlikely as *C. aureus* den openings are too small for either of these species (Jhala and Moehlman 2004). Earthen dens may have 1–3 openings and are typically about 2–3 m long and 0.5–1 m deep (Jhala and Moehlman 2004). Pups may be moved to different dens 2–4 times during their first 14 weeks of development (Moehlman 1983, 1986, 1989).

In Tanzania, *C. aureus* typically produces litters once a year during the wet season (December to February) when the large migratory herds of wildebeest (*Connochaetes taurinus*), zebra (*Equus burchelli*), and Thomson’s gazelles are present on the short-grass plains (Maddock 1979). Thus, when *C. aureus* has pups, food items ranging in size from dung beetles to Thomson’s gazelle fawns are abundant. *C. aureus* also scavenges at carcasses of wildebeest and zebra that are killed by larger predators, mainly spotted hyenas (van Lawick and van Lawick-Goodall 1970; Moehlman 1983, 1986).

In the Serengeti since mated pairs are monogamous with long-term bonds, sequential litters are full siblings as confirmed by genetic analysis of microsatellite DNA (P. D. Moehlman, in litt.). The male provisions the female while she is pregnant and both the male and the helpers (the previous years pups) provide the female with food while she is lactating (Moehlman 1983, 1986, 1989; Moehlman and Hofer 1997). The presence of helpers correlates significantly with a higher pup survival (Macdonald and Moehlman 1983; Moehlman 1986, 1989).

Communication.—Territories are maintained indirectly by scent marking and vocalizations. Both members of a pair use a raised-leg urination posture to scent mark grass tufts, bushes, and trees at nose height. Urination with a raised rear leg instead of squatting is considered to be a ‘marking posture’

and indicates that the individual is a territory holder and on its territory. Marking, for example “raised-leg” urinations indicate that an individual is a territory holder and on its territory. When a territorial male or female leaves their area they do not “mark.” Both the boundary and the area within a territory are “marked.” When foraging with their mate, territorial individuals mark at twice the frequency as when they are foraging alone. When travelling together they mark the same spot in tandem (Moehlman 1983, 1986). The presence of scent from both members of the pair may advertise to potential intruders that they are both actively in residence (Moehlman 1983). Such scent marks are considered to play an important role in territorial defense (Rosevear 1974).

Affiliative behaviors like greeting ceremonies, grooming, and group vocalizations are common in *C. aureus* social interactions (Van Lawick and van Lawick-Goodall 1970; Golani and Keller 1975). Vocalization consists of a complex howl repertoire beginning with 2–3 simple low-pitch howls and culminating in a high-pitched staccato of calls. In the Serengeti, individuals give reciprocal howls to locate their mates and family members. Families also occasionally group howl in response to neighboring family howls (Moehlman 1983). Similar functions accrue to howling in Bangladesh, for territory maintenance and advertisement of dominance (Jaeger et al. 1996). *C. aureus* has a “rumble growl” and “predator bark” when potential predators like spotted hyenas approach a den with pups. Pups respond to these vocalizations by running into the den (Moehlman 1983).

Miscellaneous behavior.—In Kenya, *Canis aureus* was active during the day, as well as at dawn and dusk but not in the middle of the night (Fuller et al. 1989). In Bangladesh, it preferred the cover of sugarcane during the day (Jaeger et al. 2007).

GENETICS

The diploid chromosome number (2N) of *Canis aureus* is 78 (Wurster and Benirschke 1968). A total of 12–24 microsatellite loci from Israeli *C. aureus* generated expected levels of heterozygosity (Cohen et al. 2013). In Serbian *C. aureus*, mitochondrial DNA has no variation and nuclear variability is low (Zachos et al. 2009). The mitochondrial DNA control regions are variable across *C. aureus* from Bulgaria, Serbia, Croatia (Dalmatia and Slavonia), and Italy and most genetically distinct in the population from Dalmatia (Fabbri et al. 2014).

A melanistic individual occurred in Turkey (Ambarli and Bilgin 2012). Other melanistic and piebald forms are known (Muller-Using 1975; Jerdon 1984). Hybridization with domestic dogs may occur in Croatia (Galov et al. 2015) but not in India (Yumnam et al. 2015).

Genetic data suggest that *C. a. lupaster* represents an ancient wolf lineage that most likely colonized Africa prior to the northern hemisphere radiation (Rueness et al. 2011). A combined analysis of nuclear and mitochondrial DNA (Bardleben et al. 2005) indicates that *C. aureus* is in a wolf-like clade that includes *Canis adustus*, *C. mesomelas*, *C. familiaris*, *C. lupus*,

C. latrans, *Cuon alpinus*, and *Lycaon pictus*. A phylogeny of living canids based on 22 nuclear and 3 mitochondrial loci produced a similar wolf-like clade (Perini et al. 2010).

CONSERVATION

Canis aureus is fairly common throughout its range and is listed on the International Union for Conservation of Nature and Natural Resources Red List as “Least Concern” with at least 80,000 individuals on the Indian continent and an unknown number in Africa (Jhala and Moehlman 2008). This species is not listed in CITES. In the European Union, *C. aureus* is listed as an Annex V species and should be legally protected (Rutkowski et al. 2015; Trouwborst et al. 2015). As the expansion of *C. aureus* is a natural long-range dispersal, it should not be considered an alien invasive species (Rutkowski et al. 2015).

In the Serengeti–Ngorongoro ecosystem the population may be in decline (Durant et al. 2011) but it is still at a relatively high density (0.5 individuals per km²). In Greece, the population has been declining from at least 1975 (Giannatos et al. 2005). In areas where *C. aureus* is perceived as a threat to livestock, it is often killed in indiscriminate predator control programs (Moehlman and Jhala 2013).

ACKNOWLEDGMENTS

We are grateful for the help of several people with diverse aspects of the manuscript. B. Glumac translated Brusina’s archaic Serbo-Croatian. J. Benkley helped with various translations from German. B. Patterson took the skull photos. I. Fielding formatted the distribution map. PDM is grateful for permission from the Tanzania Commission for Science and Technology (COSTECH) and the Tanzania Wildlife Research Institute (TAWIRI) to conduct long-term research on *C. aureus* in the Serengeti–Ngorongoro ecosystem. TAWIRI, the Tanzania National Parks Authority (TANAPA), and the Ngorongoro Conservation Area Authority (NCAA) have provided important support for this research. VH was supported by the Blakeslee Grant for Genetics Research, Smith College.

LITERATURE CITED

- ADMASU, E., S. J. THIRGOOD, A. BEKELE, AND K. M. LAURENSEN. 2004. Spatial ecology of golden jackal in farmland in the Ethiopian highlands. *African Journal of Ecology* 42:144–152.
- ALEXANDER, K. A., P. W. KAT, R. K. WAYNE, AND T. K. FULLER. 1994. Serologic survey of selected canine pathogens among free-ranging jackals in Kenya. *Journal of Wildlife Diseases* 30:486–491.
- ALLEN, G. M. 1939. A checklist of African mammals. *Bulletin of the Museum of Comparative Zoology at Harvard College* 83:1–763.
- AMBARLI, H., AND C. BILGIN. 2012. First record of a melanistic golden jackal (*Canis aureus*, Canidae) from Turkey. *Mammalia* 77:219–222.
- AMROUN, N., P. GIRAUDOUX, AND P. DELLATRE. 2006. A comparative study of the diets of two sympatric carnivores – the golden jackal (*Canis aureus*) and the common genet (*Genetta genetia*) – in Kabylia, Algeria. *Mammalia* 70:247–254.

- ARNOLD, J., A. HUMER, M. HELTAI, D. MURARIU, N. SPASSOV, AND K. HACKLANDER. 2012. Current status and distribution of golden jackals *Canis aureus* in Europe. *Mammal Review* 42:1–11.
- BARDELEBEN, C., R. L. MOORE, AND R. K. WAYNE. 2005. A molecular phylogeny of the Canidae based on six nuclear loci. *Molecular Phylogenetics and Evolution* 37:815–831.
- BERDYEVA, A. T. 1998. Dracunculiasis eradication in Turkmenistan. Country Report by the Ministry of Health, Turkmenistan.
- BLAGA, R., C. GHERMAN, D. SEUCOM, V. COZMA, AND P. BOIREAU. 2008. First identification of *Trichinella* sp. in golden jackal (*Canis aureus*) in Romania. *Journal of Wildlife Diseases* 44:457–459.
- BORKOWSKI, J., A. ZALEWSKI, AND R. MANOR. 2011. Diet composition of golden jackals in Israel. *Annales Zoologici Fennici* 48:106–118.
- BRUSINA, S. 1892. Čagalj balkanski (*Canis aureus balcanicus* Brusn. form.) iz Slavonije. *Glasnik Hrvatskoga Naravoslovnoga Društva Zagreb* 7:316–317.
- CABRERA, A. 1921. Algunos carnívoros africanos nuevos. *Boletín de la Real Sociedad Española de Historia Natural, Madrid* 21:261–264.
- ĆIROVIĆ, D., I. PAVLOVIĆ, A. PENEZIĆ, Z. KULIŠIĆ, AND S. SELAKOVIĆ. 2015. Levels of infection of intestinal helminth species in the golden jackal *Canis aureus* from Serbia. *Journal of Helminthology* 89:28–33.
- CLUTTON-BROCK, J., G. B. CORBET, AND M. HILLS. 1976. A review of the family Canidae, with a classification by numerical methods. *Bulletin of the British Museum (Natural History)* Zoology 29:1–199.
- COETZEE, C. G. 1977. Order Carnivora. Part 8. Pp. 1–42 in *The mammals of Africa: an identification manual* (J. Meester and H. W. Setzer, eds.). Smithsonian Institution Press, Washington, D.C.
- COHEN, T. M., R. KING, A. DOLEV, A. BOLDO, A. LICHTER-PELED, AND G. K. BARGAL. 2013. Genetic characterization of populations of the golden jackal and the red fox in Israel. *Conservation Genetics* 14:55–63.
- CRETZSCHMAR, P. J. 1826. Atlas zu der Reise im nördlichen Afrika von Eduard Rüppell. Erste Abtheilung Zoologie. Gedruckt und in Commission bei Heinr. Ludw. Brönnner, Frankfurt, Germany.
- CUVIER, F. 1820. Le chacal du Sénégal, femelle. Pp. 1–3 in *Histoire naturelle des mammifères, avec des figures dessinées d'après les animaux vivants, tome deuxième* (E. Geoffroy Saint-Hilaire and F. Cuvier, eds.). A. Belin, Paris, France. 1824 on title page but material on the jackal is June 1820.
- DIXON, A. F. 1995. Baculum length and copulatory behaviour in carnivores and pinnipeds (Grand Order Ferae). *Journal of Zoology (London)* 235:67–76.
- DUBEY, J. P., ET AL. 2010. Toxoplasmosis in sand cats (*Felis margarita*) and other animals in the Breeding Centre for Endangered Arabian Wildlife in the United Arab Emirates and Al Wabra Wildlife Preservation, the State of Qatar. *Veterinary Parasitology* 172:195–203.
- DURANT, S., M. E. CRAFT, R. HILBORN, S. BASHIR, J. HANDO, AND L. THOMAS. 2011. Long-term trends in carnivore abundance using distance sampling in Serengeti National Park, Tanzania. *Journal of Applied Ecology* 48:1490–1500.
- ÉHIK, J. 1938. Sakál vagy nádifarkas hazánkban [Jackal or reed-wolf from Hungary]. *Annales Musei Nationalis Hungarici, Pars Zoologica* 31:11–15.
- EISENBERG, J. F., AND M. LOCKHART. 1972. An ecological reconnaissance of Wilpattu National Park, Ceylon. *Smithsonian Contributions to Zoology* 101:1–118.
- ELLERMAN, J. R., AND T. C. S. MORRISON-SCOTT. 1951. Checklist of Palaearctic and Indian mammals 1758 to 1946. Trustees of the British Museum (Natural History), London, United Kingdom.
- EWER, R. F. 1956. The fossil carnivores of the Transvaal caves: Canidae. *Proceedings of the Zoological Society of London* 126:97–120.
- FABBRI, E., ET AL. 2014. Genetic structure and expansion of golden jackals (*Canis aureus*) in the north-western distribution range (Croatia and eastern Italian Alps). *Conservation Genetics* 15:187–199.
- FERGUSON, W. W. 1981. The systematic position of *Canis aureus lupaster* (Carnivora: Canidae) and the occurrence of *Canis lupus* in North Africa, Egypt and Sinai. *Mammalia* 45:459–465.
- FLOWER, S. S. 1932. Notes on the recent mammals of Egypt, with a list of the species recorded from that kingdom. *Proceedings of the Zoological Society of London* 1932:369–450.
- FERGUSON, W. W. 2002. *The mammals of Israel*. Gefen Publishing, Jerusalem, Israel.
- FULLER, T. K., A. R. BIKNEVICIUS, P. W. KAT, B. VAN VALKENBURGH, AND R. K. WAYNE. 1989. The ecology of three sympatric jackal species in the Rift Valley of Kenya. *African Journal of Ecology* 27:313–323.
- GALOV, A., ET AL. 2015. First evidence of hybridization between golden jackal (*Canis aureus*) and domestic dog (*Canis familiaris*) as revealed by genetic markers. *Royal Society Open Science* 2:150450.
- GAUBERT, P., ET AL. 2012. Reviving the African wolf *Canis lupus lupaster* in North and West Africa: a mitochondrial lineage ranging more than 6,000 km wide. *PLoS One* 7:e24740.
- GEOFFROY SAINT-HILAIRE, I. 1835. *Expédition scientifique de Morée. Section des sciences physiques. Tome III, 1st Partie. Zoologie. Première section, des animaux vertébrés. Mammifères et oiseaux*. F.G. Levrault, Paris, France.
- GIANNATOS, G., Y. MARINOS, P. MARAGOU, AND G. CATSADORAKIS. 2005. The status of the golden jackal (*Canis aureus*) in Greece. *Belgian Journal of Zoology* 135:145–149.
- GINSBERG, J. R., AND D. W. MACDONALD (eds.). 1990. Foxes, wolves, jackals, and dogs. An action plan for the conservation of canids. International Union for the Conservation of Nature, Gland, Switzerland.
- GOLANI, I., AND A. KELLER. 1975. A longitudinal field study of the behavior of a pair of golden jackals. Pp. 303–335 in *The wild canids* (M. W. Fox, ed.). Van Nostrand Reinhold, New York.
- GOLANI, I., AND H. MENDELSSOHN. 1971. Sequences of precopulatory behaviour of the jackal (*Canis aureus* L.). *Behaviour* 38:169–192.
- HAMILTON SMITH, C. 1839. *The natural history of dogs: Canidae or genus Canis of authors; including also the genera Hyaena and Proteles*. Pp. 77–267 in *Jardine's naturalist library. Volume 18. Mammalia. Dogs*. Volume 1 (W. Jardine, ed.). W. H. Lizars, Edinburgh, Scotland.
- HATT, R. T. 1959. *The mammals of Iraq*. Miscellaneous Publications of the Museum of Zoology, University of Michigan 106:1–113.
- HELLER, E. 1914. New subspecies of mammals from equatorial Africa. *Smithsonian Miscellaneous Collections* 63:1–12.
- HEMPRICH, F. G., AND C. G. EHRENBURG. 1830 (1833). *Symbolae physicae, seu icones et descriptiones mammalium quae ex itinere per Africam borealem et Asiam occidentalem. Zoologica I, Mammalium, pt. 2. Ex Officina Academica, venditur a Mittler, Berlin, Germany*.
- HEMPRICH, F. G., AND C. G. EHRENBURG. 1833. *Symbolae Physicae, seu icones et descriptiones corporum naturalium novorum aut minus cognitorum quae ex itinere per Libyam Aegyptum Nubiam Dongolam Syrian Arabiam et Habessianiam publico institutis sumptu. Signature ff. Ex Officina Academica, venditur a Mittler, Berlin, Germany*.
- HILZHEIMER, M. 1906a. *Papio mundamensis, Felis deliensis, Canis reissii* und andre neue Säugetiere. *Zoologischer Anzeiger* 30:109–118.
- HILZHEIMER, M. 1906b. Die geographische Verbreitung der Afrikanischen Grauschakale. *Zoologischer Beobachter* 47:363–373.
- HODGSON, B. H. 1833. Description of the wild dog of the Himalaya. *Asiatic Research* 18:221–237.
- HOOIJER, D. A. 1961. The fossil vertebrates of Ksar' Akil, a Paleolithic rock-shelter in Lebanon. *Zoologische Verhandlungen* 49:1–68.
- JAEGER, M. M., E. HAQUE, P. SULTANA, AND R. L. BRUGGERS. 2007. Daytime cover, diet and space-use of golden jackals (*Canis aureus*) in agroecosystems of Bangladesh. *Mammalia* 71:1–10.
- JAEGER, M. M., R. K. PANDIT, AND E. HAQUE. 1996. Season differences in territorial behavior by golden jackals in Bangladesh: howling versus confrontation. *Journal of Mammalogy* 77:768–775.
- JERDON, T. C. 1984. *A hand book of the mammals of India*. Mittal Publishers, Delhi, India.
- JHALA, Y., AND P. D. MOEHLMAN. 2004. Golden jackal (*Canis aureus*). Pp. 156–160 in *Canids: foxes, wolves, jackals and dogs. Status survey and conservation action plan* (C. Sillero-Zubiri, C. M. Hoffmann, and D. W. Macdonald, eds.). IUCN/SSC Canid Specialist Group, Gland, Switzerland.
- JHALA, Y., AND P. D. MOEHLMAN. 2008. *Canis aureus*. The IUCN Red List of Threatened Species 2008: e.T3744A10054631. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T3744A10054631.en>. Accessed 19 May 2016.

- KHIDAS, K. 1990. Contribution à la connaissance du chacal doré. Facteurs modulant l'organisation sociale et territoriale de la sous-espèce algérienne (*Canis aureus algirensis* Wagner, 1841). *Mammalia* 54:361–375.
- KINGDON, J. 1977. East African mammals. An atlas of evolution in East Africa. Volume IIIA: carnivores. Academic Press, London, United Kingdom.
- KOEPFLI, K.-P., ET AL. 2015. Genome-wide evidence reveals that African and Eurasian golden jackals are distinct species. *Current Biology* 25:2158–2165.
- KOLENATI, F. A. 1858. Reiseerinnerungen. Die Bereisung Hocharmeniens und Elisabethopols, der Schekinschen Provinz und des Kasbek im Central-Kaukasus. Verlagsbuchhandlung von Rudolf Kuntze, Dresden, Germany.
- KRETZOI, M. 1947. New names for mammals (Notes on nomenclature No. 3). *Annales Historico-Naturales Musei Nationalis Hungarici* 40:285–287.
- KROFEL, M. 2007. Golden jackals (*Canis aureus* L.) on the Pelješac Peninsula (southern Dalmatia, Croatia). *Natura Croatica* 16:201–204.
- KROFEL, M. 2008. Survey of golden jackals (*Canis aureus* L.) in northern Dalmatia, Croatia: preliminary results. *Natura Croatica* 17:259–264.
- KRUUK, H. 1972. The spotted hyena. University of Chicago Press, Chicago, Illinois.
- KRYŠTUFÉK, B., D. MURARIU, AND C. KURTONUR. 1997. Present distribution of the golden jackal *Canis aureus* in the Balkans and adjacent regions. *Mammal Review* 27:109–114.
- KURTEN, B. 1974. A history of coyote-like dogs (Canidae, Mammalia). *Acta Zoologica Fennica* 140:1–38.
- LAHMAR, S., ET AL. 2009. *Echinococcus* in the wild carnivores and stray dogs of northern Tunisia: the results of a pilot survey. *Annals of Tropical Medicine & Parasitology* 103:323–331.
- LANSZKI, J., M. HELTAL, AND L. SZABÓ. 2006. Feeding habits and trophic niche overlap between sympatric golden jackal (*Canis aureus*) and red fox (*Vulpes vulpes*) in the Pannonian ecoregion (Hungary). *Canadian Journal of Zoology* 84: 1647–1656.
- LAPINI, L., P. MOLINARI, L. DORIGO, G. ARE, AND P. BERALDO. 2009. Reproduction of the golden jackal (*Canis aureus moreoticus* I. Geoffroy Saint Hilaire, 1835) in Julian Pre-Alps, with new data on its range-expansion in the High-Adriatic Hinterland. *Bollettino del Museo Civico di Storia Naturale di Venezia* 60:169–186.
- VAN LAWICK, H., AND J. VAN LAWICK-GOODALL. 1970. Innocent killers. Collins, London, United Kingdom.
- LEWIS, R. E., J. H. LEWIS, AND S. I. ATALLAH. 1968. A review of Lebanese mammals. Carnivora, Pinnipedia, Hyracoidea and Artiodactyla. *Journal of Zoology (London)* 154:517–531.
- LINNAEUS, C. 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata*. Volume 1. Laurentii Salvii, Stockholm, Sweden.
- L. M. ISAEV INSTITUTE OF MEDICAL PARASITOLOGY. 1998. Dracunculiasis eradication in Uzbekistan. Country Report to the Ministry of Health, Uzbekistan.
- LORENZ, L. 1906. Über zwei neue Schakale aus Nordostafrika aus der Sammlung des Freiherrn C. von Erlanger. *Anzeiger, Kaiserliche Akademie der Wissenschaften in Wien. Mathematisch-Naturwissenschaftliche Klasse* 43:306–310.
- MACDONALD, D. W. 1979. The flexible social system of the golden jackal, *Canis aureus*. *Behavioral Ecology and Sociobiology* 5:17–38.
- MACDONALD, D. W., AND P. D. MOEHLMAN. 1983. Cooperation, altruism, and restraint in the reproduction of carnivores. Pp. 433–467 in *Perspectives in ethology* (P. Bateson and P. Klopfer, eds.). Plenum Press, New York.
- MADDOCK, L. 1979. The “migration” and grazing succession. Pp. 104–129 in *Serengeti, dynamics of an ecosystem* (A. R. E. Sinclair and M. Norton-Griffiths, eds.). University of Chicago Press, Chicago, Illinois.
- MALEZ, V. 1984. The zooarchaeological data as the base of colonizing the Markova cave on the island of Hvar. Pp. 617–621 in 9th Yugoslavian speleological congress proceedings, Yugoslavia (in Croatian).
- MARKOV, G., AND J. LANSZKI. 2012. Diet composition of the golden jackal, *Canis aureus* in an agricultural environment. *Folia Zoologica* 61:44–48.
- MATSCHE, P. 1900. Herr Matschie sprach über den Schakal des Menam-Gebietes in Siam. *Sitzungs-Berichte der Gesellschaft der Naturforschender Freunde zu Berlin* 3:144–145.
- MOEHLMAN, P. D. 1983. Socioecology of silverbacked and golden jackals (*Canis mesomelas* and *Canis aureus*). Pp. 423–453 in *Advances in the study of mammalian behavior* (J. F. Eisenberg and D. G. Kleiman, eds.). American Society of Mammalogists Special Publications, Lawrence, Kansas.
- MOEHLMAN, P. D. 1986. Ecology of cooperation in canids. Pp. 64–86 in *Ecological aspects of social evolution* (D. I. Rubenstein and R. W. Wrangham, eds.). Princeton University Press, Princeton, New Jersey.
- MOEHLMAN, P. D. 1989. Intraspecific variation in canid social systems. Pp. 143–163 in *Carnivore behavior, ecology, and evolution* (J. L. Gittleman, ed.). Chapman & Hall, London, United Kingdom.
- MOEHLMAN, P. D., AND H. HOFER. 1997. Cooperative breeding, reproductive suppression, and body mass in canids. Pp. 76–128 in *Cooperative breeding in mammals* (N. G. Solomon and J. A. French, eds.). Cambridge University Press, Cambridge, United Kingdom.
- MOEHLMAN, P. D., AND Y. V. JHALA. 2013. *Canis aureus* golden jackal (Asiatic jackal, common jackal). Pp. 35–38 in *Mammals of Africa, volume 5* (J. Kingdon, D. Happold, T. Butynski, M. Hoffman, M. Happold, and J. Kalina, eds.). Bloomsbury Publishing, London, United Kingdom.
- MOFARIED, A. L. 2013. Macro-anatomical investigation of the skull of golden jackal (*Canis aureus*) and its clinical application during regional anesthesia. *Global Veterinaria* 10:547–550.
- MOJSISOVICO, A. 1897. Das Thierleben der Österreichisch-Ungarischen Tiefebene. Biologische und thiergeographische Skizzen und Bilder. A. Hölder, Wien, Austria.
- MORRISON-SCOTT, T. C. S. 1939. Some Arabian mammals collected by Mr. H. St. J. B. Philby, C. I. E. *Novitates Zoologicae* 41:181–211.
- MUKHERJEE, S., S. P. GOYAL, A. J. T. JOHNSINGH, AND M. R. P. LEITE PITMAN. 2004. The importance of rodents in the diet of jungle cat (*Felis chaus*), caracal (*Caracal caracal*) and golden jackal (*Canis aureus*) in Sariska Tiger Reserve, Rajasthan, India. *Journal of Zoology (London)* 262:405–411.
- MULLER-USING, D. 1975. Jackals. Pp. 236–242 in *Grzimek's animal life encyclopedia* (B. Grzimek, ed.). Van Nostrand Reinhold Company, New York.
- NOACK, T. 1896. I. Wissenschaftliche Mittheilungen. I. Ein neuer Steinbock und ein neuer (?) [sic] Canide aus Arabien. *Zoologischer Anzeiger* 19:353–356.
- NOACK, T. 1897. Ostafrikanische Schakale. *Zoologischer Anzeiger* 20:517–521.
- OLARTE-CASTILLO, X. A., ET AL. 2015. Molecular characterization of canine kobuvirus in wild carnivores and the domestic dog in Africa. *Virology* 477:89–97.
- OSBORN, D. J., AND I. HELMY. 1980. The contemporary land mammals of Egypt (including Sinai). *Fieldiana Zoology, New Series* 5:1–579.
- PALMQVIST, P., M. MENDOZA, A. ARRIBAS, AND D. R. GRÖCKE. 2002. Estimating the body mass of Pleistocene canids: discussion of some methodological problems and a new ‘taxon free’ approach. *Lethaia* 35:358–360.
- PATTON, W. S. 1910. Preliminary report on a new piroplasm (*Piroplasma gibsoni* sp. nov.) found in the blood of the hounds of the Madras Hunt and subsequently discovered in the blood of the jackal *Canis aureus*. *Bulletin de la Société de Pathologie Exotique* 3:274–280.
- PERINI, F. A., C. A. M. RUSSO, AND C. G. SCHRAGO. 2010. The evolution of South American endemic canids: a history of rapid diversification and morphological parallelism. *Journal of Evolutionary Biology* 23:311–322.
- POCHÉ, R. M., S. J. EVANS, P. SULTANA, M. E. HAGUE, R. STERNER, AND M. A. SIDDIQUE. 1987. Notes on the golden jackal (*Canis aureus*) in Bangladesh. *Mammalia* 51:259–270.
- POCOCK, R. I. 1938. The jackals of S.W. Asia and S.E. Europe. *Proceedings of the Zoological Society of London* 108:37–40.
- PRATER, S. 1971. The book of Indian animals. Bombay Natural History Society, Bombay, India.
- QUMSIYEH, M. B. 1996. *Mammals of the Holy Land*. Texas Tech University Press, Lubbock.

- RAICHEV, E. 2011. Effect of shooting on the structure of population of golden jackal (*Canis aureus* L.) in Sarnena Sredna Gora Mountain. *Agricultural Science and Technology* 3:276–280.
- ROELKE-PARKER, M. E., ET AL. 1996. A canine distemper virus epidemic in Serengeti lions (*Panthera leo*). *Nature* 379:441–445.
- ROSEVEAR, D. R. 1974. The carnivores of West Africa. Trustees of the British Museum (Natural History), London, United Kingdom.
- ROTEM, G., H. BERGER, R. KING, P. B. (KUTIEL) [SIC], AND D. SALTZ. 2011. The effect of anthropogenic resources on the space-use patterns of golden jackals. *Journal of Wildlife Management* 78:132–136.
- RUENESS, E. K., ET AL. 2011. The cryptic African wolf: *Canis aureus lupaster* is not a golden jackal and is not endemic to Egypt. *PLoS One* 6:e16385.
- RUTKOWSKI, R., ET AL. 2015. A European concern? Genetic structure and expansion of golden jackals (*Canis aureus*) in Europe and the Caucasus. *PLoS One* 10:e0141236.
- SALEH, M. A., AND M. I. BASUONY. 2014. Mammals of the genus *Canis* Linnaeus, 1758 (Canidae, Carnivora) in Egypt. *Egyptian Journal of Zoology* 62:49–92.
- SAVAGE, R. J. 1977. Evolution in carnivorous mammals. *Paleontology* 20:237–271.
- SEITZ, V. A. 1959. Beobachtungen an Handaufgezogenen Goldschakalen (*Canis aureus aligirensis*). *Zeitschrift für Tierpsychologie* 16:747–771.
- SHAMIR, M., ET AL. 2001. Antibodies to selected canine pathogens and infestation with intestinal helminths in golden jackals (*Canis aureus*) in Israel. *Veterinary Journal* 162:66–72.
- SHAW, G. 1800. General zoology or systematic natural history. Volume 1, part 2. G. Kearsley, London, United Kingdom.
- SINGH, A. N. 1983. A study of diverse prey species of python (*Python molurus*) with special reference to its interaction with jackal (*Canis aureus*). *Tigerpaper* 10:31–32.
- SHELDON, J. W. 1992. Wild dogs: the natural history of the non-domestic Canidae. Academic Press, Chicago, Illinois.
- SOMMER, R., AND N. BENECKE. 2005. Late-Pleistocene and early Holocene history of the canid fauna of Europe (Canidae). *Mammalian Biology* 70:227–241.
- SPASSOV, N. 1989. The position of jackals in the *Canis* genus and life-history of the golden jackal (*Canis aureus* L.) in Bulgaria and the Balkans. *Historia Naturalis Bulgarica*, Sofia 1:44–56.
- STOYANEV, S. 2012. Craniometric differentiation of golden jackals (*Canis aureus* L., 1758) in Bulgaria. Pp. 39–47 in International symposium on hunting, modern aspects of sustainable management of game populations, Zemun-Belgrade, Serbia.
- TALMI-FRANK, D., ET AL. 2010. *Leishmania tropica* infection in golden jackals and red foxes, Israel. *Emerging Infectious Diseases* 16:1973–1975.
- TEMU, S. E., C. L. NAHONYO, AND P. D. MOEHLMAN. 2016. Comparative foraging efficiency of two sympatric jackals, silver-backed jackals (*Canis mesomelas*) and golden jackals (*Canis aureus*), in the Ngorongoro Crater, Tanzania. *International Journal of Ecology*, Article ID 6178940, 5 pages. doi:10.1155/2016/6178940
- TEMU, S. E., C. L. NAHONYO, AND P. D. MOEHLMAN. 2017. Hunting of a bat-eared fox (*Otocyon megalotis*) and an adult Thomson's gazelle (*Eudorcas thomsoni*) by golden jackals (*Canis aureus*) in the Ngorongoro Crater, Tanzania. *African Journal of Ecology* 55:386–387.
- THOMAS, O. 1903. On some mammals collected by Capt. H. N. Dunn, R. A. M. C., in the Soudan. *Proceedings of the Zoological Society of London* 103:294–300.
- THOMAS, O. 1911. The mammals of the tenth edition of Linnaeus; an attempt to fix the types of the genera and the exact bases and localities of the species. *Proceedings of the Zoological Society of London* 1911:120–158.
- TÓTH, T., L. KRECSÁK, E. SZÜCS, M. HELTAI, AND G. HUSZÁR. 2009. Records of the golden jackal (*Canis aureus* Linnaeus, 1758) in Hungary from 1800th until 2007, based on a literature survey. *North-Western Journal of Zoology* 5:386–405.
- TROUWBORST, A., M. KROFEL, AND J. D. C. LINNELL. 2015. Legal implications of range expansions in a terrestrial carnivore: the case of the golden jackal (*Canis aureus*) in Europe. *Biodiversity and Conservation* 24:2593–2610.
- VAN VALKENBURGH, B. 2009. Costs of carnivory: tooth fracture in Pleistocene and Recent carnivores. *Biological Journal of the Linnean Society* 96:68–81.
- VAN VALKENBURGH, B., AND R. K. WAYNE. 1994. Shape divergence associated with size convergence in sympatric East African jackals. *Ecology* 75:1567–1581.
- WAGNER, A. 1829. Oken's Isis 9:986–989.
- WAGNER, J. A. 1841. Schreber's Die Säugethiere in Abbildungen nach der Natur mit Beschreibungen, Supplementary Volume 2. Wolfgang Walther, Erlangen, Germany.
- WANDRY, R. 1975. Contribution to the study of the social behaviour of captive golden jackals (*Canis aureus* L.). *Tierpsychol* 39:365–402.
- WALTON, O. R., AND D. O. JOLY. 2003. *Canis mesomelas*. *Mammalian Species* 715:1–9.
- WEIGL, R. 2005. Longevity of mammals in captivity; from the living collections of the world. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, Germany.
- WOZENCRAFT, W. C. 2005. Order Carnivora. Pp. 532–628 in *Mammal species of the world: a taxonomic and geographic reference* (D. E. Wilson and D. M. Reeder, eds.). 3rd ed. Johns Hopkins University Press, Baltimore, Maryland.
- WROUGHTON, R. C. 1916. Scientific results of the mammal survey. (E) The Indian jackals. *Journal of the Bombay Natural History Society* 24:649–655.
- WURSTER, D. H., AND K. BENIRSCHKE. 1968. Comparative cytogenetic studies in the order Carnivora. *Chromosoma* 24:336–382.
- WYMAN, J. 1967. The jackals of the Serengeti. *Animals* 10:79–83.
- YOM-TOV, Y., S. ASHKENAZI, AND O. VINER. 1995. Cattle predation by the golden jackal *Canis aureus* in the Golan Heights, Israel. *Biological Conservation* 73:19–22.
- YUMNAM, B., T. NEGI, J. E. MALDONADO, R. C. FLEISCHER, AND Y. V. JHALA. 2015. Phylogeography of the golden jackal (*Canis aureus*) in India. *PLoS One* 10:e0138497.
- ZACHOS, F. E., ET AL. 2009. Genetic variability, differentiation, and founder effect in golden jackals (*Canis aureus*) from Serbia as revealed by mitochondrial DNA and nuclear microsatellite loci. *Biochemical Genetics* 47:241–250.

Associate Editor of this account was Jamie M. Harris. Editor was MEREDITH J. HAMILTON.