



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Commentary

Preparing nurses for COVID-19 response efforts through involvement in antimicrobial stewardship programmes

M. Courtenay^{a,*}, E. Burnett^b, E. Castro-Sánchez^c, B. Du Toit^d,
R.M. Figueiredo^e, R. Gallagher^f, F. Gotterson^g, H. Kennedy^h, E. Maniasⁱ,
J. McEwen^h, V. Ness^j, R. Olans^k, M.C. Padoveze^l

^a School of Health Sciences, Cardiff University, Cardiff, UK

^b School of Health Sciences, University of Dundee, Scotland, UK

^c NIHR Health Protection Unit in Healthcare Associated Infection and Antimicrobial Resistance at Imperial College London, London, UK

^d Mediclinic Southern Africa, Stellenbosch, South Africa

^e Nursing Department, University of Sao Carlos, Brazil

^f Royal College of Nursing, London, UK

^g The University of Melbourne, National Centre for Antimicrobial Stewardship, Australia

^h NHS Tayside, Dundee, UK

ⁱ School of Nursing and Midwifery, Centre for Quality and Patient Safety, Institute for Health Transformation, Deakin University, Australia

^j Glasgow Caledonian University, Glasgow, UK

^k School of Nursing, MGH Institute of Health Professions, Boston, MA, USA

^l School of Nursing, University of São Paulo, Brazil

ARTICLE INFO

Article history:

Received 30 May 2020

Accepted 3 June 2020

Available online 10 June 2020



Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) continues to spread internationally with 5,593,631 cases reported globally including 353,334 deaths [1]. Its rapid emergence and dissemination have highlighted multiple areas in which competencies in antimicrobial stewardship (AMS) (the safe and effective use of antimicrobials), specifically by nurses,

can support response efforts. There have been calls for nurses to be recognized as legitimate contributors to AMS team efforts [2–5]. Unfortunately, nurses' roles in these efforts have received minimal mention in international and national policies [4–10]. It is reported that nurses' knowledge of antibiotics is poor [11,12], and that AMS taught in nurse undergraduate programmes is disparate or lacking [13]. To address this gap, AMS consensus-based international competency statements have been developed, focused on six domains (infection prevention and control, antimicrobials and antimicrobial resistance, the diagnosis of infection and use of antibiotics, antimicrobial prescribing practice, person-centred care, interprofessional collaborative practice), which are (seen as) priorities/minimum requirements for nurses [14,15].

Nurses are often the first healthcare workers to come into contact with patients infected by SARS-CoV-2, and thus need to be familiar with the signs and symptoms of the disease in order to ensure prompt implementation of additional precautions to stop transmission [16]. Differentiating between viral and bacterial pneumonia can be challenging [17] and thus awareness of how these symptoms differ is crucial. This awareness is central to reducing unnecessary prescriptions of not just antibiotics, as COVID-19 is caused by a virus, but other anti-infective drugs

* Corresponding author. Address: School of Health Sciences, Cardiff University, Cardiff, CF24 OAB, UK.

E-mail address: courtenaym@cardiff.ac.uk (M. Courtenay).

associated with AMS such as antifungal and antiviral medications. AMS requires nurses to be able to distinguish between symptoms and ask questions about the use of antibiotics, should symptoms be more consistent with a viral infection, or when microbiology results do not indicate a bacterial cause [14]. Such actions will help to ensure that antimicrobial resistance is not an unintended consequence of COVID-19.

Myths about COVID-19 circulated through social media and disreputable websites, the plethora of ever-evolving data and information, and the guidance on COVID-19 prevention presented daily, have made it difficult for the public to understand how the infection is transmitted and if antibiotics are necessary. Increasingly, the importance of engaging with patients and involving them in decision making to optimize AMS efforts, has become recognized [14]. Nurses, with their focus on patient-centred care, are in an excellent position to engage with the public and communicate messages about interrupting the spread of infection and the appropriate use of antimicrobials, including the importance of prudent use for patients who really need them.

It is also important to recognize nurses' engagement through interprofessional collaboration, which involves nurses and other health professionals from various disciplines working together with shared goals, mutual trust, respect, and understanding about each other's roles, along with acceptance that patients are team members [18,19]. There is an association between interprofessional collaboration and patient safety [20,21]. In AMS, interprofessional collaboration requires shared understanding about antimicrobial treatment decisions and plans, and about the expected outcomes of antimicrobial therapy [14].

Infection prevention and control quality-assurance frameworks [22] highlight that arrangements around AMS should be maintained during the COVID-19 pandemic, however, the outbreak of SARS-CoV-2 presents enormous challenges for AMS and for teamwork and interprofessional collaboration [23]. Traditional AMS teams may have fallen apart during the pandemic, through repurposing of staff and the impact on microbiology services of supporting the COVID-19 workload. Furthermore, the rapid emergence and dissemination of COVID-19 has resulted in reconfiguration of healthcare services, redeployment of healthcare workers into areas where they may have less expertise, and employment of retired practitioners who have less experience of antibiotic stewardship. This reshaping of interprofessional relations, with the addition of different professionals, with various professional backgrounds and experiences, potentially dilutes existing team cohesions, and could affect discussions and decisions about antimicrobial prescribing. Nurses are pivotal as leaders and influencers and this pandemic throws a spotlight on their communication and coordination of holistic patient care, clinical practice and facilitation of interprofessional collaboration. Nurses, however, may lack confidence in this endeavour; with the reshaping of healthcare teams increasing uncertainty and the need for clinical and professional supervision to overcome challenges. Applying AMS competencies will help nurses to consolidate their appreciation of interprofessional collaboration, and its contribution to AMS, and empower them to enact their AMS role, which in turn would enable sustainability of AMS activities during challenging times [14]. In applying AMS competencies, nurses will know the prerequisites for meaningful collaboration, including effective communication, negotiation

and assertiveness skills [5,24,25]. Consequently, they are in a better position to confidently discuss antimicrobial-related issues with various health professionals, such as a switch from intravenous to oral therapy, and would recognize when specific communication techniques and tools are needed to convey and act on critical messages [24,26]. Building competence to practise collaboratively enables nurses to be active participants in AMS policy decisions, and involves nurses taking advantage of opportunities to discuss antimicrobial treatment decisions and management plans with their colleagues, patients, and carers. Through the application of competencies [14], nurses can work deliberately to build rapport and trust amongst team members, knowing that rapport and trust are necessary for genuine interprofessional collaborative practice to happen.

COVID-19 has focused attention on nurse leaders' power and potential to promote AMS. A key AMS strategy in the current pandemic is to promote the message that antibiotics should not be prescribed for viral infections, and that these medicines must only be prescribed for those for whom serious bacterial (primary or secondary) infection is suspected. By involving nurses in AMS leadership positions, role modelling the importance of AMS behaviours, and advocating for, and supporting nurses in enacting their AMS role, a strong signal will be sent out to nurses about the importance of responsible antimicrobial management.

Conflict of interest statement

None declared.

Funding sources

None.

References

- [1] WHO. Coronavirus disease (COVID-2019) situation reports. 2020. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/> [last accessed May 2020].
- [2] Edwards R, Drumright LN, Kiernan M, Holmes A. Covering more territory to fight resistance: considering nurses' role in antimicrobial stewardship. *J Infect Prevent* 2011;12:6–10.
- [3] Olans RN, Olans RD, Demaria A. The critical role of the staff nurse in antimicrobial stewardship – unrecognized, but already there. *Clin Infect Dis* 2016;62:84–9.
- [4] American Association of Nurses (AANs). White Paper: Redefining the Antibiotic Stewardship Team: Recommendations from the AANs/Centers for Disease Control (CDC) and Prevention workgroup on the role of registered nurses in hospital antibiotic stewardship practices. 2017. <https://www.cdc.gov/antibiotic-use/healthcare/pdfs/ANA-CDC-whitepaper.pdf> [last accessed May 2020].
- [5] Castro-Sánchez E, Gilchrist M, McEwen J, Smith M, Holmes A. Antimicrobial Stewardship: Widening the Collaborative Approach. *J Antimicrob Stewardship* 2017;1(1):29–37.
- [6] European Commission. EU Guidelines for the prudent use of antimicrobials in human health (2017/C 212/01). https://ec.europa.eu/health/amr/sites/amr/files/amr_guidelines_prudent_use_en.pdf [last accessed May 2020].
- [7] European Federation of Nurses Associations (EFN). Nurses are frontline combating antimicrobial resistance. <http://www.efnweb.be/wp-content/uploads/EFN-AMR-Report-Nurses-are-frontline-combating-AMR-07-11-2017.pdf> [last accessed May 2020].

- [8] CDC. Core elements of hospital antibiotic stewardship programs. Atlanta, GA: US Department of Health and Human Services, CDC; 2014 [cited 02/04/2019]. Available at: <http://www.cdc.gov/getsmart/healthcare/implementation/core-elements> [last accessed May 2020].
- [9] RCN. Antimicrobial resistance. 2020. <https://www.rcn.org.uk/clinical-topics/infection-prevention-and-control/antimicrobial-resistance> [last accessed May 2020].
- [10] National Health Services Scotland. Scottish management of antimicrobial resistance action plan 2014–2018 (ScotMARAP 2). Edinburgh: The Scottish Government.
- [11] McEwen J, Burnett E. Antimicrobial, stewardship and pre-registration student nurses: evaluation of teaching. *J Infect Prev* 2019;19:80–6.
- [12] McGregor W, Walker G, Bayne G, McEwen J. Assessing knowledge of antimicrobial stewardship. *Nurs Times* 2015;111:15–7.
- [13] Castro-Sánchez E, Drumright LN, Gharbi M, Farrell S, Holmes AH. Mapping antimicrobial stewardship in undergraduate medical, dental, pharmacy, nursing and veterinary education in the United Kingdom. *PLoS One* 2016;11(2):e0150056.
- [14] Courtenay M, Castro-Sanchez E, Gallagher R, McEwen J, Bulabula ANH, Carre Y, et al. Development of consensus-based international antimicrobial stewardship competencies for undergraduate nurse education. *J Hosp Infect* 2019;103:244–250.
- [15] Courtenay M, Castro-Sanchez E, editors. *Antimicrobial stewardship for nursing practice*. Oxfordshire: CABI; 2020.
- [16] PHE. Reducing the risk of transmission of COVID-19 in the hospital setting. 2020. <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/reducing-the-risk-of-transmission-of-covid-19-in-the-hospital-setting> [last accessed May 2020].
- [17] NICE. COVID 19 Rapid guideline: antibiotics for pneumonia in adults in hospital. 2020 [NG 173]. Available at: <https://www.nice.org.uk/guidance/ng173> [last accessed May 2020].
- [18] D'Amour D, Ferrada-Videla M, San Martin Rodriguez L, Beaulieu M-D. The conceptual basis for interprofessional collaboration: Core concepts and theoretical frameworks. *J Interprof Care* 2005;19:116–31.
- [19] Petri L. Concept Analysis of Interdisciplinary Collaboration. *Nurs Forum* 2010;45:73–82.
- [20] Levett-Jones T, Burdett T, Chow YL, Jonsson L, Lasater K, Rae Mathews L, et al. Case studies of interprofessional education initiatives from five countries. *J Nurs Scholarsh* 2018;50:332–324.
- [21] Reeves S, Pelone F, Harrison R, Goldman J, Zwarenstein M. Interprofessional collaboration to improve professional practice and healthcare outcomes. *Cochrane Database Syst Rev* 2017;6:Cd000072.
- [22] Infection NHS. Prevention and control board assurance framework. NHS; 2020.
- [23] Nembhard IM, Burns LR, Shortell SM. Responding to covid-19: lessons from management research. 2020. Catalyst non-issue content. 1.
- [24] Curtis K, Tzannes A, Rudge T. How to talk to doctors – a guide for effective communication. *Int Nurs Rev* 2011;58:13–20.
- [25] Wilson AJ, Palmer L, Levett-Jones T, Gilligan C, Outram S. Interprofessional collaborative practice for medication safety: nursing, pharmacy, and medical graduates' experiences and perspectives. *J Interprof Care* 2016;30:649–54.
- [26] Foronda C, MacWilliams B, McArthur E. Interprofessional communication in healthcare: An integrative review. *Nurse Ed Pract* 2016;19:36–40.