

MEETING ABSTRACT

Open Access

EHMTI-0069. The low prevalence of “forehead and facial flushing” and “sensation of fullness in the ear” in Japanese cluster headache patients

N Imai

From 4th European Headache and Migraine Trust International Congress: EHMTIC 2014
Copenhagen, Denmark. 18-21 September 2014

Background

New ICHD-3 beta criteria for cluster headache added two accompanying autonomic symptoms: “forehead and facial flushing” and “sensation of fullness in the ear”. Previous reports showed that several clinical features of cluster headache were different between Caucasian and East Asian.

Aim

We evaluated the presence of “forehead and facial flushing” and “sensation of fullness in the ear” in Japan.

Method

Subjects comprised 23 consecutive cluster headache patients who had been diagnosed with cluster headache according to ICHD-2 (14 males, 9 females; mean age, 41.4 ±13.8 years; range, 20-71 years). Information was collected by face to face interview.

Results

All patients were episodic cluster headache. Two patients (9%) had “forehead and facial flushing” and 4 patients (17%) had “sensation of fullness in the ear”. The patients with “forehead and facial flushing” had also “sensation of fullness in the ear”. The prevalence of “forehead and facial flushing” and “sensation of fullness in the ear” in our study were lower than that of previous Caucasian study (20.0% reported “forehead and facial flushing” and 33.3% “sensation of fullness in the ear”).

Conclusion

The low prevalence of “forehead and facial flushing” and “sensation of fullness in the ear” may be clinical characteristic in Japanese cluster headache patients.

No conflict of interest.

Published: 18 September 2014

doi:10.1186/1129-2377-15-S1-C24

Cite this article as: Imai: EHMTI-0069. The low prevalence of “forehead and facial flushing” and “sensation of fullness in the ear” in Japanese cluster headache patients. *The Journal of Headache and Pain* 2014 **15** (Suppl 1):C24.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com

Department of Neurology, Japanese Red Cross Shizuoka Hospital, Shizuoka, Japan