

# Health Status and Working Condition of Migrant Workers: Major Public Health Problems

## Abstract

**Background:** Very little research and survey have been performed on the occupational health, hazards, and working condition of urban and rural of workers in Qatar. The aim of the current study is to identify the health status, lifestyle condition, working-related problems, and accidents experienced by Indian subcontinental migrants (ISCM) in Qatar and their access to health-care facilities. **Methods:** This is a cross-sectional study based on a representative sample of 1186 workers and study covering sociodemographic characteristics, medical conditions, and health-seeking behaviors and personal experience. **Results:** There were statistically significant differences between semi-urban and urban migrant workers in terms of educational, occupation, income, working hours, and accommodation type ( $P < 0.001$ ). Furthermore, there were statistically significant differences between migrant workers in terms of body mass index, delay in receiving salary, the right to medical insurance and sick day entitlement, cigarettes and shisha smoking, amenities, having on-site safety measure, and sleeping hours ( $P < 0.01$ ). Moreover, there were statistically significant differences between semi-urban and urban migrant workers in terms of pain, cardiopulmonary, gastrointestinal, and pseudoneurologic and medical symptoms ( $P < 0.01$ ). Multiple logistic regression was used for predictors' health problems in migrant workers such as the absence of drinking water, tap water availability, safety facility tools, occupational status, shisha smoking, toilet facility, working hours, and accommodation type were considered the strong predictors. **Conclusions:** The current study revealed that there is a lack of insufficient information for the migrants about their lifestyle, medical conditions, health risks, injury, and rights privilege in relation to legal working condition and health services.

**Keywords:** Access health care, health status, lifestyle, migrant workers, safety

## Introduction

Although the State of Qatar is a relatively new nation in the Middle East, Qatar has rapid developments, especially in the petroleum, airline, maritime, commercial, and construction industries.<sup>[1,2]</sup> The economic and industrial developments have caused numerous social and environmental problems during the past two decades.<sup>[1]</sup> Exposure to lead is considered a primary health issue in the industrial and oil-producing countries. In Qatar, the increase in the amount of oil and gas has been associated with the growth of urbanization and industry and increasing number of motor vehicles in transport.<sup>[2]</sup>

Low-skilled or unskilled workers are employed from poor countries with usually short-term contracts by most of the oil-rich countries.<sup>[1-5]</sup> They are recruited usually in projects which need workforce such

as construction and service industries.<sup>[1,2]</sup> Majority of the workers usually do not have enough access to health-care facilities. The main reasons can be language and cultural differences to access health care.<sup>[6-8]</sup> Overall, South Asian countries and North Africa are the main suppliers of migrant workers to the State of Qatar and other Gulf countries.<sup>[2,9-14]</sup>

More recently, it was asked by Guardian Newspaper that "Have Qatar's work conditions could be improved?"<sup>[15]</sup> Numerous workers still complain about low payment, long work hours, and limitation of migrant workers' rights and freedoms despite better accommodation and new payment rules.<sup>[15,16]</sup> Migrant workers' salary and payment conditions are not good, for example, the salaries of carpenters and steel fixers working on a World Cup stadium<sup>[17]</sup> site ranging from \$250 to \$300 US dollars with overtime. Qatari government commissioned detailed

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**How to cite this article:** Bener A. Health status and working condition of migrant workers: Major public health problems. *Int J Prev Med* 2017;8:68.

## Abdulbari Bener<sup>1,2,3</sup>

<sup>1</sup>Department of Biostatistics and Medical Informatics, Cerrahpaşa Faculty of Medicine, Istanbul University, Istanbul, Turkey, <sup>2</sup>Department of Evidence for Population Health Unit, School of Epidemiology and Health Sciences, University of Manchester, Manchester, UK, <sup>3</sup>Department of Endocrinology, Regenerative and Restorative Medicine Research Center, International School of Medicine, Istanbul Medipol University, Istanbul, Turkey

## Address for correspondence:

Prof. Abdulbari Bener,  
Department of Biostatistics  
and Medical Informatics,  
Cerrahpaşa Faculty of  
Medicine, Istanbul University,  
34098 Cerrahpasa, Istanbul,  
Turkey.  
E-mail: abdulbari.bener@istanbul.edu.tr

## Access this article online

**Website:**  
[www.ijpvmjournal.net/www.ijpvm.ir](http://www.ijpvmjournal.net/www.ijpvm.ir)

**DOI:**  
10.4103/ijpvm.IJPVM\_396\_16

## Quick Response Code:



report<sup>[18]</sup> for improving life conditions for migrant laborers with suggested substantial reforms.

During 2014, over 80% of all migrant workers in Qatar were expatriated. More than 1.5 million migrant workers were from South Asian countries such as India, Nepal, Pakistan, Bangladesh, Sri Lanka, Pilipino, and Korea.<sup>[19,20]</sup> It is estimated that one million migrant workers will be recruited in many projects including construction, soccer stadiums, urban–rural roads, tunnel, and metro railway systems for the preparation of World Cup football matches in 2022.

The aim of the current study is to identify the health status, lifestyle condition, work problems, and accidents experienced by Indian subcontinental migrants (ISCM) migrant workers in Qatar and their access to health-care facilities in urban and semi-urban areas.

## Methods

This is a cross-sectional study of the self-reported health condition of the migrant workers during their stay in Qatar. As of December 2013, there were an estimated 1,466,851 migrant workers in Nepalese, Indian, Bangladeshi, Sri Lankan, Arabs, and Pakistani workforce (ISCM) in Qatar. Of these, 1,466,851 were semi-skilled or unskilled workers including over 1.2 million in construction.<sup>[19,20]</sup>

The data were collected in January 2013 to June 2014 from various types of location: the International Airport, shopping malls, bus parking area, and working sites. The inclusion criteria of the present study were being adult people who had work experience of at least 12 months in Qatar; meanwhile, in most cases, working contract is about minimum 18 months. We have developed and partially used the previous questionnaire on the basis of author experience in this area which is one of the demographic and clinical questionnaires were adopted from a previous study.<sup>[12]</sup> We computed the sample size based on 50% prevalence of migrants might have access to health-care services, assuming 3.2% bound on error estimation, and taking 99% confidence level, the required minimum sample size for this study was  $n = 1623$ .

$$n = \frac{z_{\alpha/2}^2 pq}{d^2}$$

We have obtained a list of representative sample of 1623 workers, and we have approached workers, but 1186 participants agreed to participate in this study (73%). Urban area is mostly thickly populated area, usually greater than a town and larger congested area, having fixed boundaries and certain local powers of government. Semi-urban population is not thickly populated area with homogeneous people, and it is quite place and has less economic development.

The variables in the analysis were the migrants' self-reported health problems and the accidents during their past 12 months of working period in Qatar. The

self-reported questions were did you suffer from any health problem during your past 12 months of stay in that region? Did you have any type of injury or accident at your workplace during past 12 months? Variables such as age, marital status, occupation, educational status, ethnicity, working hours, accommodation type, body mass index (BMI), and types of health problems were recorded.

The data were analyzed using the Statistical Package for the Social Sciences (IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp). Student's *t*-test was used to ascertain the significance of differences between mean values of two continuous variables. Chi-square and Fisher's exact test (two-tailed) were used to test for differences in proportions of categorical variables between two or more groups. Multiple logistic regression analysis method was used to assess the potential confounders and order the importance of risk factors for the health problems.  $P < 0.05$  was considered as the cutoff value for significance.

## Results

Table 1 shows comparison of sociodemographic characteristics among participants from semi-urban and urban locations. There were statistically significant differences between semi-urban and urban migrant workers in terms of educational level, occupation status, household income, working hours, and accommodation type ( $P < 0.001$ ).

Table 2 shows some important comparison of working and lifestyle characteristics of migrant workers among participants from semi-urban and urban locations. There were statistically significant differences between semi-urban and urban migrant workers in terms of BMI, delay with receiving salary, having medical insurance, having sick day entitlement, smoking cigarettes and shisha smoking, amenities, having on-site safety measure, number of hours sleeping, and self-reported health status ( $P < 0.01$ ).

Table 3 shows some important comparison of lifetime prevalence of medical symptoms among migrant workers from semi-urban and urban locations. There were statistically significant differences between semi-urban and urban migrant workers in terms of pain, cardiopulmonary, gastrointestinal (GI), and pseudoneurologic and medical symptom conditions grouping ( $P < 0.01$ ).

Figure 1 shows the type of occupational accidents according to locations. As can be seen from this figure that there were statistically significant differences between semi-urban and urban migrant workers in terms of over half of the fatal injuries to workers were of three kinds of most common falls from height (18.6%), then slip/trip (8.5%), followed by machinery accidents (7.3%) ( $P = 0.014$ ).

Multiple logistic regression method was used to investigate the potential confounding risk factors in the multivariable analyses. Table 4 identified the predictors for health problems in migrant workers such as absence of drinking

**Table 1: Sociodemographic characteristics of the studied participants (n=1186)**

Variables	Semi-urban (n=564), n (%)	Urban (n=622), n (%)	P
Age group, (years)			
18-24	58 (10.3)	83 (13.4)	0.549
25-34	188 (33.3)	206 (33.1)	
35-44	150 (26.6)	164 (26.4)	
45-54	115 (20.4)	116 (18.6)	
≥55	53 (9.4)	53 (8.5)	
Sex			
Male	518 (91.8)	581 (93.4)	0.317
Female	46 (8.2)	41 (6.6)	
Ethnicity			
Sri Lankan	53 (9.4)	52 (8.4)	0.559
Bangladeshi	62 (11.0)	64 (10.3)	
Nepalese	126 (22.4)	154 (24.7)	
Indian	111 (19.7)	113 (18.1)	
Pakistani	71 (12.6)	71 (11.4)	
Philippines	43 (7.6)	47 (7.6)	
Arabs/Egypt/Syrian/Palestinian	73 (12.9)	102 (16.4)	
North Korean	25 (4.4)	19 (3.1)	
Marital status			
Single	64 (11.3)	82 (13.2)	0.376
Married	500 (88.7)	540 (86.8)	
Educational level			
Primary	297 (52.7)	281 (45.2)	<0.001
Intermediate	161 (28.5)	145 (23.3)	
Secondary	64 (11.3)	116 (18.6)	
University	42 (7.4)	80 (12.9)	
Occupation			
Construction works	269 (47.7)	210 (33.8)	<0.001
General laborer	83 (14.7)	53 (8.5)	
Carpentry and repair	77 (13.7)	88 (14.1)	
Agricultural works	48 (8.5)	79 (12.7)	
House service/driver	34 (6.0)	89 (14.3)	
Engineer/sedentary/clerical	53 (9.4)	103 (16.6)	
Household income (QRs)			
<1500	3534 (62.6)	263 (42.3)	<0.001
1500-2999	118 (21.0)	160 (25.7)	
3000-4500	34 (6.0)	89 (14.3)	
>5000	59 (10.5)	110 (17.7)	
Working hours			
<8	178 (31.6)	232 (37.3)	<0.001
9-10	259 (45.9)	277 (44.5)	
>10	127 (22.5)	113 (18.2)	
Working days			
5	124 (22.0)	135 (21.7)	0.144
6	313 (55.5)	374 (60.1)	
7	127 (22.5)	113 (18.2)	
Accommodation type			
Dormitory room	310 (55.0)	293 (47.1)	<0.001
Prefabric house	142 (25.2)	123 (19.8)	
Traditional house	61 (10.8)	116 (18.6)	
Apartment/flat	51 (9.0)	90 (14.5)	

QRs=\$ 0.275 US Dollars

**Table 2: Some important characteristics lifestyle habits among the studied participants (n=1186)**

Variables	Semi-urban (n=564), n (%)	Urban (n=622), n (%)	P
BMI			
Normal (<25 kg/m <sup>2</sup> )	174 (30.1)	161 (25.9)	0.058
Over weight (25-30 kg/m <sup>2</sup> )	228 (40.4)	289 (46.4)	0.036
Obese (30+ kg/m <sup>2</sup> )	166 (29.5)	172 (27.7)	0.497
Delay with receiving salary			
Yes	216 (38.3)	200 (32.1)	0.027
No	348 (61.7)	422 (67.9)	
Having medical insurance			
Yes	70 (12.4)	52 (8.4)	0.022
No	494 (87.6)	570 (91.6)	
Having sick day entitlement			
Yes	132 (23.4)	266 (19.5)	<0.001
No	432 (67.0)	356 (57.2)	
Have you ever smoked shisha			
Yes	83 (14.7)	55 (8.8)	0.002
No	432 (67.6)	501 (80.5)	
Have you ever smoked cigarette			
Never	367 (65.1)	453 (72.8)	0.014
Current smoker	127 (22.5)	105 (16.9)	
Ex-smoker	70 (12.4)	64 (10.3)	
Amenities			
Inside tap water	288 (51.1)	384 (61.7)	<0.001
Drinking water available	327 (58.1)	400 (64.3)	0.023
Inside toilet	340 (60.3)	433 (69.6)	<0.001
Air condition available	175 (31.0)	238 (38.3)	0.004
Have on-site safety			
Yes	176 (31.2)	256 (41.2)	<0.001
No	388 (68.8)	366 (58.2)	
Hours of sleep (mean±SD)	6.1±1.2	6.8±1.6	<0.001
Self-reported health status			
Excellent/very good	169 (28.4)	190 (30.5)	<0.001
Good	280 (49.6)	336 (54.0)	
Poor	124 (22.0)	96 (15.4)	

BMI=Body mass index, SD=Standard deviation

water (odds ratio [OR] = 2.73; 95% confidence interval [CI]: 2.48–3.95,  $P < 0.001$ ), inside tap water availability (OR = 2.69; 95% CI: 2.24–3.58,  $P = 0.003$ ), safety facility and tools (OR = 2.73; 95% CI: 2.48–3.95,  $P < 0.001$ ), occupational status (OR = 2.13; 95% CI: 1.65–3.22,  $P < 0.001$ ), shisha smoking (OR = 1.96; 95% CI: 1.27–3.14,  $P = 0.031$ ), toilet facility, (OR = 1.94; 95% CI: 1.36–3.02,  $P = 0.044$ ), working hours (OR = 1.82; 95% CI: 1.53–2.16,  $P = 0.034$ ), and accommodation type (OR = 1.66; 95% CI: 1.38–2.49,  $P = 0.008$ ) were considered the strong predictors and significant contributing risk factors for health problems after adjusting for age, gender, and other variables.

## Discussion

In Arabian Gulf countries, Qatar is a top country which is recruited a great number of migrant workers. As of

**Table 3: Prevalence of medical conditions among studied participants (n=1186)**


Symptom	Semi-urban (n=564), n (%)	Urban (n=622), n (%)	P
<b>Pain</b>			
Back pain	223 (39.5)	176 (28.3)	<0.001
Joint pain	117 (20.7)	89 (14.3)	<0.001
Pain in arms and legs	125 (22.0)	95 (15.2)	<0.001
Headache	224 (39.7)	155 (24.9)	<0.001
<b>Cardiopulmonary</b>			
Chest pains	156 (27.7)	79 (12.7)	0.006
Shortness of breath	94 (16.7)	72 (11.6)	<0.001
Heart disease	156 (27.7)	140 (22.5)	0.044
Dizziness	110 (19.50)	87 (14.0)	0.012
<b>Gastrointestinal</b>			
Abdominal pain	135 (23.9)	113 (18.2)	0.015
Nausea	92 (16.3)	73 (11.7)	0.024
Gas or indigestion	88 (15.6)	70 (11.3)	0.019
Diarrhea	153 (27.1)	133 (21.4)	0.025
Irritability	59 (10.5)	38 (6.1)	0.008
Vomiting	102 (18.1)	97 (15.6)	0.276
Peptic ulcer	129 (22.9)	108 (17.4)	0.020
<b>Pseudoneurologic</b>			
Feeling tired	156 (27.6)	116 (18.6)	<0.001
Trouble sleeping	149 (26.5)	118 (18.9)	<0.001
Weakness/fainting	129 (22.8)	117 (18.8)	0.043
Blurred vision	141 (25.0)	143 (23.0)	0.323
Depression	126 (22.3)	106 (17.0)	0.023
Anxiety	74 (13.1)	55 (8.8)	0.020
Stress	124 (22.0)	99 (15.9)	0.009
Fatigue	136 (24.1)	104 (16.7)	0.002
<b>Medical condition</b>			
Diabetes	103 (18.3)	109 (17.5)	0.399
Hypertension	130 (23.0)	104 (16.7)	0.007
Asthma and allergy	77 (13.7)	59 (9.5)	0.028

**Table 4: Multivariable logistic regression analysis predictors for determinants of health problem (n=1186)**

Independent variables	OR	95% CI	P
Absence of drinking water	2.73	2.48-3.95	<0.001*
Inside tap water availability	2.69	2.24-3.58	0.003
Safety facility and tools absence	2.30	1.49-3.91	0.002
Occupation status	2.13	1.65-3.22	<0.001
Sheehsa smoking	1.96	1.27-3.14	0.031
Toilet facility	1.94	1.36-3.02	0.044
Working hours	1.82	1.53-2.16	0.034
Accommodation type	1.66	1.38-2.49	0.008

\*Two-sided P value based on -2 log-likelihood ratio statistics. Dependent variable=Health problem. Hosmer-Lemeshow test, Chi-square P=0.440 shows our model is quite a good fit. OR=Odds ratio, CI=Confidence interval

December 2013, it was estimated that 1,466,851 migrants were working in Qatar and over 1.2 million were semi-skilled or unskilled workers including construction.<sup>[19,20]</sup> Those



**Figure 1: Type of common occupational accidents**

increasing number of migrant workers and long working hours will lead an increase in the risk of occupational health conditions, occupational, and traffic accidents.<sup>[15,17,19]</sup> Migrant workers are also at risk from occupational medical condition and health problems such as pain, cardiopulmonary heart and respiratory diseases, GI abdominal, diarrhea, vomiting and peptic ulcer, pseudoneurologic-like fatigue, depression, anxiety and stress, and medical condition, i.e., diabetes, hypertension, and asthma and allergic diseases. Furthermore, it is reported that busy schedule and unsanitary lifestyle cause outbreaks of infectious diseases such as tuberculosis and pneumonia.<sup>[21]</sup> The suitable and timely access to health-care facilities may help to eliminate the high risk of health problems, particularly among low-income workers. The results of the current study are consistent with the Singapore study<sup>[22]</sup> and other neighboring Gulf Arab Countries' study.<sup>[23]</sup>

The risk of occupation and poor knowledge about health and safety rules lead to occupational-related health problems in Qatar. The present study found that there are many differences in terms of socioculture, religion, and education between the ethnic groups.<sup>[23]</sup> A greater part of the Qatari's workforce is at risk during the summer months where mostly working 10–12 h with over 45°C. About over 12,000 cases of heat-related illness were treated in health centers at the Hamad Medical Corporation, accident emergency departments (AED) during 2013.<sup>[24,25]</sup>

In fact, immigration and health care are controversial concerns politically, socially, economically, and ethically. Immigration has determined as a deep subject with more than 2 million migrant labors who are working and living in Qatar now. Over 80% of these labors have arrived from ISCM (Nepalese, Indian, Bangladeshi, Sri Lankan, and Pakistani) and Arab countries.<sup>[19,20,24,25]</sup> Many of those immigrants are not able to totally use the health-care facilities. They do not access to emergency services through hospital AED. Although over 8% of the national gross domestic product is spent on health care each year, uninsured labors are not utilize such services.<sup>[24,25]</sup>

## Conclusions

The migrant workers in Qatar have suffered from medical condition and health problems such as back pain, headache, cardiopulmonary as heart and respiratory diseases, GI abdominal, diarrhea, peptic ulcer, pseudoneurologic such as fatigue, depression, anxiety, stress, and diabetes, hypertension, and asthma and allergic diseases. Meanwhile, the most common occupational accident was falls, slip, and struck by object. Insurance for health services should be provided to all migrant workers during their working contracts. Almost 80% of migrant workers had no medical insurance.

## Acknowledgments

The author would like to thank the Hamad Medical Corporation for their support and ethical approval research project studies (HMC RC/36012/2012 and RP# 12025/12).

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

**Received:** 04 Dec 16 **Accepted:** 21 Apr 17

**Published:** 06 Sep 17

## References

- Bener A, Al Mazroei A. Health services management in Qatar. *Croat Med J* 2010;51:85-8.
- Bener A, Abdul Rahman YS, Abdel Aleem EY, Khalid MK. Trends and characteristics of injuries in the state of Qatar: Hospital-based study. *Int J Inj Contr Saf Promot* 2012;19:368-72.
- Joshi S, Simkhada P, Prescott GJ. Health problems of Nepalese migrants working in three Gulf countries. *BMC Int Health Hum Rights* 2011;11:3.
- WHO. Health of Migrants. Geneva: World Health Organisation; 2007. Available from: [http://www.apps.who.int/gb/ebwha/pdf\\_files/EB122/B122\\_11en.pdf](http://www.apps.who.int/gb/ebwha/pdf_files/EB122/B122_11en.pdf). [Last accessed on 2009 Feb 19].
- Weiner M. International Migration and Development: Indians in the Persian Gulf. *Population and Development Review* 1982;8:1-36.
- WHO. International Migration Health and Human Rights. Geneva: Health and Human Right Publication Series, World Health Organisation; 2003. p. 1-40. Available from: [http://www.who.int/hhr/activities/en/intl\\_migration\\_hhr.pdf](http://www.who.int/hhr/activities/en/intl_migration_hhr.pdf). [Last accessed on 2009 Feb 19].
- Lowell B, Kemper Y. Transatlantic roundtable on low-skilled migration in the twenty-first century. Prospects and policies. *Int Migr* 2004;42:118-40.
- Wolffers I, Verghis S, Marin M. Migration, human rights, and health. *Lancet* 2003;362:2019-20.
- Kronfol Z, Saleh M, Al-Ghafry M. Mental health issues among migrant workers in Gulf cooperation council countries: Literature review and case illustrations. *Asian J Psychiatr* 2014;10:109-13.
- IOM. Situation Report on International Migration in East and South-East Asia. Geneva: International Organisation for Migration; 2008. Available from: [http://www.publications.iom.int/system/files/pdf/situation\\_report.pdf](http://www.publications.iom.int/system/files/pdf/situation_report.pdf). [Last accessed on 2016 Apr 05].
- Szczepura A. Access to health care for ethnic minority populations. *Postgrad Med J* 2005;81:141-7.
- Bener A, Obineche E, Gillett M, Pasha MA, Bishawi B. Association between blood levels of lead, blood pressure and risk of diabetes and heart disease in workers. *Int Arch Occup Environ Health* 2001;74:375-8.
- Bhattarai R. Globalisation and flow of international labour migration. In Population Association of Nepal. *Nepal Population Journal* 2007;13:144-7.
- Kapiszewski A. Arab Versus Asian Migrant Workers in the GCC Countries' Paper Presented at the United Nations Expert Group Meeting on the International Migration and Development in the Arab Region, Beirut, May 15-17 2006, UN/POP/EGM/2006/02. United Nations: Dept. of Economic and Social Affairs; 2006.
- Black I. Modern-Day Slavery in Focus. Have Qatar's Work Conditions Improved? Migrant Laborers Tell Their Stories. *Guardian Newspaper*; Thursday 14 May, 2015 12.00 BST.
- Longva AN. Keeping migrant workers in check: The kafala system in the Gulf. *Middle East Rep* 1999;211:20-2.
- Weaver M. Modern-Day Slavery in Focus. Fifa to Investigate Arrest of BBC News Team in Qatar. *Guardian Newspaper*; Monday 18 May, 2015 11.30 BST.
- Gibson O. Modern-Day Slavery in Focus. Qatar: How Have Conditions for Migrant Workers Changed in Nine Key Areas? *Guardian Newspaper*; Monday 14 May, 2015 12.00 BST.
- Qatar Information Exchange. Labour Market, Qatar Statistics Authority; 2014. Available from: <http://www.qsa.gov.qa/>. [Last accessed on 2015 Aug 19].
- United Nations- Economic and Social Commission for Western Asia. The Demographic Profile of Qatar; 2011. Available from: <http://www.escwa.un.org/popin/members/qatar.pdf>. [Last accessed on 2015 Aug 15].
- Al-Khal AL, Bener A, Enarson DA. Tuberculosis among garment workers in an Arabian developing country: State of Qatar. *Arch Environ Occup Health* 2005;60:295-8.
- Lee W, Neo A, Tan S, Cook AR, Wong ML, Tan J, *et al.* Health-seeking behaviour of male foreign migrant workers living in a dormitory in Singapore. *BMC Health Serv Res* 2014;14:300.
- Loney T, Cooling RF, Aw TC. Lost in translation? Challenges and opportunities for raising health and safety awareness among a multinational workforce in the United Arab Emirates. *Saf Health Work* 2012;3:298-304.
- Annual Health Report Qatar, Department of Epidemiology and Medical Statistics, Doha, Qatar. Available from: <http://www.site.hmc.org.qa/msrc/ahr.htm>. [Last accessed on 2015 Aug 10].
- Bener A, Abdul Rahman YS, Abdel Aleem EY, Khalid MK. Trends and characteristics of head and neck injury from falls: A hospital based study, Qatar. *Sultan Qaboos Univ Med J* 2011;11:244-51.