

Poster presentation

Open Access

## **PI8-10. Ability of HIV antigens-pulsed monocyte-derived dendritic cells to induce HIV-specific T cell response: potential use in therapeutic vaccine**

GG Silveira\*, TM Oshiro, A Almeida, C Finazzo, RM Oliveira, AJ Duarte and G Benard

Address: Laboratory of Investigation in Dermatology and Immunodeficiencies, School of Medicine of the University of São Paulo, São Paulo, Brazil

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P319 doi:10.1186/1742-4690-6-S3-P319

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P319>

© 2009 Silveira et al; licensee BioMed Central Ltd.

### **Background**

Despite 20 years of effort, the design of an effective HIV-1 vaccine remains an enormous challenge. In this scenario, new immunological approaches must be considered. Dendritic cells-based vaccines have been widely used in cancer therapy showing interesting successful results. More recently, Lu et al. (2004) described large reductions on virus load in untreated HIV-infected patients immunized with autologous DC pulsed with inactivated viruses. Although encouraging, these results showed sustained virus reduction in only half of vaccinees. Besides, isolating virus from each patient is a rather laborious and expensive process. Based on that monocyte-derived dendritic cell (Mo-DC) vaccine model we evaluated the use of alternative HIV products in order to simplify the process of vaccine production, reduce costs and achieve protective immune responses on a greater number of individuals.

### **Methods**

Monocytes from HIV-serodiscordant couples were differentiated *in vitro* into dendritic cells, pulsed with a pool of aldrithiol-2-inactivated HIV-1 subtypes or the HIV-1IIIIB p55Gag protein and then cultured with autologous lymphocytes for 7 days. At day 7, the culture received a boost of fresh Mo-DCs pulsed with the same antigens. The T lymphocyte immunological response was determined by proliferation assay, IFN $\gamma$  production and cellular activation. The Mo-DC phenotype was also evaluated.

### **Results**

Mo-DCs were shown to be fully matured and activated (CD11c<sup>+</sup>HLA-DR<sup>hi</sup>CD86<sup>hi</sup>CD80<sup>hi</sup>CD83<sup>+</sup>) in both antigen-pulsing protocols. IFN $\gamma$  production by T CD4<sup>+</sup> and CD8<sup>+</sup> lymphocytes, as well as the percentage of CD38<sup>+</sup> cells, was always higher when stimulated with pulsed Mo-DCs compared to non-pulsed cells. The T lymphocyte proliferation of pulsed Mo-DCs was also significant greater compared to the non-pulsed condition, being slightly but not significantly higher in p55Gag-pulsed cells.

### **Conclusion**

This easier and cheaper Mo-DC vaccine model elicited *in vitro* detectable cellular immune responses in HIV-uninfected subjects, which suggests a broader range of action and represents a viable and promising alternative of therapeutic vaccination.