

A case of “pseudo science”? A study claiming effects of the Cry1Ab protein on larvae of the two-spotted ladybird is reminiscent of the case of the green lacewing

Stefan Rauschen

Published online: 30 June 2009
© Springer Science+Business Media B.V. 2009

A recent report on the potential negative impact in a laboratory setting of the Cry1Ab protein on larvae of the two spotted ladybird *Adalia bipunctata* (Schmidt et al. 2009) has gained notoriety. It was used in Germany, along with some other studies supposedly showing a negative impact of the transgenic MON810 maize on non-target organisms, to temporarily ban the cultivation of this Bt-maize under a safeguard clause conforming with Article 23 of the EU directive 2001/18/EC. This decision, although officially communicated as based on new evidence, was in fact based on flawed science and has been recognized to be politically motivated by a number of the stakeholders involved (Sinha 2009). The present temporary ban of MON810 by the German Government will now be considered by the EU commission, which will consult the European Food Safety Authority (EFSA), and then decide on the merits of the scientific data. Whether the ban will then be lifted or not lies within the committological decision making process within the European Union. The data on *A. bipunctata* was first published in the Proceedings of the German Society for General and Applied Entomology half a decade ago (Schmidt et al. 2004). Thus, it does not even constitute “new evidence”.

In their experiment, Schmidt et al. fed *A. bipunctata* larvae with eggs of the flour moth *Ephestia kuehniella*

that had been sprayed with solutions of Cry1Ab and Cry3Bb of different concentrations. Mortality in the treatments was higher than in the control groups. Larval developmental time and adult biomass were also measured, but there were no differences between the experimental groups. The authors concluded that the Cry proteins, and especially Cry1Ab, had a negative impact on *A. bipunctata* larvae, based on some unknown mode of action.

There are some obvious methodological flaws and inconsistencies in the Schmidt et al. (2009) paper, however:

Firstly, the quantity of stock solutions applied and the actual quantity of test proteins taken up by the ladybird larvae is not reported. It is only written that the eggs of the flour moth *E. kuehniella* used as food were “sprayed” with the different Cry protein dilutions. There is also no information given on the food consumption by the larvae during their development. The authors thus did not know, at least they did not state, the actual dose that the tested individuals took up during the experiment. Thus, the first important criterion defining risk, i.e., the degree of exposure, was not properly quantified.

Secondly, the mortality rates in identical control groups for the three separate treatments varied greatly between 7.5 and 20.8% for the 1st larval instars. These differences are neither explained nor addressed in the statistical analysis and the interpretation of the data, but strongly suggest methodological problems.

S. Rauschen (✉)
Department of Plant Physiology, RWTH Aachen
University, Worringerweg 1, 52074 Aachen, Germany
e-mail: Rauschen@Bio3.RWTH-Aachen.de

Thirdly, the treatment groups with the highest Bt-protein concentrations did not show an increase in mortality rates over the groups treated with lower concentrations. This was the case for both Cry1Ab and Cry3Bb, and is a very astonishing result that contradicts classical dose-response relationship models. It is also interesting to note in this context that the Coleopteran-specific Cry3Bb protein had a smaller effect than the Lepidopteran-specific Cry1Ab, whereas one might expect the opposite to be the case for this test organism.

Fourthly, none of the treatments had any impact on either larval developmental time, or on the body-weight of adult beetles. This means that the larvae either died, or seemed to be totally unaffected by the treatments with Cry proteins. This is also surprising, since it is well established that susceptible organisms suffer from sub-lethal effects long before direct toxic effects (i.e. mortality) can be observed (e.g. Sears et al. 2001).

Schmidt et al. also totally neglect relevant recent literature. For example, Bai et al. (2005) reported the absence of any effects of rice pollen containing Cry1Ab on *Propylea japonica*. Similarly, a study by Álvarez-Alfageme et al. (2008) did not report any effects of Cry1Ab on another coccinellid, *Stethorus punctillum*. The discussion in the paper by Schmidt et al. would also have benefited from published results from a large number of field studies with Bt-maize reporting on a variety of beetle families encompassing Coccinellidae, Carabidae and Lathridiidae (de la Poza et al. 2005; Ahmad et al. 2006; Eckert et al. 2006; Toschki et al. 2006; Floate et al. 2007; Leslie et al. 2007). A recent report by Dhillon and Sharma (2009) that did show effects of Cry1Ab on larvae of *Cheilomenes sexmaculatus* in a laboratory setting, but using an exposure regime more relevant to the field situation, showed that “direct exposure to Bt toxins expressed in transgenic plants [...] will have little effect on the activity and abundance of the ladybird”. The claims and interpretations by Schmidt et al. need to be put into the context of the vast body of evidence showing that genetically modified plants expressing Cry1Ab do not have a negative impact on ladybirds.

An absence of ecological considerations in the interpretation of data is another striking flaw of the Schmidt et al. paper:

Adalia bipunctata are aphid predators and it is unclear to what extent these beetles would take up Bt-proteins in the field. For it is well documented that Bt-maize varieties do not carry Cry proteins in their phloem sap, and that aphid predators are unlikely to be exposed to Bt-proteins via their prey (Head et al. 2001; Raps et al. 2001; Dutton et al. 2002; Lundgren and Wiedenmann 2005). Therefore, exposure will only occur if these beetles consume maize pollen (see Cottrell and Yeargan 1998; Lundgren et al. 2005). This consumption of plant materials in comparison to that of aphid prey needs to be quantified in the field to assess the potential risk. Schmidt et al. touch these issues only superficially, without reaching reasonable conclusions on the relevance of their own data. In this instance, it is important to note that the expression level for Cry1Ab in pollen from MON810 (the only Bt-maize cultivated in the European Union) is extremely low, ranging to a maximum of 97 ng Cry1Ab/g fresh weight (Nguyen and Jehle 2007). Interestingly, Schmidt et al. give false expression levels of 7.93–10.34 µg Cry1Ab/g fresh weight (which is the expression level in leaves according to their source, AGBIOS), that exaggerate the potential exposure in the field and thus give a false impression on the relevance of this exposure pathway.

Adalia bipunctata cannibalise their own eggs and suffer from intra-guild predation by other Coccinellids (Schellhorn and Andow 1999; Burgio et al. 2002). The authors fail to give a perspective on the quantitative relevance of a potential additional mortality based on the effects of Bt-proteins in relation to these two alternative common mortality factors.

Moreover, *A. bipunctata* is in fact not a very common ladybird in maize, at least not in Germany (Rauschen et al. 2009). The significance of a potential negative impact on the limited populations occurring in maize cropping systems for the population as a whole on a larger regional scale remains doubtful.

Overall, the findings and interpretations in Schmidt et al. and the consideration of this paper for the justification of the ban of MON810 in Germany appear erroneous.

In the past, similar results on potential negative impacts of Cry1Ab expressing Bt-maize varieties on

another beneficial insect, the green lacewing *Chrysoperla carnea*, had been published. After about 10 years of research, we now recognize that these early claims were flawed, and that effects measured in the original investigations were based on poor prey quality, and not on direct toxic effects of Cry1Ab (Shelton et al. 2009; for a detailed account see Romeis et al. 2009). The green lacewing case has been very prevalent in discussions concerning the potential risks of Bt-plants for the environment and is still often cited by non-governmental organisations and certain ill-informed regulatory bodies, although scientifically has been thoroughly refuted. Sadly, it appears that results from the flawed experiments utilizing the two-spotted ladybird might be used to justify an indefensible political position and misinform the lay public. In the overall context of non-target risk assessment (Romeis et al. 2008), the Schmidt et al. paper at best indicates a potential hazard of an insecticidal compound to larvae of *A. bipunctata* under certain artificial conditions. The next step must be to assess the relative importance of different exposure pathways, and to assess whether the effects are still visible under more realistic exposure regimes. If they do, it remains to be assessed whether these effects have wider implications for the exposed population of this species and for its function as a biological control agent. Only if these questions are thoroughly and reasonably addressed, in a scientifically sound manner, can reliable conclusions on real potential risks be drawn. And only then can political decision makers reach judgments that are not clouded by the results of ill-conceived and shoddy research.

References

- Ahmad A, Wilde GE, Whitworth RJ, Zolnerowich G (2006) Effect of corn hybrids expressing the coleopteran-specific Cry3Bb1 protein for corn rootworm control on above-ground insect predators. *J Econ Entomol* 99:1085–1095
- Álvarez-Alfageme F, Ferry N, Castañera P, Ortego F, Gatehouse AMR (2008) Prey mediated effects of Bt maize on fitness and digestive physiology of the red spider mite predator *Stethorus punctillum* Weise (Coleoptera: Coccinellidae). *Transgenic Res* 17:943–954
- Bai YY, Jiang MX, Cheng JA (2005) Effects of transgenic rice pollen on fitness of *Propylea japonica*. *J Pest Sci* 78: 123–128
- Burgio G, Santi F, Maini S (2002) On intra-guild predation and cannibalism in *Harmonia axyridis* (Pallas) and *Adalia bipunctata* L. (Coleoptera: Coccinellidae). *Biol Control* 24(2):110–116
- Cottrell TE, Yeargan KV (1998) Effect of pollen on *Coleomegilla maculata* (Coleoptera : Coccinellidae) population density, predation, and cannibalism in sweet corn. *Environ Entomol* 27:1402–1410
- de la Poza M, Pons X, Farinós GP, López C, Ortego F, Eizaguirre M, Castañera P, Albajes R (2005) Impact of farm-scale Bt maize on abundance of predatory arthropods in Spain. *Crop Prot* 24:677–684
- Dhillon MK, Sharma HC (2009) Effects of *Bacillus thuringiensis*-endotoxins Cry1Ab and Cry1Ac on the coccinellid beetle, *Cheilomenes sexmaculatus* (Coleoptera, Coccinellidae) under direct and indirect exposure conditions. *Biocontrol Sci Techn* 19:407–420
- Dutton A, Klein H, Romeis J, Bigler F (2002) Uptake of Bt-toxin by herbivores feeding on transgenic maize and consequences for the predator *Chrysoperla carnea*. *Ecol Entomol* 27:441–447
- Eckert J, Schuphan I, Hothorn LA, Gathmann A (2006) Arthropods on maize ears for detecting impacts of Bt maize on nontarget organisms. *Environ Entomol* 35: 554–560
- Floate KD, Cárcamo HA, Blackshaw RE, Postman B, Bourassa S (2007) Response of ground beetle (Coleoptera: Carabidae) field populations to four years of lepidoptera-specific Bt corn production. *Environ Entomol* 36:1269–1274
- Head G, Brown CR, Groth ME, Duan JJ (2001) Cry1Ab protein levels in phytophagous insects feeding on transgenic corn: implications for secondary exposure risk assessment. *Entomol Exp Appl* 99:37–45
- Leslie TW, Hoheisel GA, Biddinger DJ, Rohr JR, Fleischer SJ (2007) Transgenes sustain epigeal insect biodiversity in diversified vegetable farm systems. *Environ Entomol* 36:234–244
- Lundgren JG, Wiedenmann RN (2005) Tritrophic interactions among Bt (Cry3Bb1) corn, aphid prey, and the predator *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Environ Entomol* 34:1621–1625
- Lundgren JG, Huber A, Wiedenmann RN (2005) Quantification of consumption of corn pollen by the predator *Coleomegilla maculata* (Coleoptera: Coccinellidae) during anthesis in an Illinois cornfield. *Agric Forest Entomol* 7:53–60
- Nguyen HT, Jehle JA (2007) Quantitative analysis of the seasonal and tissue-specific expression of Cry1Ab in transgenic maize MON810. *J Plant Dis* 114:82–87
- Raps A, Kehr J, Gugerli P, Moar WJ, Bigler F, Hilbeck A (2001) Immunological analysis of phloem sap of *Bacillus thuringiensis* corn and of the nontarget herbivore *Rhopalosiphum padi* (Homoptera: Aphididae) for the presence of Cry1Ab. *Mol Ecol* 10:525–533
- Rauschen S, Schaarschmidt F, Gathmann A (2009) Occurrence and field densities of Coccinellidae in the maize herb layer: implications for Environmental Risk Assessment. *IOBC/WPRS Bulletin* (in press)
- Romeis J, Bartsch D, Bigler F, Candolfi MP, Gielkens MMC, Hartley SE, Hellmich RL, Huesing JE, Jepson PC, Layton R, Quemada H, Raybould A, Rose RI, Schiemann J, Sears MK, Shelton AM, Sweet J, Vaituzis Z, Wolt JD (2008)

- Assessment of risks of insect-resistant transgenic crops to nontarget arthropods. *Nat Biotechnol* 26:203–208
- Romeis J, Meissle M, Raybould A, Hellmich RL (2009) Impact of insect-resistant transgenic crops on above-ground nontarget arthropods. In: Ferry N, Gatehouse AMR (eds) *Environmental impact of genetically modified crops*. CABI, Wallingford, pp 165–198
- Schellhorn NA, Andow DA (1999) Cannibalism and interspecific predation: role of oviposition behavior. *Ecol Appl* 9(2):418–428
- Schmidt JEU, Braun CU, L'Abate C, Whitehouse LP, Hilbeck A (2004) Untersuchungen zu Effekten von *Bacillus thuringiensis*-Toxinen aus transgenen insektenresistenten Pflanzen auf räuberische Marienkäfer (Coleoptera: Coccinellidae). *Mitt Dtsch Ges Allg Angew Ent* 14:419–422 (Abstract in English)
- Schmidt JEU, Braun CU, Whitehouse LP, Hilbeck A (2009) Effects of activated Bt transgene products (Cry1Ab, Cry3Bb) on immature stages of the ladybird *Adalia bipunctata* in laboratory ecotoxicity testing. *Arch Environ Contam Toxicol* 56(2):221–228
- Sears MK, Hellmich RL, Stanley-Horn DE, Oberhauser KS, Pleasants JM, Mattila HR, Siegfried BD, Dively GP (2001) Impact of Bt corn pollen on monarch butterfly populations: a risk assessment. *PNAS* 98(21):11937–11942
- Shelton AM, Naranjo SE, Romeis J, Hellmich RL, Wolt JD, Federici BA, Albajes R, Bigler F, Burgess EPJ, Dively GP, Gatehouse AMR, Malone LA, Roush R, Sears M, Sehnal F (2009) Setting the record straight: a rebuttal to an erroneous analysis on transgenic insecticidal crops and natural enemies. *Transgenic Res* 18:317–322
- Sinha G (2009) Up in arms. *Nat Biotechnol* 27(7) (in press)
- Toschki T, Hothorn LA, Roß-Nickoll M (2006) Effects of cultivation of genetically modified Bt maize on epigeic arthropods (Araneae; Carabidae). *Environ Entomol* 36: 967–981