


# Cost Determinants of Continuum-Care Episodes for Hip Fracture

Davide Golinelli<sup>1</sup> , Erik Boetto<sup>2</sup>, Antonio Mazzotti<sup>3</sup>, Simona Rosa<sup>1</sup>, Paola Rucci<sup>1</sup>, Elena Berti<sup>4</sup>, Cristina Ugolini<sup>5</sup> and Maria Pia Fantini<sup>1</sup>

<sup>1</sup>Department of Biomedical and Neuromotor Sciences (DIBINEM), Alma Mater Studiorum–University of Bologna, Italy. <sup>2</sup>School of Hygiene and Preventive Medicine, Alma Mater Studiorum–University of Bologna, Italy. <sup>3</sup>1<sup>st</sup> Orthopedic and Traumatologic Clinic, IRCCS–Istituto Ortopedico Rizzoli, Bologna, Italy. <sup>4</sup>Regional Agency for Health and Social Care, Emilia-Romagna Region - ASSR, Bologna, Italy. <sup>5</sup>Department of Economics and CRIFSP-School of Advanced Studies in Health Policy, University of Bologna, Italy.

Health Services Insights  
Volume 14: 1–11  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1178632921991122



**ABSTRACT:** Many factors affect the healthcare costs and outcomes in patients with hip fracture (HF). Through the construction of a Continuum-Care Episode (CCE), we investigated the costs of CCEs for HF and their determinants. We used data extracted from administrative databases of 5094 consecutive elderly patients hospitalized in 2017 in Emilia Romagna, Italy, to evaluate the overall costs of the CCE. We calculated the acute and post-acute costs from the date of the hospital admission to the end of the CCE. The determinants of costs by type of surgical intervention (total hip replacement, partial hip replacement, open reduction, and internal fixation) were investigated using generalized linear regression models. Regardless of the type of surgical intervention, hospital bed-based rehabilitation in public or private healthcare facilities either followed by rehabilitation in a community hospital/temporary nursing home beds or not were the strongest determinants of costs, while rehabilitation in intermediate care facilities alone was associated with lower costs. CCE's cost and its variability is mainly related to the rehabilitation setting. Cost-wise, intermediate care resulted to be an appropriate setting for providing post-acute rehabilitation for HF, representing the one associated with lower overall costs. Intermediate care organizational setting should be privileged when planning integrated care HF pathways.

**KEYWORDS:** Hip fractures, healthcare costs, episode of care, rehabilitation, intermediate care facilities

**RECEIVED:** October 11, 2020. **ACCEPTED:** January 8, 2021.

**TYPE:** Original Research

**FUNDING:** The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The project is the result of the partnership "Economic evaluation of clinical pathways and prescribing appropriateness" between the School of Health Policies-University of Bologna and the Regional Agency for Health and Social Care of the Emilia-Romagna Region (ASSR-RER), with the cooperation of the Local Health Authority of Parma. The 3 institutions have jointly concurred to the

project with their own resources with the ASSR-RER assuring the financial resources necessary for research assistant contracts.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Davide Golinelli, Department of Biomedical and Neuromotor Sciences (DIBINEM), Alma Mater Studiorum–University of Bologna, Via San Giacomo 12, Bologna 40126, Italy. Email: [davide.golinelli@unibo.it](mailto:davide.golinelli@unibo.it)

## Introduction

Hip fracture (HF) is a common event in the elderly that causes a growing burden on healthcare systems worldwide and requires an intensive healthcare resources utilization, especially in the first year after the fracture.<sup>1–3</sup> From an epidemiological point of view, more than 300 000 hospitalizations for HF occur annually in the United States, with a 1 year mortality of almost 25%.<sup>4</sup> In the United Kingdom, HF produces £ 1.1 billion of hospital costs per year, which are estimated to increase to £1.5 billion by 2025, when 104 000 annual cases are expected.<sup>5</sup> Assuming that the age-related incidence will increase by just 1% per year, the number of HF in the world will reach approximately 8.2 million in 2050.<sup>6</sup> HF also represents an important public health concern in Italy: in 2016, it accounted for 120 845 hospitalizations and 92 624 surgical procedures, with an annual incidence of 189.5 per 100 000 in men and of 498.4 per 100 000 in women  $\geq 50$  years old<sup>7,8</sup> and with relevant functional impairments in the individuals who experience it, especially the elderly.<sup>9–11</sup>

Data on HF in Italy reflect its aging population. In Emilia Romagna, one of the largest administrative regions of North-Eastern Italy (4.4 million inhabitants), during 2011 6368

patients aged  $\geq 65$  years were treated in regional hospitals for HF (equal to 142.8 cases per 100 000) with a clear predominance of females (75.3% of cases) and with a median age of 84 years in both sexes.<sup>12</sup>

One of the most problematic expression of population aging is frailty,<sup>13</sup> which plays an important role in reduced functional recovery and overall health status decline.<sup>4,14–16</sup> HF in frail elderly are associated with multiple comorbidities,<sup>17–19</sup> and several studies indicate poor functional outcomes leading to decreased mobility, less self-dependence in daily living, decreased life quality, and increased dependency and mortality rates.<sup>20–22</sup> To reduce complications and improve functional outcomes, international guidelines recommend to perform surgery within 48 hours from hospital admission, ensure early mobilization the day after the procedure and provide a post-acute rehabilitation plan involving a multidisciplinary team in different healthcare settings.<sup>3,23–25</sup>

The debate is open on what the most appropriate integrated healthcare pathways for HF patients are. In order to define the most effective and efficient rehabilitation pathway, a Cochrane systematic review pointed to the need to evaluate all components of the process of acute care and rehabilitation together



rather than separately.<sup>26</sup> Nonetheless, there is currently no definite framework that specifies the appropriate healthcare pathway after HF.

Among the possible healthcare settings addressing HF patients' needs, attention was drawn to intermediate care.<sup>27</sup> Intermediate care has been developed since the late 1990s, to foster the integration between acute and primary care in several European countries.<sup>28</sup> Intermediate care has both "preventive" aims (avoiding unnecessary hospitalizations and "delayed discharges") and "rehabilitation" aims (supporting discharge and access to rehabilitation services to enforce care close to home).<sup>29</sup> In fact, it is essential to ensure continuity of care by developing services and healthcare pathways that integrate hospital and primary care services.<sup>30</sup> Accordingly, intermediate care has been introduced in countries with a national and universalistic healthcare service, such as the UK and Italy, in order to provide an integrated multi-professional healthcare pathway for patients with complex/chronic needs who can be treated outside the hospital setting.<sup>31-33</sup>

To evaluate that, Sheehan et al.<sup>34,35</sup> designed a conceptual framework for an hip fracture integrated episode of care, defined as Continuum-Care Episode (CCE), that encompasses the hospital admission for the surgical procedure for HF and the following bed-based rehabilitation phase performed in different organizational settings and suggested its usefulness also for comparative quality and costs analyses.

Pitzul et al.<sup>36</sup> described HF patient characteristics and the most common post-acute pathways within a 1-year episode of care in Ontario, Canada, and found that similar HF patients were discharged to different post-acute settings, calling into question both the appropriateness of care delivered in the post-acute period and health expenditures. Braithwaite et al.<sup>4</sup> pointed out that almost half of the costs attributable to HF are mainly related to the costs of rehabilitation. Nonetheless, many clinical, social, and organizational factors may affect the costs and outcomes of episodes of care in patients with HF. It seems therefore of interest to investigate whether rehabilitation setting and other organizational factors may determine the variability of costs in the CCE of patients with HF.<sup>34</sup>

The aim of this study is to investigate the costs of CCEs for hip fracture and to identify their determinants in Emilia-Romagna region, Italy.

## Methods

### *Study setting*

In Italy, the National Health System (NHS) is responsible for providing "essential levels of care" to the population, through autonomous planning and delivery of healthcare services in the 21 Italian Regions.

Emilia-Romagna Regional Health System includes 8 Local Health Trusts (LHT), 4 University Hospitals, 1 General Hospital Trust, and 4 Research Hospitals. In the last decade,

Emilia-Romagna Region has improved the management of patients with HF, issuing policies and guidelines aimed at reducing the delay of surgery and designing specific modalities for postoperative rehabilitation.<sup>12</sup> Each of the 8 LHTs is responsible for adopting regional policies and guidelines and defining predetermined healthcare pathways (or CCEs) to patients with HF. However, each LHTs is autonomous in defining specific CCEs, relying on the characteristics of its territory and on the offer of available services. In particular, a determinant of the variability in CCEs for HF patients is the availability of intermediate care facilities, such as nursing homes and community hospitals (CH). CHs are bed-based intermediate care services that may be defined as "local hospitals," staffed mainly by general practitioners (GPs), nurses, and also rehabilitation professionals (such as physiatrists and physical therapists).


### *Data and study population*

We conducted a retrospective observational cohort study using administrative databases of Emilia-Romagna Region that were linked and anonymized for the analysis, as reported in Supplemental Table 1 (Supplemental Files).

The study population consisted of consecutive patients hospitalized for HF over the year 2017, aged  $\geq 65$  years and resident in Emilia Romagna. The inclusion and exclusion criteria were those adopted by the Italian National Outcomes Program<sup>7</sup> in protocol n.42 ("Surgery within 2 days after HF in the elderly"). Accordingly, all patients hospitalized with a primary or secondary diagnosis of HF (ICD-9-CM codes 820.0-820.9), in Emilia Romagna from January 1, 2017 to December 31, 2017, were included in the study with the following exceptions: admissions preceded by hospitalization with a diagnosis of HF in the previous 2 years; hospitalizations of patients aged under 65 and over 100; hospitalizations of patients not resident in Italy; patients hospitalized after transfer from another facility; hospitalizations of polytraumatized patients (Diagnosis Related Groups—DRG 484-487); patients who died within 2 days without any surgical intervention (difference between the date of death and the date of entry to the hospital equal to 0–1 days); admissions with a primary or secondary diagnosis of malignant tumor (codes ICD-9-CM 140.0-208.9) in the index hospitalization or in the 2 previous years.

### *Continuum-care episode*

According to Sheehan et al.,<sup>34,35</sup> definition the CCE represents an integrated episode of care consisting of an acute hospital admission phase for HF (early surgery, multidimensional evaluation, and early mobilization) and a post-acute phase (rehabilitation in appropriate settings and follow-up), ending with a specific event (eg, death of patient, long-term hospitalization in elderly care facility, or hospital discharge at home) or the end date (6 months) of the time window (Figure 1).<sup>34,35</sup>



**Figure 1.** Continuum-care episode for hip fracture, including acute hospital admission and post-acute rehabilitation. Types of post-acute rehabilitation: (1) Type 1, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital); (2) Type 2, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital), followed by rehabilitation in intermediate care facilities (community hospital [CH], or temporary nursing home beds); (3) Type 3, rehabilitation in intermediate care facilities (CH, or temporary nursing home beds).

Therefore, the CCE is a framework describing an integrated healthcare pathways that include the acute hospitalization phase for the surgical intervention for HF (coded using the Diagnosis Related Groups—DRGs—related to HF) and continues with 3 different types of post-acute rehabilitation: (1) *Type 1*, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital); (2) *Type 2*, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital), followed by rehabilitation in intermediate care facilities (community hospital [CH], or temporary nursing home beds); and (3) *Type 3*, Rehabilitation in intermediate care facilities (CH, or temporary nursing home beds).

The CCE (Figure 1) termination rules are met when the patient: (a) is deceased; (b) is hospitalized in a long-term elderly care facility; (c) has no other hospitalizations within 2 days from the last discharge date. The CCE (Figure 1) summarizes the healthcare and rehabilitation pathways that patients with HF can face. The CCE represents the conceptual framework that allows to describe the variability in terms of services provisions and costs in Emilia Romagna. Each patient was followed up for 6 months after the first hospital admission for HF.

### Cost analysis

The study population for the cost analysis included only patients undergoing surgery. To evaluate the overall costs of the CCE, we calculated the direct medical costs incurred by the regional healthcare system from the date of the first hospital admission to the end of the CCE, as illustrated in Figure 1.

The healthcare provisions included in the CCE costs account for the acute hospitalization (ie, DRGs) plus other

costs related to: post-acute re-admissions (public or private hospital) in Orthopedics and Traumatology beds (Italian NHS bed code 036), Recovery and Rehabilitation beds (code 056), Long-Term Care beds (code 060), admissions to CHs, temporary admissions to nursing homes, orthopedic check-ups, and outpatient rehabilitation (code R20).

We have also calculated the costs from 3 up to 180 days after the end of CCE, in order to describe the costs of possible early rehospitalizations of patients within 6 months.

Cost data for patients who died before the end of the CCE included all costs incurred by the healthcare services until the death date.

### Analysis strategy

The socio-demographic and clinical characteristics of patients and costs were summarized using descriptive statistics such as mean and standard deviation for continuous variables, and median and interquartile range for continuous asymmetric variables.

All the descriptive statistics are provided for the overall study population and by type of surgery. This was done because the type of surgical intervention is mainly influenced by the surgeons' decision. Following the regional guidelines for surgery after HF,<sup>12</sup> the decision is based on the anatomic-pathological characteristics of the fracture and on patient age and clinical conditions. This defines 4 different sub-categories of population, namely: (1) no intervention, (2) total hip replacement (THR), (3) hemiarthroplasty or partial hip replacement (PHR), and (4) open reduction and internal fixation (ORIF).

The determinants of costs for patients undergoing ORIF, PHR, and THR surgery were investigated using generalized

**Table 1.** Characteristics of patients by type of intervention.

	NO INTERVENTION N=257	TOTAL HIP ARTHROPLASTY N=531	PARTIAL HIP ARTHROPLASTY N=1793	OPEN REDUCTION AND INTERNAL FIXATION N=3323	TOTAL N=5904
Mean age $\pm$ SD, y	83.7 $\pm$ 7.9	75.4 $\pm$ 5.9	85.9 $\pm$ 6.0	84.4 $\pm$ 7.4	84.0 $\pm$ 7.5
Sex, n (col %)					
Female	174 (67.7)	390 (73.4)	1275 (71.1)	2617 (78.8)	4456 (75.5)
Male	83 (32.3)	141 (26.6)	518 (28.9)	706 (21.2)	1448 (24.5)
Elixhauser (classes), n (col %)					
0 morbidities	112 (43.6)	363 (68.4)	1009 (56.3)	1878 (56.5)	3362 (56.9)
1–2 morbidities	75 (29.2)	133 (25.0)	499 (27.8)	923 (27.8)	1630 (27.6)
3+ morbidities	70 (27.2)	35 (6.6)	285 (15.9)	522 (15.7)	912 (15.4)
Total	257 (4.4)	531 (9.0)	1793 (30.4)	3323 (56.3)	5904 (100)

Abbreviations: SD, standard deviation; col, column.

linear multiple regression models (GLM) with a gamma probability distribution and log link, to take into account the skewness of the cost distribution. The dependent variables were the costs of CCE, and the predictors of costs were chosen according to the literature and included: gender, age, comorbidities (Elixhauser index<sup>37</sup>), timing of the surgery (over 48 hours vs within 48 hours), an active home care program before the hospital admission (yes/no), residing in nursing home before hospital admission (yes/no), hospital length of stay (below or above the DRG threshold days), rehabilitation type (Type 1, Type 2, Type 3, vs no rehabilitation), length of rehabilitation (from 1 to 21 days vs more than 22 days),<sup>38</sup> and Local Health Trust (7 dummy variables for each LHT vs the LHT of the capital town of Emilia Romagna).

Predictors of costs included “non-modifiable” variables (age, gender, multimorbidity, and long-term HA or residing in a nursing home before hospital admission for HF), “modifiable” variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and the LHT of residence.

This selection was made in order to account for well-known predictors of costs, such as age and number of comorbidities,<sup>10,13,15,21</sup> and other organizational factors that can mitigate the effect of hip fracture on patient outcomes and costs, such as timing of surgery or length or setting of rehabilitation.<sup>18,19,23,24</sup> The impact of modifiable factors was tested after adjusting for “non-modifiable” factors.

The goodness of fit of the models was assessed using the following indices: the Akaike information criterion (AIC) and Bayesian information criterion (BIC). These indices have no predefined cut-offs and can only be interpreted when comparing 2 different models. Lower indices denote a better model fit.

*Ethics approval.* The study has been approved by the AVEC (Comitato Etico Indipendente di Area Vasta Emilia Centro) research ethics committee board, on April 17, 2019. Data used in this research was obtained from the Regional Healthcare Information System which includes detailed information on the use of healthcare services by all regional patients, with the patient as our unit of observation. The study, based on routine administrative information, was carried out in conformity with the regulations on data management of the Emilia-Romagna Region and with Italian privacy law. Administrative data were anonymized by an ad hoc service of the Regional Department of Health and the University of Bologna have no possibility to retrospectively identify individuals included in the study.

## Results

The study population included 5094 patients aged  $\geq 65$  years with a hospital admission for HF in 2017. Table 1 shows the characteristics of the study population, by type of surgical intervention: no intervention (n=257), THR (n=531), PHR (n=1793), and ORIF (n=3323).

Patients with THR were on average 10 years younger (75.4  $\pm$  7.9 years) than those with PHR (85.9  $\pm$  6.0), ORIF (84.4  $\pm$  7.4), and no intervention (83.7  $\pm$  7.9). Female patients were predominant in all types of surgery (from 71.1% in THR to 78.8% in ORIF). Patients with comorbidities were 68.4% of those with THR, 56.3% and 56.5% respectively of those with PHR or ORIF and only 43.6% of those not undergoing any surgery. Patients with 3 or more comorbidities were less frequent among those undergoing THR (6.6%), while representing a large proportion of patients without any surgical intervention (27.2%). Fifteen percent died over 6 months from HF (31.5% among those without an intervention, 1.9% THR, 17.1% PHR, 14.6% ORIF).



Table 2 shows the empirical costs of the CCEs in € currency by type of surgery. The costs were broken down into the 2 components of the CCE (acute hospitalization + bed-based rehabilitation) and are provided for the total CCE (see also Figure 1). The median cost for the acute phase of patients undergoing THR and PHR intervention (531 and 1793 patients, respectively) was the same (€ 10079.43), corresponding to the DRG reimbursement, whereas the median costs for bed-based rehabilitation are slightly different in patients undergoing THR and PHR, respectively € 3234.00 and € 3198.74. While the costs for the acute phase are lower in patients undergoing ORIF (€ 6526.13), the bed-based rehabilitation phase has a higher median cost (€ 3622.08) compared with patients with THR and PHR.

Costs of HF patients at 180 days after end of CCE resulted in 4.3% incremental costs (Table 2).

Tables 3 to 5 show the results of 3 GLM models predicting the costs incurred by the regional healthcare system for patients with HF undergoing ORIF, THR, and PHR surgery in 2017, respectively. The **first model (model A)** includes “non-modifiable” variables such as age, gender, multimorbidity, and long-term HA or residing in a nursing home before hospital admission for HF. The **second model (model B)** included, in addition to model A variables, the “modifiable” variables that depend on healthcare organizational models, that is, timing of surgery, the rehabilitation type, the hospital length of stay, the length of rehabilitation, and the LHT of residence.

In model A, including 3323 patients undergoing ORIF (Table 3), the number of comorbidities was associated with increased costs ( $b=0.026, P<.001$ ), while age and gender were unrelated with costs. Receiving home care or residing in nursing home before the hospitalization were associated with lower costs ( $b=-0.096, P=.040$  and  $b=-0.179, P<.001$ , respectively). The AIC and the BIC for this model were respectively 62580 and 62623. After adjusting for “non-modifiable” variables (model B, Table 3), the ORIF surgical intervention after 48 hours from the hospitalization was unrelated with costs ( $b=-0.004, P=.650$ ). Type 2 rehabilitation (hospital bed-based rehabilitation in public or private healthcare facilities, followed by bed-based rehabilitation in intermediate care facilities) was associated with higher costs ( $b=0.456, P<.001$ ). Lastly, a hospital length of stay over the DRG threshold was associated with higher costs ( $b=0.209, P<.001$ ), as well as a rehabilitation lasting 22 days or more ( $b=0.281, P<.001$ ). The inclusion of organizational variables led to a substantive change in the goodness of fit of the model to the data, with an AIC index of 58664 and a BIC index of 58786.

Model A includes “non-modifiable” variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, the “modifiable” variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and LHT of residence.

**Table 2.** Costs (in € currency) for acute hospitalization, rehabilitation, and total costs of CCE by type of intervention and total costs at 3 and 6 months after CCE.

TYPE OF INTERVENTION	ACUTE COSTS (€)			REHABILITATION COSTS (€)			TOTAL COSTS OF CCE (€)			TOTAL COSTS AT 3MO AFTER CCE			TOTAL COSTS AT 6MO AFTER CCE			
	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	N	MEDIAN (95%CI)	PERCENTAGE INCREASE IN TOTAL COSTS (%)	
No intervention	257	2278.4 (2278.4–2437.9)	111	3640.0 (3212.4–4158.0)	257	3921.0 (2437.9–5555.5)	257	3921.01 (2437.86–5804.39)	257	3921.01 (2437.86–5804.39)	257	4005.2 (2740.4–5804.4)	257	4005.2 (2740.4–5804.4)	0.3	1.6
Total hip arthroplasty	531	10079.4 (10079.4–10284.3)	324	3234.0 (2964.9–3542.0)	531	11927.4 (11633.1–12335.4)	531	12029.43 (11662.43–12419.43)	531	12029.43 (11662.43–12419.43)	531	12210.8 (11791.4–12549.4)	531	12210.8 (11791.4–12549.4)	1.4	3.8
Partial hip arthroplasty	1793	10079.4 (10079.4–10284.3)	1224	3198.7 (2990.0–3277.1)	1793	12494.2 (12356.8–12643.4)	1793	12543.43 (12419.43–12697.43)	1793	12543.43 (12419.43–12697.43)	1793	12679.4 (12537.3–12813.4)	1793	12679.4 (12537.3–12813.4)	1.4	3.2
Open reduction and internal fixation	3323	6526.1 (6526.1–6526.1)	2099	3622.1 (3542.0–3770.0)	3323	8686.4 (8566.2–8871.2)	3323	8723.19 (8614.11–8926.47)	3323	8723.19 (8614.11–8926.47)	3323	9000.1 (8789.4–9168.2)	3323	9000.1 (8789.4–9168.2)	2.2	5.3
Total	5904	7404.5 (7113.3–7404.5)	3758	3449.6 (3449.6–3542.0)	5904	10079.4 (10079.4–10284.3)	5904	10079.43 (10079.43–10238.71)	5904	10079.43 (10079.43–10238.71)	5904	10303.9 (10190.4–10429.2)	5904	10303.9 (10190.4–10429.2)	1.8	4.3

Abbreviations: N, number of patients; CI, Confidence Interval.

**Table 3.** Results of generalized linear multiple regression models.

VARIABLE	MODEL A		MODEL B	
	"NOT MODIFIABLE" PREDICTORS AIC=62 580, BIC=62 623		"NOT MODIFIABLE" AND "MODIFIABLE" PREDICTORS AIC=58 664, BIC=58 786	
	B	P VALUE	B	P VALUE
Male gender	-0.026	.069	-0.014	.075
Age	-0.001	.305	-0.002	<.001
Number of comorbid conditions	0.026	<.001	0.012	<.001
Active home care before hospitalization (yes vs no)	-0.096	.040	-0.020	.225
Nursing home stay before hospitalization (yes vs no)	-0.179	<.001	-0.022	.119
Surgery after $\geq$ 48h from admission (yes vs no)			-0.004	.650
Length of hospital stay over DRG threshold			0.209	<.001
Type 1 rehabilitation vs no rehabilitation			0.356	<.001
Type 2 rehabilitation vs no rehabilitation			0.456	<.001
Type 3 rehabilitation vs no rehabilitation			0.184	<.001
Rehabilitation duration over 22 d			0.281	<.001
Rehabilitation duration between 1 and 21 d			0	
LHT 8 vs LHT 5			-0.074	<.001
LHT 7 vs LHT 5			-0.021	.101
LHT 6 vs LHT 5			-0.147	<.001
LHT 4 vs LHT 5			-0.073	<.001
LHT 3 vs LHT 5			-0.093	<.001
LHT 2 vs LHT 5			0.034	.007
LHT 1 vs LHT 5			0.060	<.001

Dependent variable: CCE costs for patients undergoing ORIF surgery (N=3323). Model A includes "non-modifiable" variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, "modifiable" variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and LHT of residence.

Types of post-acute rehabilitation: (1) Type 1, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital); (2) Type 2, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital), followed by rehabilitation in intermediate care facilities (community hospital [CH], or temporary nursing home beds); (3) Type 3, rehabilitation in intermediate care facilities (CH, or temporary nursing home beds).

Abbreviations: AIC, Akaike information criterion; BIC, Bayesian information criterion; b, coefficient; LHT, local health trust; THR, total hip replacement; PHR, hemiarthroplasty or partial hip replacement (PHR); ORIF, open reduction and internal fixation; CH, community hospital.

In model A, for patients undergoing PHR (Table 4), the number of comorbidities was associated with increased CCE costs ( $b=0.018$ ,  $P<.001$ ), while gender was unrelated with CCE costs. Age was associated with costs ( $b=-0.003$ ,  $P<.001$ ). Receiving home care was unrelated with costs, while residing in nursing home before the hospitalization was associated with lower costs ( $b=-0.104$ ,  $P<.001$ , respectively). The AIC and the BIC for this model were respectively 33 602 and 33 640. After adjusting for case-mix variables (model B, Table 4), the surgical intervention after 48 hours from the hospitalization was related with higher costs ( $b=0.017$ ,  $P=.042$ ). Type 2 rehabilitation (hospital bed-based rehabilitation in public or private

healthcare facilities, followed by bed-based rehabilitation in intermediate care facilities) was associated with higher costs ( $b=0.235$ ,  $P<.001$ ). Lastly, a hospital length of stay over the DRG threshold was associated with higher costs ( $b=0.074$ ,  $P<.001$ ), as well as a rehabilitation lasting 22 days or more ( $b=0.202$ ,  $P<.001$ ). The inclusion of organizational variables improved the goodness of fit of the model to the data, with an AIC index of 32 059 and a BIC index of 32 169.

Model A includes "non-modifiable" variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, "modifiable"

**Table 4.** Results of generalized linear multiple regression models.

VARIABLE	MODEL A		MODEL B	
	"NOT MODIFIABLE" PREDICTORS AIC = 33 602, BIC = 33 640		"NOT MODIFIABLE" AND "MODIFIABLE" PREDICTORS AIC = 32 059, BIC = 32 169	
	B	P VALUE	B	P VALUE
Male gender	0.005	.641	0.006	.421
Age	-0.003	.001	-0.001	.012
Number of comorbid conditions	0.018	<.001	0.008	<.001
Active home care before hospitalization (yes vs no)	-0.042	.083	-0.019	.220
Nursing home stay before hospitalization (yes vs no)	-0.104	<.001	-0.029	.048
Surgery after ≥48 h from admission (yes vs no)			0.017	.042
Length of hospital stay over DRG threshold			0.074	<.001
Type 1 rehabilitation vs no rehabilitation			0.218	<.001
Type 2 rehabilitation vs no rehabilitation			0.235	<.001
Type 3 rehabilitation vs no rehabilitation			0.100	<.001
Rehabilitation duration over 22 d			0.202	<.001
Rehabilitation duration between 1 and 21 d			0	
LHT 8 vs LHT 5			-0.020	.061
LHT 7 vs LHT 5			0.028	.042
LHT 6 vs LHT 5			-0.045	.035
LHT 4 vs LHT 5			-0.030	.007
LHT 3 vs LHT 5			-0.031	.020
LHT 2 vs LHT 5			0.041	.005
LHT 1 vs LHT 5			0.057	.002

Dependent variable: CCE costs for patients undergoing PHR surgery (N = 1793). Model A includes "non-modifiable" variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, "modifiable" variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and LHT of residence.

Types of post-acute rehabilitation: (1) Type 1, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital); (2) Type 2, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital), followed by rehabilitation in intermediate care facilities (community hospital [CH], or temporary nursing home beds); (3) Type 3, rehabilitation in intermediate care facilities (CH, or temporary nursing home beds).

Abbreviations: AIC, Akaike information criterion; BIC, Bayesian information criterion; b, coefficient; LHT, local health trust; THR, total hip replacement; PHR, hemiarthroplasty or partial hip replacement (PHR); ORIF, open reduction and internal fixation; CH, community hospital.

variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and LHT of residence.

In model A, for patients undergoing THR (Table 5), the number of comorbidities was associated with increased CCE costs ( $b=0.040$ ,  $P<.001$ ), while gender was unrelated with CCE costs. Age was associated with costs ( $b=0.006$ ,  $P<.001$ ). Receiving home care was associated with lower costs ( $b=-0.156$ ,  $P=.040$ ), while residing in nursing home before the hospitalization was not associated with CCE costs ( $b=0.025$ ,  $P=.785$ ). The AIC and the BIC for this model were respectively 9927 and 9957. After adjusting for "non-modifiable" variables (model B, Table 5), the surgical intervention after 48 hours

from the hospitalization was unrelated with costs ( $b=-0.018$ ,  $P=.203$ ). Type 2 rehabilitation was again associated with higher costs ( $b=0.284$ ,  $P<.001$ ). Lastly, a hospital length of stay over the DRG threshold was associated with higher costs ( $b=0.101$ ,  $P<.001$ ), as well as a rehabilitation lasting 22 days or more ( $b=0.204$ ,  $P<.001$ ). The inclusion of organizational variables improved the goodness of fit of the model to the data, with an AIC index of 9398 and a BIC index of 9482.

Model A includes the "non-modifiable" variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, the "modifiable" variables that depend on organizational models (timing of

**Table 5.** Results of generalized linear multiple regression models.

VARIABLE	MODEL A		MODEL B	
	"NOT MODIFIABLE" PREDICTORS AIC=9927, BIC=9957		"NOT MODIFIABLE" AND "MODIFIABLE" PREDICTORS AIC=9398, BIC=9482	
	B	P VALUE	B	P VALUE
Male gender	0.050	.024	0.021	.108
Age	0.006	<.001	0.000	.904
Number of comorbid conditions	0.040	<.001	0.014	.012
Active home care before hospitalization (yes vs no)	-0.156	.040	-0.039	.406
Nursing home stay before hospitalization (yes vs no)	0.025	.785	-0.075	.184
Surgery after ≥48 h from admission (yes vs no)			-0.018	.203
Length of hospital stay over DRG threshold			0.101	<.001
Type 1 rehabilitation vs no rehabilitation			0.258	<.001
Type 2 rehabilitation vs no rehabilitation			0.284	<.001
Type 3 rehabilitation vs no rehabilitation			0.168	<.001
Rehabilitation duration over 22 d			0.204	<.001
Rehabilitation duration between 1 and 21 d			0	
LHT 8 vs LHT 5			-0.040	.033
LHT 7 vs LHT 5			-0.002	.931
LHT 6 vs LHT 5			-0.055	.079
LHT 4 vs LHT 5			-0.052	.014
LHT 3 vs LHT 5			-0.066	.004
LHT 2 vs LHT 5			0.087	.000
LHT 1 vs LHT 5			0.028	.311

Dependent variable: CCE costs for patients undergoing THR surgery (N=531). Model A includes "non-modifiable" variables (age, gender, multimorbidity, long-term home assistance, and residing in a nursing home before hospital admission for HF). Model B also includes, in addition to Model A variables, "modifiable" variables that depend on organizational models (timing of surgery, rehabilitation type, hospital length of stay, length of rehabilitation) and LHT of residence.

Types of post-acute rehabilitation: (1) Type 1, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital); (2) Type 2, Hospital bed-based rehabilitation in public or private healthcare facilities (including the same hospital), followed by rehabilitation in intermediate care facilities (community hospital [CH], or temporary nursing home beds); (3) Type 3, rehabilitation in intermediate care facilities (CH, or temporary nursing home beds).

Abbreviations: AIC, Akaike information criterion; BIC, Bayesian information criterion; b, coefficient; LHT, local health trust; THR, total hip replacement; PHR, hemiarthroplasty or partial hip replacement (PHR); ORIF, open reduction and internal fixation; CH, community hospital.

surgery, rehabilitation type, hospital length of stay, length of rehabilitation and LHT of residence).

## Discussion

Our study uses a theoretical conceptual framework for the evaluation of cost determinants of integrated healthcare pathways for HF. Investigating the costs of integrated CCEs for patients with HF and identifying their determinants is highly relevant for different healthcare stakeholders, given that HF is one of the conditions with the highest impact on health.<sup>1-5</sup> Despite progress in surgery, HF is characterized by high mortality and a marked reduction in the quality of life in at least half of the surviving subjects; the literature suggests that HF's

clinical outcomes are more strongly associated with the demographic and clinical characteristics of the affected subjects than with the type of trauma or surgical and rehabilitative modality.<sup>39,40</sup> However, less is known about the determinants of costs related to episodes of care. The definition of the CCE allowed us to describe the integrated healthcare pathway of patients with HF and to estimate their cost determinants.

We found that the characteristics of HF patients undergoing different types of surgery have a wide variability. First, patients with THR were 10 years younger than those undergoing PHR. Moreover, we found a high prevalence of female patients in the population with HF.<sup>40</sup> This results, in line with previous literature,<sup>41-43</sup> confirms that older women are at higher



risk of HF due to reduced bone density and other risk factors such as familial predisposition or obesity.

Our results also confirm that the choice of the type of surgery depends not only on the fracture type but also largely on patients' clinical characteristics. When a HF leads to hip replacement surgery, PHR is the preferred option in patients with a more complicated health profile and whose physical activities are limited, while THR is usually applied in younger and healthier patients.<sup>44</sup> Notably, our results indicate that the costs of patients undergoing THR and PHR do not show a substantial difference, neither for the acute part of the CCE, where the cost corresponds to the DRG, nor for the bed-based post-acute rehabilitation. As expected, ORIF costs less than hip replacement (both PHR and THR) in the acute part and in the total costs, while in these patients (and in those who are not subjected to surgery) the rehabilitation part of the CCE is more expensive.

Our findings confirmed that much of the variability in total costs is mainly attributable to differences in post-acute care costs, particularly to rehabilitation.<sup>45-47</sup> Moreover, these costs are in line with the range found in the literature.<sup>48</sup>

The analysis of the determinants of CCE's costs shows that some patients' demographic and clinical characteristics are associated with differential costs. Gender is unrelated to CCE costs in patients undergoing ORIF, PHR, or THR surgery. This is interesting, as it underlines that no gender differences are present in this specific context of care. Age is weakly associated with costs. Notably, as patients' age increases, CCE costs are reduced. This association is statistically significant for patients undergoing ORIF and PHR, but not THR, and confirms what reported in literature and guidelines.<sup>25,45,48,49</sup> In fact, older patients usually undergo less complex and therefore less expensive surgery, thus reducing overall costs.<sup>43</sup>

Increasing spending is correlated with the co-morbidity burden of geriatric HF patients, in line with previous literature.<sup>50-53</sup> Johnson et al.<sup>54</sup> showed that only 4.95% of HF patients presented without associated comorbidities. Many authors demonstrated that a higher comorbidity burden is associated with higher costs of care and longer length of stay for geriatric HF. Data presented here corroborates Casaletto and Gatt's<sup>55</sup> findings that patients with more comorbidities (HF) account for higher cost of care.

Current geriatric HF literature supports surgical stabilization within 48 hours as a pathway to reduce morbidity, reduce mortality, and preserve function.<sup>56-59</sup> It is widely reported in the literature, and underlined by the guidelines, that performing the surgery within 48 hours from the hospital admission provides better clinical outcomes.<sup>3,48</sup> However, our results underscore that carrying out each of the surgical interventions (ORIF, PHR, and THR) after 48 hours from the hospital admission does not determine an increase in the overall costs of the CCE. We can therefore argue that, while the majority of the literature identifies the early intervention as an instrument to achieve

better clinical and functional outcomes, it cannot be considered associated with a reduction of the overall costs of the CCE.

On the contrary, active home care or being resident in a nursing home before the hospital admission for HF is associated with cost reduction, particularly for patients undergoing ORIF and PHR. A possible explanation is that patients with these characteristics are already being treated by healthcare and social services, thus requiring less contacts with other healthcare facilities, and are also older, which has been proved to be associated with lower cost interventions.<sup>44</sup>

After adjusting for other variables, hospital bed-based rehabilitation in public or private healthcare facilities, either followed by rehabilitation in a community hospital/temporary nursing home beds or not, is the rehabilitation type associated with higher overall costs of CCE, regardless of the type of surgical intervention. Conversely, bed-based rehabilitation performed exclusively in intermediate care facilities (CH or nursing home) is associated with lower overall costs. The duration of the rehabilitation has also an impact on costs, in line with existing literature.<sup>60,61</sup>

These results are mainly due to the lower costs attributable to the patients' stay in intermediate care beds, which reflects the lower intensity of care required by HF patients in the post-acute phase. Net of patients' case-mix and surgical intervention, it appears that the intermediate care organizational model contributes to the integrated episode of care for HF with a lower economic weight. Therefore, in designing appropriate integrated care pathways for this type of patient, this organizational option should certainly be taken into consideration.

This study has some limitations. Our analyses are based on HF occurring in 1 calendar year; it is possible that variations of costs occur over time because of changes in reimbursement policies. Moreover, it is possible that the differences in costs we found reflect unmeasured differences in patient complexity, such as frailty degree, vitamin D status, and fall dynamics. In fact, many factors can determine health outcomes and costs in elderly patients with hip fracture, as health outcomes and costs could vary based on presence of certain conditions that impact bone quality as well as the risk of falls. However, our analyses are based on data extracted from the hospital discharge records database, including patients with HF resident in Emilia Romagna in 2017. In order to account for patients' comorbidity in the predictive models of costs, we used an internationally validated comorbidity indicator, the ECI.<sup>37</sup> Unfortunately, characteristics as individual frailty, drug use and concomitant conditions that may affect the risk of hip fracture are not recorded in the hospital discharge records database.

Non-medical costs were not included in the analyses because they were not available. Also, in the present study we did not focus on the clinical outcomes of HF patients, as evaluating HF functional outcomes on a large retrospective cohort such as the one present here was not within the specific objectives.

## Conclusion

In this study, we investigated the determinants of costs of continuum-care episodes for elderly patients with hip fracture. Our findings suggest that, after adjusting for patients' case-mix, the type of rehabilitation results to be the main determinant of overall costs, and that bed-based rehabilitation performed in intermediate care facilities is associated with lower overall spending. Cost-wise, intermediate care organizational settings should be privileged when planning hip fracture rehabilitation pathways.

## Author Contributions

Davide Golinelli, Erik Boetto, Elena Berti, Cristina Ugolini, and Maria Pia Fantini conceived and designed the study, reviewed the literature, and wrote the manuscript. Simona Rosa and Paola Rucci acquired the data, carried out the statistical analyses, and reviewed the manuscript. Davide Golinelli, Erik Boetto, and Antonio Mazzotti interpreted the data and reviewed and edited the final manuscript in collaboration with Maria Pia Fantini for intellectual contents. All authors have read and approved the final manuscript.

## Ethics Approval

The study has been approved by the AVEC (Comitato Etico Indipendente di Area Vasta Emilia Centro) research ethics committee board, ID 233/2019/OSS/AOUBo on April 17, 2019. Data used in this research was obtained from the Regional Healthcare Information System which includes detailed information on the use of healthcare services by all regional patients, with the patient as our unit of observation. The study, based on routine administrative information, was carried out in conformity with the regulations on data management of the Emilia-Romagna Region and with Italian privacy law. Administrative data were anonymized by an ad hoc service of the Regional Department of Health and the University of Bologna have no possibility to retrospectively identify individuals included in the study.

## ORCID iD

Davide Golinelli  <https://orcid.org/0000-0001-7331-9520>

## Supplemental Material

Supplemental material for this article is available online.

## Availability of Data and Material

The data analyzed are property of the Emilia Romagna Regional Department of Health.

## REFERENCES

- Hansen L, Mathiesen AS, Vestergaard P, Ehlers LH, Petersen KD. A health economic analysis of osteoporotic fractures: who carries the burden? *Arch Osteoporos*. 2013;8:126. doi:10.1007/s11657-013-0126-3
- Canton G, Moghnie A, Cleva M, Kostoris FM, Murena L. Dual mobility total hip arthroplasty in the treatment of femoral neck fractures: a retrospective evaluation at mid-term follow-up. *Acta Biomed*. 2019;90:98-103. doi:10.23750/abm.v90i1-S.8070
- Moja L, Piatti A, Pecoraro V, et al. Timing matters in hip fracture surgery: patients operated within 48 hours have better outcomes. A meta-analysis and meta-regression of over 190,000 patients. *PLoS One*. 2012;7:e46175. doi:10.1371/journal.pone.0046175
- Braithwaite RS, Col NF, Wong JB. Estimating hip fracture morbidity, mortality and costs. *J Am Geriatr Soc*. 2003;51:364-370. doi:10.1046/j.1532-5415.2003.51110.x
- Leal J, Gray AM, Hawley S, et al. Cost-effectiveness of orthogeriatric and fracture liaison service models of care for hip fracture patients: a population-based study. *J Bone Miner Res*. 2017;32:203-211. doi:10.1002/jbmr.2995
- Sambrook P, Cooper C. Osteoporosis [published correction appears in *Lancet*. 2006 Jul 1;368(9529):28]. *Lancet*. 2006;367:2010-2018. doi:10.1016/S0140-6736(06)68891-0
- Programma Nazionale Esiti 2018. [Outcomes National Program] AGENAS. <https://pne.agenas.it/>
- Svedbom A, Hernlund E, Ivergård M, et al. Osteoporosis in the European Union: a compendium of country-specific reports. *Arch Osteoporos*. 2013;8:137. doi:10.1007/s11657-013-0137-0
- Piscitelli P, Chitano G, Johansson H, Brandi ML, Kanis JA, Black DM. Updated fracture incidence rates for the Italian version of FRAX®. *Osteoporos Int*. 2013;24:859-866. doi:10.1007/s00198-012-2021-y
- Lahtinen A, Leppilähti J, Vähänikkilä H, et al. Costs after hip fracture in independently living patients: a randomised comparison of three rehabilitation modalities. *Clin Rehabil*. 2017;31:672-685. doi:10.1177/0269215516651480
- Cheng SY, Levy AR, Lefavre KA, Guy P, Kuramoto L, Sobolev B. Geographic trends in incidence of hip fractures: a comprehensive literature review. *Osteoporos Int*. 2011;22:2575-2586. doi:10.1007/s00198-011-1596-z
- Regione Emilia-Romagna, Deliberazione della Giunta Regionale 2 Agosto 2013, N. 1133. Organizzazione dell'assistenza integrata al paziente anziano con fratture di femore - Approvazione linee guida per le Aziende Sanitarie della Regione Emilia-Romagna. [Organization of integrated care for elderly patients with femur fractures - Approval of guidelines for Healthcare Organizations of the Emilia-Romagna Region.] Bollettino Ufficiale della Regione Emilia-Romagna n.247 del 22.08.2013. Accessed November 15, 2020. <http://bur.regione.emilia-romagna.it/dettaglio-inserzione?i=eb0819fb559aa40449f54cf9cf32bc39>
- Clegg A, Young J, Illiffe S, Rikkert MO, Rockwood K. Frailty in elderly people [published correction appears in *Lancet*. 2013 Oct 19;382(9901):1328]. *Lancet*. 2013;381:752-762. doi:10.1016/S0140-6736(12)62167-9
- Zidén L, Wenestam CG, Hansson-Scherman M. A life-breaking event: early experiences of the consequences of a hip fracture for elderly people. *Clin Rehabil*. 2008;22:801-811. doi:10.1177/0269215508090204
- Auais M, Morin S, Nadeau L, Finch L, Mayo N. Changes in frailty-related characteristics of the hip fracture population and their implications for health-care services: evidence from Quebec, Canada. *Osteoporos Int*. 2013;24:2713-2724. doi:10.1007/s00198-013-2390-x
- Loizzo M, Gallo F, Caruso D. Reducing complications and overall healthcare costs of hip fracture management: a retrospective study on the application of a Diagnostic Therapeutic Pathway in the Cosenza General Hospital. *Ann Ig*. 2018;30:191-199. doi:10.7416/ai.2018.2210
- Kistler EA, Nicholas JA, Kates SL, Friedman SM. Frailty and short-term outcomes in patients with hip fracture. *Geriatr Orthop Surg Rehabil*. 2015;6:209-214. doi:10.1177/2151458515591170
- Endo A, Baer HJ, Nagao M, Weaver MJ. Prediction model of in-hospital mortality after hip fracture surgery. *J Orthop Trauma*. 2018;32:34-38. doi:10.1097/BOT.0000000000001026
- Kalmet PH, Koc BB, Hemmes B, et al. Effectiveness of a multidisciplinary clinical pathway for elderly patients with hip fracture: a multicenter comparative cohort study. *Geriatr Orthop Surg Rehabil*. 2016;7:81-85. doi:10.1177/2151458516645633
- Petros RSB, Ferreira PEV, Petros RSB. Influence of proximal femur fractures in the autonomy and mortality of elderly patients submitted to osteosynthesis with cephalomedullary nail. *Rev Bras Ortop*. 2017;52:57-62. doi:10.1016/j.rboe.2017.08.014
- Aarden JJ, van der Esch M, Engelbert RHH, van der Schaaf M, de Rooij SE, Buurman BM. Hip fractures in older patients: trajectories of disability after surgery. *J Nutr Health Aging*. 2017;21:837-842. doi:10.1007/s12603-016-0830-y
- de Jooe SGCJ, Kalmet PHS, Fiddlers AAA, Poeze M, Blokhuis TJ. Long-term functional outcome after a low-energy hip fracture in elderly patients. *J Orthop Traumatol*. 2019;20:20. doi:10.1186/s10195-019-0529-z
- Shiga T, Wajima Z, Ohe Y. Is operative delay associated with increased mortality of hip fracture patients? Systematic review, meta-analysis, and meta-regression. *Can J Anaesth*. 2008;55:146-154. doi:10.1007/BF03016088
- Carretta E, Bochicchio V, Rucci P, Fabbri G, Laus M, Fantini MP. Hip fracture: effectiveness of early surgery to prevent 30-day mortality. *Int Orthop*. 2011;35:419-424. doi:10.1007/s00264-010-1004-x
- Scottish Intercollegiate Guidelines Network. Management of hip fracture in older people. *A national clinical guideline*. 2009. [https://www.shfa.scot.nhs.uk/docs/20161109\\_SSC\\_for\\_Hip\\_Fracture\\_Patients.pdf](https://www.shfa.scot.nhs.uk/docs/20161109_SSC_for_Hip_Fracture_Patients.pdf)

26. Handoll HH, Cameron ID, Mak JC, Finnegan TP. Multidisciplinary rehabilitation for older people with hip fractures. *Cochrane Database Syst Rev.* 2009;4:CD007125. doi:10.1002/14651858.CD007125.pub2
27. Melis RJ, Olde Rikkert MG, Parker SG, van Eijken MI. What is intermediate care? *BMJ.* 2004;329:360-361. doi:10.1136/bmj.329.7462.360
28. Young J, Stevenson J. Intermediate care in England: where next? *Age Ageing.* 2006;35:339-341. doi:10.1093/ageing/af026
29. Young J. The development of intermediate care services in England. *Arch Gerontol Geriatr.* 2009;49:S21-S25. doi:10.1016/S0167-4943(09)70008-1
30. Lenzi J, Mongardi M, Rucci P, et al. Sociodemographic, clinical and organisational factors associated with delayed hospital discharges: a cross-sectional study. *BMC Health Serv Res.* 2014;14:128. doi:10.1186/1472-6963-14-128
31. Young J, Gladman JR, Forsyth DR, Holditch C. The second national audit of intermediate care. *Age Ageing.* 2015;44:182-184. doi:10.1093/ageing/afu174
32. Compagni A, Tediosi F, Tozzi VD. L'integrazione tra ospedale e territorio nelle Aziende Sanitarie. In: Cantù E, ed. *Rapporto OASI 2010: L'aziendalizzazione della sanità in Italia.* EGEA; 2010:519-538.
33. Morando V, Tozzi VD. Lo zibaldone delle cure intermedie: analisi delle esperienze regionali, in Italia e in Europa. In: CERGAS - Bocconi, ed. *Rapporto OASI 2016: Osservatorio sulle Aziende e sul Sistema sanitario Italiano.* EGEA, 2010. [The zibaldone of intermediate care: analysis of regional experiences, in Italy and in Europe. In: CERGAS - Bocconi, ed. *OASI Report 2016: Observatory on Companies and the Italian Health System*].
34. Sheehan KJ, Sobolev B, Guy P, et al. Constructing an episode of care from acute hospitalization records for studying effects of timing of hip fracture surgery. *J Orthop Res.* 2016;34:197-204. doi:10.1002/jor.22997
35. Sheehan KJ, Smith TO, Martin FC, et al. Conceptual framework for an episode of rehabilitative care after surgical repair of hip fracture. *Phys Ther.* 2019;99:276-285. doi:10.1093/ptj/pzy145
36. Pitzul KB, Wodchis WP, Carter MW, Kreder HJ, Voth J, Jaglal SB. Post-acute pathways among hip fracture patients: a system-level analysis. *BMC Health Serv Res.* 2016;16:275. doi:10.1186/s12913-016-1524-1
37. Elixhauser A, Steiner C, Harris DR, et al. Comorbidity measures for use with administrative data. *Med Care.* 1998;36:8-27.
38. Cecchi F, Pancani S, Antonioli D, et al. Predictors of recovering ambulation after hip fracture inpatient rehabilitation. *BMC Geriatr.* 2018;18:201. doi:10.1186/s12877-018-0884-2
39. Veronese N, Maggi S. Epidemiology and social costs of hip fracture. *Injury.* 2018;49:1458-1460. doi:10.1016/j.injury.2018.04.015
40. Williamson S, Landeiro F, McConnell T, et al. Costs of fragility hip fractures globally: a systematic review and meta-regression analysis. *Osteoporos Int.* 2017;28:2791-2800. doi:10.1007/s00198-017-4153-6
41. Cummings SR, Nevitt MC, Browner WS, et al. Risk factors for hip fracture in white women. Study of Osteoporotic Fractures Research Group. *N Engl J Med.* 1995;332:767-773. doi:10.1056/NEJM199503233321202
42. Sterling RS. Gender and race/ethnicity differences in hip fracture incidence, morbidity, mortality, and function. *Clin Orthop Relat Res.* 2011;469:1913-1918. doi:10.1007/s11999-010-1736-3
43. Farmer ME, White LR, Brody JA, Bailey KR. Race and sex differences in hip fracture incidence. *Am J Public Health.* 1984;74:1374-1380. doi:10.2105/ajph.74.12.1374
44. Guyen O. Hemiarthroplasty or total hip arthroplasty in recent femoral neck fractures? *Orthop Traumatol Surg Res.* 2019;105:S95-S101. doi:10.1016/j.otsr.2018.04.034
45. Weeks WB, Schoellkopf WJ, Ballard DJ, Kaplan GS, James B, Weinstein JN. Episode-of-care characteristics and costs for hip and knee replacement surgery in hospitals belonging to the high value healthcare collaborative compared with similar hospitals in the same health care markets. *Med Care.* 2017;55:583-589. doi:10.1097/MLR.0000000000000710
46. Weeks WB, Schoellkopf WJ, Sorensen LS, Masica AL, Nesse RE, Weinstein JN. The high value healthcare collaborative: observational analyses of care episodes for hip and knee arthroplasty surgery. *J Arthroplasty.* 2017;32:702-708. doi:10.1016/j.arth.2016.09.009
47. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987;40:373-383. doi:10.1016/0021-9681(87)90171-8
48. Burgers PT, Hoogendoorn M, Van Woensel EA, et al. Total medical costs of treating femoral neck fracture patients with hemi- or total hip arthroplasty: a cost analysis of a multicenter prospective study. *Osteoporos Int.* 2016;27:1999-2008. doi:10.1007/s00198-016-3484-z
49. Tedesco D, Gibertoni D, Rucci P, et al. Impact of rehabilitation on mortality and readmissions after surgery for hip fracture. *BMC Health Serv Res.* 2018;18:701. doi:10.1186/s12913-018-3523-x
50. Kirkland LL, Kashiwagi DT, Burton MC, Cha S, Varkey P. The Charlson comorbidity index score as a predictor of 30-day mortality after hip fracture surgery. *Am J Med Qual.* 2011;26:461-467. doi:10.1177/1062860611402188
51. Neuhaus V, King J, Hageman MG, Ring DC. Charlson comorbidity indices and in-hospital deaths in patients with hip fractures. *Clin Orthop Relat Res.* 2013;471:1712-1719. doi:10.1007/s11999-012-2705-9
52. de Luise C, Brimacombe M, Pedersen L, Sorensen HT. Comorbidity and mortality following hip fracture: a population-based cohort study. *Aging Clin Exp Res.* 2008;20:412-418. doi:10.1007/BF03325146
53. Nikkel LE, Fox EJ, Black KP, Davis C, Andersen L, Hollenbeak CS. Impact of comorbidities on hospitalization costs following hip fracture. *J Bone Joint Surg Am.* 2012;94:9-17. doi:10.2106/JBJS.J.01077
54. Johnson DJ, Greenberg SE, Sathiyakumar V, et al. Relationship between the Charlson Comorbidity Index and cost of treating hip fractures: implications for bundled payment. *J Orthop Traumatol.* 2015;16:209-213. doi:10.1007/s10195-015-0337-z
55. Casaletto JA, Gatt R. Post-operative mortality related to waiting time for hip fracture surgery. *Injury.* 2004;35:114-120. doi:10.1016/s0020-1383(03)0210-9
56. Simunovic N, Devereaux PJ, Sprague S, et al. Effect of early surgery after hip fracture on mortality and complications: systematic review and meta-analysis. *CMAJ.* 2010;182:1609-1616. doi:10.1503/cmaj.092220
57. Moran CG, Wenn RT, Sikand M, Taylor AM. Early mortality after hip fracture: is delay before surgery important? *J Bone Joint Surg Am.* 2005;87:483-489. doi:10.2106/JBJS.D.01796
58. Kenzora JE, McCarthy RE, Lowell JD, Sledge CB. Hip fracture mortality. Relation to age, treatment, preoperative illness, time of surgery, and complications. *Clin Orthop Relat Res.* 1984;186:45-56.
59. Bohm E, Loucks L, Wittmeier K, Lix LM, Oppenheimer L. Reduced time to surgery improves mortality and length of stay following hip fracture: results from an intervention study in a Canadian health authority. *Can J Surg.* 2015;58:257-263. doi:10.1503/cjs.017714
60. Huusko TM, Karppi P, Avikainen V, Kautiainen H, Sulkava R. Intensive geriatric rehabilitation of hip fracture patients: a randomized, controlled trial. *Acta Orthop Scand.* 2002;73:425-431. doi:10.1080/00016470216324
61. Fliess E, Weinstein O, Sherf M, Dreier J. Healthcare services utilization following admission for hip fracture in elderly patients. *Int J Qual Health Care.* 2018;30:104-109. doi:10.1093/intqhc/mzx178