

Japanese Nationwide Study on the Association Between Short-term Exposure to Particulate Matter and Mortality

Takehiro Michikawa^{1,2}, Kayo Ueda^{3,4}, Akinori Takami⁵, Seiji Sugata⁵,
Ayako Yoshino⁵, Hiroshi Nitta¹, and Shin Yamazaki¹

¹Centre for Health and Environmental Risk Research, National Institute for Environmental Studies, Ibaraki, Japan

²Department of Environmental and Occupational Health, School of Medicine, Toho University, Tokyo, Japan

³Environmental Health Sciences, Kyoto University Graduate School of Global Environmental Studies, Kyoto, Japan

⁴Department of Environmental Engineering, Graduate School of Engineering, Kyoto University, Kyoto, Japan

⁵Centre for Regional Environmental Research, National Institute for Environmental Studies, Ibaraki, Japan

Received June 19, 2018; accepted October 8, 2018; released online October 27, 2018

ABSTRACT

Background: From around 2012, the use of automated equipment for fine particulate matter (PM_{2.5}) measurement with equivalence to a reference method has become popular nationwide in Japan. This enabled us to perform a national health effect assessment employing PM_{2.5} concentrations based on the standardized measurement method. We evaluated the association between non-accidental mortality and short-term exposure to PM_{2.5} and coarse particulate matter (PM), with the latter estimated as the difference between suspended particulate matter and PM_{2.5}, for the fiscal years 2012–2014.

Methods: This was a time-stratified case-crossover study in 100 highly-populated Japanese cities. Mortality data was obtained from the Ministry of Health, Labour and Welfare. City-specific estimates of PM-mortality association were calculated by applying a conditional logistic regression analysis, and combined with a random-effects meta-analysis.

Results: The respective averages of daily mean concentration were 14.6 µg/m³ for PM_{2.5} and 6.4 µg/m³ for coarse PM. A 10 µg/m³ increase in PM_{2.5} concentrations for the average of the day of death and the previous day was associated with an increase of 1.3% (95% confidence interval (CI), 0.9–1.6%) in total non-accidental mortality. For cause-specific mortality, PM_{2.5} was positively associated with cardiovascular and respiratory mortality. After adjustment for PM_{2.5}, we observed a 1.4% (95% CI, 0.2–2.6%) increase in total mortality with a 10 µg/m³ increase in coarse PM.

Conclusion: The study revealed that short-term exposure to PM_{2.5} had adverse effects on total non-accidental, cardiovascular, and respiratory mortality in Japan. Coarse PM exposure also increased the risk of total mortality.

Key words: fine particulate matter; coarse particle; mortality; case-crossover; Japanese

Copyright © 2018 Takehiro Michikawa et al. This is an open access article distributed under the terms of Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

INTRODUCTION

The health effects of particulate matter, particularly fine particulate matter (PM_{2.5} that passes through a size-selective inlet with a 50% cut-off level of 2.5 µm in aerodynamic diameter), are an international concern,¹ and epidemiologic evidence increases year by year. For instance, with respect to mortality as a classical outcome, a PM_{2.5}-daily mortality association has been established, and the effect size, on a national or multi-country scale, has been considered from the viewpoint of public health. One nationwide study in 112 American cities revealed that an increase of 10 µg/m³ in PM_{2.5} exposure was associated with a 0.98% increase in all-cause mortality,² and another nationwide study, in 272 Chinese cities, estimated a similar increase of 0.22% in mortality.³ Further, a multi-country study in 10 European cities estimated a 0.55% increase in mortality.⁴ Nonetheless, evidence of the effects of PM_{2.5} exposure on health in Japan is extremely limited because the routine measurement of PM_{2.5} in Japanese air

pollution monitoring stations only began after the Air Quality Standard for PM_{2.5} was established in 2009.⁵ Previously, we reported that short-term exposure to PM_{2.5} was positively associated with all-cause mortality in those aged 65 years or older in 20 regions between 2002 and 2004⁶; however, PM_{2.5} concentration measurement equivalence was not ensured among the 20 automated measuring devices in the study.

To perform PM_{2.5} concentration comparison among cities, systematic measurement error based on the measuring equipment should be avoided. Therefore, the Japanese Air Quality Standards have defined the PM_{2.5} measurement method as follows: “mass measurement with filter sample collection, which is designated as a reference method, or alternative automated methods, designated as equivalent methods, which are proved to have measurement performance comparable to the corresponding reference method”.⁵ From around 2012, use of automated equipment enabling PM_{2.5} measurement with equivalence to a reference method has become popular nationwide. This allowed us to perform a

Address for correspondence. Takehiro Michikawa, Department of Environmental and Occupational Health, School of Medicine, Toho University, 5-21-16 Omori-nishi, Ota-ku, Tokyo 143-8540, Japan (e-mail: takehiro.michikawa@med.toho-u.ac.jp).

national health effect assessment that employed PM_{2.5} concentrations based on the standardized measurement method.

We conducted a multi-city case-crossover study of the association between short-term exposure to PM_{2.5} and mortality in Japan. Additionally, with regard to exposure to coarse particulate matter (coarse PM), for which there is still limited health effects evidence globally, we explored the association with mortality using an exposure surrogate for coarse PM, estimated as the difference between the concentrations of suspended particulate matter (SPM that passes through a size-selective inlet with a 100% cut-off level of 10 µm in aerodynamic diameter) and PM_{2.5}.

METHODS

Study area

In the 2015 Japanese Population Census, there were 110 cities with a total population exceeding 200,000. Of these 110 cities, 99 had one or more ambient air pollution monitoring stations with automated PM_{2.5} measuring equipment with measurement equivalence to the reference method. Additionally, we included in the study area Tsukuba city (population: roughly 227,000 people in the 2015 Population Census), which has no monitoring station but is the home of the Japan National Institute for Environmental Studies, by combining Tsukuba City and adjacent Tsuchiura City (which has a monitoring station) into one city. Our 100 target cities were located throughout the length of Japan, from Hokkaido Prefecture (Sapporo, Asahikawa, and Hakodate) to Okinawa Prefecture (Naha). The population and size of the 100 target cities are presented in eTable 1. The study protocol was reviewed and approved by the Ethics Committee of the National Institute for Environmental Studies (2017-004), and the Faculty of Medicine, Toho University (A18051).

Environmental data

The hourly measurement of PM_{2.5} concentrations using the reference method-equivalent automated PM_{2.5} measuring equipment, measured at one ambient air pollution monitoring station within each city from April 2012 to March 2015, were obtained from the Japan National Institute for Environmental Studies' atmospheric environment database. The PM_{2.5} measurement periods varied by city (eTable 1). Among 100 cities, 66 cities had only one ambient air monitoring station measuring PM_{2.5}. According to the pollutant monitoring manual of the Ministry of the Environment, Japan,⁷ such stations are located at the points that seem to be representative of pollutant concentrations in the community. For the remaining cities, which all had two or more ambient air pollution monitoring stations, we selected one that satisfied the following criteria: (1) the station measured other co-pollutants, including SPM, photochemical oxidants (Ox), nitrogen dioxide (NO₂), and sulphur dioxide (SO₂), in addition to PM_{2.5}; and/or (2) the station was located nearer to the center of the city than other stations. However, in such cases, we confirmed a strong correlation between the PM_{2.5} concentrations measured at the selected station and those of the other station(s) in the same city (mean correlation coefficient = 0.95). The locations of the monitoring stations in the 100 target Japanese cities are presented in eFigure 1. The daily mean concentrations of PM_{2.5} were calculated based on hourly measurements from 0 to 23 h, and no daily mean concentrations were calculated for days when more than four hourly measurements were missing. The median %

missing days in the PM_{2.5} data was 1.2% (interquartile range [IQR], 0.6–2.2%), and these days were excluded in the analysis.

The hourly measurements of other co-pollutants were also obtained, and the daily mean concentrations of SPM, NO₂ and SO₂, as well as the maximum 8-h concentrations of Ox, were calculated. In this study, the difference between the respective concentrations of SPM and PM_{2.5} was used as an exposure surrogate for coarse PM. Internationally, coarse PM is defined as particulate matter measuring between 2.5 µm and 10 µm (PM_{10-2.5}) in diameter.⁸ However, the monitoring stations in Japan do not routinely measure PM₁₀ (particles that pass through a size-selective inlet with a 50% cut-off level of 10 µm in aerodynamic diameter). For the coarse PM-mortality association, we included 77 cities where concentrations of coarse PM had been reasonably distributed.

Meteorological data (daily mean ambient temperature and relative humidity), measured at the nearest meteorological observatory to each city, were collected from the Japan Meteorological Agency. Finally, information on the weekly influenza counts was collected from the Japan National Institute of Infectious Diseases, and an influenza epidemic was considered to persist for the number of weeks above the 90 percentile of the distribution during the study period.⁹

Mortality data

For all the study areas, we applied to the Ministry of Health, Labour and Welfare, Japan, for use of the data from its Vital Statistic Survey, and obtained daily mortality records, including the age and sex of the deceased, and the date and location of death, as well as the primary cause of death according to the 10th edition of the International Classification of Diseases (ICD-10). In this study, our primary outcome was total non-accidental death (ICD-10: A00 through R99), and specific causes were defined as: cardiovascular disease (I00 through I99); coronary heart disease (I20 through I25); stroke (I60 through I69); and respiratory disease (J00 through J99).

Statistical analysis

We used a time-stratified case-crossover design and compared an individual's exposure on the "case" day with his or her exposure on the "control" days, when he or she did not become a case.¹⁰ We defined the "case" day as the day of death, and selected three or four "control" days from the same days of the week, in the same month and year, as the "case" day. For instance, if an individual died on Mar 15, 2013, four control days were assigned: March 1, 8, 22, and 29 of that year. We excluded cases on national holidays (not including weekends), to avoid bias in selection control due to the difference between pollutant concentrations on holidays and on non-holidays.

To investigate the association between exposure to PM_{2.5} and mortality, we first applied city-specific conditional logistic regression models. Based on past studies,^{2,3,11} the choice of the lag0-1 (2-day moving average of the case-day and previous-day) concentrations of PM_{2.5} was made before commencement of the study. The PM_{2.5} concentrations at lag2 and lag3 were used to investigate the lag pattern of the PM_{2.5}-mortality association. We constructed a distributed lag model, simultaneously including PM_{2.5} concentrations at lag0-1, lag2, and lag3, and adjusted for ambient temperature at lag0-1 using a five-knot natural cubic spline, relative humidity at lag0-1 using a three-knot cubic spline, and influenza epidemics. Odds ratios (ORs) with 95% confidence

intervals (CIs) of mortality based on an increase of 10 µg/m³ in PM_{2.5} concentration at lag0-1 were estimated because a linear model was not inferior to a non-linear model with respect to estimation of the PM_{2.5}-mortality association. Subsequently, the city-specific estimates were used to obtain pooled estimates on the PM_{2.5}-mortality association using random-effects meta-analysis. The ORs were converted into excess risks approximated by [(OR - 1) × 100]. The proportion of total variation in the city-specific estimates, attributable to heterogeneity, was assessed by computing the I² statistic. The same two-stage approach was followed in estimating the association between coarse PM and mortality. We mutually adjusted for PM_{2.5} and coarse PM after estimating each PM-mortality association in the single-pollutant model.

For sensitivity analysis, we constructed two-pollutant models; and another model additionally adjusted for temperature at lag2-21 to investigate the long delay of the cold effects.¹² With respect to PM_{2.5} exposure, to estimate the health effects at PM_{2.5} concentrations not exceeding the daily World Health Organization (WHO) guideline values,¹³ we analyzed after restricting to days with PM_{2.5} concentrations at lag0-1 ≤25 µg/m³. We also performed stratified analyses by age (<75 or ≥75 years) and sex, and examined seasonal variation in the PM-mortality association. Since the health effects of PM_{2.5} seem likely to differ with the composition and sources of PM_{2.5},¹⁴ we also explored region-specific associations: East Japan, where domestic pollution is the major contributor to PM_{2.5} concentrations; Central Japan; and West Japan, where transboundary pollution makes a large contribution to PM_{2.5} (eTable 1).¹⁵ All analyses were performed using STATA version 14 (Stata Corporation, College Station, TX, USA).

RESULTS

In this nationwide analysis, the average daily mean PM_{2.5} concentration was 14.6 (standard deviation [SD], 8.3) µg/m³ (Table 1), and the 98th percentile of daily mean concentrations was 37.8 µg/m³. In terms of region-specific statistics, the respective averages of PM_{2.5} concentration were 13.6 (SD, 8.0) µg/m³ in East Japan, 14.6 (SD, 8.3) µg/m³ in Central Japan, and 17.9 (SD, 9.5) µg/m³ in West Japan. The average daily mean concentration of coarse PM was 6.4 (SD, 5.1) µg/m³. The city-specific results for environmental factors are presented in eTable 1. The Pearson’s correlation coefficients with PM_{2.5} were 0.33 for coarse PM, 0.37 for Ox, 0.47 for NO₂, and 0.42 for SO₂ (eTable 2). Among 1,347,152 non-accidental deaths, cardiovascular mortality accounted for 28.1% and respiratory mortality for 16.7% (Table 2).

When we explored the city-specific association between exposure to PM_{2.5} and total non-accidental mortality, the point estimates of percentage increase for 10 µg/m³ increase in PM_{2.5} at lag0-1 showed a direction of increased mortality risk in 72 cities (eFigure 2). We did not observe significant heterogeneity (I² = 9.8%). Pooled estimates for the association between PM_{2.5} exposure and mortality are presented in Table 3. Exposure to PM_{2.5} at lag0-1 was associated with an increase in total mortality of 1.3% per 10 µg/m³ (95% CI, 0.9–1.6). The respective percentage increases for the association of exposure to PM_{2.5} and cause-specific mortality were 1.6% (95% CI, 0.8–2.4%) for cardiovascular disease, 2.7% (95% CI, 1.0–4.4%) for coronary heart disease, 1.3% (95% CI, -0.2 to 2.8%) for stroke, and 1.5%

Table 1. Overall statistics of environmental factors in 100 Japanese cities, April 2012 to March 2015

Environmental factors, daily mean	Number of cities	Mean (SD)	Percentile		
			25th	50th	75th
PM _{2.5} , µg/m ³	100	14.6 (8.3)	8.6	13.1	18.8
Coarse PM, µg/m ³ ^a	77	6.4 (5.1)	2.9	5.6	8.9
Ox, ppb ^b	95	42.3 (15.2)	31.8	40.6	51.7
NO ₂ , ppb	99	12.4 (6.1)	7.9	11.2	15.7
SO ₂ , ppb	91	2.1 (1.2)	1.3	1.9	2.7
Ambient temperature, °C	100	15.7 (8.5)	7.7	16.0	23.0
Relative humidity, %	100	67 (13)	58	68	76

NO₂, nitrogen dioxide; Ox, photochemical oxidants; PM, particulate matter; SD, standard deviation; SO₂, sulphur dioxide.

^aConcentrations of coarse PM were calculated by subtracting PM_{2.5} concentrations from those of suspended particulate matter.

^bDaily maximum 8-h mean concentrations.

Table 2. Overall characteristics of non-accidental deaths (1,347,152 deaths) in 100 Japanese cities, April 2012 to March 2015

Outcome	Percent
Total non-accidental (ICD-10: A00–R99)	100
Age, years	
<75	27.9
≥75	72.1
Sex	
Men	51.9
Women	48.1
Cardiovascular disease (ICD-10: I00–99)	28.1
Coronary heart disease (ICD-10: I20–25)	6.4
Stroke (ICD-10: I60–69)	9.2
Respiratory disease (ICD-10: J00–99)	16.7

ICD, international classification of diseases.

(95% CI, 0.7–2.3%) for respiratory disease. PM_{2.5} at lag2 and lag3 was not associated with either total or cause-specific mortality.

We performed sensitivity and stratified analyses of the association between PM_{2.5} exposure and total non-accidental mortality (Figure 1). After adjustment for exposure to co-pollutants, or for temperature at long lag, the percentage increase point estimates were attenuated; however, the positive association between PM_{2.5} and mortality persisted. Even based on WHO guideline values, the total mortality increased by 1.1% (95% CI, 0.6–1.7%) with a 10 µg/m³ increase in PM_{2.5} at lag0-1. When we performed the stratified analyses, the point estimates of percentage increase tended to be slightly greater in adults aged 75 years or older and in spring and autumn. The PM_{2.5}-mortality association varied somewhat by region: there was a positive association in East Japan (percentage increase, 1.5%; 95% CI, 0.9–2.0%) and in Central Japan (1.7%; 95% CI, 1.1–2.3%), but no clear positive association was found in West Japan (0.5%; 95% CI, -0.5 to 1.4%).

We analyzed the association between exposure to coarse PM and non-accidental mortality in 77 cities (eFigure 3), and the results of pooling the city-specific estimates are summarized in Table 4 and eTable 3. The coarse PM concentrations at lag0-1 were positively associated with an increased risk of total mortality (percentage increase per 10 µg/m³, 2.3%; 95% CI, 1.4–3.3%) without heterogeneity (I² = 5.5%). After adjustment for exposure to PM_{2.5}, though the pooled estimate of percentage increase was attenuated, the coarse PM-mortality association was positive (percentage increase, 1.4%; 95% CI, 0.2–2.6%). In the two-

Table 3. Pooled estimates for the association between PM_{2.5} exposure and mortality in 100 Japanese cities, April 2012 to March 2015

Outcome	Lag0-1		Lag2		Lag3	
	Percentage increase for 10 µg/m ³ increase in PM _{2.5} ^a	(95% CI)	Percentage increase for 10 µg/m ³ increase in PM _{2.5} ^a	(95% CI)	Percentage increase for 10 µg/m ³ increase in PM _{2.5} ^a	(95% CI)
Total non-accidental (ICD-10: A00–R99)	1.3	(0.9 to 1.6)	0.2	(–0.2 to 0.5)	–0.3	(–0.7 to 0.1)
Cardiovascular disease (ICD-10: I00–99)	1.6	(0.8 to 2.4)	0.5	(–0.1 to 1.1)	–0.4	(–1.1 to 0.4)
Coronary heart disease (ICD-10: I20–25)	2.7	(1.0 to 4.4)	1.0	(–0.6 to 2.6)	1.0	(–0.1 to 2.2)
Stroke (ICD-10: I60–69)	1.3	(–0.2 to 2.8)	–0.3	(–1.4 to 0.8)	–0.3	(–1.5 to 0.9)
Respiratory disease (ICD-10: J00–99)	1.5	(0.7 to 2.3)	–0.3	(–1.3 to 0.8)	0.1	(–1.0 to 1.1)

CI, confidence interval; ICD, international classification of diseases; PM, particulate matter.

^aAdjusted for PM_{2.5} at lag0-1, lag2, and lag3 simultaneously, ambient temperature at lag0-1, relative humidity at lag0-1, and influenza epidemics.




Figure 1. Sensitivity and stratified analyses of the association between PM_{2.5} exposure and total non-accidental mortality in 100 Japanese cities, April 2012 to March 2015

CI, confidence interval; ICD, international classification of diseases; NO₂, nitrogen dioxide; Ox, photochemical oxidants; PM, particulate matter; SO₂, sulphur dioxide

^aAdjusted for PM_{2.5} at lag2 and lag3, ambient temperature at lag0-1, relative humidity at lag0-1, and influenza epidemics.

^bThe sum of the region was 99, because we did not include Naha, Okinawa, in West Japan due to geographical differences (see eTable 1).

pollutant model, which included both coarse PM and PM_{2.5}, the point estimates of percentage increase for the association of coarse PM with cardiovascular and respiratory mortality showed a positive direction but were not statistically significant (percentage increase for cardiovascular mortality, 1.3%; 95% CI, –0.9 to 3.5%; and percentage increase for respiratory mortality, 0.9%; 95% CI,

–1.8 to 3.8%). With regard to PM_{2.5}, adjustment for coarse PM did not substantially affect the pattern of the PM_{2.5}-mortality association (Table 4).

For coarse PM-mortality association, the elevated risk was more evident in West Japan than in East and Central Japan (eFigure 4).

DISCUSSION

Under environmental conditions, in which the average (14.6 µg/m³) and 98th percentile (37.8 µg/m³) of daily mean PM_{2.5} concentration were similar to those of the Japan Air Quality Standard (annual mean ≤15 µg/m³, daily mean ≤35 µg/m³), an overall 10 µg/m³ increase in PM_{2.5} concentrations on the day of death and the previous day was associated with a 1.3% increase in total non-accidental mortality. This positive association was observed even when we restricted to days with PM_{2.5} concentrations at lag0-1 ≤25 µg/m³. The adverse effects of PM_{2.5} in this study were consistent with the previous, 20-cities study in Japan, which observed that a 10 µg/m³ increase on the previous day was associated with a 0.88% increase in non-accidental all-cause mortality for adults aged 65 years or older,⁶ though direct comparison of the effect size of the PM_{2.5}-mortality association was difficult due to the difference in the number of cities, age category, and PM_{2.5} measurement method involved.

Our effect estimate of PM_{2.5} (a 1.3% increase in total mortality) tended to be higher than the estimates in the studies of other countries: 0.98% (95% CI, 0.75–1.22%) in 112 American cities from 1999–2005,² 0.55% (95% CI, 0.27–0.84%) in 10 Mediterranean metropolitan areas from 2001–2010,⁴ and 0.8% (95% CI, 0.3–1.2%) in a nationwide Dutch study from 2008–2009,¹⁶ and 0.22% (95% CI, 0.15–0.28%) in 272 Chinese cities from 2013–2015.³ This discrepancy may be partly explained by differences in population structure. In Japan, the proportion of older population tends to be higher than in other countries.¹⁷ As we observed, the effect estimate of PM_{2.5} tended to be higher in those 75 years old or more, than in those less than 75 years old; and older individuals are likely more vulnerable to PM exposure.¹⁸ In addition, the difference may be influenced by regional differences in the composition and sources of PM_{2.5}. Even in Japan, though a positive direction in the association between PM_{2.5} and mortality was observed in all three regions that were distinguished based on the composition and sources of PM_{2.5}, the

Table 4. Pooled estimates for non-accidental mortality associated with PM_{2.5} and coarse PM in 77 Japanese cities, April 2012 to March 2015

Outcome	PM _{2.5}		Coarse PM ^a	
	Percentage increase for 10 µg/m ³ increase at lag0-1 ^b	(95% CI)	Percentage increase for 10 µg/m ³ increase at lag0-1 ^b	(95% CI)
Single pollutant model				
Total non-accidental (ICD-10: A00–R99)	1.2	(0.8 to 1.7)	2.3	(1.4 to 3.3)
Cardiovascular disease (ICD-10: I00–99)	1.5	(0.6 to 2.4)	2.6	(0.7 to 4.4)
Coronary heart disease (ICD-10: I20–25)	1.8	(–0.2 to 3.9)	3.8	(0 to 7.7)
Stroke (ICD-10: I60–69)	1.0	(–0.6 to 2.7)	2.7	(–0.9 to 6.4)
Respiratory disease (ICD-10: J00–99)	1.5	(0.6 to 2.4)	1.4	(–0.8 to 3.7)
Two pollutant model^c				
Total non-accidental (ICD-10: A00–R99)	1.2	(0.6 to 1.7)	1.4	(0.2 to 2.6)
Cardiovascular disease (ICD-10: I00–99)	1.4	(0.3 to 2.4)	1.3	(–0.9 to 3.5)
Coronary heart disease (ICD-10: I20–25)	2.5	(0.3 to 4.6)	2.5	(–1.7 to 6.7)
Stroke (ICD-10: I60–69)	0.6	(–1.0 to 2.3)	2.9	(–0.9 to 6.9)
Respiratory disease (ICD-10: J00–99)	2.1	(1.0 to 3.2)	0.9	(–1.8 to 3.8)

CI, confidence interval; ICD, international classification of diseases; PM, particulate matter.

^aConcentrations of coarse PM were calculated by subtracting PM_{2.5} concentrations from those of suspended particulate matter.

^bAdjusted for PM_{2.5} or coarse PM at lag2 and lag3, ambient temperature at lag0-1, relative humidity at lag0-1, and influenza epidemics.

^cWe included both PM_{2.5} and coarse PM in the model.

mortality effect of PM_{2.5} appeared to vary somewhat by region. In East Japan, domestic traffic-related pollution, as a source of elemental carbon (EC), made a major contribution to PM_{2.5} concentrations^{15,19}; while sulfate (SO₄²⁻) concentrations, influenced by transboundary pollution, were lower than in West Japan.^{20,21} Both EC and SO₄²⁻ were associated with mortality, but the largest association per unit mass was observed for EC.^{14,22} Such regional differences in the composition and sources of PM_{2.5} may help interpret the apparent association in East Japan rather than in West Japan. Thus, a related health effect assessment, focused on the composition and sources of PM_{2.5},²³ is required in the future. Among the possible reasons for seasonal difference, spring and autumn tend to witness an increase in outdoor activities, and decrease in air conditioner usage, meaning increased exposure to outdoor air pollution.

We found that exposure to PM_{2.5} was positively associated with both cardiovascular and respiratory mortality, and this was consistent with the results for the PM_{2.5}-mortality association in past studies.^{2-4,16} With respect to cardiovascular disease, we found that PM_{2.5} exposure was associated with mortality due to coronary heart disease but not clearly associated with stroke mortality. The existence of acute cardiac effects due to PM_{2.5} exposure has been internationally accepted,^{24,25} and we believe that the supposition of a positive PM_{2.5}-stroke association is not far from gaining consensus. Indeed, there is steadily increasing evidence suggesting a positive association between short-term exposure to PM_{2.5} and mortality due to stroke.^{2,3,26} In Tokyo, for example, PM_{2.5} exposure was associated with an elevated risk of elderly mortality related to stroke as well as coronary heart disease.²⁷ In fact, the non-statistical elevated risk of stroke-related mortality we here observed may partially explain the fact that the acute fatality rate of coronary heart disease is higher than that of stroke.²⁸

The health effects of coarse PM are still under investigation.^{8,29} In the American multi-city study,² exposure to coarse PM, estimated as the difference between PM₁₀ and PM_{2.5} (PM_{10-2.5}), was associated with an increase in all-cause mortality; however, no such association was observed in the European multi-city study.⁴ In the present study, we used the difference between the

respective concentrations of SPM and PM_{2.5} as a surrogate marker of coarse PM, and observed a 1.4% increase in total mortality with a 10 µg/m³ increase in coarse PM at lag0-1, after adjustment for PM_{2.5}. This finding was similar to the Japanese single-city (ie, 23 wards of Tokyo) result for the association between coarse PM (defined as in our study) and all-cause mortality among adults aged 65 years or older between 2002 and 2013 (PM_{2.5}-adjusted percentage increase, 1.6%; 95% CI, 1.1–2.2%).²⁷ The clear association between coarse PM and mortality in West Japan might be partially related to the health effects of Asian dust originating in the deserts of China and Mongolia, which is mainly composed of coarse particles.³⁰⁻³² There were more days of Asian dust in West Japan than in East and Central Japan.³³ Our results did not show a clear association between coarse PM exposure and cause-specific mortality but did suggest a positive direction in the association. Coarse particles cause inflammation with deposition in the bronchial tubes, but, unlike fine particles, do not penetrate to the alveolus.^{34,35} Any significant conclusion regarding the health effects of coarse PM exposure in consideration of PM_{2.5} exposure would require further evidence.

The main strength of the present study is that it provides the first nationwide evidence of an association between mortality and short-term exposure to PM_{2.5} or coarse PM in Japan. Further, data for a large number of deaths were analyzed, and the effects of PM exposure on mortality were estimated with narrow CIs. Another strength of the study is its use of PM_{2.5} data measured using automated PM_{2.5} measuring equipment with measurement equivalence to a reference method. This enabled evaluation of the PM_{2.5}-mortality association without allowance for systematic PM_{2.5} measurement error due to the measuring equipment. On the other hand, we should acknowledge some limitations in this study. There was random measurement error in the exposure assessment, and coarse PM (difference between SPM and PM_{2.5}) was more likely to be susceptible to such error than PM_{2.5}. Since this classical error appears to be unrelated to outcome, we must interpret the results of the association between coarse PM and mortality with allowance for an underestimate on the effect. As in past environmental epidemiologic studies investigating acute

health effects of air pollution, another limitation of the study was the inevitability of exposure misclassification, giving that the results were based on the use of pollutant data measured at a single monitoring station in each city. It was difficult to estimate the biasing effect of a combination of this Berkson error and classical error on our estimates of the PM-mortality association.³⁶

In conclusion, we observed that short-term exposure to PM_{2.5} had adverse effects on total non-accidental, cardiovascular, and respiratory mortality in Japan. Coarse PM exposure also increased the risk of total mortality. Our findings suggest that we must continue monitoring for exposure to PM, including the coarse component, and continue assessing the health effects of such exposure.

ACKNOWLEDGEMENTS

The authors thank Mr Minoru Soga, Ms Satomi Takahashi, Ms Yayoi Kida, and Ms Noriko Shidai (National Institute for Environmental Studies) for data-collection and secretarial assistance.

Funding: This work was supported by the Environment Research and Technology Development Fund of the Ministry of the Environment, Japan [Number 5-1751, Akinori Takami]. The funders were not involved in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

Conflicts of interest: None declared.

APPENDIX A. SUPPLEMENTARY DATA

Supplementary data related to this article can be found at <https://doi.org/10.2188/jea.JE20180122>.

REFERENCES

- Cohen AJ, Brauer M, Burnett R, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet*. 2017;389:1907–1918.
- Zanobetti A, Schwartz J. The effect of fine and coarse particulate air pollution on mortality: a national analysis. *Environ Health Perspect*. 2009;117:898–903.
- Chen R, Yin P, Meng X, et al. Fine particulate air pollution and daily mortality. A nationwide analysis in 272 Chinese cities. *Am J Respir Crit Care Med*. 2017;196:73–81.
- Samoli E, Stafoggia M, Rodopoulou S, et al; MED-PARTICLES Study Group. Associations between fine and coarse particles and mortality in Mediterranean cities: Results from the MED-PARTICLES Project. *Environ Health Perspect*. 2013;121:932–938.
- Ministry of the Environment, Japan. Environmental quality standards in Japan - Air quality. <http://www.env.go.jp/en/air/aaq/aaq.html>. Accessed 18.09.02.
- Ueda K, Nitta H, Ono M, Takeuchi A. Estimating mortality effects of fine particulate matter in Japan: a comparison of time-series and case-crossover analyses. *J Air Waste Manag Assoc*. 2009;59:1212–1218.
- Continuous monitoring manual for atmosphere environment, 6th edition; 2010. <http://www.env.go.jp/air/osen/manual.6th/index.html>. Accessed 18.09.02.
- Adar SD, Filigrana PA, Clements N, Peel JL. Ambient coarse particulate matter and human health: a systematic review and meta-analysis. *Curr Environ Health Rep*. 2014;1:258–274.
- Ng CF, Ueda K, Takeuchi A, et al. Sociogeographic variation in the effects of heat and cold on daily mortality in Japan. *J Epidemiol*. 2014;24:15–24.
- Janes H, Sheppard L, Lumley T. Case-crossover analyses of air pollution exposure data: referent selection strategies and their implications for bias. *Epidemiology*. 2005;16:717–726.
- Di Q, Dai L, Wang Y, et al. Association of short-term exposure to air pollution with mortality in older adults. *JAMA*. 2017;318:2446–2456.
- Gasparri A, Guo Y, Hashizume M, et al. Mortality risk attributable to high and low ambient temperature: a multicountry observational study. *Lancet*. 2015;386:369–375.
- World Health Organization. Ambient (outdoor) air quality and health 2016. <http://www.who.int/mediacentre/factsheets/fs313/en/>. Accessed 18.09.02.
- Achilleos S, Kioumourtzoglou MA, Wu CD, Schwartz JD, Koutrakis P, Papatheodorou SI. Acute effects of fine particulate matter constituents on mortality: a systematic review and meta-regression analysis. *Environ Int*. 2017;109:89–100.
- Morino Y, Ueda K, Takami A, et al. Sensitivities of simulated source contributions and health impacts of PM_{2.5} to aerosol models. *Environ Sci Technol*. 2017;51:14273–14282.
- Janssen NA, Fischer P, Marra M, Ameling C, Cassee FR. Short-term effects of PM_{2.5}, PM₁₀ and PM_{2.5-10} on daily mortality in The Netherlands. *Sci Total Environ*. 2013;463–464:20–26.
- United Nations, Department of Economic and Social Affairs, Population Division, 2015. World Population Prospects: The 2015 Revision, World Population 2015 Wallchart. <https://esa.un.org/unpd/wpp/Publications/>. Accessed 18.09.02.
- Bell ML, Zanobetti A, Dominici F. Evidence on vulnerability and susceptibility to health risks associated with short-term exposure to particulate matter: a systematic review and meta-analysis. *Am J Epidemiol*. 2013;178:865–876.
- Ikeda K, Yamaji K, Kanaya Y, et al. Source region attribution of PM_{2.5} mass concentrations over Japan. *Geochem J*. 2015;49:185–194.
- Morino Y, Nagashima T, Sugata S, et al. Verification of chemical transport models for PM_{2.5} chemical composition using simultaneous measurement data over Japan. *Aerosol Air Qual Res*. 2015;15:2009–2023.
- Aikawa M, Ohara T, Hiraki T, et al. Significant geographic gradients in particulate sulfate over Japan determined from multiple-site measurements and a chemical transport model: impacts of transboundary pollution from the Asian continent. *Atmos Environ*. 2010;44:381–391.
- Atkinson RW, Mills IC, Walton HA, Anderson HR. Fine particle components and health—a systematic review and meta-analysis of epidemiological time series studies of daily mortality and hospital admissions. *J Expo Sci Environ Epidemiol*. 2015;25:208–214.
- Ueda K, Yamagami M, Ikemori F, Hisatsune K, Nitta H. Associations between fine particulate matter components and daily mortality in Nagoya, Japan. *J Epidemiol*. 2016;26:249–257.
- Newby DE, Mannucci PM, Tell GS, et al; ESC Working Group on Thrombosis, European Association for Cardiovascular Prevention and Rehabilitation; ESC Heart Failure Association. Expert position paper on air pollution and cardiovascular disease. *Eur Heart J*. 2015;36:83–93b.
- Mustafić H, Jabre P, Caussin C, et al. Main air pollutants and myocardial infarction: a systematic review and meta-analysis. *JAMA*. 2012;307:713–721.
- Shah AS, Lee KK, McAllister DA, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *BMJ*. 2015;350:h1295.
- Yorifuji T, Kashima S, Doi H. Associations of acute exposure to fine and coarse particulate matter and mortality among older people in Tokyo, Japan. *Sci Total Environ*. 2016;542:354–359.
- Rumana N, Kita Y, Turin TC, et al. Acute case-fatality rates of stroke and acute myocardial infarction in a Japanese population: Takashima stroke and AMI registry, 1989–2005. *Int J Stroke*. 2014;9(Suppl A100):69–75.
- U.S. Environmental Protection Agency. Integrated science assessment for particulate matter. Research Triangle Park, NC; 2009.

30. Kamouchi M, Ueda K, Ago T, Nitta H, Kitazono T; Fukuoka Stroke Registry Investigators. Relationship between Asian dust and ischemic stroke: a time-stratified case-crossover study. *Stroke*. 2012; 43:3085–3087.
31. Kashima S, Yorifuji T, Tsuda T, Eboshida A. Asian dust and daily all-cause or cause-specific mortality in western Japan. *Occup Environ Med*. 2012;69:908–915.
32. Kojima S, Michikawa T, Ueda K, et al. Asian dust exposure triggers acute myocardial infarction. *Eur Heart J*. 2017;38:3202–3208.
33. Nakamura T, Hashizume M, Ueda K, et al. The relationship between Asian dust events and out-of-hospital cardiac arrests in Japan. *J Epidemiol*. 2015;25:289–296.
34. Lippmann M, Chen LC. Health effects of concentrated ambient air particulate matter (CAPs) and its components. *Crit Rev Toxicol*. 2009;39:865–913.
35. Graff DW, Cascio WE, Rappold A, Zhou H, Huang YC, Devlin RB. Exposure to concentrated coarse air pollution particles causes mild cardiopulmonary effects in healthy young adults. *Environ Health Perspect*. 2009;117:1089–1094.
36. Armstrong BG. Effect of measurement error on epidemiological studies of environmental and occupational exposures. *Occup Environ Med*. 1998;55:651–656.