

ERRATUM

Open Access



# Erratum to: The effect of Katsura-uri (Japanese pickling melon, *Cucumis melo* var. *conomon*) and its derived ingredient methylthioacetic acid on energy metabolism during aerobic exercise

Wataru Aoi<sup>1\*</sup>, Kazuya Takeda<sup>2</sup>, Azusa Sasaki<sup>2</sup>, Yuki Hasegawa<sup>2</sup>, Yasushi Nakamura<sup>2</sup>, Eun Young Park<sup>2</sup>, Kenji Sato<sup>3</sup>, Masayo Iwasa<sup>1</sup>, Airi Nakayama<sup>1</sup>, Mizuki Minamikawa<sup>4</sup>, Yukiko Kobayashi<sup>4</sup>, Koji Shiota<sup>5</sup> and Noboru Suetome<sup>5</sup>

## Erratum to: SpringerPlus (2015) 4:377 DOI 10.1186/s40064-015-1144-y

It has come to our attention that during production of the original article, an error was introduced into Table 1 during copyediting. The corrected Table 1 can be found below. The publisher apologises for inconvenience caused.

### Author details

<sup>1</sup> Laboratory of Health Science, Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto 606-8522, Japan. <sup>2</sup> Laboratory of Food Science, Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto 606-8522, Japan. <sup>3</sup> Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan. <sup>4</sup> Laboratory of Nutrition Science, Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto 606-8522, Japan. <sup>5</sup> Horticultural Division, Kyoto Prefectural Agriculture, Forestry and Fisheries Technology Center, Kameoka 621-0806, Japan.

**Table 1 Blood metabolic parameters in mice**

	Sedentary	Exercise		
		Control	MTA-25	MTA-250
Blood glucose (mM)	6.6 ± 0.7	7.8 ± 1.4	7.5 ± 2.0	6.7 ± 1.0
Plasma NEFA (μEq/L)	880 ± 281	916 ± 360	883 ± 529	889 ± 268
Blood ammonia (μM)	110 ± 43	135 ± 37	125 ± 68	107 ± 33
Blood lactate (mM)	–	3.2 ± 1.2	2.5 ± 0.7	2.1 ± 0.3 <sup>#</sup>

Values are presented as mean ± SD. Control, exercise group administered water; MTA-25, exercise group receiving 25 ppm MTA supplementation; MTA-250, exercise group receiving 250 ppm MTA supplementation

MTA methylthioacetic acid

<sup>#</sup> P < 0.05 vs. control

Published online: 21 September 2015

The online version of the original article can be found under doi:[10.1186/s40064-015-1144-y](https://doi.org/10.1186/s40064-015-1144-y).

\*Correspondence: waoui@kpu.ac.jp

<sup>1</sup> Laboratory of Health Science, Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto 606-8522, Japan

Full list of author information is available at the end of the article