



MEETING ABSTRACT

Open Access

Selenium and gastrointestinal cancers risk

Marcin Lener^{1*}, Anna Jakubowska¹, Anna Wiechowska-Kozłowska², Józef Kładny³, Magdalena Muszyńska¹, Grzegorz Sukiennicki¹, Jan Lubiński¹

From Annual Conference on Hereditary Cancers 2012
Szczecin, Poland. 30-31 August 2012

Research suggests that selenium may influence the behavior of the cancer risk in two ways. As an antioxidant, selenium helps to protect the body against free radicals. Selenium may also prevent or slow tumor growth, as some breakdown products of selenium can inhibit tumor growth by enhancing immune cell activity and inhibition of tumor blood vessel development.

Aim

The aim of this study was to determine the level of selenium in blood serum as a potential marker of risk for cancers of the colon, stomach or pancreas.

Material and methods

The research material was a total of 110 samples of blood serum from people with cancer, diagnosed and confirmed in one of the organs: colon (67 cases), pancreas (30 cases) or stomach (13 cases) and 110 samples of blood serum derived from healthy individuals representing paired control group. The criteria adopted for pairing included: gender, year of birth (+/- 3 years),

history of the occurrence of cancers in the family among first degree relatives and smoking status expressed in pack-years.

Selenium concentration in blood serum was determined using inductively coupled plasma mass spectrometry (ICP-MS) – Perkin Elmer. Validation: SeronormTM, Nycomed Pharma AS, Oslo, Norway. The measurement accuracy was +/- 5% µg Se/l.

Results

Association between Se concentration and frequency of cancers in quartiles are presented in table 1. Statistical analyses are summarized in table 2.

Conclusions

1. There is a very strong correlation between the level of selenium in serum and the risk of gastrointestinal cancers (pancreas, colon, stomach) in the Polish population.

2. Due to the risk of gastrointestinal cancers evaluated most people from the Polish population should increase

Table 1 Association between Se plasma concentration and risk of cancers analyzed.

Cancer site	Quartile	Se concentration range (µg/l)	No. of Cases/ Controls
Pancreas	I	25,69-50,09	15/0
	II	50,72-65,58	9/6
	III	66,34-73,30	6/9
	IV	74,07-113,89	0/15
Colorectal and stomach	I	15,92-56,75	35/5
	II	57,3-67,91	22/18
	III	68,13-78,07	12/28
	IV	78,38-155,72	11/29

¹Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland

Full list of author information is available at the end of the article

Table 2 Results of statistical analyses of cancer site depending on Se concentration.

Cancer site	Quartiles compared	Se concentration range ($\mu\text{g/l}$)	Cases/controls in compared groups	Fisher's Exact Test		
				P	OR	CI
Pancreas	I vs II	25,69-50,09 vs 50,72-65,58	15/0 vs 9/6	0,017	21,2	1,07-421,11
	I vs III	25,69-50,09 vs 66,34-73,30	15/0 vs 6/9	0,0007	45,3	2,2-899,53
	I vs IV	25,69-50,09 vs 74,07-113,89	15/0 vs 0/15	<0,0001	961	7,9-51,617
Colorectal and stomach	I vs II	15,92-56,75 vs 57,3-67,91	(5)*35/5 vs (4)*22/18	0,0026	5,7	1,86-17,66
	I vs III	15,92-56,75 vs 68,13-78,07	(5)*35/5 vs (1)*12/28	<0,0001	16,33	5,14-51,89
	I vs IV	15,92-56,75 vs 78,38-155,72	(5)*35/5 vs (3)*11/29	<0,0001	18,46	5,76-59,25

*stomach cancer cases

the level of selenium in serum to approximately 80-100 $\mu\text{g/l}$.

3. Prospective studies can elucidate:

a) the use of selenium measurements as markers of risk of above cancers,

b) possibility of lowering risk of the cancers of the colon, pancreas and stomach by supplementation of diet with selenium.

4. Assessment of the level of selenium may increase the effectiveness of many screening programs of gastrointestinal cancers, for example, the cost of detecting asymptomatic colorectal cancer by colonoscopy can be reduced dozen times.

Author details

¹Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland. ²Laboratory of Endoscopy, Division of Health Care Ministry of Internal Affairs and Administration, Szczecin, Poland. ³Department of General and Oncological Surgery, Pomeranian Medical University, Szczecin, Poland.

Published: 10 December 2012

doi:10.1186/1897-4287-10-S4-A3

Cite this article as: Lener et al.: Selenium and gastrointestinal cancers risk. *Hereditary Cancer in Clinical Practice* 2012 **10**(Suppl 4):A3.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

