

Research article

Open Access

Expression of cytokine and chemokine mRNA and secretion of tumor necrosis factor- α by gallbladder epithelial cells: Response to bacterial lipopolysaccharides

Christopher E Savard¹, Thane A Blinman², Ho-Soon Choi¹, Sung-Koo Lee¹, Stephen J Pandol² and Sum P Lee*¹

Address: ¹Department of Medicine, University of Washington and VA Puget Sound Health Care System, Seattle, Washington USA and ²Department of Medicine, University of California at Los Angeles and West Los Angeles VA Medical Center, Los Angeles, California USA

E-mail: Christopher E Savard - cesavard@u.washington.edu; Thane A Blinman - blinman@email.chop.edu; Ho-Soon Choi - choihs@hanyang.ac.kr; Sung-Koo Lee - sklee@amc.seoul.kr; Stephen J Pandol - Stephen.Pandol@med.va.gov; Sum P Lee* - splee@u.washington.edu

*Corresponding author

Published: 11 October 2002

Received: 13 August 2002

BMC Gastroenterology 2002, **2**:23

Accepted: 11 October 2002

This article is available from: <http://www.biomedcentral.com/1471-230X/2/23>

© 2002 Savard et al; licensee BioMed Central Ltd. This article is published in Open Access: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: In addition to immune cells, many other cell types are known to produce cytokines. Cultured normal mouse gallbladder epithelial cells, used as a model system for gallbladder epithelium, were examined for their ability to express the mRNA of various cytokines and chemokines in response to bacterial lipopolysaccharide. The synthesis and secretion of the tumor necrosis factor- α (TNF- α) protein by these cells was also measured.

Results: Untreated mouse gallbladder cells expressed mRNA for TNF- α , RANTES, and macrophage inflammatory protein-2 (MIP-2). Upon treatment with lipopolysaccharide, these cells now produced mRNA for Interleukin-1 β (IL-1 β), IL-6, monocyte chemoattractant protein-1 (MCP-1), and showed increased expression of TNF- α and MIP-2 mRNA. Untreated mouse gallbladder cells did not synthesize TNF- α protein; however, they did synthesize and secrete TNF- α upon treatment with lipopolysaccharide.

Methods: Cells were treated with lipopolysaccharides from 3 strains of bacteria. Qualitative and semi-quantitative RT-PCR, using cytokine or chemokine-specific primers, was used to measure mRNA levels of TNF α , IL-1 β , IL-6, IL-10, KC, RANTES, MCP-1, and MIP-2. TNF- α protein was measured by immunoassays.

Conclusion: This research demonstrates that gallbladder epithelial cells in response to lipopolysaccharide exposure can alter their cytokine and chemokine RNA expression pattern and can synthesize and secrete TNF α protein. This suggests a mechanism whereby gallbladder epithelial cells in vivo may mediate gallbladder secretory function, inflammation and diseases in an autocrine/paracrine fashion by producing and secreting cytokines and/or chemokines during sepsis.

Background

The biliary system is more than a passive conduit transporting bile from the liver to the intestines. Hepatic bile is actively modified by the epithelial cells that line the bile ducts and gallbladder. Biliary epithelial function is partially under the control of secretin and vasoactive intestinal peptide [1]. Other extracellular agonists that influence biliary function include neuropeptides [2,3], ATP [4,5], and bile salts [6]. In addition to the well-documented role that cytokines play in the immune response and inflammation [7,8], cytokines also modify the functioning of epithelial cells. For instance, tumor necrosis factor- α (TNF- α) can modify intracellular signal transduction and regulate ionic channel activities in various epithelial cells [9]. A few studies have shown that cytokines can alter biliary epithelial cell function [10,11].

Biliary epithelial cells are capable of producing a great variety of physiologically active compounds. For example, it has been shown that biliary epithelial cells synthesize nitric oxide, a strong vasodilator [12], and endothelin-1 [13], a potent vasoconstrictor. Various cytokines are also produced by biliary epithelial cells [14]. As biliary epithelial cells are capable of both producing and responding to cytokines, these cells may use cytokines for the autocrine/paracrine regulation of their own function as well as that of other cells.

Bile in the biliary ducts and the gallbladder often contains bacteria and the bacterial product lipopolysaccharide (LPS). This may be due to ascending infection from the intestine or descending infection via portal venous blood [15,16]. Bacterial infection usually leads to activation of the immune system followed by inflammation. While inflammation is an important element in the defense against infection, it can also contribute to cholestatic biliary diseases and cholelithiasis. Therefore, the control of the immune and inflammatory response of biliary tissue to bacteria is important in determining disease progres-

sion. While the biliary epithelium can act as a physical barrier to limit bacterial infection of the underlying tissue [16], its physiological response to bacteria and bacterial products is not well characterized. Like pulmonary and intestinal epithelial cells, biliary epithelial cells might respond to bacteria in bile by producing cytokines and other factors. To test this hypothesis, we used cultured normal mouse gallbladder epithelial cells (MGBE) as an *in vitro* model system to examine the ability of gallbladder epithelium to produce mRNA for various cytokines and chemokines. The synthesis and secretion of TNF- α protein was also measured. These cells demonstrated distinct changes in their cytokine and chemokine mRNA expression profile and TNF- α protein production in response to LPS.

Methods

Materials

Analytical grade chemicals and tissue culture supplies were obtained from Sigma Chemical Co. (St. Louis, MO), except where noted. Vitrogen (bovine dermal collagen) was purchased from Collagen Corp. (Palo Alto, CA). Transwell inserts (24 mm diameter, 3 μ m pore size) were purchased from Corning Corp. (Acton, MA).

Cell culture

MGBE were grown as previously described [17]. In brief, MGBE were grown on Vitrogen-coated Transwell inserts suspended above human gallbladder myo-fibroblast feeder cells. The cells were fed with Eagle's minimum essential medium containing 10% fetal bovine serum, 2 mM L-glutamine, 20 mM HEPES, 100 IU/mL penicillin, 100 μ g/mL streptomycin, plus Insulin-Transferrin-Sodium Selenite Media Supplement, MEM-vitamins, and MEM-non-essential amino acid solution (Sigma Chemical Co.). Confluent MGBE have well-differentiated morphology and high transepithelial electrical resistance (163 Ω /cm²) [17]. Prior to each experiment, Transwell inserts with confluent MGBE were transferred to new plates without feeder cells.

Table 1: Murine specific primer sequences used in PCR

mRNA	Forward Primer	Reverse Primer	Expected product size (bp)	GenBank accession number
IL-1 β	5'-gtgtgatgttcccattagacagc	5'-tgagaggtgctgatgtaccagtt	383	
IL-6	5'-gaaatctgctctggtcttctgg	5'-ttttctgaccacagtgaggaatg	357	M26744
IL-10	5'-tcagtcaagtgaagactttcttc	5'-gcagttgatgaagatgtcaaactc	358	X60675
TNF α	5'-tgaacttcggggatgatcggtc	5'-agccttgcctctgaagagaac	295	X66539
KC (CXCL1)	5'-ttgaaggtgatgccgaccag	5'-cccagactctcatctctcc	205	
RANTES (CCL5)	5'-ccccatagggctcgacacca	5'-ctagctcatctcctaatagttgat	205	
MCP-1 (CCL2)	5'-cactatgcaggtctctgtcacg	5'-gatctcaacttggtctggtcca	294	M57441
MIP-2 (CXCL2)	5'-cagtgcagctgcgctgtccaatg	5'-cagttagccttgcctttgttcag	210	X65647
GAPDH	5'-atgtcagatccacaacggatacat	5'-actccctcaagattgtcagcaat	307	
ARP	5'-gttgaacatctccccctcttc	5'-atgtcctcatcggattcctcc	402	Z29530




Figure 1
RT-PCR for cytokine mRNAs Treated MGBE were exposed to 100 $\mu\text{g/mL}$ *E. coli* LPS on the apical surface for two hours. GAPDH was used as an internal control to standardize starting mRNA quantity. This experiment was repeated twice with equivalent results.

RT-PCR assay of cytokine and chemokine mRNA

Confluent MGBE were fed serum-free medium. Cells were treated with LPS from *E. coli* (serotype 0111:B4) or *Pseudomonas aeruginosa* (serotype 10) at 100 $\mu\text{g/mL}$, added to the apical compartment of the insert. After two hours, the cells were harvested by trypsinization, and total RNA was extracted using TRIzol reagent (Gibco BRL, Gaithersburg, MD). Five μg of total RNA was reverse-transcribed according to the manufacturer's protocol (Superscript II Preamplification System, Gibco BRL) using oligo (dT) as primer. cDNA prepared from 0.5 μg RNA was subjected to PCR using murine gene-specific primers for 8 different cytokines and chemokines (Table 1) as previously detailed [18]. All of the primer pairs spanned at least one intron in the corresponding genomic DNA. Positive RNA controls were performed previously to confirm specificity of primer pairs (data not shown). Negative controls were performed by omitting the RT step or the cDNA template from PCR amplification. For qualitative PCR, target sequences were amplified at 56°C for 25 cycles; for semi-quantitative PCR, target sequences for TNF- α , IL-1 β , and MIP-2 (CXCL2) were amplified at 56°C between 22 and 32 cycles in order to yield visible products within the linear amplification range. PCR products were separated by electrophoresis on a 2% agarose gel and stained with ethidium bromide. All RT-PCR bands at the expected size were also directly sequenced to confirm their identity. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or acidic ribosomal phosphoprotein (ARP) were used as internal controls.

TNF- α Immunoassay

Confluent MGBE were fed serum-free medium, with or without LPS added to the apical side (LPS concentration = 2 or 100 $\mu\text{g/mL}$). Medium from the apical and basolateral compartment was collected separately at measured intervals and immediately frozen at -70°C. The cells were lysed with 1% NP-40 buffer (150 mM NaCl, 50 mM Tris, pH 8.0) containing 100 $\mu\text{g/mL}$ phenylmethylsulfonyl fluoride and 1 $\mu\text{g/mL}$ aprotinin, sonicated for 30 seconds, then frozen. TNF- α was measured in cell lysate and undiluted medium using the Mouse TNF- α Immunoassay Kit from R&D Systems (Minneapolis, MN). Statistical analysis on the TNF- α immunoassay data was performed using Student's t-test. Cell viability was assessed by measuring lactate dehydrogenase release into the medium, as previously described [6]. Changes in transepithelial permeability was measured using ^3H -mannitol as a small molecular weight probe. Tritiated mannitol (2 μCi) with cold mannitol (0.15 mM) was added to the apical medium and samples of the basolateral medium harvested and counted at 1, 2, 4, and 24 hours after LPS treatment.

Results

RT-PCR

Using cytokine/chemokine-specific primers with RT-PCR, untreated MGBE showed distinct bands representing mRNA of TNF- α , RANTES (Regulated on activation, normal T cell expressed and secreted) (CCL5), and macrophage inflammatory protein-2 (MIP-2/CXCL2) (Figure 1). Upon apical treatment with LPS from *E. coli* for two hours, these cells additionally produced mRNA for interleukin-1 β (IL-1 β), IL-6, and monocyte chemoattractant protein-1 (MCP-1/CCL2). There was no detectable expression of IL-10 or KC (murine GRO- α /CXCL1) mRNA before or after LPS treatment. Semi-quantitative RT-PCR using specific primers for TNF- α , MIP-2 (CXCL2) and IL-1 β showed that LPS treatment caused the amount of TNF- α mRNA to slightly increase compared to the already abundant levels found in untreated cells (figure 2 and 3). In contrast, there was a substantial increase in the expression of MIP-2 (CXCL2) mRNA and IL-1 β mRNA following LPS treatment compared to control. Similar responses were seen when the cells were treated with LPS from *P. aeruginosa* (Figure 2 and 3). All visualized PCR product bands were of the expected size and nucleotide sequence confirming proper identification. Although the RANTES (CCL5) sample had extra bands of unknown origin, the band at the expected size of 205 bp had a nucleotide sequence that matched RANTES.

TNF- α Immunoassay

The immunoassay experiments showed no detectable TNF- α in untreated MGBE or their surrounding media after 5 hours of incubation (limit of detection = 5.1 pg/mL). In contrast, following treatment with 100 $\mu\text{g/mL}$ *E. coli*




Figure 2
Semi-quantitative RT-PCR Semi-quantitative RT-PCR products from cytokine mRNA using MIP-2, TNF- α , and IL-1 β specific primers. C is untreated control cells. Treated cells were exposed to 100 μ g/mL LPS from either *E. coli* or *P. aeruginosa* on the apical surface for two hours. ARP was used as an internal control to standardize starting mRNA quantity. PCR cycle number used was between 22 for ARP and 31 for TNF- α .

LPS the concentration of TNF- α protein in the cells was 33.6 pg per 10^6 cells in one hour (Figure 4). Intracellular TNF- α concentration reached a peak two hours after LPS treatment with a subsequent decrease by five hours. After two and five hours of LPS treatment, there was measurable release of TNF- α into the media of both the apical and basolateral compartments (figure 5). Three different species of LPS (*E. coli*, *P. aeruginosa*, and *Klebsiella pneumoniae*) stimulated TNF- α release in a similar fashion (figure 6). Treatment with 2 μ g/mL *E. coli* LPS showed comparable results. None of the treatments increased lactate dehydrogenase in the medium compared with controls showing




Figure 3
Densitometry analysis of semi-quantitative RT-PCR results Densitometry of PCR bands for MIP-2, TNF- α , and IL-1 β (from figure 2) based on % of ARP control bands using ImageJ Program (Version 1.20s).




Figure 4
Intracellular TNF- α following LPS treatment Intracellular TNF- α concentration of MGBE over time following apical exposure to 100 μ g/mL *E. coli* LPS was measured by immunoassay. Values are the means \pm standard error of results from three to four assays.




Figure 5
TNF-α secretion into medium Release of TNF-α into the apical and basolateral medium by cultured MGBE exposed to 100 μg/mL *E. coli* LPS over time. TNF-α secretion was calculated as the total pg amount of TNF-α present in the apical or basolateral medium. Values are the means of triplicates ± standard error from a single assay. This assay was repeated twice more with similar results. Statistical differences between the concentration of TNF-α in the basal and apical medium were measured by paired Student's t-test (* = p < 0.01).

that the LPS treatments did not alter epithelial cell integrity (data not shown). LPS treatment also had no effect on the transepithelial permeability of these cells for up to 24 hours as measured by ³H-mannitol diffusion experiments when compared to controls (data not shown).

Discussion

Bacteria are found in the bile of the gallbladder and bile ducts [15,16], the presence of which may have relevance to clinical diseases such as stone formation [19–21], primary sclerosing cholangitis [22], and cholestasis [23,24]. Direct exposure of sub-epithelial tissue to LPS from gram-negative bacteria causes release of cytokines by resident leukocytes and activation of macrophages [25]. The biliary fluid of patients with cholangitis has increased concentrations of the inflammatory cytokines TNF-α and IL-6 suggesting that bacterial infection leads to an increase in cytokines released into the bile [26]. Adding LPS to the bile of guinea pig gallbladders *in vivo* led to increased IL-1 concentration in the bile [27]. What role biliary epithelial cells play in the response to bacteria in bile is not known.

While most cytokine research has focused on cells of the immune system, other cell types produce cytokines including the epithelial cells of the lung [28,29], retina [30], and intestines [31,32]. Among the few studies involving the biliary system, it was demonstrated that cultured human intrahepatic duct epithelial cells produce functional IL-8 and MCP-1 (CCL2) [14,33]. Isolated human intrahe-




Figure 6
Effects of LPS from different bacteria on TNF-α secretion Release of TNF-α into the apical and basolateral medium by cultured MGBE exposed to 100 μg/mL LPS from various bacteria for two hours. TNF-α was measured by immunoassay and calculated as the total pg present in the apical and basolateral medium. Each treatment was done in triplicate and assay repeated twice more with *E. coli* LPS and once more with *P. aeruginosa* and *K. pneumoniae* LPS. Values represent the mean of each treatment (n = 9 for *E. coli* LPS and n = 6 for *P. aeruginosa* and *K. pneumoniae* LPS) ± standard error. There was a statistical difference between the amount of TNF-α in the basal versus apical medium as measured by paired Student's t-test (* = p < 0.01).

patic epithelial ducts expressed immunoreactive IL-6 when treated with IL-1β or phorbol myristate acetate [34]. Intracellular TNF-α mRNA and protein was detected in intrahepatic biliary duct cells in tissue sections [35], but not in cultured intrahepatic epithelial cells; therefore, there was no clear demonstration that biliary epithelium secretes TNF-α. Little research has focused on the ability of the epithelial cells of the extrahepatic biliary system to produce cytokines or chemokines.

As described in this paper, MGBE were used as an *in vitro* model system of gallbladder epithelium to measure the endogenous production of cytokines and chemokines by extrahepatic biliary epithelial cells. These cultured cells consist exclusively of normal, non-neoplastic epithelial cells [17], without other cell types such as leukocytes ordinarily found in biliary tissues. The experiments showed that unstimulated MGBE express mRNA for TNF-α, MIP-2, and RANTES (CCL5). After treating the apical surface of the cells with LPS, the mRNA levels for TNF-α and MIP-2 (CXCL2) increased, while mRNA for IL-1β, IL-6, and MCP-1 (CCL2) appeared for the first time. Therefore, cytokines (TNF-α, IL-1β, and IL-6), a C-C type chemokine (MCP-1/CCL2), and a C-X-C type chemokine (MIP-2/CXCL2) are all up regulated at the RNA level by bacterial product interaction with the epithelial cells. The effects of LPS treatment for two hours on cytokine and chemokine mRNA expression clearly suggests that biliary epithelial

cells can modify their own cytokine and chemokine production after bacterial challenge at the mRNA level. As with all *in vitro* cell systems there is the possibility that the cells in culture may have phenotypic changes to their cytokine profile compared to *in vivo* cells. In other parameters tested (i.e., morphology, mucin secretion, etc.), these cells retain the normal phenotype of cells *in vivo*.

Measurements of TNF- α protein showed that MGBE synthesize and secrete TNF- α when exposed to LPS, while untreated cells do not. This demonstrates that gallbladder epithelial cells themselves secrete a cytokine in response to bacterial products. The results from RT-PCR (showing a relatively high TNF- α mRNA expression levels in untreated cells which is only slightly increased by LPS treatment) along with the immunoassay results suggest that TNF- α mRNA is constantly present in these cells, but is not translated into protein until certain triggers are provided. This is consistent with the post-transcriptional regulation of TNF- α observed in other cells [36]. In contrast, the mRNA of other factors, such as IL-1 β , IL-6, and MCP-1 (CCL2) are not synthesized until the cells are stimulated, suggesting that their production is under transcriptional control. The protein expression of these other cytokines and chemokines along with the precise nature of the interaction between LPS and cell surface receptors warrants further study.

While the exact role that gallbladder epithelial cells play in inflammatory and infectious diseases has yet to be determined, our research supports the hypothesis that gallbladder epithelium have some innate cytokine-producing capabilities in response to the presence of bacteria on their apical surface. The accumulation of TNF- α in the gallbladder bile of patients with cholangitis may be due in part to the apical release of TNF- α by biliary epithelial cells. In contrast, basolateral secretion of TNF- α could lead to a focal accumulation of TNF- α in the sub-mucosa of biliary tissue. Sufficient cytokine secretion could then stimulate the leukocytes in the sub-epithelial tissue, leading to inflammation of the tissue, without the need for the bacteria to pass through the epithelial barrier. The production of cytokines by gallbladder epithelial cells following LPS exposure demonstrates that biliary epithelial cells may be an active partner with the immune system in the detection of and defense against bacteria in bile.

During sepsis, biliary epithelial cells could mediate biliary secretory function through cytokine and chemokine production. Dog gallbladder epithelial cells demonstrated increased mucin secretion following TNF- α treatment [10]. In addition, exogenous TNF- α altered the absorptive properties of cultured human gallbladder epithelial cells [37]. It is therefore reasonable to suggest that TNF- α produced by gallbladder epithelial cells in response to LPS

could affect cellular function in an autocrine manner. Since mucin secretion and gallbladder transport properties play vital roles in cholelithiasis, changes in these parameters caused by endogenous cytokines and chemokines may promote stone formation. Epithelial-derived cytokines and chemokines also may alter biliary function leading to cholestasis.

Conclusions

Gallbladder epithelial cells demonstrated distinct patterns of expression of mRNA for various cytokines and chemokines, and responded to treatment with LPS by altering their expression of certain cytokine and chemokine RNA and increasing TNF- α protein synthesis and secretion. The ability of biliary epithelial cell to control endogenous cytokine and chemokine production has important implications for biliary inflammation and diseases, such as cholangitis, cholestasis, and cholelithiasis.

Competing interests

None declared.

Authors' contributions

CES participated in the design and coordination of the study, performed the immunoassay studies, and drafted the manuscript. TAB performed the RT-PCR studies. H-SC assisted in the immunoassay studies. S-KL initiated the immunoassay studies. SJP participated in the design of the RT-PCR studies. SPL conceived of the study, and participated in its design and coordination.

Abbreviations

IL (Interleukin), LPS (Lipopolysaccharides), MCP-1 (Monocyte chemoattractant protein-1), MGBE (Mouse gallbladder epithelial cells), MIP-2 (Macrophage inflammatory protein-2), RANTES (Regulated on activation, normal T cell expressed and secreted), TNF- α (Tumor Necrosis Factor-alpha).

Acknowledgements

Thanks to Ilya Gukovsky for assistance with the cytokine mRNA assays and Dr. Don Ostrow for helpful manuscript assistance. This work was supported by a grant from the NIH (DK50246) and in part by the Medical Research Service of the Department of Veterans Affairs.

References

1. Jansson R, Svanvik J: **Effects of intravenous secretin and cholecystokinin on gallbladder net water absorption and motility in the cat.** *Gastroenterology* 1977, **72**:639-643
2. Cho WK, Mennone A, Rydberg SA, Boyer JL: **Bombesin stimulates bicarbonate secretion from rat cholangiocytes: implications for neural regulation of bile secretion.** *Gastroenterology* 1997, **113**:311-321
3. Sand J, Tainio H, Nordback I: **Neuropeptides in pig sphincter of Oddi, bile duct, gallbladder, and duodenum.** *Dig Dis Sci* 1993, **38**:694-700
4. Elsing C, Kassner A, Stremmel W: **Sodium, hydrogen antiporter activation by extracellular adenosine triphosphate in biliary epithelial cells.** *Gastroenterology* 1996, **111**:1321-1332

5. McGill JM, Basavappa S, Mangel AW, Shimokura GH, Middleton JP, Fitz JG: **Adenosine triphosphate activates ion permeabilities in biliary epithelial cells.** *Gastroenterology* 1994, **107**:236-243
6. Klinkspoor JH, Tytgat GN, Lee SP, Groen AK: **Mechanism of bile salt-induced mucin secretion by cultured dog gallbladder epithelial cells.** *Biochemistry Journal* 1996, **316**:873-877
7. Ali H, Haribabu B, Richardson R, Snyderman R: **Mechanisms of inflammation and leukocyte activation.** *Med Clin North Am* 1997, **81**:1-28
8. Luster AD: **Chemokines – chemotactic cytokines that mediate inflammation.** *New Engl J Med* 1998, **338**:436-445
9. Papadakis KA, Targan SR: **Tumor necrosis factor: biology and therapeutic inhibitors.** *Gastroenterology* 2000, **119**:1148-1157
10. Choi J, Klinkspoor JH, Yoshida T, Lee SP: **Lipopolysaccharide from Escherichia coli stimulates mucin secretion by cultured dog gallbladder epithelial cells.** *Hepatology* 1999, **29**:1352-1357
11. Spirli C, Nathanson MH, Fiorotto R, Duner E, Denson LA, Sanz JM, Di Virgilio F, Okolicsanyi L, Casagrande F, Strazzabosco M: **Proinflammatory cytokines inhibit secretion in rat bile duct epithelium.** *Gastroenterology* 2001, **121**:156-169
12. Salomons H, Keaveny AP, Henihan R, Offner G, Sengupta A, Lamorte WV, Afdhal NH: **Nitric oxide and gallbladder motility in prairie dogs.** *Am J Physiol* 1997, **272**:G770-778
13. Housset C, Carayon A, Housset B, Legendre C, Hannoun L, Poupon R: **Endothelin-1 secretion by human gallbladder cells in primary culture.** *Lab Invest* 1993, **69**:750-755
14. Morland CM, Fear J, Joplin R, James DH: **Inflammatory cytokines stimulate human biliary epithelial cells to express interleukin-8 and monocyte chemotactic protein-1.** *Biochem Soc Trans* 1997, **25**:232s
15. Luk WK, Liu CL, Yuen KY, Wong SSY, Woo PCY, Fan ST: **Biliary tract infection due to bile-soluble bacteria: An intriguing paradox.** *Clin Infect Dis* 1998, **26**:1010-1012
16. Sung JY, Costerton JW, Shaffer EA: **Defense system in the biliary tract against bacterial infection.** *Dig Dis Sci* 1992, **37**:689-696
17. Kuver R, Savard C, Nguyen TD, Osborne WR, Lee SP: **Isolation and long-term culture of gallbladder epithelial cells from wild-type and CF mice.** *In Vitro Cellular & Developmental Biology Animal* 1997, **33**:104-109
18. Pandolfi SJ, Periskic S, Gukovsky I, Zaninovic V, Jung Y, Zong Y, Solomon TE, Gukovskaya AS, Tsukamoto H: **Ethanol diet increases the sensitivity of rats to pancreatitis induced by cholecystokinin octapeptide.** *Gastroenterology* 1999, **117**:706-716
19. Swidsinski A, Khilkin M, Pahlig H, Swidsinski S, Priem F: **Time dependent changes in the concentration and type of bacterial sequences found in cholesterol gallstones.** *Hepatology* 1998, **27**:662-665
20. Osnes T, Sandstad O, Skar V, Osnes M: **Lipopolysaccharides and beta-glucuronidase activity in choledochal bile in relation to choledocholithiasis.** *Digestion* 1997, **58**:437-443
21. Lee D-K, Tarr PI, Haigh WG, Lee SP: **Bacterial DNA in mixed cholesterol gallstones.** *Am J Gastro* 1999, **94**:3502-3506
22. Olsson R, Bjornsson E, Backman L, Friman S, Hockerstedt K, Kaijser B, Olausson M: **Bile duct bacterial isolates in primary sclerosing cholangitis: a study of explanted livers.** *J Hepatol* 1998, **28**:426-432
23. Utili R, Abernathy CO, Zimmerman HJ: **Studies on the effects of C. coli endotoxin on canalicular bile formation in the isolated perfused rat liver.** *J Lab Clin Med* 1977, **89**:471-482
24. Nishida T, Nakahara M, Nakao K, Matsuda H: **Biliary bacterial infection decreased the secretion of bile acids and bilirubin into bile.** *Am J Surg* 1999, **177**:38-41
25. Eskandari MK, Kalf J, Billiar TR, Lee KKW, Bauer AJ: **Lipopolysaccharide activates the muscularis macrophage network and suppresses circular smooth muscle activity.** *Am J Physiol* 1997, **273**:G727-734
26. Rosen HR, Winkle PJ, Kendal BJ, Diehl DL: **Biliary interleukin-6 and tumor necrosis factor- α in patients undergoing endoscopic retrograde cholangiopancreatography.** *Dig Dis Sci* 1997, **42**:1290-1294
27. Prystowsky JB, Rege RV: **Interleukin-1 mediates guinea pig gallbladder inflammation in vivo.** *J Surg Res* 1997, **71**:123-126
28. Hedges S, Svensson M, Svanborg C: **Interleukin-6 response of epithelial cell lines to bacterial stimulation in vitro.** *Infect Immun* 1992, **60**:1295-1301
29. Khair OA, Davies RJ, Devalia JL: **Bacterial-induced release of inflammatory mediators by bronchial epithelial cells.** *Euro Resp J* 1996, **9**:1913-1922
30. Holtkamp GM, Van Rossem M, de Vos AF, Willekens B, Peek R, Kijlstra A: **Polarized secretion of IL-6 and IL-8 by human retinal pigment epithelial cells.** *Clin Exp Immunol* 1998, **112**:34-43
31. Jung HC, Eckmann L, Yang S-K, Panja A, Fierer J, E Morzycka-Wroblewska, Kagnoff MF: **A distinct array of proinflammatory cytokines is expressed in human colon epithelial cells in response to bacterial invasion.** *J Clin Invest* 1995, **95**:55-65
32. Taylor CT, Dzusz AL, Colgan SP: **Autocrine regulation of epithelial permeability by hypoxia: Role for polarized release of tumor necrosis factor α .** *Gastroenterology* 1998, **114**:657-668
33. Morland CM, Fear J, McNab G, Joplin R, Adams DH: **Promotion of leukocyte transendothelial cell migration by chemokines derived from human biliary epithelial cells in vitro.** *Proc Assoc Am Physicians* 1997, **109**:372-382
34. Matsumoto K, Fujii H, Michalopoulos G, Fung JJ, Demetris AJ: **Human biliary epithelial cells secrete and respond to cytokines and hepatocyte growth factors in vitro: interleukin-6, hepatocyte growth factor and epidermal growth factor promote DNA synthesis in vitro.** *Hepatology* 1994, **20**:376-382
35. Yasoshima M, Kono N, Sugawara H, Katayanagi K, Harada K, Nakamura Y: **Increased expression of interleukin-6 and tumor necrosis factor-alpha in pathologic biliary epithelial cells: in situ and culture study.** *Lab Invest* 1998, **78**:89-100
36. Keshav S, Lawson L, Chung LP, Stein M, Perry VH, Gordon S: **Tumor necrosis factor mRNA localized to paneth cells of normal murine intestinal epithelium by in situ hybridization.** *J Exp Med* 1990, **171**:327-332
37. Rege RV: **Inflammatory cytokines alter human gallbladder epithelial cell absorption/secretion.** *J Gastrointest Surg* 2000, **4**:185-192

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-230X/2/23/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedCentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com