

The Notch and TGF- β Signaling Pathways Contribute to the Aggressiveness of Clear Cell Renal Cell Carcinoma

Jonas Sjölund^{1‡}, Anna-Karin Boström^{1,9}, David Lindgren^{1,9}, Sugata Manna¹, Aristidis Moustakas², Börje Ljungberg³, Martin Johansson¹, Erik Fredlund¹, Håkan Axelsson^{1*}

1 Department of Laboratory Medicine, Center for Molecular Pathology, Lund University, University Hospital MAS, Malmö, Sweden, **2** Ludwig Institute for Cancer Research, Biomedical Center, Uppsala University, Uppsala, Sweden, **3** Department of Surgical and Perioperative Sciences, Urology and Andrology, Umeå University, Umeå, Sweden

Abstract

Background: Despite recent progress, therapy for metastatic clear cell renal cell carcinoma (CCRCC) is still inadequate. Dysregulated Notch signaling in CCRCC contributes to tumor growth, but the full spectrum of downstream processes regulated by Notch in this tumor form is unknown.

Methodology/Principal Findings: We show that inhibition of endogenous Notch signaling modulates TGF- β dependent gene regulation in CCRCC cells. Analysis of gene expression data representing 176 CCRCCs showed that elevated TGF- β pathway activity correlated significantly with shortened disease specific survival (log-rank test, $p = 0.006$) and patients with metastatic disease showed a significantly elevated TGF- β signaling activity (two-sided Student's t -test, $p = 0.044$). Inhibition of Notch signaling led to attenuation of both basal and TGF- β 1 induced TGF- β signaling in CCRCC cells, including an extensive set of genes known to be involved in migration and invasion. Functional analyses revealed that Notch inhibition decreased the migratory and invasive capacity of CCRCC cells.

Conclusion: An extensive cross-talk between the Notch and TGF- β signaling cascades is present in CCRCC and the functional properties of these two pathways are associated with the aggressiveness of this disease.

Citation: Sjölund J, Boström A-K, Lindgren D, Manna S, Moustakas A, et al. (2011) The Notch and TGF- β Signaling Pathways Contribute to the Aggressiveness of Clear Cell Renal Cell Carcinoma. PLoS ONE 6(8): e23057. doi:10.1371/journal.pone.0023057

Editor: Andrew Yeudall, Virginia Commonwealth University, United States of America

Received: November 2, 2010; **Accepted:** July 12, 2011; **Published:** August 3, 2011

Copyright: © 2011 Sjölund et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the Swedish Cancer Society (grant no 09 0643), the Swedish Research Council (grant no 90348001), Åke Wiberg's Foundation, Ollie and Elof Ericsson's Foundation, the Crafoord Foundation, Gunnar Nilsson's Cancer Foundation, The Royal Physiographic Society in Lund, and Malmö University Hospital Research Funds. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: hakan.axelsson@med.lu.se

‡ These authors contributed equally to this work.

‡ Current address: Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, California, United States of America

Introduction

Clear cell renal cell carcinoma (CCRCC) is the most common malignancy of the kidney [1]. About a quarter of the CCRCC patients have metastatic disease at the time of diagnosis and eventually one-third of the patients presented with localized tumors at diagnosis relapse. Despite recent advances using multi-kinase inhibitors, disseminated CCRCC remains inherently treatment resistant [2]. Consequently, studies leading to a better understanding of the factors that determines the metastatic phenotype of CCRCC are warranted [3]. The tumor suppressor gene *VHL* is lost in approximately 80% of all CCRCCs and represents a hallmark feature of CCRCC, but additional oncogenic events are required for both tumor formation and progression [4].

Notch signaling is an evolutionarily conserved signaling pathway of fundamental importance during development and post-natal life, regulating cell fate decisions, proliferation and survival. Dysregulated Notch signaling has been implicated in a wide variety of pathological conditions, including cancer [5].

Ligand (Jagged and Delta-like families) binding leads to two proteolytic cleavages of the receptors, the latter being dependent on the γ -secretase complex. Upon cleavage, the intracellular domain of the Notch receptor (icNotch) translocates to the nucleus where it converts the transcriptional repressor CSL to an activator [6]. Small molecule inhibitors that are capable of inhibiting Notch activation by targeting the γ -secretase complex are being tested for treatment of tumor types characterized by elevated Notch signaling, such as breast cancer and T-ALL [7]. In a recent study, we showed that Notch signaling components were elevated in primary CCRCC specimens compared to normal kidney and inhibition of Notch signaling attenuated growth of CCRCC cells, both *in vitro* and *in vivo* [8]. Thus, we have postulated that Notch signaling might represent a novel, clinically targetable oncogenic pathway in this pathological context.

The TGF- β pathway has a dual role in tumorigenesis: the growth inhibiting function at early stages of tumor formation is breached during tumor progression and at later stages TGF- β signaling can promote cell migration and invasion [9]. TGF- β elicits its cellular responses by binding to TGF- β type I and type II

serine/threonine kinase receptors (TGFBR1 and TGFBR2) that phosphorylate intracellular messengers SMAD2 and SMAD3, which in complex with SMAD4 transcriptionally induce or repress a diverse array of genes. In CCRCC, loss of TGFBR2 has been reported, which has been associated with tumor progression and also suggested to be the mechanism responsible for the escape from TGF- β -mediated growth repression [10,11,12,13]. However, there are also studies showing that loss of TGFBR2 expression is associated with improved CCRCC patient survival and that the TGF- β cascade promotes CCRCC bone metastasis *in vivo* [14,15].

Here we sought to identify downstream targets of the Notch pathway in CCRCC by employing transcriptome analyses of γ -secretase treated CCRCC cells. Our data indicate that inhibition of Notch signaling attenuates the TGF- β transcriptional output and that elevated TGF- β signaling activity in primary CCRCC is associated with decreased survival. This study thus provides additional rationale for targeting the Notch pathway for treatment of CCRCC.

Results

Notch inhibition in CCRCC cells affects TGF- β gene signatures

Our previous work established that active Notch signaling is an inherent property of CCRCC cells [8]. To further confirm this observation, we performed Western blot experiments on extracts from 786-O and SKRC-10 cells using an antibody that specifically recognizes activated Notch1 (icNotch1). As anticipated, icNotch1 was detected in control treated cells whereas treatment with the γ -secretase inhibitor DAPT completely abolished the levels of icNotch1 in both cell lines (Figure 1A). We next analyzed global gene expression changes following Notch inhibition using microarrays. The Notch target genes *HES1* and *IL7R* [8,16] were both strongly downregulated in both 786-O and SKRC-10 cells (Table 1), thus validating our approach.

Interestingly, five of the most downregulated genes common to both cell lines are known TGF- β target genes [9,17,18] (Table 1). We next asked whether this cross talk could be statistically verified in our data using gene set enrichment analysis (GSEA) [19]. DAPT modulated gene expression in the SKRC-10 microarray experiment were ranked based on Rank product analysis FDR [20]. As exemplified in Figure 1B, GSEA showed significant enrichment of several genes related to TGF- β induced transcription among the DAPT down regulated genes [21]. In a direct comparison we noted a consistent repression of several well-described TGF- β induced genes in DAPT treated samples (Figure 1C) [9,17,18]. Likewise, some previously characterized TGF- β downregulated genes (e.g. *CEPBD*, *CITED2*, *SPRY2*, and *RINT*) were significantly upregulated upon γ -secretase inhibition (Figure 1D). The downregulation of *HES1* and the TGF- β target genes *SERPINE1* and *SKIL* were confirmed using Quantitative real-time PCR (Q-PCR) in both 786-O and SKRC-10 cells (Figure 1E).

Overall, these results show that Notch inhibition not only affects prototypical Notch target genes but also modulates TGF- β dependent gene regulation in CCRCC cells.

Association between TGF- β pathway activity and prognosis in CCRCC patients

We next investigated the clinical relevance of TGF- β signaling in primary CCRCCs. It is known that TGF- β signaling acts in a highly tissue-specific manner. We therefore extracted a core set of TGF- β target genes relevant for CCRCC cells, by analyzing published TGF- β gene expression signatures derived from hepatocytes, breast cancer and fibroblasts using GSEA

[21,22,23]. We selected those genes from each of the TGF- β gene sets that contributed to the significant enrichment in the data from DAPT treated SKRC-10 cells, i.e. the leading edge subset from each GSEA analysis [19]. We thereby defined a core TGF- β gene expression signature of 145 genes, representing documented TGF- β target genes also affected in γ -secretase inhibited CCRCC cells (Table S1). This gene set was used to query a published gene expression data set of 176 CCRCCs [24] for correlations to survival. For each sample a specific TGF- β activity score was calculated based on the 145-gene signature. Survival analysis using Kaplan-Meier plots revealed that high TGF- β pathway activity score was significantly associated with a worse disease-specific survival (log-rank, $p = 0.006$; Figure 2A). Interestingly, using AJCC stage grouping, grade and performance status as covariates the TGF- β pathway activity also provided independent prognostic information when treated as a continuous variable in a multivariate Cox regression model ($p = 0.004$, HR = 4.04, 95% confidence interval (CI) = 1.55–10.53) (Table S2). Together, these analyses show that the TGF- β pathway is active and correlates to poor outcome in primary CCRCCs.

We next surveyed a collection of primary CCRCC specimens using an antibody directed against phosphorylated SMAD2 (pSMAD2), generally considered as a specific assessment of TGF- β signaling activity. We validated the anti-pSMAD2 antibody by staining paraffin-embedded 786-O cells that had been cultured in the presence or absence of the potent TGFBR1 inhibitor SB431542 [25] (Figure 2B). The primary CCRCCs generally showed strong nuclear pSMAD2 staining (Figure 2C), indicating that activation of the TGF- β cascade is a persistent feature of CCRCC.

Characterization of the TGF- β responsiveness in CCRCC cells

Since the published data regarding the role of the TGF- β receptors in CCRCC are conflicting, we analyzed the prognostic impact of these receptors in the data set from 176 CCRCCs (Figure S1). Our analysis revealed a significant association between high expression of *TGFBR1* and worse disease-specific survival (log-rank, $p = 0.030$). Low expression of *TGFBR3* was also associated with worse disease-specific survival (log-rank, $p = 0.010$). No significant association between survival and expression of *TGFBR2* could be detected.

Studies using CCRCC cells (including the 786-O cell line) suggested that intracellular TGF- β signaling is lost due to absence of the TGFBR2 receptor [10,11]. Baseline level of pSMAD2 could however be detected in 786-O cells using immunohistochemistry (Figure 2B). To further confirm the existence of a cell autonomous TGF- β signaling pathway in CCRCC cells, we analyzed the expression of TGFBR1 and TGFBR2 using Western blotting. Both 786-O and SKRC-10 cells expressed appreciable levels of the two receptors (Figure 3A). We also monitored the level of pSMAD2 in the presence or absence of exogenously added TGF- β 1 by Western blotting. We noted a baseline activity of pSMAD2 in unstimulated 786-O and SKRC-10 cells. The level of pSMAD2 in TGF- β 1 treated cells remained higher compared to control cells during the entire experiment in both cell lines (Figure 3B and 3C). Treatment with SB431542 led to a complete loss of the pSMAD2 signal (Figure 3D). Previous reports have indicated that the expression of TGF- β 1 is elevated in CCRCC due to the loss of pVHL [10,26], which are results compatible with our observations of pSMAD2 expression in primary CCRCCs and baseline expression of pSMAD2 in unstimulated CCRCC cells. In order to assess TGF- β 1 production in 786-O and SKRC-10 cells, we employed an ELISA assay. After 48 hours, TGF- β 1 could be




Figure 1. Notch inhibition in CCRCC cells attenuates TGF- β signaling. (A) Western blot analysis of icNotch1 in 786-O and SKRC-10 cells treated for 24 h with DAPT (+) or vehicle control (-). Cell extracts were separated by SDS-PAGE and after transfer the membrane was probed with an anti-cleaved Notch1 antibody. The blot was re-probed with an anti-ACTIN antibody to ensure equal loading. (B) GSEA of a TGF- β upregulated gene set classified as upregulated by TGF- β 1 treatment of skin fibroblasts at 1–4 h in the SKRC-10 microarray data. Isolated and purified RNA from SKRC-10 cells, treated with DAPT or vehicle control in 1% FCS supplemented media for 24 h, was used in oligomer microarray experiments. Genes in the SKRC-10 data list were ranked for downregulation upon DAPT treatment by measure of rank product analysis FDR. Gene sets with a q-value below 0.25 were considered significantly enriched. Genes upregulated by TGF- β 1 from the gene set were significantly enriched in the list ranked by downregulated by DAPT (NES = 1.75, $q = 0.09$). (C) Gene expression analysis of indicated genes reported to be activated in response to TGF- β in DAPT or vehicle control (c) treated SKRC-10 cells. Cells were treated as indicated in (B). Data represents mean log₂ ratios+95% confidence intervals of three separate experiments. (D) Gene expression analysis of indicated genes reported to be repressed in response to TGF- β in DAPT or vehicle control (c) treated SKRC-10 cells. Cells were treated as indicated in (B). Data represents mean log₂ ratios+95% confidence intervals of three separate experiments. (E) Notch inhibition in CCRCC cells decreases the mRNA expression of the Notch primary target *HES1* and the TGF- β target genes *SERPINE1* and *SKIL*. Q-PCR analyses of *HES1*, *SKIL* and *SERPINE1* mRNA levels in 786-O or SKRC-10 cells treated for 24 h with DAPT or vehicle control (c). mRNA levels were normalized to *SDHA*, *YWHAZ* and *UBC* expression and data represents mean+95% confidence intervals of three separate experiments. ***, ** and * indicates statistical significant changes (two-sided Student's t-test, $p < 0.001$, $p < 0.01$ and $p < 0.05$ respectively). doi:10.1371/journal.pone.0023057.g001

readily detected in the medium of both cell lines (Figure 3E). When 786-O and SKRC-10 cells were transfected with a plasmid containing a SMAD regulated luciferase reporter ((*CAGA*)₁₂-*Luc*), a dose-dependent increase of the luciferase reporter was detected upon TGF- β 1 stimulation (Figure 3F and 3G, respectively). This induction was dependent of TGFBR1 activation, since treatment with SB431542 abrogated both TGF- β 1 induced and basal reporter gene activity (Figure 3F–H). Q-PCR experiments showed

that the expression of the TGF- β responsive genes *JUNB* and *SERPINE1* were significantly induced upon treatment with TGF- β 1 and suppressed below baseline when the cells were treated with SB431542 alone or in combination with TGF- β 1 (Figure 3I).

Based on these experiments, we conclude that the TGF- β signaling pathway is functional in the two investigated CCRCC cell lines and that the basal activity might be a consequence of endogenous TGF- β 1 production.

Table 1. Top downregulated genes common to both 786-O and SKRC-10 cells upon γ -secretase inhibition when compared to vehicle treatment for 24 hours as evaluated from microarray data.

Gene name	786-O		SKRC-10	
	Mean ratio	Z	Mean ratio	Z
<i>HES1</i>	-1.44	-4.80	-2.98	-8.54
<i>IL7R</i>	-0.77	-2.57	-1.74	-5.02
<i>LOC284422</i>	-2.10	-6.96	-1.48	-4.30
<i>CXCL12</i>	-1.32	-4.38	-1.31	-3.80
<i>CXCR7</i>	-1.03	-3.44	-1.30	-3.78
<i>SNF1LK</i>	-0.71	-2.37	-1.29	-3.74
<i>GPR56</i>	-0.82	-2.73	-1.25	-3.63
<i>MAOB</i>	-1.15	-3.84	-1.22	-3.55
<i>SERPINE1</i>	-0.96	-3.19	-1.21	-3.53
<i>SEMA3F</i>	-1.11	-3.69	-1.19	-3.47
<i>KRTAP9-4</i>	-1.07	-3.59	-1.16	-3.38
<i>LPHN1</i>	-1.37	-4.58	-1.14	-3.34
<i>HSF2BP</i>	-0.71	-2.38	-1.13	-3.29
<i>CLTCL1</i>	-1.05	-3.49	-1.07	-3.13
<i>MN1</i>	-0.93	-3.10	-1.07	-3.13
<i>GALNT2</i>	-0.82	-2.76	-0.91	-2.68
<i>LRP4</i>	-0.71	-2.40	-0.90	-2.63
<i>CHAC1</i>	-0.70	-2.34	-0.84	-2.47
<i>FAM20C</i>	-0.63	-2.13	-0.80	-2.35
<i>INHBA</i>	-1.36	-4.54	-0.79	-2.32
<i>SR-A1</i>	-0.61	-2.06	-0.74	-2.18
<i>SKIL</i>	-0.68	-2.27	-0.69	-2.05

Genes in bold represents previously described TGF- β signaling target genes. Data represents mean \log_2 ratios (DAPT/DMSO) and Z-scores of three separate experiments. The genes are sorted according to the Z-scores of SKRC-10. doi:10.1371/journal.pone.0023057.t001

Notch inhibition perturbs both basal and induced TGF- β signaling activity

We next wanted to characterize the effects of Notch inhibition on TGF- β signaling. 786-O cells showed a modest but consistent decrease of basal pSMAD2 levels upon treatment with DAPT at all time-points analyzed, while the SMAD2 levels were unaffected all through the time course of the experiment (Figure 4A). Similar results were obtained when analyzing the SK-RC10 cell line (data not shown). Also, when transfecting the cells with siRNA against *Notch1* a down-regulation of pSMAD2 levels could be detected (Figure 4B). Notch inhibition also decreased phosphorylation of SMAD2 in cells stimulated with TGF- β 1 (Figure 4C).

We next wanted to analyze whether the suppressive effect on TGF- β pathway activity by Notch inhibition could be reversed by constitutively active, γ -secretase insensitive Notch signaling. For this purpose, we co-transfected 786-O cells with the (*CAGA*)₁₂ luciferase reporter together with an icNotch1 expression vector and treated the cells with DAPT. As shown in figure 4E, expression of icNotch1 led to a significant increase in reporter activity compared to vector control. In the presence of DAPT, expression of icNotch1 led to a partial but significant reversal of the DAPT induced suppression of reporter activity (Figure 4E). We further corroborated the diminished basal and TGF- β 1 induced TGF- β activity upon treatment with DAPT using the (*CAGA*)₁₂ reporter

(Figure 4D, 4F and 4G). Modulation of Notch signaling also affected TGF- β responsive genes (*JUNB* and *SERPINE1*) in the presence of TGF- β 1 (Figure 4H).

Altogether, our data indicate that inhibition of Notch signaling downregulates TGF- β signaling in CCRCC cells.

Notch inhibition perturbs the migratory capacity of CCRCC cells

The dual role of TGF- β signaling in cancer is well established, with a cytostatic effect in the early stages, which can be subdued to a metastatic promoting program at the later stages of tumor progression [17]. We noted however very modest effects on the cytostatic TGF- β transcripts [9] in DAPT treated SKRC-10 cells (Figure S2A). Consistent with this observation and with a previous report [10], thymidine incorporation assays confirmed that the growth capacity of CCRCC cells was not decreased by treatment with TGF- β 1 for up to 72 h (Figure 5A). In case of the SKRC-10 cells there was even a modest but significant increase in growth at 72 h of treatment with TGF- β 1.

Our microarray experiments indicated that genes regulating migration and/or invasion were downregulated (Figure S2B). Most of these genes are also directly regulated by TGF- β signaling and have been associated with aggressive and invasive cancer [17]. This observation suggested that Notch inhibition perturbs the migratory and/or invasive capacity of CCRCC cells. We functionally verified this using Boyden chamber assays and noted a significant decrease in migration when CCRCC cells were treated with DAPT (Figure 5B) or upon *Notch1* knockdown (Figure 5C) compared to control treated cells. Also, treatment with the TGF- β inhibitor SB431542 led to a significant decrease in migration of SK-RC10 cells and when combining SB431542 and DAPT treatments, no further decrease in migration was noted (Figure 5D). Addition of exogenous TGF- β 1 further stimulated the migratory capacity and this effect could be attenuated by Notch inhibition (Figure 5E). Furthermore, Notch inhibition led to a pronounced and significant decrease in invasion in both cell lines tested when compared with vehicle control (Figure 5F). To verify the clinical significance of these results, we assessed TGF- β signaling activity based on our 145-gene TGF- β signature in a previously published microarray study [27]. CCRCCs from patients with either metastatic disease at diagnosis or that later developed metastasis (n = 13) showed a significantly elevated TGF- β signaling activity as compared to tumors from patients with a localized disease and with no documented metastases during follow-up (n = 9) (two-sided Student's t-test, p = 0.044, Figure 5G).

Thus, dysregulated Notch signaling might contribute to CCRCC aggressiveness at least in part by modulating TGF- β signaling activity.

Discussion

It has been shown that loss of VHL, which is the key oncogenic event in CCRCC, leads to elevated expression of TGF- β 1 [10]. Interestingly, elevated levels of TGF- β 1 in serum from CCRCC patients are correlated with unfavorable outcome of the disease [26,28,29]. Hence, the tumor microenvironment in CCRCC is rich in TGF- β 1. These observations therefore suggest that CCRCC cells might have acquired the capacity to evade the cytostatic effects imposed by the presence of TGF- β 1. It has been postulated that structural alterations of TGF- β pathway components, such as mutations of *TGFBR2* render tumor cells insensitive to TGF- β cytostatic effects [17]. In CCRCC there are conflicting reports on such alterations and there is an apparent lack of functional analyses of signaling activity, e.g. assessment of






Figure 2. High TGF- β signaling pathway activity is associated with decreased patient survival. (A) Kaplan-Meier curves comparing disease-specific survival among 176 CCRCC patients divided into quartiles based on the TGF- β activity score. High TGF- β pathway activity score (76–100%) was significantly (log-rank, $p=0.006$) associated with a worse disease-specific survival. (B) Immunohistochemical analyses using phospho-specific SMAD2 (pSMAD2) antibody of control (c) or SB431542 treated 786-O cells. Sections were counterstained with hematoxylin-eosin. Original magnification, $\times 40$. (C) pSMAD2 is expressed *in vivo*. Immunohistochemistry of pSMAD2 in nine CCRCC tumors, that all showed positive nuclear staining. Sections were counterstained with hematoxylin-eosin. Original magnification, $\times 40$. doi:10.1371/journal.pone.0023057.g002

pSMAD2 levels. Experimental *in vitro* and *in vivo* studies have indicated that TGBR3 have antitumoral effects in CCRCC cells independent of TGF- β 1 and canonical TGFBR1/TGFBR2/SMAD signaling [11]. Our data showing that low *TGFBR3* expression in primary CCRCC is significantly associated with worse disease-specific survival is thus adding further support for this notion. Loss of TGFBR2 has been linked to CCRCC progression [10,11,12,13], whilst another investigation showed that loss of TGFBR2 improve CCRCC patient survival [15]. In favor of the latter study, the TGF- β cascade has been shown to promote CCRCC bone metastasis *in vivo* [14]. It is noteworthy that Ananth *et al*, concluded that the 786-O cells lacks a working TGF- β signaling pathway due to the absence of TGFBR2 expression [10]. In contrast, our functional assessment of the pathway in 786-O cells clearly shows that the pathway remains intact. In normal renal cells, TGF- β 1 elicits an antimitogenic response and triggers epithelial-to-mesenchymal transition [18,30,31,32,33]. While our data indicate that CCRCC cells are insensitive to TGF- β -induced growth inhibition, the cells retain an operational TGF- β pathway that directs pro-migratory and pro-metastatic functions. Consistent with the experimental data, we found evidence of SMAD2 activation in clinical specimens and an association between TGF- β signaling activity, disease-specific survival and metastatic progression in the analyses of primary CCRCCs. Our observation that elevated *TGFBR1* is significantly associated with worse disease-specific survival provides further support for a pro-metastatic function of TGF- β signaling in CCRCC. Thus, we extend previous data and suggest a pro-oncogenic role for a hyperactivated autocrine TGF- β pathway in CCRCC. This tumor-promoting effect of pathogenic TGF- β signaling could partly be manifested in an increased metastatic potential of the tumor cells, but also through paracrine angiogenic and immunosuppressive effects of TGF- β secreted by the growing tumor mass [10,34].

Different modes of cross-talk between the TGF- β and Notch signaling pathways of both synergistic and antagonistic nature have been reported in various cellular contexts [35,36,37,38,39,40,41,42,43,44]. In CCRCC cells, characterized by high activity of both pathways, Notch signaling seems superimposed on TGF- β signaling since Notch inhibition, either by siRNA targeting *Notch1* or pharmacological inhibition of Notch receptor activation, clearly perturbs important aspects of metastasis associated TGF- β signaling.

Since metastatic CCRCC has a particularly poor prognosis, with a five-year survival of about 9%, it is critical to develop treatment strategies that target the metastatic process [1]. We have recently developed a novel γ -secretase inhibition strategy, using intermittent treatment cycles that strongly inhibited the growth of xenotransplanted CCRCC cells while limiting the toxicity of the intestine, which is a major obstacle in achieving effective doses of these drugs in humans [8]. In a recent study it was also shown that glucocorticoids abrogate the gastrointestinal toxicity of γ -secretase inhibitors [45]. Thus, these studies provide alternative strategies to spare the patients from the side effects of systemic Notch inhibition.

We now provide evidence that Notch inhibition also attenuates the migratory capacity of CCRCC cells, at least in part through modulation of TGF- β signaling. In addition, it is known that

inhibition of Notch signaling perturbs tumor angiogenesis [46]. Thus, we conclude that Notch inhibition might be a particularly appealing approach for treatment of CCRCC, potentially curbing several key aspects of tumor aggressiveness.

Materials and Methods

Cell culture and reagents

The 786-O (ATCC, Rockville, MD, USA) CCRCC cell line was cultured in DMEM (Invitrogen, Stockholm, Sweden) containing 10% fetal calf serum (FCS, Invitrogen) and supplemented with 1% penicillin and streptomycin (PEST, Invitrogen). The SKRC-10 CCRCC cell line was maintained in RPMI 1640 (Invitrogen) containing 10% FCS and 1% PEST. Human recombinant TGF- β 1 was obtained from PeproTech (London, United Kingdom). Cells were treated with 2 μ M TGFBR1 inhibitor (SB431542, Sigma Aldrich, Munich, Germany), 10 μ M γ -secretase inhibitor DAPT (*N*-[*N*-(3,5-difluorophenacetyl)-L-alanyl]-*S*-phenylglycine *t*-butyl ester) from Calbiochem (Darmstadt, Germany) or the corresponding volume of DMSO (Sigma Aldrich) for indicated times. All experiments were performed in reduced serum conditions.

Microarray and data analyses

RNA from 786-O and SKRC-10 cells, treated with DAPT or vehicle control in 1% FCS supplemented media for 24 h, was used for gene expression microarray experiments with a 27 k cDNA array platform (<http://www.lth.se/sciblu>). Array production, sample labeling, hybridization and scanning were performed essentially as described previously [47]. In short, 5 μ g of total RNA (DAPT or vehicle control) was labeled with Cy3 and hybridized against 5 μ g of Cy5-labeled RNA from a pool representing nine untreated CCRCC cell lines. As the effects of DAPT treatment were of different magnitude in SKRC-10 and 786-O cells, a comparative Z-score was calculated by dividing the mean \log_2 ratio values for each gene and cell line with the standard deviation of all mean \log_2 ratios for each cell line. We performed a second round of experiments (dubbed SKRC-10 data set), that were used for GSEA [19] and extraction of gene expression signatures for pathway analysis. Rank product analysis [20] was used to create ranked gene lists based on both upregulation and downregulation. The downregulated ranked gene lists were used for correlation analyses to known gene signatures according to the GSEA method using the Molecular Signatures Database (MSigDB), and additional published TGF- β regulated gene sets [22,23].

Genes in the SKRC-10 data set contributing to a significant enrichment of the TGF- β gene sets were thereafter used to generate a DAPT/TGF- β specific signature. To investigate possible clinical significance of this obtained TGF- β gene signature, two gene expression data sets were used. The first, which comprised 177 CCRCCs, was obtained from the Stanford microarray database [48] and normalized as described in the original publication [24]. Replicate reporters were merged by gene-symbol and a presence filter was applied allowing not less than 50% presence for each gene across arrays. The second data set comprised 22 CCRCCs and 23 normal kidney samples [27].




Figure 3. The TGF- β signaling pathway is functional in CRCC cells. (A) Immunoblotting using TGFBR1 and TGFBR2 antibodies of cell lysates from 786-O and SKRC-10 cells. (B and C) pSMAD2 and SMAD-1/2/3 protein levels in cell lysates from 786-O (B) and SKRC-10 (C) cells stimulated with vehicle control (-) or 0.25 ng/ml TGF- β 1 (+) for the indicated time points. (D) Western blot analysis using pSMAD2 antibody and SMAD2 antibody, of cell lysates from 786-O and SKRC-10 cells treated with vehicle (-) or SB431542 (+) for 4 h. (E) ELISA measurement of TGF- β 1 levels in serum free media from 786-O and SKRC-10 cells grown for 48 h. Data represents mean+95% confidence intervals of three separate experiments. (F and G) 786-O (F) and SKRC-10 (G) cells were transfected with the SMAD regulated (CAGA)₁₂-Luc reporter construct, treated with vehicle control (c), increasing concentrations of TGF- β 1 or 0.25 ng/ml TGF- β 1 and SB431542 for 24 h and analyzed for relative luciferase activity. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (H) 786-O or SKRC-10 cells were transfected with the (CAGA)₁₂-Luc reporter construct, treated with vehicle control (-) or SB431542 (+) for 24 h and analyzed for relative luciferase activity. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (I) Q-PCR analyses of *JUNB* and *SERPINE1* mRNA levels in SKRC-10 cells treated with vehicle control (c), SB431542 and/or 0.25 ng/ml TGF- β 1 for 4 h. mRNA levels were normalized to *SDHA*, *YWHAZ* and *UBC* expression and data represents mean+95% confidence intervals of three separate experiments. ***, ** and * indicates statistical significant changes (two-sided Student's t-test, $p < 0.001$, $p < 0.01$ and $p < 0.05$ respectively). doi:10.1371/journal.pone.0023057.g003




Figure 4. Notch inhibition alters endogenous and TGF- β 1 induced activation of the TGF- β signaling pathway. (A) Immunoblotting using pSMAD2 antibody and SMAD2 antibody, of cell lysates from 786-O cells stimulated with vehicle control (–) or DAPT (+) for the indicated time points. (B) Immunoblotting using Notch1 antibody and pSMAD2 antibody of cell lysates from 786-O and SKRC-10 cells transfected with control siRNA (c-si) or siRNA against *Notch1* (siN-1). Cells were harvested after 24 h of transfection. (C) Western blot analysis, using pSMAD2 antibody and SMAD-1/2/3 antibody of cell lysates from 786-O and SKRC-10 cells stimulated with 0.25 ng/ml TGF- β 1 and vehicle (–) or 0.25 ng/ml TGF- β 1 and DAPT (+) for 4 h. (D) 786-O and SKRC-10 cells transfected with the (CAGA)₁₂-Luc reporter construct and treated with vehicle control (c) or DAPT for 24 h and (E) 786-O and SKRC-10 cells transfected with the Notch1 reporter construct and treated with vehicle control (c) or DAPT for 24 h and (F) 786-O cells transfected with the Notch1 reporter construct and treated with vehicle control (c) or DAPT for 24 h and (G) SKRC-10 cells transfected with the Notch1 reporter construct and treated with vehicle control (c) or DAPT for 24 h and (H) 786-O cells transfected with the Notch1 reporter construct and treated with vehicle control (c) or DAPT for 24 h and (I) SKRC-10 cells transfected with the Notch1 reporter construct and treated with vehicle control (c) or DAPT for 24 h. Relative luciferase activity and relative mRNA levels are shown. Error bars represent standard deviation. Statistical significance is indicated by asterisks: * p < 0.05, ** p < 0.01, *** p < 0.001.

followed by measurement of relative luciferase activity. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (E) Relative luciferase activity in extracts from 786-O cells transfected with the (CAGA)₁₂-Luc reporter construct and vector control (Vector C) or icNotch1 expression vector followed by treatment with DAPT or vehicle control (c) for 24 h. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated and vector control transfected cells. (F and G) 786-O (F) and SKRC-10 (G) cells were transfected with the (CAGA)₁₂-Luc reporter construct, treated with vehicle control (c), 0.25 ng/ml TGF- β 1 and vehicle control or 0.25 ng/ml TGF- β 1 and DAPT for 24 h and analyzed for relative luciferase activity. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (H) Q-PCR analyses of *JUNB* and *SERPINE1* mRNA levels in SKRC-10 cells treated with vehicle control (c), 0.25 ng/ml TGF- β 1 and vehicle (c) or 0.25 ng/ml TGF- β 1 and DAPT for 4 h. mRNA levels were normalized to *SDHA*, *YWHAZ* and *UBC* expression and data represents mean+95% confidence intervals of three separate experiments. ***, ** and * indicates statistical significant changes (two-sided Student's t-test, $p < 0.001$, $p < 0.01$ and $p < 0.05$ respectively). doi:10.1371/journal.pone.0023057.g004

Log₂ expression values for each reporter were centered according to the median expression of the normal samples and replicate reporters were merged by gene symbol. For each sample, a TGF- β pathway activity score was calculated as previously described [49]. Disease-specific survival (DSS) and American Joint Committee on Cancer (AJCC) stage grouping (I, II, III or IV) were available for 176 of the 177 patients in the Zhao *et al* cDNA gene expression data set [24]. Follow-up time was limited to five years. For Kaplan-Meier analyses, patients were divided into quartiles based on their relative TGF- β pathway activity score and interquartile differences in survival were assayed using the log-rank test. All statistical analyses were carried out using the R statistical programming environment (<http://www.r-project.org>). Specifically, for survival analyses the Survival package was used.

Quantitative real-time PCR analyses

Q-PCR analyses, total RNA extraction and quantification of gene expression using SYBR Green (Applied Biosystems, Foster City, CA, USA) were done according to previously published procedures [50]. Primer sequences are given in Table S3. Quantification of relative mRNA levels was done using the comparative C_t method and normalized to three endogenous references genes (*SDHA*, *YWHAZ* and *UBC*) [51].

Luciferase reporter assays and siRNA transfection

For siRNA experiments, cells were transfected with control siRNA or siRNA against *Notch1* (Santa Cruz Biotechnology, Santa Cruz, CA, USA) using Lipofectamine 2000 (Invitrogen) and OptiMEM I Reduced Serum Medium (Invitrogen) as described elsewhere [8].

For luciferase experiments, cells were transiently transfected with the luciferase reporter vector (CAGA)₁₂-Luc containing 12 CAGA SMAD binding sites [52]. *phRL-TK* renilla expression vector (Promega, Madison, WI, USA) was used as a control for transfection efficiency. The icNotch1 expression construct was kindly provided by J.C. Aster [53]. Cells were lysed and assayed for luciferase and renilla activities using the Dual-Luciferase Reporter Assay System (Promega).

Western blot analyses and immunohistochemistry

Cells were lysed in RIPA buffer, separated on a SDS-PAGE gel and blotted onto Immobilon-P (Millipore, Bedford, MA, USA) or Hybond-C (Amersham Biosciences, Uppsala, Sweden) membranes. The membranes were incubated with the following primary antibodies: anti-Notch1 (Santa Cruz Biotechnology), anti-cleaved Notch1 (Cell Signaling Technology, MA, USA), anti-phosphorylated SMAD2 (pSMAD2, Cell Signaling Technology), anti-SMAD2 (Cell Signaling Technology), anti-SMAD-1/-2/-3 (Santa Cruz Biotechnology), anti-TGFBR1 (Santa Cruz Biotechnology), anti-TGFBR2 (Santa Cruz Biotechnology) or anti-ACTIN (ICN Biomedicals, Aurora, OH, USA). HRP-conjugated secondary antibodies were obtained from Amersham Biosciences, Dako

(Glostrup, Denmark) and Jackson ImmunoResearch Laboratories Inc (West Grove, PA, USA). Proteins were detected by Super Signal chemiluminescence substrate (Pierce, Rockford, IL, USA).

Tumor samples collected at the University Hospital in Umeå, Sweden, including nine nephrectomy specimens were analyzed by immunohistochemistry. The tumors were classified as CCRCCs according to the Heidelberg classification system [54]. All tumor samples were obtained after permission from the patients with informed and signed consent, and the Institutional Review Board approved the study. pSMAD2 immunoreactivity was detected using the Dako EnVision system and the Dako TechMate 500 as previously described [8]. Sections were counterstained with hematoxylin-eosin. To evaluate antibody specificity, pSMAD2 immunoreactivity of vehicle control treated 786-O cells or 786-O cells in which the antigen had been eliminated by SB431542 for 24 h were performed.

TGF- β 1 ELISA assay

Cells were maintained in FCS-free media for 48 h, whereafter an ELISA was performed using the Human TGF- β 1 immunoassay (R&D Systems Inc, Minneapolis, MN, USA) according to the manufacturer's description. An ELISA microplate reader (BioTek Synergy 2, Fisher Scientific, Gothenburg, Sweden) was used to analyze the absorbance (450 nm).

Cell proliferation assays

Cells were seeded in 1% FCS media supplemented with vehicle control or TGF- β 1 and incubated for 24, 48 or 72 h. [³H]thymidine (Amersham Biosciences) was then added to the culture. Cells were harvested after 24 h of incubation. The incorporated [³H]thymidine was measured in a β -liquid scintillation counter (LKB RackBeta Wallace, Turku, Finland).

Migration and invasion assays

In γ -secretase inhibition experiments, cells were pretreated for 24 h with DAPT or vehicle control before initiation of the migration assay. Cells were then seeded in FCS-free media supplemented with DMSO, DAPT and/or TGF- β 1 into Boyden chambers (Corning, Bodenheim, Germany) with 8- μ m pore size polycarbonate membrane filters. In experiments combining SB431542 and DAPT, SKRC-10 cells were pretreated with 2 μ M SB431542 alone, in combination with 10 μ M DAPT or with vehicle control (DMSO) in 1% FCS supplemented media for 24 h. The cells were then allowed to migrate towards the lower compartment containing 10% FCS for 4 h (786-O) or 5 h (SKRC-10). The migrated cells were then fixed with 4% paraformaldehyde (Sigma Aldrich) and stained with DAPI ([4',6'-diamidino-2-phenylindole], Sigma Aldrich). The cells were thereafter counted by microscopy at 40 \times magnification. Four representative fields were counted for each filter, and each treatment condition was assayed in triplicate and repeated three times. In siRNA experiments, cells were transfected with control siRNA or siRNA against *Notch1* 24 h preceding migration assay.




Figure 5. Notch inhibition attenuates CCRCC cell migration and invasion. (A) [3 H]thymidine incorporation of 786-O and SKRC-10 cells grown for indicated hours (h) in the presence of 1.0 ng/ml TGF- β 1 or treated with vehicle control (c). Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (B) Cell migration assessed by Boyden chamber assays of 786-O and SKRC-10 cells treated with vehicle control (c) or DAPT. The cells were allowed to migrate towards the lower compartment for 4 h (786-O) or 5 h (SKRC-10). Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. (C) Cell migration as determined by Boyden chamber assays of 786-O and SKRC-10 cells transfected with non-specific control (c-si) or *Notch1* (siN-1) specific siRNAs. After 24 h of transfection, cells were counted and seeded into the Boyden chamber. The cells were allowed to migrate towards the

lower compartment for 4 h (786-O) or 5 h (SKRC-10). Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. **(D)** Cell migration as determined by Boyden chamber assays of SKRC-10 cells treated with vehicle control (c), 2 μ M SB431542, 10 μ M DAPT and control. The cells were allowed to migrate towards the lower compartment for 5 h. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. **(E)** Cell migration analyses of SKRC-10 cells treated with vehicle control (c) or treated with 0.25 ng/ml TGF- β 1 and vehicle (c) or 0.25 ng/ml TGF- β 1 and DAPT. The cells were allowed to migrate towards the lower compartment for 5 h. Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. **(F)** Cell invasion as determined by Matrigel coated Boyden chamber assays of 786-O and SKRC-10 cells treated with vehicle control (c) or DAPT. The cells were allowed to invade and migrate towards the lower compartment containing 10% FCS for 16 h (786-O) or 21 h (SKRC-10). Data represents mean+95% confidence intervals of three separate experiments. Data were normalized to vehicle control treated cells. ***, ** and * indicates statistical significant changes (two-sided Student's t-test, $p < 0.001$, $p < 0.01$ and $p < 0.05$ respectively). **(G)** TGF- β pathway activity correlates to metastatic spread of primary CCRCCs. CCRCCs from patients with either metastatic disease at diagnosis or that later developed metastasis (Metastatic, $n = 13$) showed a significantly elevated TGF- β signaling activity based on the 145-gene TGF- β signature as compared to tumors from patients with a localized disease and with no documented metastases during follow-up (Localized, $n = 9$) (two-sided Student's t-test, $p = 0.044$). doi:10.1371/journal.pone.0023057.g005

For invasion assays, 12.5% Growth Factor Reduced BD MatrigelTM Matrix (BD Biosciences, Bedford, MA, USA) diluted in FCS-free media was added on top of each Boyden chamber membrane. The cells were seeded in FCS-free media supplemented with DAPT or DMSO and were then allowed to invade through the Matrigel towards the lower compartment containing 10% FCS for 16 (786-O) or 21 h (SKRC-10) at 37°C. After incubation cells were analyzed as described for the migration assay.

Statistical Analysis

Data were calculated as the mean values with 95% confidence intervals. All statistical tests were two-sided Student's t-test and statistical significance was defined as p less than 0.05. For the statistical design and analyses of gene expression microarray data refer to "Microarray and data analyses" above.

Supporting Information

Figure S1 Disease-specific survival of 176 CCRCC patients based on the gene expression of TGFBRs. Kaplan-Meier plots of disease-specific survival of 176 CCRCC patients that were divided into four groups based on the median gene expression values of *TGFBR1*, *TGFBR2*, and *TGFBR3*. Elevated *TGFBR1* expression (log-rank, $p = 0.030$) and decreased (0–25%) *TGFBR3* expression (log-rank, $p = 0.010$) were significantly associated with worse disease-specific survival. (PDF)

Figure S2 Effect of γ -secretase inhibition on gene programs of interest in SKRC-10 cells. **(A)** Notch inhibition does not profoundly affect the TGF- β cytosolic gene program as assessed by gene expression analysis of SKRC-10 cells. Isolated and purified RNA from SKRC-10 cells treated with vehicle control (c)

or DAPT in 1% FCS for 24 hours was used in oligomer microarray experiments. Data represents mean log₂ ratios of three independent experiments+95% confidence intervals. **(B)** Notch inhibition leads to downregulation of a large set of genes (* indicates previously described TGF- β target genes) associated with cell migration and invasion as determined by gene expression analysis of SKRC-10 cells. Isolated and purified RNA from SKRC-10 cells treated with vehicle control (c) or DAPT in 1% FCS for 24 hours was used in oligomer microarray experiments. Data represents mean log₂ ratios of three independent experiments+95% confidence intervals. ***, ** and * indicates statistical significant changes (two-sided Student's t-test, $p < 0.001$, $p < 0.01$ and $p < 0.05$ respectively). (PDF)

Table S1 Core TGF- β gene expression signature of 145 genes.

(PDF)

Table S2 Multivariate COX regression analyses.

(PDF)

Table S3 Q-PCR primer sequences.

(PDF)

Acknowledgments

We are grateful to E. Oosterwijk for providing the SKRC-10 cells and J.C. Aster for providing the icNotch1 vector.

Author Contributions

Conceived and designed the experiments: JS A-KB DL EF HA. Performed the experiments: JS A-KB DL SM EF. Analyzed the data: JS A-KB DL SM EF HA MJ. Contributed reagents/materials/analysis tools: AM BL MJ. Wrote the paper: JS A-KB DL EF HA.

References

- Cohen HT, McGovern FJ (2005) Renal-cell carcinoma. *N Engl J Med* 353: 2477–2490.
- Rini BI, Atkins MB (2009) Resistance to targeted therapy in renal-cell carcinoma. *Lancet Oncol* 10: 992–1000.
- Linehan WM, Rubin JS, Bottaro DP (2009) VHL loss of function and its impact on oncogenic signaling networks in clear cell renal cell carcinoma. *Int J Biochem Cell Biol* 41: 753–756.
- Kim WY, Kaelin WG (2004) Role of VHL gene mutation in human cancer. *J Clin Oncol* 22: 4991–5004.
- Bolos V, Grego-Bessa J, de la Pompa JL (2007) Notch signaling in development and cancer. *Endocr Rev* 28: 339–363.
- Nichols JT, Miyamoto A, Weinmaster G (2007) Notch signaling—constantly on the move. *Traffic* 8: 959–969.
- Garber K (2007) Notch emerges as new cancer drug target. *J Natl Cancer Inst* 99: 1284–1285.
- Sjolund J, Johansson M, Manna S, Norin C, Pietras A, et al. (2008) Suppression of renal cell carcinoma growth by inhibition of Notch signaling in vitro and in vivo. *J Clin Invest* 118: 217–228.
- Chen CR, Kang Y, Massague J (2001) Defective repression of c-myc in breast cancer cells: A loss at the core of the transforming growth factor beta growth arrest program. *Proc Natl Acad Sci U S A* 98: 992–999.
- Ananth S, Knebelmann B, Gruning W, Dhanabal M, Walz G, et al. (1999) Transforming growth factor beta1 is a target for the von Hippel-Lindau tumor suppressor and a critical growth factor for clear cell renal carcinoma. *Cancer Res* 59: 2210–2216.
- Copland JA, Luxon BA, Ajani L, Maity T, Campagnaro E, et al. (2003) Genomic profiling identifies alterations in TGFbeta signaling through loss of TGFbeta receptor expression in human renal cell carcinogenesis and progression. *Oncogene* 22: 8053–8062.
- Miyajima A, Asano T, Seta K, Kakoi N, Hayakawa M (2003) Loss of expression of transforming growth factor-beta receptor as a prognostic factor in patients with renal cell carcinoma. *Urology* 61: 1072–1077.
- Takahashi M, Rhodes DR, Furge KA, Kanayama H, Kagawa S, et al. (2001) Gene expression profiling of clear cell renal cell carcinoma: gene identification and prognostic classification. *Proc Natl Acad Sci U S A* 98: 9754–9759.
- Kominsky SL, Doucet M, Brady K, Weber KL (2007) TGF-beta promotes the establishment of renal cell carcinoma bone metastasis. *J Bone Miner Res* 22: 37–44.
- Parker AS, Lohse CM, Wu K, Kreinest P, Copland JA, et al. (2007) Lower expression levels of the transforming growth factor beta receptor type II protein are associated with a less aggressive tumor phenotype and improved survival among patients with clear cell renal cell carcinoma. *Hum Pathol* 38: 453–461.

16. Gonzalez-Garcia S, Garcia-Peydro M, Martin-Gayo E, Ballestar E, Esteller M, et al. (2009) CSL-MAML-dependent Notch1 signaling controls T lineage-specific IL-7R α gene expression in early human thymopoiesis and leukemia. *J Exp Med* 206: 779–791.
17. Massague J, Gomis RR (2006) The logic of TGF β signaling. *FEBS Lett* 580: 2811–2820.
18. Zavadil J, Bitzer M, Liang D, Yang YC, Massimi A, et al. (2001) Genetic programs of epithelial cell plasticity directed by transforming growth factor- β . *Proc Natl Acad Sci U S A* 98: 6686–6691.
19. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, et al. (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 102: 15545–15550.
20. Breitling R, Armengaud P, Amtmann A, Herzyk P (2004) Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS Lett* 573: 83–92.
21. Verrecchia F, Chu ML, Mauviel A (2001) Identification of novel TGF- β / Smad gene targets in dermal fibroblasts using a combined cDNA microarray/promoter transactivation approach. *J Biol Chem* 276: 17058–17062.
22. Coulouarn C, Factor VM, Thorgeirsson SS (2008) Transforming growth factor- β gene expression signature in mouse hepatocytes predicts clinical outcome in human cancer. *Hepatology* 47: 2059–2067.
23. Padua D, Zhang XH, Wang Q, Nadal C, Gerald WL, et al. (2008) TGF β primes breast tumors for lung metastasis seeding through angiopoietin-like 4. *Cell* 133: 66–77.
24. Zhao H, Ljungberg B, Grankvist K, Rasmuson T, Tibshirani R, et al. (2006) Gene expression profiling predicts survival in conventional renal cell carcinoma. *PLoS Med* 3: e13.
25. Laping NJ, Grygielko E, Mathur A, Butter S, Bomberger J, et al. (2002) Inhibition of transforming growth factor (TGF)- β 1-induced extracellular matrix with a novel inhibitor of the TGF- β type I receptor kinase activity: SB-431542. *Mol Pharmacol* 62: 58–64.
26. Mitropoulos D, Kiroudi A, Christelli E, Serafetinidis E, Zervas A, et al. (2004) Expression of transforming growth factor β in renal cell carcinoma and matched non-involved renal tissue. *Urol Res* 32: 317–322.
27. Jones J, Otu H, Spentzos D, Kolia S, Inan M, et al. (2005) Gene signatures of progression and metastasis in renal cell cancer. *Clin Cancer Res* 11: 5730–5739.
28. Hegele A, Varga Z, von Knobloch R, Heidenreich A, Kropf J, et al. (2002) TGF- β 1 in patients with renal cell carcinoma. *Urol Res* 30: 126–129.
29. Sievers E, Dreimuller P, Haferkamp A, Schmidt-Wolf IG, Buchler MW, et al. (2007) Characterization of primary renal carcinoma cultures. *Urol Int* 79: 235–243.
30. Chen S, Hoffman BB, Lee JS, Kasama Y, Jim B, et al. (2004) Cultured tubule cells from TGF- β 1 null mice exhibit impaired hypertrophy and fibronectin expression in high glucose. *Kidney Int* 65: 1191–1204.
31. Dudas PL, Argentieri RL, Farrell FX (2009) BMP-7 fails to attenuate TGF- β 1-induced epithelial-to-mesenchymal transition in human proximal tubule epithelial cells. *Nephrol Dial Transplant* 24: 1406–1416.
32. Guh JY, Chuang TD, Chen HC, Hung WC, Lai YH, et al. (2003) Beta-hydroxybutyrate-induced growth inhibition and collagen production in HK-2 cells are dependent on TGF- β and Smad3. *Kidney Int* 64: 2041–2051.
33. Humes HD, Beals TF, Cieslinski DA, Sanchez IO, Page TP (1991) Effects of transforming growth factor- β , transforming growth factor- α , and other growth factors on renal proximal tubule cells. *Lab Invest* 64: 538–545.
34. Massague J (2008) TGF β in Cancer. *Cell* 134: 215–230.
35. Asano N, Watanabe T, Kitani A, Fuss IJ, Strober W (2008) Notch1 signaling and regulatory T cell function. *J Immunol* 180: 2796–2804.
36. Blokzijl A, Dahlqvist C, Reissmann E, Falk A, Moliner A, et al. (2003) Cross-talk between the Notch and TGF- β signaling pathways mediated by interaction of the Notch intracellular domain with Smad3. *J Cell Biol* 163: 723–728.
37. Fu Y, Chang A, Chang L, Niessen K, Eapen S, et al. (2009) Differential regulation of transforming growth factor β signaling pathways by Notch in human endothelial cells. *J Biol Chem* 284: 19452–19462.
38. Kennard S, Liu H, Lilly B (2008) Transforming growth factor- β (TGF- β) down-regulates Notch3 in fibroblasts to promote smooth muscle gene expression. *J Biol Chem* 283: 1324–1333.
39. Masuda S, Kumano K, Shimizu K, Imai Y, Kurokawa M, et al. (2005) Notch1 oncoprotein antagonizes TGF- β /Smad-mediated cell growth suppression via sequestration of coactivator p300. *Cancer Sci* 96: 274–282.
40. Niimi H, Pardali K, Vanlandewijck M, Heldin CH, Moustakas A (2007) Notch signaling is necessary for epithelial growth arrest by TGF- β . *J Cell Biol* 176: 695–707.
41. Ono Y, Sensui H, Okutsu S, Nagatomi R (2007) Notch2 negatively regulates myofibroblastic differentiation of myoblasts. *J Cell Physiol* 210: 358–369.
42. Samon JB, Champhekar A, Minter LM, Telfer JC, Miele L, et al. (2008) Notch1 and TGF β 1 cooperatively regulate Foxp3 expression and the maintenance of peripheral regulatory T cells. *Blood* 112: 1813–1821.
43. Sun Y, Lowther W, Kato K, Bianco C, Kenney N, et al. (2005) Notch4 intracellular domain binding to Smad3 and inhibition of the TGF- β signaling. *Oncogene* 24: 5365–5374.
44. Zavadil J, Cermak L, Soto-Nieves N, Bottinger EP (2004) Integration of TGF- β /Smad and Jagged1/Notch signalling in epithelial-to-mesenchymal transition. *Embo J* 23: 1155–1165.
45. Real PJ, Tosello V, Palomero T, Castillo M, Hernando E, et al. (2009) Gamma-secretase inhibitors reverse glucocorticoid resistance in T cell acute lymphoblastic leukemia. *Nat Med* 15: 50–58.
46. Thurston G, Noguera-Troise I, Yancopoulos GD (2007) The Delta paradox: DLL4 blockade leads to more tumour vessels but less tumour growth. *Nat Rev Cancer* 7: 327–331.
47. Lindgren D, Liedberg F, Andersson A, Chebil G, Gudjonsson S, et al. (2006) Molecular characterization of early-stage bladder carcinomas by expression profiles, FGFR3 mutation status, and loss of 9q. *Oncogene* 25: 2685–2696.
48. Hubble J, Demeter J, Jin H, Mao M, Nitzberg M, et al. (2009) Implementation of GenePattern within the Stanford Microarray Database. *Nucleic Acids Res* 37: D898–901.
49. Fredlund E, Ringner M, Maris JM, Pahlman S (2008) High Myc pathway activity and low stage of neuronal differentiation associate with poor outcome in neuroblastoma. *Proc Natl Acad Sci U S A* 105: 14094–14099.
50. Stockhausen MT, Sjolund J, Axelson H (2005) Regulation of the Notch target gene Hes-1 by TGF α induced Ras/MAPK signaling in human neuroblastoma cells. *Exp Cell Res* 310: 218–228.
51. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, et al. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034.
52. Dennler S, Itoh S, Vivien D, ten Dijke P, Huet S, et al. (1998) Direct binding of Smad3 and Smad4 to critical TGF β -inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. *EMBO J* 17: 3091–3100.
53. Aster JC, Robertson ES, Hasserrjan RP, Turner JR, Kieff E, et al. (1997) Oncogenic forms of NOTCH1 lacking either the primary binding site for RBP-Jkappa or nuclear localization sequences retain the ability to associate with RBP-Jkappa and activate transcription. *The Journal of biological chemistry* 272: 11336–11343.
54. Kovacs G, Akhtar M, Beckwith BJ, Bugert P, Cooper CS, et al. (1997) The Heidelberg classification of renal cell tumours. *J Pathol* 183: 131–133.