

RESEARCH ARTICLE

Fact boxes that inform individual decisions may contribute to a more positive evaluation of COVID-19 vaccinations at the population level

Felix G. Rebitschek^{1,2*}, Christin Ellermann¹, Mirjam A. Jenny^{1,2,3,4}, Nico A. Siegel⁵, Christian Spinner⁵, Gert G. Wagner^{1,2,6,7}

1 Harding Center for Risk Literacy, Faculty of Health Sciences Brandenburg, University of Potsdam, Potsdam, Brandenburg, Germany, **2** Max Planck Institute for Human Development, Berlin, Germany, **3** Health Communication Research Group, Implementation Science, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany, **4** Media and Communication Science, University of Erfurt, Erfurt, Thüringen, Germany, **5** Infratest dimap, Berlin, Germany, **6** German Socio-Economic Panel Study (SOEP), Berlin, Germany, **7** Federal Institute for Population Research, Wiesbaden, Hessen, Germany

* rebitschek@uni-potsdam.de



OPEN ACCESS

Citation: Rebitschek FG, Ellermann C, Jenny MA, Siegel NA, Spinner C, Wagner GG (2022) Fact boxes that inform individual decisions may contribute to a more positive evaluation of COVID-19 vaccinations at the population level. PLoS ONE 17(9): e0274186. <https://doi.org/10.1371/journal.pone.0274186>

Editor: Florent Meyniel, INSERM CEA Cognitive Neuroimaging Unit, FRANCE

Received: April 6, 2022

Accepted: August 23, 2022

Published: September 12, 2022

Copyright: © 2022 Rebitschek et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data files are contained in the [Supporting Information](#) files.

Funding: Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer 491466077.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Objective

For an effective control of the SARS-CoV-2 pandemic with vaccines, most people in a population need to be vaccinated. It is thus important to know how to inform the public with reference to individual preferences—while also acknowledging the societal preference to encourage vaccinations. According to the health care standard of informed decision-making, a comparison of the benefits and harms of (not) having the vaccination would be required to inform undecided and skeptical people. To test evidence-based fact boxes, an established risk communication format, and to inform their development, we investigated their contribution to knowledge and evaluations of COVID-19 vaccines.

Methods

We conducted four studies (1, 2, and 4 were population-wide surveys with N = 1,942 to N = 6,056): Study 1 assessed the relationship between vaccination knowledge and intentions in Germany over three months. Study 2 assessed respective information gaps and needs of the population in Germany. In parallel, an experiment (Study 3) with a mixed design (presentation formats; pre-post-comparison) assessed the effect of fact boxes on risk perceptions and fear, using a convenience sample (N = 719). Study 4 examined how effective two fact box formats are for informing vaccination intentions, with a mixed experimental design: between-subjects (presentation formats) and within-subjects (pre-post-comparison).

Results

Study 1 showed that vaccination knowledge and vaccination intentions increased between November 2020 and February 2021. Study 2 revealed objective information requirements

and subjective information needs. Study 3 showed that the fact box format is effective in adjusting risk perceptions concerning COVID-19. Based on those results, fact boxes were revised and implemented with the help of a national health authority in Germany. Study 4 showed that simple fact boxes increase vaccination knowledge and positive evaluations in skeptics and undecideds.

Conclusion

Fact boxes can inform COVID-19 vaccination intentions of undecided and skeptical people without threatening societal vaccination goals of the population.

Introduction

Effective vaccination strategies against the SARS-CoV-2 pandemic require a large proportion of the population to have the vaccine [1]. In December 2020, when the first vaccine was approved, about 37 to 45% of the adult population in Germany definitely intended to get vaccinated, according to population-wide national polls [2, 3]. In December 2021, vaccination rates in Germany approximated a plateau below 85% of people aged between 18 and 59 years, and below 90% of people 60 years and older, respectively [4]. The vaccinated population included a large share of those who were undecideds (who initially did not show a preference for or against vaccination) and skeptics (who more likely would not have had a vaccination initially but were not categorically opposed to it) before (in sum about 24 to 25% of the adult population between April and December 2020 [5, 6]). Before vaccines became available at the end of 2020, undecideds and skeptics sought information to make vaccination decisions [7, 8]. In fact, however, persuasive measures—including political, societal, moral, and social pressure—that do not serve to inform citizens played a substantial role in these decisions population-wide [9].

Health communication that informs and potentially convinces undecideds and skeptics about medical countermeasures during an emergency [10] represents an alternative to persuasive strategies that threaten trustworthiness and credibility of the communicator and of vaccinations [11]. According to international standards of evidence-based health care [12] and the patient protection law in Germany [13]—along with the ethical ideal of evidence-based decision-making [14]—every citizen should be enabled to weigh the possible benefits and harms of medical options on the basis of the best available evidence and to decide freely on this basis (informed decision-making) [15]. Accordingly, the information communicated should enable undecideds and skeptics to compare the benefits and harms of having or not having the vaccination.

The objective of our studies was to investigate how a public health information intervention (“mRNA COVID-19 vaccination fact boxes”) that was jointly developed by the Harding Center for Risk Literacy and the Robert Koch Institute (RKI) might shape the evaluation of COVID-19 vaccination by means of balanced information. Because the best available evidence clearly shows effectiveness (preventing COVID-19 and related hospitalization) and safety (with limited severe side effects), inferred benefit-harm ratios may be convincing. Thus, informed decision making, which considers individual preferences, more likely encourages vaccination and is expected to result in a net surplus of proponents over opponents at the population level. The expected surplus would be in line with the societal preference to increase the vaccinating share of the population in order to protect public health.

A “fact box” is a tabular or graphical version of a balance sheet [16] that summarizes the best available evidence on the benefits and harms of medical options and how likely these will occur [17, 18]. Fact boxes inform various health decisions, including those about medical treatments, cancer screenings, and vaccinations [17, 19, 20]. In contrast to regulation, incentives, and (invisible) nudges [21], fact boxes are not designed to enforce directed behavioral change [22]. They are boosts that have been shown to enable comprehension of medical risks and options and help acquire short-term knowledge thereof [23, 24].

We hypothesized that an intervention with a COVID-19 vaccination fact box not only conveys knowledge and corrects risk perceptions but also can support positive COVID-19 vaccination evaluations without employing persuasion. We conducted four studies to inform the further development of a COVID-19 vaccine fact box and to evaluate its impact on vaccination knowledge and evaluation. First, we assessed the relationship between vaccination knowledge and intention in Germany (Study 1) to determine whether such knowledge potentially contributes to the intention to vaccinate. To inform further development of a COVID-19 vaccination fact box, we identified information requirements and needs of the population in Germany (Study 2). Because accurate risk perception is a prerequisite for informed decision-making [25], Study 3 –conducted in parallel to study 2 –tested whether the fact box improves COVID-19 risk perception, as compared with conventional information. Study 4 evaluated whether different fact box formats are effective for enabling informed vaccination intentions. Studies 1, 2, and 4 were population-wide surveys, each sampling from a multi-level stratified sample from a pool of 72,000 panelists who belonged to the 25 million members of the German PAYBACK customer loyalty program (S1 Fig).

Study 1

To provide a foundation for our plan to promote COVID-19 vaccination knowledge, we assessed the relationship between vaccination knowledge and acceptance in Germany over a three-month period that included introduction of the first vaccines. According to the 5C model [26], which describes the five psychological antecedents of vaccination (confidence in vaccines, structural and psychological constraints, complacency in disease risk perception, collective responsibility in protecting others), the motivation to search for vaccination information is often linked to negative vaccination intention (calculation). On the other hand, vaccination knowledge can be linked to higher vaccination intentions [27] and can close the gap between intentions and actual vaccination decisions [28].

Materials and methods

Survey data from the company Infratest dimap served as the basis for a secondary analysis of vaccination intentions between November 2020 and February 2021. As part of the company’s representative Coronavirus Online Panel Survey Special (Corona-Online-Meinungs-Panel-Survey Spezial, COMPASS), German citizens’ attitudes toward COVID-19 policy, perceptions of constraints, and general attitudes during the pandemic were surveyed. The survey was supplemented by questions on vaccination knowledge. Data collection by the company was exempt from institutional review board approval according to the guidelines of the German Psychological Society DGPS (Deutsche Gesellschaft für Psychologie) [29]. It was reasonable to assume that participation in the commercial survey did not produce harm or discomfort beyond everyday experience. Participants gave written consent to participate and also provided written consent when they registered for repeated multi-topic surveys (e.g. topics such as policy or health), receiving obligatory information on data protection. The study was conducted consistent with the Declaration of Helsinki. The study was not pre-registered.

Participants. The cross-sectional data were obtained from daily surveys of members of the German population [30]. For Study 1, independent samples with (T₁, T₃) and without (T₀, T₂, T₄) knowledge assessment were conducted between 25 November 2020 and 16 February 2021. We do not provide a longitudinal sample analysis because there were different samples across assessments (S1 Fig). Between 13,664 and 13,816 German-speaking respondents with German citizenship were invited per assessment (about 300 per day). Between 14 and 46 of them were not eligible. Among those who were eligible, between 25.4% and 28.0% were non-responders in each case, between 1.0% and 1.5% did not complete the survey, and N = 2,037 (T₀), N = 2,090 (T₁), N = 4,021 (T₂), N = 6,056 (T₃), and N = 1,942 (T₄) were presented with our items. Of the participants, 50.4% to 50.5% were female. They had an average age of 47.2 to 52.1 years, between 31.9% and 36.8% had a university entrance qualification, and about the half of them lived in places with less than 50,000 inhabitants and earned less than 3,500 euros per household per month (S1 Table). Our samples are representative of German citizens who are active online (for details see [31]): The samples were drawn by simple random sampling. To take the varying commitment of different groups in study participation into account (non-response bias), the samples were weighted according to demographic population features. Only data of participants who completed the survey were analyzed. Participants received remuneration worth about 1 euro.

Procedure. After their informed consent to multi-topic study participation, participants received demographic questions, questions about pandemic conditions, an inquiry on their experiences with COVID-19, and a request to evaluate non-pharmaceutical interventions, (e.g. mask wearing obligations) followed by questions on vaccination intention, evaluation, and knowledge items.

Measures. The outcomes of interest in Study 1 were vaccination intention and vaccination knowledge.

Vaccination intention was assessed with a single-option choice ("Vaccines against the coronavirus (COVID-19) are now available. If you get the chance, will you get vaccinated against COVID-19?": Definitely yes, probably yes, probably not, definitely not, I cannot yet say / am still undecided, I am already vaccinated; the last option was introduced when vaccines were available, from T₂ onwards).

Vaccination knowledge was assessed with a focus on vaccination decisions: four items on potential harms (e.g. headache with and without vaccine), and uncertainty (e.g. reduction of contagiousness). The responses were scored according to the best available evidence in December 2020 (S3 Table). Respondents' estimates of how many people get sick with COVID-19 if vaccinated or not after meeting an infected person were elicited using a normalized frequency format (out of 1,000 people) to show understanding of vaccine efficacy. Resulting inferences reflected an underlying risk ratio between the estimations. To avoid zero in the denominator, the division numerator and denominator were adjusted with +1 out of 1,000 (88% to 98% vaccine efficacy was scored as correct based on the results of a meta-analysis, published on 4 February 2021) [32]. For correct answers, we tolerated a +-2% margin of error.

Analysis. Descriptive analyses, linear and logistic regressions, and Spearman rank correlation analyses of vaccination intentions and knowledge scores were conducted.

Results

The cross-sectional analysis of five assessments showed increasing vaccination intentions between November 2020 and February 2021 in Germany. The cumulative proportion of proponents (probably or definitely intending to get the vaccination) grew from 54.4% (T₀, end of Nov) to 65.1% (T₄, mid-February, including 2.2% already vaccinated) (Fig 1). Each assessment




Fig 1. Proportion of respondents according to their intention to get vaccinated against COVID-19 or indicating that they already had. Independent samples were weighted at the time of their assessment. Error bars show 95% confidence intervals.

<https://doi.org/10.1371/journal.pone.0274186.g001>

after T_0 made a proponent on average 1.1 times (odds ratio [OR]) more likely (95%-confidence interval [CI] [1.07, 1.12]), $\chi^2(1) = 48.55, p < .001$, although the proportion of those who probably intended to get vaccinated decreased ($p < .001$). At the same time, the proportion of undecideds declined ($p = .044$). The proportion of opponents (“definitely not”) and skeptics (“probably not”) decreased until actual vaccinations began ($p < .001$), but remained stable in January and February ($p = .228$), at about 12% and 9%, respectively.

Furthermore, between the end of December 2020 and the beginning of February 2021 (when only mRNA vaccines were available in Germany) the samples surveyed showed increasing vaccination knowledge, $F(1,5034) = 40.81, p < .001, \beta_{\text{standardized}} = 0.09$ (Fig 2). Over this period, those with low educational qualification (no formal qualification, primary, or secondary school) showed a similar increase in vaccination knowledge (from mean (M) = 4.40, standard error (SE) = 0.08 to M = 4.67, SE = 0.07; $p < .001$) compared with those with higher educational qualification (university entrance qualification) (from M = 4.68, SE = 0.04 to 4.96, SE = 0.03, $p = .001$).

More positive vaccination intentions were associated with higher knowledge scores (correlation coefficients $r_{T1} = .36$ and $r_{T3} = .36$, each $p < .001$). Among those who were undecided in February, little improvement in knowledge could be observed when compared with undecideds at the turn of the year (+0.1 correct out of 4 test items about vaccine-related uncertainty; compared with +0.3 among proponents).




Fig 2. Number of correct items increased over two months according to respondents' intention to get vaccinated against COVID-19. The independent samples were weighted at the time of their assessment. Error bars show the standard error of the mean.

<https://doi.org/10.1371/journal.pone.0274186.g002>

Discussion

From one month before first vaccine's approval in Germany to two months afterwards, knowledge about the benefits and harms of vaccination increased, as did vaccination intentions. Although this finding cannot be extrapolated to the population's knowledge development over the further course of the pandemic, it supports our assumption that improving vaccination knowledge could contribute to increasing vaccination rates. To improve vaccination knowledge, together with the RKI, the national health authority for disease surveillance and prevention, we developed evidence-based health information in the form of fact boxes.

Study 2

The development process of evidence-based health information ideally involves patients and consumers [33]. First, it is essential to identify and to consider objective informational requirements, as identified by experts [34]: What information do recipients require? What do they not know? What false beliefs do they hold? Second, subjective information needs must be elicited and considered, not only with regard to decision-making but also with regard to developing trustworthy information, which may be unrelated to the expert's view on which information is crucial. On the basis of a secondary analysis of data on perceived reasons for and against vaccination, we assessed the respective information requirements and needs of the population in Germany.

Materials and methods

The study was conducted according to the conditions described under Study 1. The study was not pre-registered.

Participants. Study 2 was based on the sample T_0 ($N = 2,037$) from the end of November 2020, described under Study 1 ([S1 Table](#)).

Design. We used the cross-sectional data from the survey described under Study 1.

Procedure. In the multi-topic survey, reasons were elicited after an inquiry on experiences with COVID-19 and a request to evaluate non-pharmaceutical interventions.

Measures. To identify required and needed information about the COVID-19 vaccination, outcomes of interest in Study 2 were reasons for and against having the vaccination.

Reasons in favor of or against COVID-19 vaccination were asked, with single-choice and open-response items depending on vaccination intention (e.g. for skeptics and opponents: “Why would you not want to get vaccinated if necessary?”).

Analysis. To analyze open responses about reasons in favor of and against COVID-19 vaccination, category systems starting with the subgroup of 18- to 39-year-olds were inductively developed independently by two researchers (one was the author C.E.). Successively, generated codes were reduced and summarized according to the 5C scale [26], a tool to monitor psychological antecedents of vaccination that describes five key elements: confidence (e.g. in the effectiveness and safety of vaccination, of the health care system), complacency (perception of risk), constraints (barriers to execution), calculation (extent of information seeking), and collective responsibility (sense of responsibility for the community). Afterward, the raters compared and consensually agreed on a combined category system with consistent codes for each item ([S5 Table](#)) and coded the responses of the three items again independently from each other. Interrater reliability was high, Cohen’s kappa = .92 (motivation), kappa = .90 (against vaccination), and kappa = .87 (undecided), respectively. Discrepancies in the coding of the individual answers were discussed, a uniform coding was jointly decided upon, and the codes were quantified.

Results

Beyond the answers to closed questions about reasons against vaccination (36%: no threat of a severe COVID-19 course, 20%: pandemic passing without larger harm; [S6 Table](#)), we identified mitigators and facilitators for COVID-19 vaccination communication using open-ended questions: objective information requirements (experts agree that these appertain to everyone making an informed decision or in need of correction, e.g. false beliefs) and subjective information needs of the target population (these are personal and can be absolutely unrelated to experts’ views on informed decision-making, e.g. desired facts about who regulates the vaccine manufacturers).

Predominantly, undecided respondents’ motivations to get vaccinated were related to confidence in the vaccines and the delivering system, such as more medical research (36%), exclusion of harms (14%; more often from 40 years of age onward), and long-lasting high efficacy (7%). Besides those requirements, about 11% of undecideds below 40 years of age explicitly claimed their motivation to depend on more or better information. Motivations of those below 40 were more likely extrinsic (e.g., no contact restrictions, freedom to travel) but also reflected collective responsibility (18% stated that they would agree to vaccination if that would protect others, but only 7% of those aged 40–59 and 4% over 60 mentioned the same).

Nearly all reasons against vaccination named by undecideds, skeptics, and opponents showed information needs related to confidence and trust ([S7 Table](#)): belief in insufficient research on the vaccine and uncertainty about its efficacy and safety (28–52%, increasing with

age), fear of harms (34–49%, decreasing with age), and distrust of policies or the vaccine (11–21%). Personal requirements (8%) and low disease risk perception (7%) played minor roles.

Discussion

Our analysis of open-ended responses showed that confidence-related information needs concerning vaccine efficacy, safety, short- and long-term reactions and harms, and uncertainty are the most essential targets of vaccination communication. This is directly in line with information requirements of health information guidelines [35] and with the aim of building trust.

In parallel to Study 2, a pilot format of the fact box was tested in Study 3 with regard to its effect on risk perception. Insights from both studies were used to inform the further development of a COVID-19 vaccination fact box (from S2A, S2B Fig to Fig 3 and S4–S6 Figs, e.g. by collapsing typical vaccination reactions).

Study 3

Accurate risk perceptions of having or not having an intervention is a prerequisite of informed choices [25]. Although higher disease risk perceptions can be associated with higher vaccination intentions [36–38], overstating the risk of COVID-19 in vaccination communication can not only create fears but can also be detrimental to trustworthiness of medical and scientific experts, which in turn predicts uptake [36]. By communicating transparently and correcting false risk perceptions, fact boxes theoretically can inform the public without undermining trust in the communicating institution. Our online study with a convenience sample controlled whether fact boxes—as evidence-based health information that helps understand the risks of medical options [23, 24]—improve COVID-19 risk perception.

Materials and methods

Ethical approval was obtained from the ethics committee of the University of Potsdam (Germany; 80/2020). The study was not pre-registered.

How safe and effective is the vaccine (mRNA method) against COVID-19 for younger adults?		
The following data refer to vaccinated and non-vaccinated adults between 18 and 60 years of age , some of them with and some of them without pre-existing conditions, using the example of a region with many cases.		
	Out of every 1,000 non-vaccinated younger adults	Out of every 1,000 vaccinated younger adults
Benefits		
How many got sick with COVID-19?	20	1
How many have to be treated at a hospital due to a severe COVID-19 course?	1 – 3 (depending on age and pre-existing conditions)	About 0 (depending on age and pre-existing conditions)
How many suffer from long-term complications of COVID-19, e.g. shortness of breath and memory impairment?	This is still uncertain.	
Harms		
Within one week (after a vaccination appointment), how many suffer temporarily from... ...fatigue and exhaustion?	8	60
...chills?	1	8
How many need fever or pain medication temporarily within one week (after a vaccination appointment)?	140	370
How many suffer severe health damage, e.g. a heart attack, within one month (after a vaccination appointment)?	No difference: about 10 in both groups.	
How many suffer from long-term complications of a COVID-19 vaccine?	There is currently no evidence on long-term complications caused by the vaccine.	
Note: Typical vaccine reactions, e.g. redness or fever, usually subside after one to two days. The potential relationships between the vaccine and rare (e.g. allergies) and atypical reactions (e.g. insomnia, enlarged lymph nodes) as well as transient facial paralysis are currently under examination.		
Sources on the vaccines Comirnaty and Moderna: Baden 2020. NEJM; BioNtech & Pfizer 2020. www.comirnatyeducation.de; CDC 2021. MMWR; EMA 2020. www.ema.europa.eu; FDA 2020. FDA Briefing Document; Polack 2020. NEJM; RKI 2020. Meldedaten; STIKO 2021. Epidemiologisches Bulletin, 2.		
Last update: 24 January 2021 Latest version available at www.hardingcenter.de/en/fact-boxes		

Fig 3. English translation of the simple fact box for people between the ages of 18 and 60 from Study 4.

<https://doi.org/10.1371/journal.pone.0274186.g003>

Participants. Seven hundred and nineteen adults aged 18 to 68 years ($M = 28.8$, $SD = 8.8$), who were recruited as a convenience sample through the online portal Prolific.co, completed Study 3 (demographics in [S2 Table](#)). Participants were eligible if their mother tongue was German and if Germany was their current country of residence. Participants were contacted by email with information about the study and a link to the online survey. They were remunerated with 2.65 British pounds (about 3 euros).

Design. Study 3 had a mixed experimental design: between-subjects variations (random assignment of each participant to one out of six presentation formats) and within-subjects analysis (pre- and post-presentation). The three presentation conditions were (1) vaccine fact box pilot ($n = 120$, “In case of contact with the coronavirus,” [S2A Fig](#)), (2) vaccine fact box pilot with social framing ($n = 123$, “If you spread the coronavirus,” [S2B Fig](#)), and (3) a control condition ($n = 116$; website from www.helios-gesundheit.de [39]), in which standard information on SARS-CoV-2, COVID-19 (without vaccination) and influenza from the internet was presented (between November 20 and February 21 among the top three German search findings comparing SARS-CoV-2 and influenza, [S9 File](#)). Three further presentation conditions for another experiment on disease risks are not reported here. Participants were not aware of the alternative formats. The same introductory text was provided for each condition.

Procedure. After giving informed consent and responding to demographic questions, participants answered items on disease risk perception and fear, on previous adherence to COVID-19 measures, and on vaccination intention and evaluation (baseline assessment). Then they were presented with the intervention (e.g. fact boxes). After reading the presentation formats with evaluation items (e.g. trust in the information presented), the intervention was removed and questions on fear, risk perception, adherence to non-pharmaceutical interventions (e.g. wearing masks, not reported here), and vaccination intentions and evaluations were repeated (post-assessment).

Measures. Outcomes of interest in Study 3 were risk perception, fear of getting or spreading COVID-19, and vaccination intention.

Risk perception and fear. Two items measured the fear of getting or spreading COVID-19 (“How greatly do you fear contracting COVID-19/infecting other adults?: Please use a scale from 0 to 10. A value of 0 means no fear at all and a value of 10 means a lot of fear. You can also use one of the values in between to indicate your rating.”). One item measured perceived severity (“How severe would an illness with COVID-19 be for you personally? Please use a scale from 0 to 10. A value of 0 means not at all severe and a value of 10 means very severe. You can also use one of the values in between to indicate your rating.”). Numerical risk perception was measured with one item employing a frequency format (Please imagine 1,000 adults like yourself. How many of these adults will contract COVID-19 as a result of close contact with someone infected with the virus?): Scoring of correct responses (5–28%) considered the range of the best available evidence, which can be found in [S4 Table](#) together with the medical references. For control, all items were presented with numbers for influenza as well, but were not the subject of this study.

Vaccination intention. Participants were asked if they planned to get vaccinated within the next six months.

Analysis. Besides descriptive statistics, repeated measures analyses of variance were conducted to compare the presentation conditions.

Results

Studying the fact boxes decreased ([Table 1](#)) numeric COVID-19 risk perception (ANOVA interaction effect: $F(1,357) = 10.05$, $p = .002$, $\eta_p^2 = 0.03$) compared with control presentation,

Table 1. Personal and social fear, subjective and numeric risk perception across conditions, diseases, pre- and post-presentation.

Presentation format	Disease	Personal fear		Social fear		Subjective risk perception		Numeric risk perception (out of 1,000)	
		Pre	Post	Pre	Post	Pre	Post	Pre	Post
		M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
Vaccine fact box	COVID-19	5.8 (2.6)	6.0 (2.7)	6.1 (2.8)	6.2 (2.8)	6.3 (2.5)	6.4 (2.6)	327 (294)	248 (222)
	Influenza	3.4 (2.4)	3.5 (2.5)	3.6 (2.5)	3.7 (2.7)	3.6 (2.4)	3.7 (2.6)	257 (246)	190 (192)
Social framing box	COVID-19	5.5 (2.5)	5.7 (2.6)	6.3 (2.4)	6.4 (2.4)	5.9 (2.4)	6.0 (2.3)	345 (323)	259 (245)
	Influenza	2.8 (2.1)	2.9 (2.1)	3.1 (2.0)	3.3 (2.2)	3.6 (2.0)	3.4 (2.2)	288 (294)	220 (242)
Standard information	COVID-19	5.5 (2.5)	6.0 (2.5)	6.3 (2.4)	6.9 (2.4)	5.5 (2.5)	6.2 (2.4)	317 (323)	314 (281)
	Influenza	2.9 (2.0)	3.4 (2.4)	3.2 (2.1)	4.0 (2.4)	3.3 (2.2)	3.8 (2.5)	207 (220)	171 (190)

<https://doi.org/10.1371/journal.pone.0274186.t001>

leading to more accurate estimates (see [S3 Fig](#), $p < .001$). Only control presentation increased ([Table 1](#)) both personal fear (ANOVA interaction effect of time and presentation format, with $F(1,357) = 4.17$, $p = .042$, $n_p^2 = 0.01$) and perceived severity of developing COVID-19 (interaction, with $F(1,357) = 19.90$, $P < .001$, $n_p^2 = 0.05$).

More positive evaluations ($F(1,357) = 12.55$, $p < .001$, $n_p^2 = 0.03$) and increasing intentions to get vaccinated ($F(1,357) = 7.63$, $p = .006$, $n_p^2 = 0.02$) were shown as ANOVA main effects, but there were no format-specific interaction effects ([S8 Table](#)). Here, control information may have been as effective as fact boxes, albeit by promoting fear and perceived severity.

Discussion

Presenting the vaccination fact box pilot decreased numerical disease risk perception and promoted accurate risk perceptions, without affecting vaccination intentions. At the same time, perceived severity of the disease and fear of contracting or spreading COVID-19 were not stimulated by the fact box presentation. Prompting fear responses would have posed the risk of interfering with informed decision-making [40]. Findings supported the further development and implementation of the fact box format and appeared to confirm the assumed relationship between fact box presentation and vaccination evaluation and intention.

Study 4

Based on the insights of Studies 1, 2, and 3, Study 4 examined whether different fact box formats are effective for enabling informed vaccination intentions in a population-wide sample. Undecideds and skeptics needed and desired information (Study 2) to weigh potential benefits and harms [7]; otherwise they were hesitant to get vaccinated [8]. Study 4 tested whether both complex and simple fact boxes—based on the implemented version—would improve vaccination knowledge and modify vaccination evaluation or intention.

Materials and methods

The study was conducted according to the conditions described under Study 1. Despite having received ethical approvals for similar fact box studies before [23, 24], we sought and received approval for the study design on mRNA fact boxes (IRB of the Max Planck Institute for Human Development, A 2021–07). The study was not pre-registered.

Participants. Study 4 was based on the sample T₃ ([S1 Table](#)), described under Study 1. We excluded respondents ($n = 182$, 6.1%) who spent more than fivefold the average time (> about 18 minutes) on reading a fact box or on completing a question, assuming that they had likely turned to other activities.

Design. The experiment had a mixed experimental design, with between-subjects variation (random assignment of participants to one of three presentation conditions) and with repeated assessments of vaccination evaluation within-subjects (pre- and post-presentation). The three presentation conditions were (1) simple fact box ($n = 984$) (see Fig 3 and S4 Fig), complex fact box ($n = 974$) (see S5 and S6 Figs), and (3) one group without information presentation ($n = 991$). The formats varied for separate age groups (18–59 years vs. 60 onward). Participants were not aware of the alternative conditions. The same introductory text was provided for each condition.

Procedure. The procedure for the surrounding survey was described under Study 1. In Study 4, the vaccine efficacy item pair was presented either together with fact boxes (complex or simple) or without a stimulus (control group). Knowledge recall and a second vaccination evaluation were prompted after intervention.

Measures. Outcomes of interest in Study 4 were vaccination intention, evaluation, and vaccination knowledge.

Vaccination evaluation was assessed with participants' ratings of the benefit-harm ratio of the COVID-19 vaccination in question on a 11-point rating scale from 0 (harms clearly outweigh the benefits) to 10 (benefits clearly outweigh the harms).

Vaccination intention and knowledge were assessed as described under Study 1. After information presentation, five items with true-false statements tested participants' recall of vaccination safety (fatigue, serious adverse events), uncertainty (later harm, facial paresis), and efficacy.

Study time (in seconds) was the time that participants took for responding to the vaccine efficacy item pair, while some of them were presented with the interventions.

Analyses. We used variance analyses of intervention effects on knowledge and comprehension, and additional (logistic) regression analyses to examine factors influencing knowledge of and the evaluation of the vaccination (age group-specific formats, fact box formats, education, and income).

Results

First, compared with no treatment, the sum score of vaccination knowledge was higher after fact box presentations (ANOVA main effect, $F(1,3101) = 36.58$, $p < .001$, $\eta_p^2 = 0.01$). Respondents of both observed age groups (below 60 and 60 onward) recalled vaccine efficacy, safety, and related uncertainties differentially when presented with a fact box instead of nothing (S9 Table). Multiple logistic regressions including the factors presentation format and age group showed that respondents of both age groups, who were presented with the fact box instead of nothing, were more likely aware of the side effect of fatigue (Odds Ratio (OR) = 1.85 with a 95%-confidence interval (CI) [1.69, 2.01]). Compared to those without fact box, respondents with fact box presentation, aged 60 and above, more likely confirmed a potential risk of facial paresis (OR = 1.16 [1.04, 1.28]), and younger respondents more likely recalled vaccine efficacy (OR = 1.13 [1.02, 1.24]).

Finally, vaccine efficacy was (OR = 1.85 [1.53, 2.17], $p < .001$) more likely correctly inferred when a fact box was presented (logistic regression controlling for the level of education).

Second, logistic regressions ($\chi^2(1) = 14.70$, $p < .001$ and $\chi^2(1) = 5.50$, $p = .020$) revealed that fact boxes more likely prompted both any change (OR = 1.36 [1.20, 1.52], $p < .001$) and a positive change (OR = 1.25 [1.06, 1.44], $p = .020$) of the evaluation of the vaccination compared with no intervention. For a distinct consideration of the different formats: Whereas 18.6% of respondents without any intervention changed their evaluation of a COVID-19 vaccination positively, 20.3% did so if studying a complex fact box, and 24.2% if studying a simple fact box.

At the same time, however, 14.5%, 19.8%, and 16.3%, respectively, evaluated the evaluation more negatively when asked a second time (at post-assessment). Compared with no intervention, presenting a simple fact box ($\chi^2(1) = 9.52, p = .002$) more likely led to a positive than to a non-positive change (OR = 1.40 [1.19, 1.62], $p = .002$) in the evaluation of the vaccination. This could not be confirmed for the complex box ($\chi^2(1) = 0.90, p = .343$).

The shift in vaccination evaluation after being presented simple fact boxes (+7.9%) could, to a substantial extent, be related to the skeptical and undecided respondents' comprehension of the information presented. Those who drew correct inferences about vaccine efficacy after having seen simple fact boxes showed a positive change in evaluation, but not those who drew incorrect inferences (main effect of an univariate ANOVA, $F(1,467) = 3.88, p < .050, n_p^2 = 0.01$). Explorative sub-analyses highlighted that this effect is due to the younger skeptics and undecideds ($F(1,387) = 5.65, p = .018, n_p^2 = 0.01$), not to those aged 60 and above ($F(1,76) = 0.04, p = .836$). In contrast, younger skeptics' and undecideds' recall, with at least 80% of correct responses after information presentation, was not related to positive changes in evaluation ($F(1,387) = 0.59, p = .445$).

Taking into consideration participants with different levels of education, we compared simple and complex fact boxes with no treatment. Respondents with the lowest educational qualification (16.5% with no school degree or degree after 9 years of schooling) recalled less than those with higher qualifications (degree after 10 years, university entrance qualification) (ANOVA main effect of sum score, $F(1,3086) = 11.44, p = .001, n_p^2 < 0.01$). But no interaction indicated an education-related recall benefit over the control condition ($F(2,3086) = 0.20, p = .820$).

Additionally, we compared the two types of fact boxes. Recall of information from simple (59%) but not from complex fact boxes (61%) was significantly lower for respondents with low to moderate levels of education compared with those with higher levels (64% and 62%, respectively, ANOVA interaction effect, $F(1,2048) = 4.10, p = .043, n_p^2 < 0.01$). An additional analysis that split the low to moderately educated people according to their income median confirmed, across both age groups, that higher education, not income, was associated with recall of information from simple fact boxes (Fig 4A and 4B).




Fig 4. Proportion of correct responses to five knowledge items according to different levels of education and household net income for respondents younger than 60 years of age (A) and 60 years of age and older (B). The sample is weighted. Error bars show the standard error of the mean.

<https://doi.org/10.1371/journal.pone.0274186.g004>

For complex fact boxes, the proportion of quick responders (below median response time of the control condition) among those with the lowest educational attainment was lower (7%; compared to those with higher education: 15%), which indicates that they needed more time for reading and/or for deciding to skip reading. In contrast, for simple fact boxes, the proportion of quick responders did not appear to depend on the level of education (in both groups about 11–12%).

Discussion

People who read the fact boxes can be expected to gain a better comprehension of and short-term knowledge about the mRNA vaccination. Furthermore, their evaluation of the mRNA vaccination changes positively, particularly if those who were initially undecided and skeptical comprehend vaccine efficacy, e.g. via the simple fact box. A separate analysis could not confirm this net surplus for complex fact boxes. Furthermore, people with lower educational attainment seem to be less likely enabled to make an informed decision than higher educated groups, even when they receive this more detailed and complex version of the fact box. This indicates a potential risk of increasing health inequality by using evidence-based fact boxes, from which people with higher educational attainment benefit disproportionately.

General discussion

The association of vaccination knowledge and uptake implies one crucial mechanism with respect to the goal of an adequately immunized society. Besides showing a differential increase in vaccination knowledge in proponents, undecideds, and opponents around the start of COVID-19 vaccination in Germany, Study 1 showed increasing intentions to have the (mRNA) vaccinations in the early phase of vaccine distribution [2]. Study 2 found that both undecideds and skeptics lacked various pieces of information for making a decision, paired with false beliefs and a lack of trust in vaccine safety and efficacy. Study 3 revealed that a COVID-19 vaccination fact box improved disease risk perception without increasing COVID-19 fears (unlike in the control condition). Based on these findings, a previously implemented vaccination fact box was amended. Applied at the population level in Study 4, fact boxes boosted knowledge of undecideds and skeptics together with a more positive evaluation of the vaccination's benefit-harm ratio. Studying simple or complex fact boxes instead of nothing was 1.3 times more likely to lead to any positive change in evaluation. In the case of complex fact boxes, however, any changes seem to be balanced.

Although a common factor (e.g. peer behaviors) admittedly may underlie information acquisition and intentions—while stable vaccination preferences may prompt differential information acquisition—knowledge about benefits and harms of COVID-19 vaccinations, which can reduce vaccine hesitancy [41], may lead to informed intentions for readers of fact boxes if they base their intentions on these facts. This mechanism, in line with the standards of health care systems in many democratic countries, also contributes to the trustworthiness of authorities engaged in vaccination risk communication. For example, in face of the fact that denying information gaps can undermine perceived credibility [42, 43], fact boxes contain epistemic uncertainty disclaimers (e.g. verbal statements or ranges as a result of incomplete knowledge about the risk numbers) [18]. Transparently informing the public about limitations of vaccinations, which considers individual preferences, does not necessarily reduce vaccination intentions [44], which can support the societal preference to increase the vaccinating share of the population. By contrast, persuasive communications, e.g. messages framed in relative risks, can increase vaccination intentions [45] but are both misinformative and incomprehensible [46], colliding with the rights and needs of undecided and skeptical citizens. In addition, incomprehensible information may provoke a backlash effect. Future research could

investigate the causal link between the grade of comprehensibility of information about benefits and harms and people's vaccination intentions.

Limitations of our study: In our population-wide sample, simple fact boxes appear to be more beneficial for those with higher education. This contradiction to the design intention could be due to a superficial reading of the simple box by participants with lower levels of education, as indicated by their brief reading times. Although information needs and requirements were elicited, the target group with lower educational attainment did not actively participate in the development process of the fact box intervention (e.g., Study 3 included a highly educated convenience sample, separate piloting was missing). However, the potential of shared decision-making interventions to reduce health inequities [47] is tied to the involvement of disadvantaged groups in their development [48]. If neglected, there is a risk of increasing health inequity [49], e.g. due to the heterogeneity in the audience's ability to comprehend given information. Those who comprehend the least, potentially benefit the least [45], e.g. low-numerate breast cancer patients [50]. Also, cancer screening fact boxes let high-numerate people update their benefit perceptions more strongly than low-numerates [22]. This could indicate distorted interpreting of fact boxes. Thus, more research needs to examine under which conditions fact boxes can strengthen informed decision-making of disadvantaged groups and reduce health inequities. Furthermore, concepts such as critical health literacy or digital health literacy should be taken into account to examine determinants that influence who benefits or does not benefit from fact boxes and how this affects health inequities [51]. Additional factors associated with formal education can also lead to inequality (e.g. working and living conditions). Future studies should thus incorporate approaches such as the PROGRESS Plus framework, which describes inequity-generating factors at multiple levels.

An additional limitation of our work concerns the set of vaccination knowledge items that covered certain requirements of health information guidelines (benefit, harms, uncertainty) but were not a validated scale of vaccination knowledge.

The results are hardly transferable to the non-democratic contexts, because, alongside the differences in media, political handling of the pandemic, and access to vaccination, the needs and preferences for the content and design of health information depend on a population's risk perception, numerical literacy, and the opportunity for informed decision-making. This suggests that the success of a complex public health intervention, e.g. the provision of fact boxes by a central health authority, depends on a number of factors related to the societal characteristics of a country or region.

We conclude that in Germany, the implementation of fact boxes [52] supports evidence-based communication and, thus, empowerment at the population level. Dissemination of the mRNA fact boxes via the RKI media channels included their webpage with about 130 million visits in 2020, an RKI Twitter account, and the cellphone app of the Permanent Vaccination Commission in Germany. Our figures imply that presenting the about 11 million undecided and skeptical adults under 60 years of age in Germany with a simple fact box for about 90 seconds would familiarize more than 600,000 people with vaccine efficacy. A majority of them would evaluate vaccinations more positively. Also, by abstaining from persuasion, reluctance and distrust concerning the sender can be prevented or alleviated. Loss of trustworthiness is a relevant risk of alternative interventions that are nontransparent (e.g., nudges), persuasive (e.g., advertisement), or that exert pressure (e.g., for authorized access to public activities).

Finally, informed decision-making is not a threat to the goal of an adequately immunized population. To achieve this, ensuring the right of each individual to decide about their own health might be sufficient, while other types of social contracts [53] could be less important. The standard of evidence-based health care, legally binding in Germany [12] and benchmark of a democratic society, would ensure responsible vaccination decisions and future

commitment when refreshments of the vaccination might be necessary or when individuals might have to decide to get vaccinated against other diseases.

Supporting information

S1 Fig. Overview of used data sets and study samples. *Data from the Waves 0 to 5 were not analyzed for this paper.

(TIF)

S2 Fig. Experimental fact boxes from Study 3 about benefits and harms of the mRNA vaccine “In case of contact with coronavirus” (A), with social framing “If you spread coronavirus” (B).

(TIF)

S3 Fig. Accuracy of COVID-19 risk perception (responding with a 5–28% probability of developing COVID-19 given close contact to an infected person; evidence from October 2020). Error bars show 95% confidence intervals.

(TIF)

S4 Fig. English translation of the simple fact box for people 60 years and older from Study 4.

(TIF)

S5 Fig. English translation of the complex fact box for people between the ages of 18 and 60 from Study 4.

(TIF)

S6 Fig. English translation of the complex fact box for people 60 years and older from Study 4.

(TIF)

S1 Table. Descriptions of the quasi-representative samples (Studies 1, 2, and 4) underlying T₀, T₁, T₂, T₃, and T₄. The data are not weighted.

(DOCX)

S2 Table. Sample descriptions for Study 3, according to fact box conditions vs. control condition and across the study conditions).

(DOCX)

S3 Table. Responses to open questions of undecideds (pro and contra vaccination), skeptics, and opponents (contra only).

(DOCX)

S4 Table. Intention to get vaccinated across presentation formats and pre- and post-assessment in Study 3.

(DOCX)

S5 Table. Proportion of correct responses to five knowledge items (true-false statements) after information presentation (recall).

(DOCX)

S6 Table. Reasons for and against COVID-19 vaccination based on multiple-choice items (Study 2) that were presented depending on prior vaccination intentions. The data are weighted.

(DOCX)

S7 Table. Items, plus best available evidence in December 2020 for the studies 1, 2, and 4.

(DOCX)

S8 Table. Best available evidence in October 2020 on transmissibility in a close-contact setting and manifestation rate given an infection of SARS-CoV-2 and influenza virus, focusing on the age groups 20–39, 40–59, and 60–79 years. Min and max represent a range that includes dark figures, vaccine efficacy, and conflicting study estimates.

(DOCX)

S9 Table. Category system about information requirements (knowledge gaps) and (subjective) needs of vaccination proponents, undecideds, skeptics, and opponents.

(DOCX)

S10 Table. Sample descriptions for Study 4, according to simple vs complex fact box conditions vs. control condition and across the study conditions. The data are not weighted.

(DOCX)

S1 File. Data file of the quasi-representative sample underlying T₀.

(SAV)

S2 File. Data file of the quasi-representative sample underlying T₁.

(SAV)

S3 File. Data file of the quasi-representative sample underlying T₂.

(SAV)

S4 File. Data file of the quasi-representative sample underlying T₃.

(SAV)

S5 File. Data file of the quasi-representative sample underlying T₄.

(SAV)

S6 File. Syntax file of the analyses of Study 1.

(SPS)

S7 File. Syntax file of the analyses of Study 3.

(SPS)

S8 File. Syntax file of the analyses of Study 4.

(SPS)

S9 File. Stimulus of the control condition in Study 3.

(PDF)

S10 File. Data file of the convenience sample of Study 3.

(SAV)

S11 File. Intention data file including samples underlying T₀ to T₄.

(SAV)

S12 File. Knowledge data file combining samples underlying T₀ to T₄.

(SAV)

Acknowledgments

We thank our proofreader Rona Unrau for her critical eye and Carolin Harder for her coding of open-ended item responses. Furthermore, we thank Gudrun Keller for programming the COMPASS items.

Author Contributions

Conceptualization: Felix G. Rebitschek, Nico A. Siegel, Gert G. Wagner.

Data curation: Christian Spinner.

Formal analysis: Felix G. Rebitschek, Christin Ellermann.

Investigation: Felix G. Rebitschek, Christin Ellermann, Nico A. Siegel, Christian Spinner, Gert G. Wagner.

Methodology: Felix G. Rebitschek.

Writing – original draft: Felix G. Rebitschek, Christin Ellermann, Mirjam A. Jenny, Gert G. Wagner.

Writing – review & editing: Felix G. Rebitschek, Christin Ellermann, Mirjam A. Jenny, Nico A. Siegel, Christian Spinner, Gert G. Wagner.

References

1. Bartsch SM, O'Shea KJ, Ferguson MC, Bottazzi ME, Wedlock PT, Strych U, et al. Vaccine efficacy needed for a COVID-19 coronavirus vaccine to prevent or stop an epidemic as the sole intervention. *American Journal of Preventive Medicine*. 2020; 59(4):493–503. <https://doi.org/10.1016/j.amepre.2020.06.011> PMID: 32778354
2. Haug S, Schnell R, Weber K. Impfbereitschaft mit einem COVID-19-Vakzin und Einflussfaktoren. Ergebnisse einer telefonischen Bevölkerungsbefragung. *Das Gesundheitswesen*. 2021; 83(10):789–96.
3. Robert Koch Institute. COVID-19 Impfquoten-Monitoring in Deutschland (COVIMO) -1. Report. 2021.
4. Bundesministerium für Gesundheit. Impfdashboard 2022 [5 August 2022]. Available from: <https://impfdashboard.de/>.
5. Neumann-Böhme S, Varghese NE, Sabat I, Barros PP, Brouwer W, van Exel J, et al. Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *The European Journal of Health Economics*. 2020; 21(7):977–82. <https://doi.org/10.1007/s10198-020-01208-6> PMID: 32591957
6. Wissenschaft im Dialog. Wissenschaftsbarometer 2020—Report. 2020.
7. Quinn SC, Kumar S, Freimuth VS, Kidwell K, Musa D. Public willingness to take a vaccine or drug under Emergency Use Authorization during the 2009 H1N1 pandemic. *Biosecurity and Bioterrorism: Biodefense strategy, Practice, and Science*. 2009; 7(3):275–90. <https://doi.org/10.1089/bsp.2009.0041> PMID: 19775200
8. Betsch C, Korn L, Felgendreff L, Eitze S, Schmid P, Sprengholz P, et al. German COVID-19 Snapshot Monitoring (COSMO)-Welle 15 (23.06.2020). PrePrint. 2020.
9. Thießen M. Immunity as Relativity: German Vaccination Campaigns and Debates in Times of COVID-19. *Historical Social Research/Historische Sozialforschung*. 2021; 46(4):316–38.
10. Schoch-Spana M, Brunson E, Chandler H, Gronvall GK, Ravi S, Sell TK, et al. Recommendations on how to manage anticipated communication dilemmas involving medical countermeasures in an emergency. *Public Health Reports*. 2018; 133(4):366–78. <https://doi.org/10.1177/0033354918773069> PMID: 29847750
11. Blastland M, Freeman AL, van der Linden S, Marteau TM, Spiegelhalter D. Five rules for evidence communication. *Nature*. 2020; 587(7834):362–4. <https://doi.org/10.1038/d41586-020-03189-1> PMID: 33208954
12. Braddock CH III, Edwards KA, Hasenberg NM, Laidley TL, Levinson W. Informed decision making in outpatient practice: time to get back to basics. *JAMA*. 1999; 282(24):2313–20. <https://doi.org/10.1001/jama.282.24.2313> PMID: 10612318
13. Bundesministerium für Gesundheit. Gesetz zur Verbesserung der Rechte von Patientinnen und Patienten. Bundesanzeiger Verlag, 2013.
14. Kerridge I, Lowe M, Henry D. Ethics and evidence based medicine. *BMJ*. 1998; 316(7138):1151–3. <https://doi.org/10.1136/bmj.316.7138.1151> PMID: 9552959

15. Elmore JG, Ganschow PS, Geller BM. Communication between patients and providers and informed decision making. *Journal of the National Cancer Institute Monographs*. 2010; 2010(41):204–9. <https://doi.org/10.1093/jncimonographs/lgq038> PMID: 20956831
16. Eddy DM. Comparing benefits and harms: the balance sheet. *JAMA* 1990; 263(18):2493–505. <https://doi.org/10.1001/jama.263.18.2493> PMID: 2329639
17. Schwartz LM, Woloshin S, Welch HG. Using a drug facts box to communicate drug benefits and harms: two randomized trials. *Annals of Internal Medicine*. 2009; 150(8):516–27. <https://doi.org/10.7326/0003-4819-150-8-200904210-00106> PMID: 19221371
18. McDowell M, Rebitschek FG, Gigerenzer G, Wegwarth O. A simple tool for communicating the benefits and harms of health interventions. *MDM Policy & Practice*. 2016; 1(1):2381468316665365.
19. AOK-Bundesverband, Harding Center for Risk Literacy. AOK-Faktenboxen 2016 [8 August 2022]. Available from: <http://www.aok.de/faktenboxen>.
20. Harding Center for Risk Literacy. Fact Boxes 2021 [8 August 2022]. Available from: <https://www.hardingcenter.de/en/transfer-and-impact/fact-boxes>.
21. Hertwig R, Grüne-Yanoff T. Nudging and boosting: Steering or empowering good decisions. *Perspectives on Psychological Science*. 2017; 12(6):973–86. <https://doi.org/10.1177/1745691617702496> PMID: 28792862
22. Eber MR, Sunstein CR, Hammitt JK, Yeh JM. The modest effects of fact boxes on cancer screening. *Journal of Risk and Uncertainty*. 2021; 62(1):29–54. <https://doi.org/10.1007/s11166-021-09344-x> PMID: 34385760
23. McDowell M, Gigerenzer G, Wegwarth O, Rebitschek FG. Effect of tabular and icon fact box formats on comprehension of benefits and harms of prostate cancer screening: A randomized trial. *Medical Decision Making*. 2019; 39(1):41–56. <https://doi.org/10.1177/0272989X18818166> PMID: 30799691
24. Brick C, McDowell M, Freeman AL. Risk communication in tables versus text: a registered report randomized trial on 'fact boxes'. *Royal Society Open Science*. 2020; 7(3):190876. <https://doi.org/10.1098/rsos.190876> PMID: 32269779
25. Trevena LJ, Zikmund-Fisher BJ, Edwards A, Gaissmaier W, Galesic M, Han PK, et al. Presenting quantitative information about decision outcomes: a risk communication primer for patient decision aid developers. *BMC Medical Informatics and Decision Making*. 2013; 13(2):1–15. <https://doi.org/10.1186/1472-6947-13-S2-S7> PMID: 24625237
26. Betsch C, Schmid P, Heinemeier D, Korn L, Holtmann C, Böhm R. Beyond confidence: Development of a measure assessing the 5C psychological antecedents of vaccination. *PLoS ONE*. 2018; 13(12): e0208601. <https://doi.org/10.1371/journal.pone.0208601> PMID: 30532274
27. Zheng H, Jiang S, Wu Q. Factors influencing COVID-19 vaccination intention: The roles of vaccine knowledge, vaccine risk perception, and doctor-patient communication. *Patient Education and Counseling*. 2022; 105(2):277–83. <https://doi.org/10.1016/j.pec.2021.09.023> PMID: 34565643
28. Wegwarth O, Kurzenhäuser-Carstens S, Gigerenzer G. Overcoming the knowledge–behavior gap: The effect of evidence-based HPV vaccination leaflets on understanding, intention, and actual vaccination decision. *Vaccine*. 2014; 32(12):1388–93. <https://doi.org/10.1016/j.vaccine.2013.12.038> PMID: 24486360
29. Deutsche Gesellschaft für Psychologie. Ethisches Handeln in der psychologischen Forschung. Empfehlungen der Deutschen Gesellschaft für Psychologie für Forschende und Ethikkommissionen. Göttingen (Germany): Hogrefe Verlag; 2018.
30. Siegel N, Wagner GG. infratest: Corona Online Meinungs Panel Survey Spezial (COMPASS) 2021 [8 August 2022]. Available from: www.infratest-dimap.de/umfragen-analysen/bundesweit/coronacompass/coronacompass/.
31. Wegwarth O, Wagner GG, Spies C, Hertwig R. Assessment of German public attitudes toward health communications with varying degrees of scientific uncertainty regarding COVID-19. *JAMA network open*. 2020; 3(12):e2032335–e. <https://doi.org/10.1001/jamanetworkopen.2020.32335> PMID: 33301021
32. STIKO. Beschluss der STIKO zur 2. Aktualisierung der COVID-19-Impfempfehlung und die dazugehörige wissenschaftliche Begründung. *Epidemiologisches Bulletin*. 2021;5.
33. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ*. 2008; 337:a1655. <https://doi.org/10.1136/bmj.a1655> PMID: 18824488
34. Bunge M, Mühlhauser I, Steckelberg A. What constitutes evidence-based patient information? Overview of discussed criteria. *Patient Education and Counseling*. 2010; 78(3):316–28. <https://doi.org/10.1016/j.pec.2009.10.029> PMID: 20005067

35. Lühnen J, Albrecht M, Hanßen K, Hildebrandt J, Steckelberg A. Leitlinie evidenzbasierte Gesundheit-sinformation: Einblick in die Methodik der Entwicklung und Implementierung. *Zeitschrift für Evidenz, Fortbildung und Qualität im Gesundheitswesen*. 2015; 109(2):159–65. <https://doi.org/10.1016/j.zefq.2015.03.004> PMID: 26028454
36. Kerr JR, Schneider CR, Recchia G, Dryhurst S, Sahlin U, Dufouil C, et al. Correlates of intended COVID-19 vaccine acceptance across time and countries: results from a series of cross-sectional surveys. *BMJ Open*. 2021; 11(8):e048025. <https://doi.org/10.1136/bmjopen-2020-048025> PMID: 34341047
37. Ling M, Kothe EJ, Mullan BA. Predicting intention to receive a seasonal influenza vaccination using Protection Motivation Theory. *Social Science & Medicine*. 2019; 233:87–92. <https://doi.org/10.1016/j.socscimed.2019.06.002> PMID: 31195194
38. Mesch GS, Schwirian KP. Social and political determinants of vaccine hesitancy: Lessons learned from the H1N1 pandemic of 2009–2010. *American Journal of Infection Control*. 2015; 43(11):1161–5. <https://doi.org/10.1016/j.ajic.2015.06.031> PMID: 26521933
39. Helios Kliniken. Corona versus Grippe: Der Unterschied liegt im Detail [15 December 2020]. Available from: <https://www.helios-gesundheit.de/magazin/corona/news/corona-versus-grippe-was-ist-gefaehrlicher/>.
40. Zikmund-Fisher BJ, Fagerlin A, Ubel PA. Risky feelings: why a 6% risk of cancer does not always feel like 6%. *Patient Education and Counseling*. 2010; 81:87–93.
41. Palm R, Bolsen T, Kingsland JT. The effect of frames on COVID-19 vaccine resistance. *Frontiers in Political Science*. 2021; 3(41):661257.
42. Frewer LJ, Miles S, Brennan M, Kuznesof S, Ness M, Ritson C. Public preferences for informed choice under conditions of risk uncertainty. *Public Understanding of Science*. 2002; 11(4):363–72.
43. Renn O, Levine D. Credibility and trust in risk communication. In: Kasperson RE, Stallen PJM, editors. *Communicating Risks to the Public: International Perspectives*. Dordrecht (Netherlands): Springer; 1991. p. 175–217.
44. Kerr JR, Freeman AL, Marteau TM, van der Linden S. Effect of information about COVID-19 vaccine effectiveness and side effects on behavioural intentions: two online experiments. *Vaccines*. 2021; 9(4):379. <https://doi.org/10.3390/vaccines9040379> PMID: 33924542
45. Lawes-Wickwar S, Ghio D, Tang MY, Keyworth C, Stanescu S, Westbrook J, et al. A rapid systematic review of public responses to health messages encouraging vaccination against infectious diseases in a pandemic or epidemic. *Vaccines*. 2021; 9(2):72. <https://doi.org/10.3390/vaccines9020072> PMID: 33498395
46. Akl EA, Oxman AD, Herrin J, Vist GE, Terrenato I, Sperati F, et al. Using alternative statistical formats for presenting risks and risk reductions. *The Cochrane Database of Systematic Reviews*. 2011;(3): CD006776. <https://doi.org/10.1002/14651858.CD006776.pub2> PMID: 21412897
47. Durand M-A, Carpenter L, Dolan H, Bravo P, Mann M, Bunn F, et al. Do interventions designed to support shared decision-making reduce health inequalities? A systematic review and meta-analysis. *PloS ONE*. 2014; 9(4):e94670. <https://doi.org/10.1371/journal.pone.0094670> PMID: 24736389
48. Enard KR, Dolan Mullen P, Kamath GR, Dixon NM, Volk RJ. Are cancer-related decision aids appropriate for socially disadvantaged patients? A systematic review of US randomized controlled trials. *BMC Medical Informatics and Decision Making*. 2016; 16(1):1–15. <https://doi.org/10.1186/s12911-016-0303-6> PMID: 27267490
49. Lorenc T, Petticrew M, Welch V, Tugwell P. What types of interventions generate inequalities? Evidence from systematic reviews. *J Epidemiol Community Health*. 2013; 67(2):190–3. <https://doi.org/10.1136/jech-2012-201257> PMID: 22875078
50. Lipkus IM, Peters E, Kimmick G, Liotcheva V, Marcom P. Breast cancer patients' treatment expectations after exposure to the decision aid program adjuvant online: the influence of numeracy. *Medical Decision Making*. 2010; 30(4):464–73. <https://doi.org/10.1177/0272989X09360371> PMID: 20160070
51. O'Neill J, Tabish H, Welch V, Petticrew M, Pottie K, Clarke M, et al. Applying an equity lens to interventions: using PROGRESS ensures consideration of socially stratifying factors to illuminate inequities in health. *Journal of Clinical Epidemiology*. 2014; 67(1):56–64. <https://doi.org/10.1016/j.jclinepi.2013.08.005> PMID: 24189091
52. Harding Center for Risk Literacy, Robert Koch Institute. *Faktenboxen zur mRNA-Schutzimpfung gegen COVID-19*. 2021.
53. Korn L, Böhm R, Meier NW, Betsch C. Vaccination as a social contract. *Proceedings of the National Academy of Sciences*. 2020; 117(26):14890–9. <https://doi.org/10.1073/pnas.1919666117> PMID: 32541033