



ELSEVIER

Journal of Optometry

www.journalofoptometry.org



ORIGINAL ARTICLE

Visual outcome of penetrating keratoplasty, deep anterior lamellar keratoplasty and Descemet membrane endothelial keratoplasty

Clàudia Garrido^a, Genís Cardona^{b,*}, Josep L. Güell^{c,d}, Jaume Pujol^{b,e}

^a Refractive and Optometry Department, Instituto de Microcirugía Ocular, Barcelona, Spain

^b Department of Optics and Optometry, Universitat Politècnica de Catalunya · BarcelonaTech, Terrassa, Spain

^c The Cornea, Cataract and Refractive Surgery Department, Instituto de Microcirugía Ocular, Barcelona, Spain

^d Department of Ophthalmology, Universitat Autònoma de Barcelona, Barcelona, Spain

^e Centre for Sensors, Instruments and Systems Development, Universitat Politècnica de Catalunya · BarcelonaTech, Terrassa, Spain

Received 21 March 2017; accepted 12 August 2017

Available online 13 November 2017

KEYWORDS
Deep anterior
lamellar
keratoplasty;
Descemet membrane
endothelial
keratoplasty;
Penetrating
keratoplasty;
Quality of vision;
Visual acuity

Abstract

Purpose: A single-center, cross-sectional study was designed to assess and compare objective and subjective quality of vision of patients intervened with penetrating keratoplasty (PK), deep anterior lamellar keratoplasty (DALK) and Descemet membrane endothelial keratoplasty (DMEK).

Methods: Forty-six patients previously intervened with PK (22 eyes), DALK (7 eyes) and DMEK (17 eyes) were recruited. Visual evaluation included spherical and cylindrical refraction, distance corrected visual acuity (DCVA), photopic contrast sensitivity (CS), optical quality, measured with the HD Analyzer (objective scattering index [OSI], MTF cut-off and Strehl ratio), and ocular and corneal aberrometry, measured with the KR-1W Wavefront Analyzer.

Results: Statistically significant between-group differences were found in age ($p=0.006$, DMEK patients were older) and time since surgery ($p<0.001$, longest time for PK patients). No statistically significant differences were found in DCVA between the techniques. Between-group differences were encountered in CS at 12 ($p=0.007$) and 18 ($p<0.001$) cycles per degree, with DMEK and DALK obtaining the best and worst outcomes, respectively. Differences in optical quality were found between the techniques (OSI, $p=0.004$; MTF cut-off, $p=0.048$; Strehl ratio $p=0.022$), with DMEK displaying the best outcomes. Highest and lowest values in ocular and corneal aberrations were for DALK and DMEK patients, respectively. Between-group differences were found in corneal astigmatism ($p<0.001$; -3.31 ± 2.00 D in PK; -2.68 ± 0.94 D in DALK; -1.09 ± 0.62 D in DMEK).

* Corresponding author at: Terrassa School of Optics and Optometry, Violinista Vellsolà, 37, E08222 Terrassa, Catalonia, Spain.
E-mail address: genis.cardona@upc.edu (G. Cardona).

Conclusion: Overall, DMEK proved superior over PK and DALK in terms of quality of vision, with PK offering slightly better outcomes than DALK in most visual function parameters under evaluation.

© 2017 Spanish General Council of Optometry. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Queratoplastia
lamelar anterior
profunda;
queratoplastia
endotelial de
membrana de
Descemet;
queratoplastia
penetrante;
calidad de visión;
agudeza visual.

Resultados visuales de la queratoplastia penetrante, la queratoplastia lamelar anterior profunda y la queratoplastia endotelial de membrana de Descemet

Resumen

Objetivo: Se diseñó un estudio en un único centro y transversal para evaluar y comparar la calidad de la visión subjetiva de los pacientes intervenidos de queratoplastia penetrante (QP), queratoplastia lamelar anterior profunda (QLAP) y queratoplastia endotelial de membrana de Descemet (DMEK).

Métodos: Se incluyó a cuarenta y seis pacientes previamente intervenidos de QP (22 ojos), QLAP (7 ojos) y DMEK (17 ojos). La evaluación visual incluyó la refracción esférica y cilíndrica, la agudeza visual de lejos corregida (AVDC), la sensibilidad al contraste fotópica (SC), la calidad óptica, medidas con el HD Analyzer (índice de dispersión objetiva [OSI], punto de corte MTF y ratio Strehl), así como la aberrometría ocular y corneal, medidas con el KR-1W Wavefront Analyzer.

Resultados: Se encontraron diferencias estadísticamente significativas entre los grupos en cuanto a edad ($p = 0,006$, los pacientes de DMEK eran mayores) y el tiempo transcurrido desde la cirugía ($p < 0,001$, tiempo superior para los pacientes de QP). No se encontraron diferencias estadísticamente significativas en cuanto a AVDC entre las técnicas. Se encontraron diferencias entre los grupos en cuanto a SC en 12 ($p = 0,007$) y 18 ($p < 0,001$) ciclos por grado, obteniendo DMEK y QLAP los mejores y peores resultados, respectivamente. Se encontraron diferencias en cuanto a calidad óptica entre las técnicas (OSI, $p = 0,004$; punto de corte MTF, $p = 0,048$; ratio Strehl $p = 0,022$), reflejando DMEK los mejores resultados. Los mejores y peores valores en cuanto a aberraciones oculares y corneales fueron registrados por los pacientes de QLAP y DMEK, respectivamente. Se hallaron diferencias entre grupos en cuanto a astigmatismo corneal ($p < 0,001$; $-3,31 \pm 2$ D en QP; $-2,68 \pm 0,94$ D en QLAP; $-1,09 \pm 0,62$ D en DMEK).

Conclusión: En general, DMEK demostró ser superior a QP y QLAP en términos de calidad de visión, y QP ofreció mejores resultados que QLAP en la mayoría de los parámetros de la función visual evaluados.

© 2017 Spanish General Council of Optometry. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Nowadays, many options are available to the corneal transplant surgeon for the treatment of corneal disorders. The keratoplasty surgical procedure has been modified to improve drawbacks associated with penetrating keratoplasty (PK), including delayed wound healing, major vulnerability to trauma, risk of immune rejection, unpredictable refractive outcomes, prolonged visual rehabilitation and high or irregular astigmatism.^{1,2} Although PK, in which the full corneal thickness is replaced, remains the gold standard,³ current keratoplasty procedures focus on replacing only the essential amount of tissue (also known as "selective keratoplasty"). Thus, in deep anterior lamellar keratoplasty (DALK),⁴ the whole corneal tissue is replaced with the exception of Descemet membrane and endothelium; in Descemet membrane endothelial keratoplasty (DMEK),^{5,6} the Descemet membrane and endothelium

are replaced by the corresponding layers from the donor cornea; and in Descemet stripping (automated) endothelial keratoplasty (DSEK or DSAEK)^{7,8} the Descemet membrane and endothelium are replaced by a thin layer of donor stroma, Descemet membrane and endothelium. Lamellar corneal grafts have proved superior in terms of fast visual rehabilitation and more predictable refractive outcomes, requiring only partial rather than full-thickness incisions.⁹ Besides, DMEK has been reported to provide similar endothelial cell count but higher rebubbling rate than other endothelial keratoplasty techniques, while also restoring physiologic pachymetry.^{10,11}

The present study aimed at describing and comparing visual outcomes of PK, DALK and DMEK. Previous researchers have assessed subjective and objective quality of vision of DALK with reference to PK. Thus, Güell and colleagues reported better best-corrected visual acuity (VA) and optical quality with PK than DALK, also noting a good correlation

between these parameters.¹² Arjomand and co-workers also observed better spectacle-corrected VA in PK than DALK, with VA in DALK increasing in those eyes with lower (<20 µm) thickness of the residual recipient stromal bed.¹³ These authors did not find any difference between PK and DALK in terms of contrast sensitivity (CS) and higher order aberrations (HOA). Pantanelli and colleagues reported better spectacle-corrected VA in PK than DALK, a difference that persisted even with full HOA correction through adaptive optics (HOA were worse in PK), although no differences in CS between PK and DALK were found.¹⁴ The same authors also evidenced the superiority of PK versus DSAEK in VA, but not in HOA, which they attributed to the longer time PK patients had for neural adaptation, and to the possible presence of corneal haze in DSAEK patients. Discrepancies amongst researchers persist, with reports describing similar best-corrected VA^{15,16} and HOA^{3,17} in PK and DALK, while others note a better performance of PK over DALK in both visual function parameters.¹⁸⁻²⁰ Fewer studies have explored the visual outcome of DMEK, probably because this challenging technique is not yet widely spread. However, all these reports suggest that DMEK may be a superior technique in terms of fast visual recovery, less refractive changes, and best-corrected visual acuity when compared with DSAEK, as well as presenting with reduced posterior HOA than PK and DSAEK.^{10,21}

Given the relatively small volume of published research addressing the objective and subjective quality of vision of patients following PK, DALK and DMEK, the purpose of the present single-center, cross-sectional study was to evaluate and compare the visual outcomes of these techniques in terms of visual acuity, contrast sensitivity, optical quality and higher-order aberrations. Although it may be argued that the decision of the corneal surgeon to select a particular technique is commonly governed by the characteristics of the condition (*i.e.*, affecting the whole cornea or being limited to certain layers), in some instances this choice may be influenced by other factors such as expertise with a technique or expected visual outcomes. The findings of the present research may provide information to assist these decisions.

Material and methods

Study sample

Forty-six consecutive patients (23 females) who had previously undergone PK (22 eyes), DALK (7 eyes) or DMEK (17 eyes) surgery were included in the study. Twenty-two eyes were right eyes (48%) and 24 left eyes (52%). All patients attended the Cornea, Cataract and Refractive Surgery Unit, Instituto de Microcirugía Ocular (IMO), Barcelona, Spain, between January and June 2015 for follow-up visits of their procedures. Age of patients ranged between 23 and 78 years, with a mean age (\pm standard deviation) of 53.2 (\pm 14.7) years. Exclusion criteria were concomitant ocular conditions known to influence visual outcome such as corneal opacities, uveitis, manifest cataracts, posterior capsular opacities, vitreous or retinal abnormalities and vitrectomized eyes. Patients who had undergone post-keratoplasty corneal refractive surgery were also excluded, as were those unable

to understand or comply with the instructions required for objective and subjective visual function assessment.

All patients received information regarding the procedures and associated possible complications and written informed consent was obtained. The study was conducted in accordance with the Declaration of Helsinki tenets of 1975 (as revised in Tokyo in 2004) and received the approval of an institutional ethical board (IMO).

Surgical procedures

All corneal surgeries had been conducted by the same surgeon (JLG) at the Cornea and Refractive Surgery Unit, Instituto de Microcirugía Ocular (IMO), Barcelona, Spain. Surgeries were executed according to the previously described procedure for PK, DALK and DMEK.²²⁻²⁴ At the time of measurements all sutures inserted during the surgical intervention had been removed.

Visual function evaluation

Following retinoscopy and subjective refraction, corrected distance high-contrast monocular visual acuity (DCVA) was measured in logMAR units with the Early Treatment Diabetic Retinopathy Study (ETDRS) charts (Optec 6500, Stereo Optical Co., Inc., Chicago, IL) at a viewing distance of 4 m and under photopic conditions (85 cd/m²). Contrast sensitivity (CS) was assessed with the CSV-1000 test (Vector Vision, Inc, Greenville, OH) placed at 2.5 m. This test presents a translucent chart divided into four sine-wave grating stimuli at spatial frequencies of 3, 6, 12 and 18 cycles per degree (cpd) and eight levels of contrast. A two-alternative forced choice paradigm was implemented for CS measures. The background illumination of the translucent chart is provided by the fluorescent luminance source of the instrument, that is, it is independent of room illumination. Best distance correction was employed during all CS measurements.

Optical quality was assessed with the HD Analyzer (Visiometrics S.L., Spain). The HD Analyzer employs a double-pass system to evaluate the retinal image degradation of a point-source object.²⁵ Double-pass images contain information about aberrations and scattering.²⁶ Explored parameters were the objective scattering index (OSI), calculated as the ratio between the light reaching an annular area ranging from 2 to 20 minute of arc from the central peak of maximum intensity and that of the central 1 minute of arc,²⁷ the modulation transfer function cut-off frequency (MTF cut-off), which is the spatial frequency corresponding to a 0.01 MTF value (the MTF value is calculated from the root square of the modulus of the Fourier transform of the double pass image), and the Strehl ratio, defined as the ratio under the measured MTF curve and the curve that would be obtained if the eye was free of aberrations.²⁸ All measurements were performed with full spherical and cylindrical correction and for a 4 mm pupil.

Finally, aberrometry data was obtained with the KR-1W Wavefront Analyzer (Topcon Medical Systems, Inc., Oakland, NJ). This instrument combines a Hartmann-Shack wavefront analyzer with a Placido disc-based anterior corneal topographer to provide ocular and anterior corneal aberrometry information, respectively (corneal aberrations are

calculated from elevation data and subsequently transformed to wavefront Zernike polynomials). A pupil diameter configuration of 4 mm was used to determine total ocular and anterior corneal HOA as the root mean square (RMS) values of all Zernike coefficients of order 3rd to 8th. The same instrument was employed as a topographer to obtain anterior corneal astigmatism, objective refraction, as well as to determine pupil diameter under mesopic conditions (5 cd/m^2). Internal aberrations were obtained by subtracting corneal aberrations from ocular aberrations.

All procedures were conducted by an experienced optometrist (CG), masked to the type of surgery being evaluated. Aberrometry and optical quality measurements were repeated three times and the average was used for statistical analysis.

Data analysis

Statistical analysis was performed with the IBM SPSS software for Windows (IBM Corporation, Armonk, NY). All data were examined for normality with the Kolmogorov-Smirnov test, which revealed compliance with a normal distribution. Accordingly, descriptive statistics are summarized as mean \pm SD. An analysis of variance test (ANOVA) was employed to investigate the statistical significance of the differences between PK, DALK and DMEK for each variable under evaluation and, when differences reached statistical significance, the Tukey *post hoc* test was employed for pair-wise analysis, thus taking into consideration family-wise error-rate. A *p*-value of 0.05 or less was considered to denote statistical significance throughout the study.

Results

Demographic data for PK, DALK and DMEK groups is summarized in Table 1, which also shows time (in months) since surgery. An ANOVA analysis revealed statistically significant differences both in age ($F=5.857$; $p=0.006$) and time since surgery ($F=17.666$; $p<0.001$). When submitted to a Tukey *post hoc* test, pair-wise differences in age were found between DMEK and PK and between DMEK and DALK, with DMEK patients (61.59 ± 11.34 years) being older than PK (49.60 ± 15.46 years) and DALK patients (44.00 ± 9.45 years). Differences were also found in time since surgery between PK and DALK and between PK and DMEK, with time since PK being much longer (84.82 ± 52.21 months) than time since DALK (21.28 ± 8.42 months) and time since DMEK (18.37 ± 12.46 months). Main reasons for intervention were Fuchs endothelial dystrophy (15 patients) and keratoconus (15 patients).

Refractive sphere and cylinder outcomes, as well as DCVA and CS for each group of patients, are shown in Table 2. Statistically significant between-group differences in both refractive sphere ($F=3.682$; $p=0.033$) and cylinder ($F=9.129$; $p<0.001$) were encountered which, when examined pair-wise, were found to originate in the differences between PK and DALK in spherical refraction and between PK and DMEK in cylinder. The largest and smallest refractive astigmatism corresponded to eyes intervened of PK ($-2.85 \text{ D} \pm 2.05 \text{ D}$) and DMEK ($-0.76 \text{ D} \pm 0.70 \text{ D}$), respectively. In terms of CS, statistically significant between-group differ-

ences were encountered at high spatial frequencies: 12 cpd ($F=5.570$; $p=0.007$) and 18 cpd ($F=11.541$; $p<0.001$). In particular, at 18 cpd, all pair-wise analyses revealed significant differences, with DMEK (0.89 ± 0.25) and DALK (0.30 ± 0.25) patients achieving the best and worst CS outcomes, respectively. No statistically significant differences were found in DCVA.

Table 3 displays a summary of the HDA measurements. An ANOVA test disclosed statistically significant differences in OSI ($F=6.231$; $p=0.004$), MTF cut-off ($F=3.272$; $p=0.048$) and Strehl Ratio ($F=4.182$; $p=0.022$) among the three procedures. A *post hoc* pair-wise analysis revealed statistically significant differences between PK and DMEK in all these parameters, as well as between DALK and DMEK in OSI. Overall, DMEK presented the best outcome in terms of optical quality and scattering, with PK performing slightly better than DALK.

The measurements of the KR-1W are shown in Table 4. Statistically significant between-group differences were found in all parameters under evaluation. The highest and lowest ocular and corneal HOA values corresponded to DALK and DMEK, respectively, with statistically significant differences between PK and DMEK and between DALK and DMEK, and a similar performance of PK and DALK. Calculated mean internal HOAs were $-0.57 \mu\text{m}$ in PK, $-0.92 \mu\text{m}$ in DALK and $-0.12 \mu\text{m}$ in DMEK. Corneal astigmatism was also different between the three procedures ($F=11.842$; $p<0.001$), with statistically significant pair-wise differences between PK and DMEK and between DALK and DMEK. Corneal astigmatism was highest in PK patients ($-3.31 \pm 2.00 \text{ D}$), followed by DALK patients ($-2.68 \pm 0.94 \text{ D}$), and DMEK patients ($-1.09 \pm 0.62 \text{ D}$). Finally, between-group differences in pupil diameter were found ($F=12.812$; $p<0.001$). Patients intervened with DALK had larger pupils than PK patients and DMEK patients, although statistically significant differences were only found between the pairs PK – DMEK and DMEK – DALK.

Discussion

Published literature is inconclusive regarding the superiority of a particular keratoplasty technique in regards to optical quality. Differences in study design, instrumentation and follow-up time, amongst other factors, may account for these discrepancies. In the present study, we aimed at examining and comparing objective and subjective visual quality in patients intervened with PK, DALK and DMEK. Although all procedures were conducted by the same experienced surgeon and in the same clinical setting, some study design limitations remained unresolved, mainly regarding between-group differences in age of patients and time since surgery. Indeed, whereas most DMEK patients continue to wear their own glasses after surgery, on account of the minor change in refraction associated with this technique, stability of visual acuity and refraction in PK may require a few months.²⁹ During this time, PK patients may have time for neural adaptation to their post-operative aberrations.¹⁴ In summary, different results may have been obtained with a sample better matched in time since surgery.

Overall, patients intervened with DMEK were found to outperform PK and DALK patients in most of the examined

Table 1 Demographic details, presented as mean \pm standard deviation (SD), of PK, DALK and DMEK patients. Results of the ANOVA test and *post hoc* Tukey pair-wise analysis are shown. Any *p*-value < 0.05 (in bold) denotes statistical significance.

	Procedure			ANOVA		Tukey pair-wise analysis	
	PK	DALK	DMEK	F	<i>p</i>	Pairs	<i>p</i>
Age (years)	49.60 \pm 15.46	44.00 \pm 9.45	61.59 \pm 11.34	5.857	0.006	PK-DMEK	0.021
						PK-DALK	0.601
						DMEK-DALK	0.014
Sex (M/F)	13/9	3/4	7/10	0.675	0.515		
Time since intervention (months)	84.82 \pm 82	21.28 \pm 8.42	18.37 \pm 12.46	17.666	<0.001	PK-DMEK	<0.001
						PK-DALK	0.001
						DMEK-DALK	0.984

Table 2 Refractive and visual outcomes, presented as mean \pm standard deviation (SD), of PK, DALK and DMEK patients. Results of the ANOVA test and *post hoc* Tukey pair-wise analysis are shown. Any *p*-value < 0.05 denotes statistical significance.

	Procedure			ANOVA		Tukey pair-wise analysis	
	PK	DALK	DMEK	F	<i>p</i>	Pairs	<i>p</i>
DCVA ^a (logMAR)	0.05 \pm 0.06	0.12 \pm 0.12	0.04 \pm 0.05	2.993	0.061		
Sphere (D)	1.06 \pm 1.60	-0.64 \pm 1.70	0.66 \pm 1.09	3.682	0.033	PK-DMEK	0.675
						PK-DALK	0.025
						DMEK-DALK	0.121
Cylinder (D)	-2.85 \pm 2.05	-2.28 \pm 0.82	-0.76 \pm 0.70	9.129	<0.001	PK-DMEK	<0.001
						PK-DALK	0.671
						DMEK-DALK	0.079
CS ^b 3 cpd ^c (log CS)	1.38 \pm 0.23	1.44 \pm 0.23	1.47 \pm 0.17	0.984	0.382		
CS 6 cpd (log CS)	1.54 \pm 0.25	1.50 \pm 0.23	1.66 \pm 0.24	1.686	0.197		
CS 12 cpd (log CS)	1.09 \pm 0.36	0.90 \pm 0.32	1.33 \pm 0.24	5.570	0.007	PK-DMEK	0.051
						PK-DALK	0.354
						DMEK-DALK	0.010
CS 18 cpd (log CS)	0.65 \pm 0.30	0.30 \pm 0.25	0.89 \pm 0.25	11.541	<0.001	PK-DMEK	0.023
						PK-DALK	0.019
						DMEK-DALK	<0.001

^a DCVA: distance corrected monocular visual acuity.^b CS: contrast sensitivity.^c cpd: cycles per degree.

visual function and refractive parameters, including refractive and corneal astigmatism, contrast sensitivity at high spatial frequencies, optical quality, scattering and corneal and ocular HOAs. It must be noted, however, that DMEK patients were older than PK and DALK patients, resulting in statistically significant differences in pupil diameter that may partially account for the superiority of this procedure in terms of CS. In contrast, albeit optical quality and aberrations are influenced by pupil diameter, all these measurements were conducted with an artificial pupil of 4 mm, that is, pupil diameter may not be considered as a confounding variable for these parameters.

No statistically significant differences in DCVA were found between the techniques. These results are in agreement with previous reports comparing PK and endothelial keratoplasty procedures,^{30,31} as well as PK and DALK,^{15,16} although

other authors describe a better performance of PK over DALK in DCVA.^{18–20} It is interesting to note that many of these authors report that, in general, patients intervened with any type of keratoplasty fail to reach their full visual potential. In this regards, 7 out of 22 PK, 3 out of 7 DALK and 3 out of 17 DMEK patients had DCVA worse than 0.1 logMAR (0.8 decimal) at the time of the follow-up appointment. Increased scattering and HOA may account for the inability of some of these patients to overcome a particular visual acuity threshold. However, the encountered between-group differences in HOA and optical quality were not accompanied by corresponding differences in DCVA. This finding may suggest that either different mechanisms may be governing the relationship between these parameters in PK, DALK and DMEK or that DCVA is not as sensitive as CS at high spatial frequencies, in which DMEK patients scored better results than PK and

Table 3 Outcomes from the HD Analyzer, presented as mean \pm standard deviation (SD) of PK, DALK and DMEK patients. Results of the ANOVA test and post hoc Tukey pair-wise analysis are shown. Any p-value < 0.05 denotes statistical significance.

	Procedure			ANOVA		Tukey pair-wise analysis	
	PK	DALK	DMEK	F	p	Pairs	p
OSI ^a	3.11 \pm 1.37	3.20 \pm 1.38	1.81 \pm 0.94	6.231	0.004	PK-DMEK PK-DALK DMEK-DALK	0.006 0.984 0.040
MTF cut-off ^b (cpd ^c)	17.20 \pm 7.37	19.27 \pm 10.67	24.02 \pm 8.44	3.272	0.048	PK-DMEK PK-DALK DMEK-DALK	0.038 0.833 0.418
Strehl ratio	0.112 \pm 0.041	0.103 \pm 0.045	0.144 \pm 0.052	4.182	0.022	PK-DMEK PK-DALK DMEK-DALK	0.027 0.989 0.109

^a OSI: objective scattering index.^b MTF cut-off: modulation transfer function cut-off frequency.^c cpd: cycles per degree.**Table 4** Results of the KR-1W Wavefront Analyzer, presented as mean \pm standard deviation (SD) of PK, DALK and DMEK patients. Results of the ANOVA test and post hoc Tukey pair-wise analysis are shown. Any p-value < 0.05 denotes statistical significance.

	Procedure			ANOVA		Tukey pair-wise analysis	
	PK	DALK	DMEK	F	p	Pairs	p
Corneal HOA ^a RMS ^b (μm)	0.534 \pm 0.18	0.732 \pm 0.46	0.244 \pm 0.07	10.760	<0.001	PK-DMEK PK-DALK DMEK-DALK	0.001 0.103 <0.001
Ocular HOA ^a RMS ^b (μm)	0.477 \pm 0.16	0.640 \pm 0.50	0.232 \pm 0.105	7.214	0.002	PK-DMEK PK-DALK DMEK-DALK	0.038 0.004 0.222
Corneal astigmatism (D)	3.31 \pm 2.00	2.68 \pm 0.94	1.09 \pm 0.62	11.842	<0.001	PK-DMEK PK-DALK DMEK-DALK	<0.001 0.600 0.057
Pupil diameter (mm)	5.68 \pm 0.77	6.39 \pm 0.57	4.90 \pm 0.72	12.812	<0.001	PK-DMEK PK-DALK DMEK-DALK	0.005 0.079 <0.001

^a HOA: high order aberrations.^b RMS: root mean square.

DALK patients, to compare visual quality between keratoplasty techniques. It may be worth noting that CS at middle and high spatial frequencies has been documented to be particularly useful for target detection and identification tasks, even in patients in whom visual acuity is not excellent.³²

Regarding the superiority of DMEK in terms of both ocular and corneal HOAs, it must be noted that the Placido disk/Hartmann-Shack configuration of the KR-1W allows for the measurement of ocular, that is, wavefront, aberrations, and of anterior corneal aberrations, which are determined from topographical height data. Internal corneal aberrations, including those originating at the posterior cornea and at the lens, may be estimated by subtracting anterior corneal from ocular HOAs. Thus, given mean internal HOAs of $-0.57 \mu\text{m}$ in PK, $-0.92 \mu\text{m}$ in DALK and $-0.12 \mu\text{m}$ in DMEK, the actual impact of posterior HOAs on visual quality in DMEK patients was probably not significant.^{33,34} These

results, which are in agreement with previous reports,²¹ are not unexpected, as the change in refractive index between the posterior cornea and the aqueous humor is smaller than the change occurring at the anterior surface between air and the cornea/tear film. Further research, including the evaluation of posterior height data, is needed to better understand this finding, however. Overall, DMEK, as the less invasive procedure of the three, was also found to present the best outcome in terms of optical quality and scattering.

Finally, it may be worth mentioning that between-group differences in age (DMEK patients were older) and particularly in time since surgery (more time in PK) may have resulted in an underestimation of the differences between the techniques. In effect, the loss with age of the ability of the posterior cornea to compensate anterior corneal HOA has been previously described,³⁵ as has the positive effect of neural adaptation on visual quality, particularly

in PK patients.¹⁴ In addition, given the recruitment difficulties encountered with DALK patients, with only 7 subjects in this group, the possibility of Type II error may not be ruled out, i.e., it may be speculated whether with a larger, more balanced sample, other statistically significant differences between techniques may have been uncovered.

In conclusion, as far as we know the present study is one of the first reports comparing quality of vision in PK, DALK and DMEK patients, and one of the few exploring ocular wavefront, as well as, anterior corneal height-data based aberrations.³⁶ Although some study limitations remain that warrant further investigation, our findings give support to the overall superiority of DMEK over PK and DALK. The lack of statistically significant differences in DCVA between the procedures highlights the advantages of exploring other visual function parameters, such as CS, optical quality, scattering and HOAs, to assess the subtle differences in visual quality of these patients. Finally, in view of the present findings, we would advise eye care providers to manage these patients with contact lenses, instead of glasses, to improve post-operative VA, contrast sensitivity and to reduce HOA.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Riddle HK Jr, Parker DA, Price FW Jr. Management of postkeratoplasty astigmatism. *Curr Opin Ophthalmol*. 1998;9:15–28.
- Pineros O, Cohen EJ, Rapuano CJ, Laibson PR. Long-term results after penetrating keratoplasty for Fuchs' endothelial dystrophy. *Arch Ophthalmol*. 1996;114:15–18.
- Koh S, Maeda N, Nakagawa T, et al. Characteristic higher-order aberrations of the anterior and posterior corneal surfaces in 3 corneal transplantation techniques. *Am J Ophthalmol*. 2012;153:284–290.
- Archila EA. Deep lamellar keratoplasty dissection of host tissue with intrastromal air injection. *Cornea*. 1984–1985;3:217–218.
- Melles GR, Lander F, Rietveld FJ. Transplantation of Descemet's membrane carrying viable endothelium through a small scleral incision. *Cornea*. 2002;21:415–418.
- Melles GR. Posterior lamellar keratoplasty: DLEK to DSEK to DMEK. *Cornea*. 2006;25:879–881.
- Melles GR, Wijdh RH, Nieuwendaal CP. A technique to excise the descemet membrane from a recipient cornea (descemetothenesis). *Cornea*. 2004;23:286–288.
- Price FW Jr, Price MO. Descemet's stripping with endothelial keratoplasty in 50 eyes: a refractive neutral corneal transplant. *J Refract Surg*. 2005;21:339–345.
- Price FW Jr, Feng MT, Price MO. Evolution of endothelial keratoplasty: where are we headed? *Cornea*. 2015;34(suppl 10):S41–S47.
- Guerra FP, Anshu A, Price MO, Giebel AW, Price FW. Descemet's membrane endothelial keratoplasty. Prospective study of 1-year visual outcomes, graft survival, and endothelial cell loss. *Ophthalmology*. 2011;118:2368–2373.
- Price MO, Giebel AW, Fairchild KM, Price FW. Descemet's membrane endothelial keratoplasty. Prospective multicenter study of visual and refractive outcomes and endothelial survival. *Ophthalmology*. 2009;116:2361–2368.
- Güell JL, Pujol J, Arjona M, Diaz-Doutón F, Luque SO, Artal P. Objective Optical quality after penetrating keratoplasty and deep anterior lamellar keratoplasty. *Invest Ophthalmol Vis Sci*. 2005;46: E-Abstract 2712.
- Ardjomand N, Hau S, McAlister JC, et al. Quality of vision and graft thickness in deep anterior lamellar and penetrating corneal allografts. *Am J Ophthalmol*. 2007;143:228–235.
- Pantanelli SM, Sabesan R, Ching SST, Yoon G, Hindman HB. Visual performance with wave aberration correction after penetrating, deep anterior lamellar, or endothelial keratoplasty. *Invest Ophthalmol Vis Sci*. 2012;53:4797–4804.
- Coombes AG, Kirwan JF, Rostron CK. Deep lamellar keratoplasty with lyophilised tissue in the management of keratoconus. *Br J Ophthalmol*. 2001;85:788–791.
- Shimazaki J, Shimmura S, Ishioka M, Tsubota K. Randomized clinical trials of deep anterior lamellar keratoplasty vs penetrating keratoplasty. *Am J Ophthalmol*. 2002;134:159–165.
- Javadi MA, Feizi S, Yazdani S, Mirbabaei F. Deep anterior lamellar keratoplasty versus penetrating keratoplasty for keratoconus: a clinical trial. *Cornea*. 2010;29:365–371.
- Watson SL, Ramsay A, Dart JK, Bunce C, Craig E. Comparison of deep lamellar keratoplasty and penetrating keratoplasty in patients with keratoconus. *Ophthalmology*. 2004;111:1676–1682.
- Panda A, Bageshwar LM, Ray M, Singh JP, Kumar A. Deep lamellar keratoplasty versus penetrating keratoplasty for corneal lesions. *Cornea*. 1999;18:172–175.
- Bahar I, Kaiserman I, Srinivasan S, Ya-Ping J, Slomovic AR, Rootman DS. Comparison of three different techniques of corneal transplantation for keratoconus. *Am J Ophthalmol*. 2008;146:905–912.
- Rudolph M, Laaser K, Bachmann BO, Cursiefen C, Epstein D, Kruse FE. Corneal higher-order aberrations after Descemet's membrane endothelial keratoplasty. *Ophthalmology*. 2012;119:528–535.
- Melles GR, Lander F, Rietveld FJ, Remeijer L, Beekhuis WH, Binder PS. A new surgical technique for deep stromal, anterior lamellar keratoplasty. *Br J Ophthalmol*. 1999;83:327–333.
- Anwar M, Teichmann KD. Big-bubble technique to bare Descemet's membrane in anterior lamellar keratoplasty. *J Cataract Refract Surg*. 2002;28:398–403.
- Güell JL, Morral M, Gris O, Elies D, Manero F. Bimanual technique for insertion and positioning of endothelium-Descemet membrane graft in Descemet membrane endothelial keratoplasty. *Cornea*. 2013;32:1521–1526.
- Güell JL, Pujol J, Arjona M, Diaz-Doutón F, Artal P. Optical quality analysis system; instrument for objective clinical evaluation of ocular optical quality. *J Cataract Refract Surg*. 2004;30:1598–1599.
- Díaz-Doutón F, Benito A, Pujol J, Arjona M, Güell JL, Artal P. Comparison of the retinal image quality with a Hartmann-Shack wavefront sensor and a double-pass instrument. *Invest Ophthalmol Vis Sci*. 2006;47:1710–1716.
- Artal P, Benito A, Pérez GM, et al. An objective scatter index based on double-pass retinal images of a point source to classify cataracts. *PLoS ONE*. 2011;6:e16823.
- Vilaseca M, Padilla A, Ondategui JC, Arjona M, Güell JL, Pujol J. Effect of laser in situ keratomileusis on vision analyzed using preoperative optical quality. *J Cataract Refract Surg*. 2010;36:1945–1953.
- Hayashi K, Hayashi H. Long-term changes in corneal surface configuration after penetrating keratoplasty. *Am J Ophthalmol*. 2006;141:241–247.
- Kosker M, Suri K, Duman F, Hammersmith KM, Nagra PK, Rapuano CJ. Long-term outcomes of penetrating keratoplasty and Descemet stripping endothelial keratoplasty for Fuchs endothelial dystrophy: fellow eye comparisons. *Cornea*. 2013;32:1083–1088.

31. Nanavaty MA, Wang X, Shortt AJ. Endothelial keratoplasty versus penetrating keratoplasty. *Cochrane Database Syst Rev*. 2014;CD008420.
32. Ginsburg AP, Evans DW, Sekule R, Harp SA. Contrast sensitivity predicts pilots' performance in aircraft simulators. *Am J Optom Physiol Opt*. 1982;59:105–109.
33. Hindman HB, Huxlin KR, Pantanelli SM, et al. Post-DSAEK optical changes: a comprehensive prospective analysis on the role of ocular wavefront aberrations, haze and corneal thickness. *Cornea*. 2013;32:1567–1577.
34. Yamaguchi T, Negishi K, Yamaguchi K, et al. Comparison of anterior and posterior corneal surface irregularity in Descemet stripping automated endothelial keratoplasty and penetrating keratoplasty. *Cornea*. 2010;29:1086–1090.
35. Dubbelman M, Sicam VA, van der Heijde RG. The contribution of the posterior surface to the coma aberration on the human cornea. *J Vis*. 2007;7:101–108.
36. Turnbull AMJ, Tsatsos M, Hossain PN, Anderson DF. Determinants of visual quality after endothelial keratoplasty. *Surv Ophthalmol*. 2016;61:257–271.