

MEETING ABSTRACT

Open Access

# Activation of kappa opioid receptors reduces seizure activity in a dose-dependent way

Luca Zangrandi, Christoph Schwarzer\*

From 18th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Croatian, Serbian and Slovenian Pharmacological Societies.

Graz, Austria. 20-21 September 2012

## Background

Neuropsychiatric disorders are one of the main challenges of human medicine with epilepsy as one of the most frequent. Temporal lobe epilepsy represents the most common type of epilepsies and is often accompanied by marked neuronal degeneration. One main factor that causes neural loss is the excitotoxicity of glutamate, which is copiously released during seizures and hypoxia accompanying seizures. It was also shown that the deletion of prodynorphin in mice and low expression in humans is associated with increased epilepsy vulnerability. Dynorphin targets opioid receptors and in particular the  $\kappa$  opioid receptor (KOP). The KOP receptors in the hippocampal formation are located in very strategically points for the control of glutamate release and, most importantly, they are not altered under epileptic conditions. Still, the functional background of these neuroprotective effects is not fully understood. The aim of this study was to investigate the influence of KOP agonists on EEG patterns of epileptic mice.

## Methods

Kainic acid (KA; 1 nmol in 50 nL saline) was injected unilaterally into the dorsal hippocampus, causing acute and delayed behavioral and EEG effects. Four-channel EEG traces were recorded from ipsi- and contralateral hippocampi and motor cortices applying depth and surface electrodes, respectively. The KOP-specific agonist U-50488H was dissolved in saline (adjusted to pH 7.4) and applied i.p.

## Results

Sharp waves and paroxysmal discharges in the ipsilateral hippocampus were recorded about 14 days after KA

injection. Paroxysmal discharges were accompanied by behavioral arrest and stereotypic behavior like head nodding. Application of KOP agonists blocked paroxysmal discharges up to 2 hours in a dose-dependent manner, which was comparable to the effect of 2.5 mg/kg diazepam. Moreover, animals treated with U-50488H were awake.

## Conclusions

Data collected so far demonstrate the anticonvulsant action of KOP agonists in the chronic phase of epilepsy, suggesting that KOP agonists may represent potential drug targets for novel anti-epileptics.

## Acknowledgements

This work was supported by the Austrian Science Fund (FWF: W1206-B05).

Published: 17 September 2012

doi:10.1186/2050-6511-13-S1-A91

**Cite this article as:** Zangrandi and Schwarzer: Activation of kappa opioid receptors reduces seizure activity in a dose-dependent way. *BMC Pharmacology and Toxicology* 2012 **13**(Suppl 1):A91.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

