

Thiazolidinedione-independent activation of peroxisome proliferator-activated receptor γ is a potential target for diabetic macrovascular complications

Takeshi Matsumura, Kayo Taketa, Seiya Shimoda, Eiichi Araki*

ABSTRACT

Macrovascular complications are responsible for the high morbidity and mortality in patients with diabetes. Peroxisome proliferator-activated receptor γ (PPAR γ) plays a central role in the process of adipocyte differentiation and insulin sensitization, and also possesses anti-atherogenic effects. Recently, some statins, angiotensin II type 1 receptor blockers and calcium channel blockers have been reported to activate PPAR γ . However, the impact of PPAR γ activation on diabetic macrovascular complications is not fully understood. It has been reported that the activation of PPAR γ by thiazolidinediones induces anti-atherogenic effects in vascular cells, including monocytes/macrophages, endothelial cells and smooth muscle cells, in atherosclerotic animal models and in clinical studies. We have reported that hydroxymethylglutaryl coenzyme A reductase inhibitors (statins), which are used for treatment of hypercholesterolemia, activate PPAR γ and mediate anti-atherogenic effects through PPAR γ activation in macrophages. Also, telmisartan, an angiotensin type I receptor blocker, has been reported to have anti-atherogenic effects through PPAR γ activation. Furthermore, we have reported that nifedipine, a dihydropyridine calcium channel blocker, can activate PPAR γ , thereby mediating anti-atherogenic effects in macrophages. Therefore, statin therapy and part of anti-hypertensive therapy might produce beneficial effects through PPAR γ activation in hypercholesterolemic and/or hypertensive patients with diabetes, and PPAR γ might be a therapeutic target for diabetic macrovascular complications. In the present review, we focus on the anti-atherogenic effects of PPAR γ and suggest potential therapeutic approaches to prevent diabetic macrovascular complications. (*J Diabetes Invest*, doi: 10.1111/j.2040-1124.2011.00182.x, 2012)

KEY WORDS: Diabetes, Macroangiopathy, Peroxisome proliferator-activated receptor γ

INTRODUCTION

Diabetic macrovascular complications, including cardiovascular disease (CVD), stroke and peripheral vascular disease, are mainly responsible for the high morbidity and mortality of diabetes patients^{1–3}. Several longitudinal epidemiological studies have shown that the risk of mortality from CVD in diabetic subjects is more than double compared with age-matched healthy subjects^{4,5}, suggesting that some features of diabetes, such as hyperglycemia, must promote an excessive tendency toward CVD.

Hyperglycemia is a diagnostic feature of diabetes, a target for antidiabetic therapy and a marker of glycemic control together with HbA_{1c}. Conserving good glycemic control has been associated with a notable reduction in the risk of developing retinopa-

thy, nephropathy and neuropathy in both type 1⁶ and type 2^{7,8} diabetic patients. Furthermore, when the therapy is started in the early stages of diabetes, intensive diabetes therapy has long-term beneficial effects on the risk of CVD in both type 1 and type 2 diabetes⁹.

There are several lines of evidence showing that the risk of cardiovascular mortality increases with the elevation in plasma glucose concentrations and HbA_{1c} levels^{10,11}. However, although the effect of plasma glucose levels appears to be preponderant in determining the microvascular risk, this does not seem to be the case for macrovascular complications.

The UK Prospective Diabetes Study (UKPDS) clearly shows that for the same degree of HbA_{1c}, particularly in its low range, the incidence of myocardial infarction is much greater than that of retinopathy¹². This apparent paradox can only be resolved by acknowledging the multifactorial nature of the cardiovascular risk in type 2 diabetic subjects. The UKPDS identified several potentially modifiable risk factors, including hypercholesterolemia and hypertension, as well as hyperglycemia, for CVD in

Department of Metabolic Medicine, Faculty of Life Sciences, Kumamoto University, Kumamoto, Japan

*Corresponding author. Eiichi Araki Tel.: +81-96-373-5169 Fax: +81-96-366-8397 E-mail address: earaki@gpo.kumamoto-u.ac.jp

Received 3 October 2011; revised 18 October 2011; accepted 19 October 2011

type 2 diabetes¹³. Furthermore, HbA_{1c} was found to be the third most important factor in determining the cardiovascular risk⁷. Indeed, several intervention studies have reported that lipid lowering with hydroxymethylglutaryl coenzyme A (HMG-CoA) reductase inhibitors (statins) reduces the incidence of cardiovascular events in patients with diabetes^{14–20}. In addition, the HOT study²¹ and UKPDS 39²² showed that lowering blood pressure decreased the risk of CVD in hypertensive patients with diabetes. Indeed, several studies have reported that treatment with angiotensin II converting enzyme inhibitor (ACEI)²³, angiotensin II type 1 receptor (AT1R) blocker (ARB)²⁴ or calcium channel blockers (CCB)^{21,25} reduced the incidence of cardiovascular events in hypertensive patients with diabetes. Therefore, it is very important for the prevention of diabetic macroangiopathy to enforce antihypercholesterolemic and antihypertensive therapy in diabetic patients.

Recently, it has been reported that some statins, ARB and CCB activate peroxisome proliferator-activated receptor γ (PPAR γ), which plays a central role in the process of adipocyte differentiation, peripheral glucose utilization and insulin sensitization, and has anti-atherogenic effects. In the present review, we examine the possibility of therapeutic approaches for the prevention of diabetic macrovascular complications through PPAR γ activation.

PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR γ

Peroxisome proliferator-activated receptor γ , a member of the nuclear hormone receptor family of ligand-dependent transcription factors²⁶, has been well characterized as a regulator of adipogenesis and is abundant in fat cells²⁷. Two major splice isoforms of PPAR γ , PPAR γ 1 and PPAR γ 2, have been identified in mice, whereas in humans and other species at least two other isoforms, PPAR γ 3 and PPAR γ 4, have been detected²⁸.

Several endogenous ligands for PPAR γ , including polyunsaturated fatty acids such as linoleic acid, arachidonic acid and eicosapentaenoic acids, have been identified. Prostaglandin-related compounds, such as 15 deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂), 9- and 13-hydroxyoctadecadienoic acid, 12- and 15-hydroxyeicosatetraenoic acid and nitro lipids, have also been identified as beneficial PPAR γ agonists^{29–32}. Thiazolidinediones (TZD), including pioglitazone and rosiglitazone, which are currently on the market, are the most notable synthetic compounds with PPAR γ activation properties that improve insulin resistance and lower blood glucose levels in type 2 diabetes.

Peroxisome proliferator-activated receptor γ forms a heterodimer with another nuclear receptor, retinoid X receptor α (RXR α), and binds to a specific DNA sequence, termed PPAR response element 5'-AAGGTCAAnAAGGTCA-3' (PPRE), in the regulatory regions of target genes³³. In addition, PPAR γ can regulate gene expression independently of PPRE, either by suppressing growth hormone protein-1, a transcription factor involved in pituitary-specific gene expression, or by interfering with the function of activator protein (AP)-1, signal transducer and activator

of transcription (STAT)-1 and nuclear factor κ B (NF- κ B)^{34–37}. Coactivators, including steroid receptor coactivator-1, p300/cAMP response element binding protein (CBP), general control non-repressed/p300/CBP-associated factor and PPAR γ coactivator, can bind PPAR γ /RXR α complexes in a ligand-dependent manner³⁸. In contrast, mitotic, stress and inflammatory signals cause PPAR γ degradation through phosphorylation on Ser84 of the mouse PPAR γ (Ser112 of the human molecule), which is in a consensus mitogen-activated protein kinase (MAPK) target motif PXSP³⁹, by extracellular signal-regulated kinases (ERK), c-Jun N-terminal kinases and p38 MAPK, which leads to ubiquitination and proteasomal degradation⁴⁰. A Ser84 to Ala84 PPAR γ mutant shows increased transcriptional activity, and a similar effect is caused by coexpression with the dual-specificity (Thr/Tyr) MAPK phosphatase, CL100³⁹.

Peroxisome proliferator-activated receptor γ plays a central role in the process of adipocyte differentiation, peripheral glucose utilization and insulin sensitization in adipocytes^{41–44}. In addition, PPAR γ is expressed in macrophages³⁶, endothelial cells (EC)⁴⁵, vascular smooth muscle cells (VSMC)⁴⁶ and in atherosclerotic lesions^{47,48}. PPAR γ agonists suppressed the progression of atherosclerosis in low-density lipoprotein (LDL)-receptor-deficient^{49,50} or apolipoprotein E-deficient (ApoE^{-/-}) mice^{51–53} (Table 1). In clinical studies, the Prospective Pioglitazone Clinical Trial in Macrovascular Events (PROactive) study suggested that pioglitazone might decrease the occurrence of all-cause mortality, non-fatal myocardial infarction (MI) and non-fatal stroke in patients with type 2 diabetes mellitus and macrovascular diseases⁵⁴. Mazzone *et al.*⁵⁵ showed that pioglitazone slowed the progression of carotid intima-media thickness compared with glimepiride in patients with type 2 diabetes. The Pioglitazone Effect on Regression of Intravascular Sonographic Coronary Obstruction Prospective Evaluation (PERISCOPE) study showed that treatment with pioglitazone resulted in a significantly lower progression rate of coronary atherosclerosis compared with glimepiride⁵⁶. Ogasawara *et al.*⁵⁷ reported that pioglitazone might stabilize coronary plaque by reducing the necrotic-core

Table 1 | Peroxisome proliferator-activated receptor γ ligands suppress the progression of atherosclerosis in animal models

Disease model	Knockout mouse	PPAR γ agonist	Decrease in lesion size, %	References
Type 2 diabetes atherosclerosis	LDL-receptor (high-fat diet)	Rosiglitazone	60–80	49
Atherosclerosis	ApoE (normal diet)	Rosiglitazone	18	53
Type 2 diabetes atherosclerosis	LDL-receptor (high-fat diet)	Troglitazone	30	51
Atherosclerosis	ApoE (high-fat diet)	Troglitazone	45	52

ApoE, apolipoprotein E; LDL, low-density lipoprotein; PPAR γ , peroxisome proliferator-activated receptor γ .

component in Japanese patients with type 2 diabetes. These findings show the usefulness of PPAR γ activator in the treatment of diabetic macroangiopathy. In contrast, meta-analysis reported that treatment with rosiglitazone, but not with pioglitazone, was associated with increased incidence of MI^{58,59}. Because meta-analysis has many limitations and the increased risk of MI is still controversial, these reports should be carefully considered. However, several differences between the effects of pioglitazone and rosiglitazone do exist. For example, pioglitazone has more beneficial effects on the lipid profile than rosiglitazone⁶⁰. Furthermore, rosiglitazone, but not pioglitazone, induced cardiac hypertrophy through a PPAR γ -independent pathway⁶¹. To clarify the mechanism(s) of the differences between the effects of pioglitazone and rosiglitazone, further studies are required.

THIAZOLIDINEDIONES

Thiazolidinediones, which are pharmacological ligands of PPAR γ , are used to treat patients with type 2 diabetes. Several anti-atherosclerotic effects of TZD have been reported in various cell types existing in atherosclerotic lesions (Figure 1). In EC, TZD inhibited the expression of vascular cell adhesion molecule-1 (VCAM-1) and intercellular adhesion molecule-1^{62,63}, decreased production of chemokines, such as interleukin-8 (IL-8) and monocyte chemoattractant protein-1 (MCP-1)⁶⁴, decreased endothelin-1, which is involved in the regulation of vascular tone and smooth muscle cell proliferation⁴⁵, and increased nitric oxide (NO) release⁶⁵. In addition, a recent report showed that adipose tissue-derived adiponectin plays an obligatory role in TZD-induced improvement of endothelial function in diabetic obese mice⁶⁶.

Vascular smooth muscle cells have pivotal roles in the progression of atherosclerosis, and recent studies have shown that they are key targets of TZD. In particular, proliferation of VSMC is one of the critical events in atherosclerosis and vascular intervention-induced restenosis. TZD inhibit these changes in VSMC and neointimal thickening after vascular injury⁶⁷. TZD inhibited VSMC proliferation^{67,68} by decreasing phosphorylation of retinoblastoma protein and increasing levels of cyclin-dependent inhibitor p27⁶⁷, and thus suppressed neointima thickening after vascular injury⁶⁹. Furthermore, TZD induced apoptosis of VSMC through p53 and Gadd45^{70,71}. In addition, TZD inhibited the expression of matrix metalloproteinase-9 (MMP-9)⁷¹ and VSMC migration^{46,68}, resulting in plaque stabilization. TZD also inhibited IL-1 β -induced inducible NO synthase (iNOS) expression and NO production in VSMC⁷². Interestingly, TZD inhibited AT1R in VSMC⁷³⁻⁷⁵, suggesting that this downregulation would be beneficial for diabetic patients with hypertension.

The involvement of PPAR γ in regulating lipid metabolism in macrophages was initially suggested by the discovery of the class B scavenger receptor CD36 as a PPAR γ target gene in macrophages⁷⁶. CD36, which is involved in the uptake of oxidized LDL (Ox-LDL), is thought to be a critical molecule in




Figure 1 | Anti-atherogenic effects of thiazolidinediones (TZD) in vascular cells. ABCA1, adenosine triphosphate-binding cassette transporter A1; ABCG1, adenosine triphosphate-binding cassette transporter G1; AT1R, angiotensin II type 1 receptor; COX-2, cyclooxygenase-2; EC, endothelial cell; eNOS, endothelial nitric oxide synthase; ET-1, endothelin-1; ICAM-1, intercellular adhesion molecule-1; IL-1 β , interleukin-1 β ; IL-6, interleukin-6; iNOS, inducible NO synthase; LXR, liver X receptor; M ϕ , macrophage; MCP-1, monocyte chemoattractant protein-1; MMP-9, matrix metalloproteinase-9; Mo, Monocyte; NO, nitric oxide; PAI-1, plasminogen activator inhibitor-1; ROS, reactive oxygen species; SMC, smooth muscle cell; TIMP-1, tissue inhibitor of metalloproteinase-1; TNF- α , tumor necrosis factor- α ; VCAM-1, vascular cell adhesion molecule-1.

macrophage-derived foam cell formation⁷⁷. In addition, CD36 is highly expressed in lipid-laden macrophages in human atherosclerotic plaques⁷⁸, and 9-hydroxyoctadecadienoic acid and 13-hydroxyoctadecadienoic acid, which are components of the Ox-LDL particle, can activate PPAR γ , suggesting that the activation of PPAR γ is involved in the formation of foam cells and the progression of atherosclerosis³¹. However, treatment with TZD has been found to increase the apolipoprotein-AI-dependent efflux of cholesterol from both mouse and human macrophages^{79,80}. Strikingly, the ability of TZD to promote cholesterol efflux is completely dependent on the expression of adenosine triphosphate-binding cassette transporter A1 (ABCA1), which is characterized as a vital molecule in the control of high-density lipoprotein and apolipoprotein A1-mediated cholesterol efflux

from macrophages^{81,82} through PPAR γ activation⁸⁰. Thus, although TZD induce the expression of CD36 in macrophages, they simultaneously induce the expression of ABCA1 in these cells, thereby enhancing the removal of cholesterol from the vessel wall through PPAR γ activation.

Classically activated macrophages, known as M1 macrophages, express pro-inflammatory cytokines, whereas alternatively activated macrophages, known as M2 macrophages, express anti-inflammatory cytokines and play a protective role in the progression of atherosclerosis. Recent studies showed that PPAR γ is one of the key regulators of M1/M2 macrophage polarization⁸², and that TZD prime monocytes into M2 macrophages and PPAR γ expression is enhanced during differentiation of M2 macrophages^{83,84}. TZD suppress the expression of M1 macrophage-associated pro-inflammatory cytokines in monocytes, including tumor necrosis factor- α (TNF- α), IL-1 β and IL-6⁸⁵, and inhibit macrophage activation *in vitro*³⁶. Meanwhile, Marx *et al.*⁴⁷ reported that TZD inhibited MMP-9 gelatinolytic activity, an enzyme responsible for plaque rupture. A similar phenomenon was also observed in monocytes⁸⁶. In association with these findings, TZD inhibited MCP-1-directed monocyte migration, which was mediated through a decrease in MMP-9 expression and/or increase in a tissue inhibitor of metalloproteinase-1 expression⁸⁷.

One of the characteristic events in atherosclerotic lesions is cell proliferation in arterial walls, including VSMC and macrophages. Indeed, previous *in vivo* studies have reported that macrophages and macrophage-derived foam cells proliferated in atherosclerotic lesions^{88–90}. We^{91,92} and other groups^{93,94} have shown that Ox-LDL enhanced macrophage proliferation and survival *in vitro*. Therefore, it is possible that macrophage proliferation promotes the progression of atherosclerosis. In relation to these findings, we have reported that TZD suppress Ox-LDL-induced macrophage proliferation⁹⁵. Thus, the suppression of macrophage proliferation might be one of the anti-atherogenic effects of TZD.

Peroxisome proliferator-activated receptor γ is thought to alter macrophage function at the molecular level through various mechanisms³⁸. PPAR γ induces a ligand and a sumoylation-dependent conformational shift that allows direct binding to NF- κ B and recruitment of the co-repressor complex, nuclear receptor corepressor, thereby suppressing transcription of NF- κ B target genes, such as iNOS^{96,97}. TZD also inhibit the activity of other transcription factors, such as AP-1 and STAT-1, both of which are involved in the induction of pro-inflammatory cytokines during differentiation of M1 macrophages^{36,85}. As aforementioned, TZD might many anti-atherogenic effects in vascular cells, and might be beneficial for diabetic macroangiopathy beyond the improvement of insulin sensitivity.

CHOLESTEROL-LOWERING THERAPY IN PATIENTS WITH DIABETES

Dyslipidemia, an established risk factor for CVD, is strikingly common in patients with type 2 diabetes, affecting almost 50%

of this population⁹⁸. In addition to hyperglycemia and hypertension, dyslipidemia is a modifiable CVD risk factor that remains largely uncontrolled in patients with diabetes⁹². HMG-CoA reductase inhibitors (statins) are known to reduce the incidence of cardiovascular events and death by functional changes of atherosclerotic lesions^{99–102}. Furthermore, because of these benefits, which were shown by many clinical studies, statin therapy is recommended as the initial pharmacological treatment for lowering LDL-cholesterol levels in patients with type 2 diabetes^{103,104}. These statin benefits are mainly achieved by its strong lipid-lowering effect. However, recent data suggest that many of the beneficial effects exerted by statins on vascular cells are independent of the cholesterol-lowering effect^{105–109}. Cholesterol is synthesized through the isoprenoid biosynthetic pathway¹¹⁰. In this pathway, isopentenyl-PP is the basic isoprene unit used for synthesis of all subsequent isoprenoids¹¹⁰. Among the isoprenoids, farnesyl pyrophosphate (FPP) and geranylgeranyl pyrophosphate (GGPP) serve as important lipid attachments for several proteins, including the small GTP-binding protein Ras and Ras-like proteins, such as Rho, Rac and Cdc42, whose proper membrane localization and function are dependent on isoprenylation¹¹⁰. The pleiotropic effects of statins, which include improving the function of EC, VSMC and macrophages^{110,111}, are thought to be mediated by blocking the synthesis of FPP and GGPP, resulting in the inhibition of small GTP-binding proteins¹¹⁰.

Several studies reported that atorvastatin and pravastatin increased PPAR γ activity in monocytes^{112,113}. We also showed that five lipophilic statins (simvastatin, cerivastatin, fluvastatin, atorvastatin and pitavastatin) can activate PPAR γ in macrophages¹¹⁴. We have shown that statins have a unique mechanism for PPAR γ activation (Figure 2). Statin-induced PPAR γ activation was suppressed by the addition of mevalonate, FPP or GGPP¹¹⁴. Moreover, farnesyl transferase inhibitor and geranylgeranyl transferase inhibitor mimicked the effects of statins. Furthermore, statins inhibited the membrane translocation of Ras, RhoA, Rac and Cdc42, and overexpression of dominant-negative mutants of RhoA and Cdc42, but not of Ras or Rac, increased PPAR γ activity¹¹⁴. Therefore, suppression of the mevalonate pathway and inactivation of RhoA and Cdc42 are involved in statin-mediated PPAR γ activation. In contrast, statins induced ERK1/2 and p38 MAPK activation, and ERK1/2- and p38 MAPK-specific inhibitors abrogated statin-induced PPAR γ activation, suggesting that statin-mediated PPAR γ activation was caused by the activation of ERK1/2 and p38 MAPK¹¹⁴. However, inhibition of RhoA and Cdc42 activated p38 MAPK, but not ERK1/2¹¹⁴. Therefore, statin-induced PPAR γ activation might be mediated by a RhoA- and Cdc42-dependent p38 MAPK pathway and by a RhoA- and Cdc42-independent ERK1/2 pathway¹¹⁴. A subsequent mechanism of PPAR γ activation by statins is a cyclooxygenase-2 (COX-2)-dependent increase in 15d-PG₂, which is a natural ligand of PPAR γ ¹¹⁴. Indeed, we showed that statins increased COX-2 expression and intracellular 15d-PG₂ levels, and statin-mediated




Figure 2 | Summary of statin-mediated peroxisome proliferator-activated receptor γ (PPAR γ) activation in macrophages. When treated with statins, farnesyl pyrophosphate (FPP) and geranylgeranyl pyrophosphate (GGPP) are downregulated by inhibition of the mevalonate pathway, thereby inhibiting geranylgeranylation and farnesylation, and the subsequent translocation of RhoA and Cdc42. The suppression of RhoA and Cdc42 translocation abrogates the RhoA and Cdc42 signaling pathways, thereby inducing p38 mitogen-activated protein kinase (MAPK)-dependent cyclooxygenase-2 (COX-2) production. In addition, small G protein-independent extracellular signal-regulated kinase 1/2 (ERK1/2) activation mediated by the suppression of geranylgeranylation and farnesylation is also involved in COX-2 production. Overexpression of COX-2 produces intracellular 15d-PGJ₂, which activates PPAR γ . The activation of PPAR γ mediates the downregulation of *TNF- α* and *MCP-1* mRNA expression by inactivating AP-1 and NF- κ B, and upregulating ABCA1. AA, arachidonic acid; ABCA1, adenosine triphosphate-binding cassette transporter A1; AP-1, activator protein-1; HMG-CoA, 3-hydroxy-3-methylglutaryl coenzyme A; 15d-PGJ₂, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂; MCP-1, monocyte chemoattractant protein-1; NF- κ B, nuclear factor- κ B; TNF- α , tumor necrosis factor- α .

PPAR γ activation was suppressed by selective COX-2 inhibitors or by downregulation of COX-2 using siRNA¹¹⁴. In relation to these findings, it has been reported that statins can induce COX-2 expression in VSMC¹¹⁵ and macrophages¹¹⁶.

Several studies have shown that statins suppressed the expression of TNF- α ¹¹⁷ and MCP-1¹¹⁸ in macrophages. We showed that lipopolysaccharide (LPS)-induced expression of TNF- α and MCP-1 was suppressed by pitavastatin in macrophages, and these effects were abrogated by PPAR γ siRNA¹¹⁴. Furthermore, pitavastatin induced ABCA1 expression, and this effect was also abrogated by PPAR γ siRNA¹¹⁴. Argmann *et al.*¹¹⁹ also reported that statins induced ABCA1 expression in macrophages, and

this effect was suppressed by the PPAR γ antagonist, GW9662. These results suggest that PPAR γ activation is one of the key mechanisms in the pleiotropic effects of statins on atherosclerosis, and statin therapy might improve the atherogenic profile by dual effects, such as improving hypercholesterolemia and activating PPAR γ , in patients with diabetes.

ANTI-HYPERTENSIVE THERAPY IN PATIENTS WITH DIABETES

Several large randomized trials have reported that ACEI²³, ARB²⁴ and dihydropyridine CCBs^{21,25} prevented cardiovascular events in hypertensive patients with type 2 diabetes. Inhibition of RAS by either ACEI or ARB reduced atherosclerotic lesion formation in experimental studies using various models of atherosclerosis^{120–123}. Among the ARB, telmisartan has been identified as a ligand for PPAR γ ^{124,125}. Indeed, we¹²⁶ and other groups^{127–130} reported that telmisartan suppressed the progression of atherosclerosis in apoE^{-/-} mice (Table 2). Similar results were shown in Watanabe heritable hyperlipidemic rabbits¹³¹ (Table 2). Furthermore, it has been reported that telmisartan prevented ischemic brain damage through PPAR γ activation in normal¹³² and diabetic mice¹³³. Several studies elucidated that telmisartan has direct anti-atherogenic effects on vascular cells (Figure 3). In EC, telmisartan suppresses VCAM-1 expression and reactive oxygen species generation¹³⁴, prevents methylglyoxal-mediated cell death¹³⁵, increases the permeability of EC¹³⁶ and induces proliferation of endothelial progenitor cells¹³⁷. In VSMC, telmisartan suppresses cell proliferation¹³⁸ and AT1R expression¹³⁹. We showed that in macrophages, telmisartan inhibits TNF- α and MCP-1 expression through inactivation of NF- κ B, induces ABCA1/ABCG1 expression through activation of liver X receptor and inhibits Ox-LDL-induced cell

Table 2 | Telmisartan suppresses the progression of atherosclerosis in animal models

Disease model	Animal models	Decrease in lesion size, %	Dose, mg/kg/day	References
Atherosclerosis	ApoE ^{-/-} mice (normal diet)	Male 30	3	126
	ApoE ^{-/-} mice (normal diet)	Female 32–66	0.3–3	127
	ApoE ^{-/-} mice (normal diet)	Male 24–57		128
	ApoE ^{-/-} mice (normal diet)	Male 38	40	128
	ApoE ^{-/-} mice (high-fat diet)	Male 43	10	129
	ApoE ^{-/-} mice (high-fat diet)	Male 50	10	130
	ApoE ^{-/-} AT1R ^{-/-} mice (high-fat diet)	Male 67	10	130
	WHHL rabbit (normal diet)	Male 84	5	131

ApoE, apolipoprotein E; AT1R, angiotensin II type 1 receptor.




Figure 3 | Anti-atherogenic effects of telmisartan in vascular cells. ABCA1, adenosine triphosphate-binding cassette transporter A1; ABCG1, adenosine triphosphate-binding cassette transporter G1; AT1R, angiotensin II type 1 receptor; EC, endothelial cell; Mφ, macrophage; MCP-1, monocyte chemoattractant protein-1; NF-κB, nuclear factor-κB; ROS, reactive oxygen species; SMC, smooth muscle cell; TNF-α, tumor necrosis factor-α; VCAM-1, vascular cell adhesion molecule-1.

proliferation through activation of PPARγ¹²⁶. Nakaya *et al.*¹⁴⁰ also reported that telmisartan induced ABCA1/ABCG1 expression through PPARγ in THP-1 macrophages. These reports suggest that telmisartan has the capacity to prevent atherosclerosis by not only lowering blood pressure, but also activating PPARγ.

Dihydropyridine CCB are relevant treatments for patients with hypertension. Beyond causing vasodilatation by inhibiting calcium channels, dihydropyridine CCB provide clinical benefits in patients with coronary artery disease that might be independent of blood pressure reductions^{141,142}. Nifedipine, one of the dihydropyridine CCB, reduced the atherosclerotic plaque area in cholesterol-fed rabbits^{143–145} (Table 3), and suppressed the progression of atherosclerosis in hypertensive patients^{146–150}. Recently, we showed that nifedipine could activate PPARγ in macrophages, which was not observed by amlodipine or Ca²⁺ chelator, suggesting that this effect was not caused by blocking calcium influx¹⁵¹. Unlike telmisartan, nifedipine does not act as a direct PPARγ ligand, but suppresses PPARγ phosphorylation

Table 3 | Nifedipine suppresses the progression of atherosclerosis in animal models

Disease model	Animal models	Decrease lesion size, %	Dose	References
Atherosclerosis	New Zealand rabbits (high-fat diet)	Male 58	16	143
	WHHL rabbits (high-fat diet)	Male 63	40	144
	Dutch-belted rabbits (high-fat diet)	Male 59	80	145
	ApoE ^{-/-} mice (normal diet)	Male 52	10	151

ApoE, apolipoprotein E.




Figure 4 | Summary of nifedipine-mediated peroxisome proliferator-activated receptor γ (PPARγ) activation in macrophages. When macrophages are treated with nifedipine, extracellular-signal regulated kinase 1/2 (ERK1/2) activity is downregulated. Inactivation of ERK1/2 suppresses PPARγ phosphorylation, leading to its activation. PPARγ activation blocks nuclear factor κB (NF-κB) activity, downregulates monocyte chemoattractant protein-1 (MCP-1) expression and upregulates adenosine triphosphate-binding cassette transporter A1 (ABCA1) expression. Nifedipine can therefore induce anti-atherogenic action.

by inhibiting ERK1/2 activity, thereby activating PPARγ (Figure 4)¹⁵¹. The exact mechanism of nifedipine-mediated inactivation of ERK1/2 is still unknown. However, Mulvaney *et al.*¹⁵² reported that nifedipine suppressed buserelin (a gonadotropin-releasing hormone agonist)-induced ERK1/2 activation through inactivation of Raf in αT3-1 cells, an immortalized mouse pituitary cell line. In this report, the authors speculated that the inactivation of Raf was caused by blocking L-type calcium channels and, subsequently, inhibiting calcium influx. However, we found that amlodipine did not activate PPARγ¹⁵². Thus, there might be mechanism(s) other than calcium influx inhibition for nifedipine-mediated inactivation of the Raf/ERK pathway. Hirata *et al.*¹⁵³ reported that nifedipine suppressed ERK1/2 activation by inhibiting proline-rich tyrosine kinase-2 (Pyk2) in VSMC. Therefore, inactivation of Pyk2 might also be involved in nifedipine-mediated inactivation of ERK1/2 in macrophages. Further studies are required to clarify the exact mechanism(s) of the nifedipine-mediated inactivation of ERK1/2.

Several studies reported multiple anti-atherogenic effects of nifedipine in vascular cells. Especially in macrophages and human peripheral blood mononuclear cells, nifedipine suppressed the LPS-induced increase of iNOS¹⁵⁴ and inflammatory cytokines, such as TNF-α, IL-1β and interferon-γ¹⁵⁵. Furthermore, nifedipine inhibited the LPS-induced NF-κB activation¹⁵⁶. In addition to these previous reports, we recently showed that

nifedipine suppressed the LPS-induced MCP-1 expression and NF- κ B activation through PPAR γ activation in macrophages¹⁵¹. These findings are supported by a previous report showing that nifedipine suppressed angiotensin II-induced MCP-1 expression through NF- κ B inhibition in rat VSMC¹⁵⁷. In contrast, Suzuki *et al.*¹⁵⁸ reported that nifedipine induced ABCA1 expression and apolipoprotein-mediated release of cellular cholesterol in macrophages. We also showed that nifedipine induced ABCA1 expression through PPAR γ activation¹⁵¹. Furthermore, our *in vivo* study showed that nifedipine suppressed the acceleration of atherosclerosis (Table 3), decreased MCP-1 expression, and increased ABCA1 expression and PPAR γ activation in apoE^{-/-} mice¹⁵¹. Our recently proposed novel mechanism of nifedipine-mediated PPAR γ activation is supported by other reports using the PPAR γ antagonist, GW9662^{159–161}.

Taking these findings into consideration, nifedipine, as well as telmisartan, might have anti-atherogenic effects through PPAR γ activation beyond the lowering of blood pressure, and treatment with such anti-hypertensive drugs might be beneficial for the prevention of atherosclerosis in hypertensive patients with diabetes.

CONCLUSIONS

The progression of diabetic macrovascular complications is caused by endothelial injury, activation of macrophages and abnormality of VSMC function. The mechanisms of anti-atherogenic effects by PPAR γ are mainly dependent on the improvement of endothelial biology, and inactivation of macrophages and VSMC. Although several studies have shown beneficial effects of TZD on experimental animal models of atherosclerosis, differences in cardiovascular risks among TZD were reported in clinical studies. Further studies are required to clarify the differences among the effects of TZD on diabetic macrovascular complications. Recently, a report showed that not the short-term use, but the use for more than 2 years of pioglitazone was weakly associated with an increased incidence of bladder cancer in a cohort study¹⁶². In contrast, statins, telmisartan and nifedipine, which possess the ability to activate PPAR γ , do not appear to mediate bodyweight gain, edema or incidence of cancers in clinical use. Thus, PPAR γ activation mediated by drugs other than TZD, such as statins, nifedipine and telmisartan, might be useful for the prevention of atherosclerosis without severe side-effects. In any case, many *in vitro*, *in vivo* and clinical studies have proposed the validity of PPAR γ activation in the progression of atherosclerosis. Therefore, a therapeutic approach using PPAR γ activation for diabetic macrovascular complications might produce beneficial effects in patients with diabetes.

ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (No. 21591144 to TM, No. 23591337 to KT and No. 20390259 to EA). None of the authors have any conflicts of interest associated with this manuscript.

REFERENCES

- Abbott RD, Donahue RP, Kannel WB, *et al.* The impact of diabetes on survival following myocardial infarction in men vs women. The Framingham Study. *JAMA* 1988; 260: 3456–3460.
- Gu K, Cowie CC, Harris MI. Mortality in adults with and without diabetes in a national cohort of the US population, 1971–1993. *Diabetes Care* 1998; 21: 1138–1145.
- Haffner SM, Lehto S, Ronnema T, *et al.* Mortality from coronary heart disease in subjects with type 2 diabetes and non-diabetic subjects with and without prior myocardial infarction. *N Engl J Med* 1998; 339: 229–234.
- Kannel WB, McGee DL. Diabetes and cardiovascular disease: the Framingham Study. *JAMA* 1979; 241: 2035–2038.
- Pan WH, Cedres LB, Liu K, *et al.* Relationship of clinical diabetes and asymptomatic hyperglycemia to risk of coronary heart disease mortality in men and women. *Am J Epidemiol* 1986; 123: 504–516.
- Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993; 329: 977–986.
- UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998; 352: 837–853.
- Ohkubo Y, Kishikawa H, Araki E, *et al.* Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995; 28: 103–117.
- Stettler C, Allemann S, Juni P, *et al.* Glycemic control and macrovascular disease in types 1 and 2 diabetes mellitus: meta-analysis of randomized trials. *Am Heart J* 2006; 152: 27–38.
- Laakso M. Glycemic control and the risk for coronary heart disease in patients with non-insulin-dependent diabetes mellitus: the Finnish Studies. *Ann Intern Med* 1996; 124: 127–130.
- Wei M, Gaskill SP, Haffner SM, *et al.* Effects of diabetes and level of glycemia on all-cause and cardiovascular mortality. *Diabetes Care* 1998; 21: 1167–1173.
- Stratton IM, Adler AI, Neil HA, *et al.* Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000; 321: 405–412.
- Turner RC, Millns H, Neil HA, *et al.* Risk factors for coronary artery disease in non-insulin dependent diabetes mellitus: United Kingdom Prospective Diabetes Study (UKPDS 23). *BMJ* 1998; 316: 823–828.
- Pyörälä K, Pedersen TR, Kjekshus J, *et al.* Cholesterol lowering with simvastatin improves prognosis of diabetic

- patients with coronary heart disease. A subgroup analysis of the Scandinavian Simvastatin Survival Study (4S). *Diabetes Care* 1997; 20: 614–620.
15. Collins R, Armitage J, Parish S, *et al.* MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003; 361: 2005–2016.
 16. Sacks FM, Pfeffer MA, Moye LA, *et al.* The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* 1996; 335: 1001–1009.
 17. Sacks FM, Tonkin AM, Shepherd J, *et al.* Effect of pravastatin on coronary disease events in subgroups defined by coronary risk factors: the Prospective Pravastatin Pooling Project. *Circulation* 2000; 102: 1893–1990.
 18. Sever PS, Dahlöf B, Poulter NR, *et al.* Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial – Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet* 2003; 361: 1149–1158.
 19. Colhoun HM, Betteridge DJ, Durrington PN, *et al.* Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 2004; 364: 685–696.
 20. Arampatzis CA, Goedhart D, Serruys PW, *et al.* Fluvastatin reduces the impact of diabetes on long-term outcome after coronary intervention – a Lescol Intervention Prevention Study (LIPS) substudy. *Am Heart J* 2005; 149: 329–335.
 21. Hansson L, Zanchetti A, Carruthers SG, *et al.* Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group. *Lancet* 1998; 351: 1755–1762.
 22. UK Prospective Diabetes Study Group. Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. *BMJ* 1998; 317: 713–720.
 23. Niskanen L, Hedner T, Hansson L, *et al.* Reduced cardiovascular morbidity and mortality in hypertensive diabetic patients on first-line therapy with an ACE inhibitor compared with a diuretic/beta-blocker-based treatment regimen: a subanalysis of the Captopril Prevention Project. *Diabetes Care* 2001; 24: 2091–2096.
 24. Dahlöf B, Devereux RB, Kjeldsen SE, *et al.* Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002; 359: 995–1003.
 25. Tuomilehto J, Rastenyte D, Birkenhäger WH, *et al.* Effects of calcium-channel blockade in older patients with diabetes and systolic hypertension. Systolic Hypertension in Europe Trial Investigators. *N Engl J Med* 1999; 340: 677–684.
 26. Spiegelman BM. Peroxisome proliferator-activated receptor γ : a key regulator of adipogenesis and systemic insulin sensitivity. *Eur J Med Res* 1997; 2: 457–464.
 27. Kliewer SA, Lenhard JM, Willson TM, *et al.* A prostaglandin J₂ metabolite binds peroxisome proliferator-activated receptor γ and promotes adipocyte differentiation. *Cell* 1995; 83: 813–819.
 28. Desvergne B, Wahli W. Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 1999; 20: 649–688.
 29. Forman BM, Tontonoz P, Chen J, *et al.* 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ is a ligand for the adipocyte determination factor PPAR- γ . *Cell* 1995; 83: 803–812.
 30. Huang JT, Welch JS, Ricote M, *et al.* Interleukin-4-dependent production of PPAR- γ ligands in macrophages by 12/15-lipoxygenase. *Nature* 1999; 400: 378–382.
 31. Nagy L, Tontonoz P, Alvarez JG, *et al.* Oxidized LDL regulates macrophage gene expression through ligand activation of PPAR- γ . *Cell* 1998; 93: 229–240.
 32. Schopfer FJ, Lin Y, Baker PR, *et al.* Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor γ ligand. *Proc Natl Acad Sci USA* 2005; 102: 2340–2345.
 33. Willson TM, Lambert MH, Kliewer SA. Peroxisome proliferator-activated receptor γ and metabolic disease. *Annu Rev Biochem* 2001; 70: 341–367.
 34. Feige JN, Gelman L, Michalik L, *et al.* From molecular action to physiological outputs: peroxisome proliferator-activated receptors are nuclear receptors at the crossroads of key cellular functions. *Progr Lipid Res* 2006; 45: 120–159.
 35. Tolon RM, Castillo AI, Aranda A. Activation of the prolactin protein gene by peroxisome proliferator-activated receptor- α appears to be DNA binding-independent. *J Biol Chem* 1998; 273: 26652–26661.
 36. Ricote M, Li AC, Willson TM, *et al.* The peroxisome proliferator-activated receptor- γ is a negative regulator of macrophage activation. *Nature* 1998; 391: 79–82.
 37. Feinstein DL, Spangolo A, Akar C, *et al.* Receptor-independent actions of PPAR thiazolidinedione agonists: is a mitochondrial function the key? *Biochem Pharm* 2005; 70: 177–188.
 38. Ricote M, Glass CK. PPARs and molecular mechanisms of transrepression. *Biochim Biophys Acta* 2007; 1771: 926–935.
 39. Diradourian C, Girard J, Pegorier JP. Phosphorylation of PPARs: from molecular characterization to physiological relevance. *Biochimie* 2005; 87: 33–82.
 40. Genini D, Catapano CV. Block of nuclear receptor ubiquitination: a mechanism of ligand-dependent control of peroxisome proliferator-activated receptor δ activity. *J Biol Chem* 2007; 282: 11776–11785.
 41. Kliewer SA, Forman BM, Blumberg B, *et al.* Differential expression and activation of a family of murine peroxisome

- proliferator-activated receptors. *Proc Natl Acad Sci USA* 1994; 91: 7355–7359.
42. Chawla A, Schwarz EJ, Dimaculangan DD, *et al.* Peroxisome proliferator activated receptor (PPAR) γ : adipose-predominant expression and induction early in adipocyte differentiation. *Endocrinology* 1994; 135: 798–800.
 43. Braissant O, Fougère F, Scotto C, *et al.* Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR- α , - β , and - γ in the adult rat. *Endocrinology* 1996; 137: 354–366.
 44. Jain S, Pulikuri S, Zhu Y, *et al.* Differential expression of the peroxisome proliferator-activated receptor γ (PPAR γ) and its coactivators steroid receptor coactivator-1 and PPAR-binding protein PBP in the brown fat, urinary bladder, colon, and breast of the mouse. *Am J Pathol* 1998; 153: 349–354.
 45. Delerive P, Martin-Nizard F, Chinetti G, *et al.* Peroxisome proliferator-activated receptor activators inhibit thrombin-induced endothelin-1 production in human vascular endothelial cells by inhibiting the activator protein-1 signaling pathway. *Circ Res* 1999; 85: 394–402.
 46. Marx N, Schonbeck U, Lazar MA, *et al.* Peroxisome proliferator-activated receptor γ activators inhibit gene expression and migration in human vascular smooth muscle cells. *Circ Res* 1998; 83: 1097–1103.
 47. Marx N, Sukhova G, Murphy C, *et al.* Macrophage in human atheroma contain PPAR γ : differentiation-dependent peroxisomal proliferator-activated receptor γ (PPAR γ) expression and reduction of MMP-9 activity through PPAR γ activation in mononuclear phagocytes *in vitro*. *Am J Pathol* 1998; 153: 17–23.
 48. Ricote M, Huang J, Fajas L, *et al.* Expression of the peroxisome proliferator-activated receptor γ (PPAR γ) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci USA* 1998; 95: 7614–7619.
 49. Li AC, Brown KK, Silvestre MJ, *et al.* Peroxisome proliferator-activated receptor γ ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *J Clin Invest* 2000; 106: 523–531.
 50. Li AC, Binder CJ, Gutierrez A, *et al.* Differential inhibition of macrophage foam-cell formation and atherosclerosis in mice by PPAR α , β/δ , and γ . *J Clin Invest* 2004; 114: 1564–1576.
 51. Collins AR, Meehan WP, Kintscher U, *et al.* Troglitazone inhibits formation of early atherosclerotic lesions in diabetic and nondiabetic low density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 2001; 21: 365–371.
 52. Chen Z, Ishibashi S, Perrey S, *et al.* Troglitazone inhibits atherosclerosis in apolipoprotein E-knockout mice: pleiotropic effects on CD36 expression and HDL. *Arterioscler Thromb Vasc Biol* 2001; 21: 372–377.
 53. Claudel T, Leibowitz MD, Fievet C, *et al.* Reduction of atherosclerosis in apolipoprotein E knockout mice by activation of the retinoid X receptor. *Proc Natl Acad Sci USA* 2001; 98: 2610–2615.
 54. Dormandy JA, Charbonnel B, Eckland DJ, *et al.* Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAZone Clinical Trial In macroVascular Events): a randomised controlled trial. *Lancet* 2005; 366: 1279–1289.
 55. Mazzone T, Meyer PM, Feinsein SB, *et al.* Effect of pioglitazone compared with glimepiride on carotid intima-media thickness in type 2 diabetes: a randomized trial. *JAMA* 2006; 296: 2572–2581.
 56. Nissen SE, Nicholls SJ, Wolski K, *et al.* Comparison of pioglitazone vs glimepiride on progression of coronary atherosclerosis in patients with type 2 diabetes: the PERISCOPE randomized controlled trial. *JAMA* 2008; 299: 1561–1573.
 57. Ogasawara D, Shite J, Shinke T, *et al.* Pioglitazone reduces the necrotic-core component in coronary plaque in association with enhanced plasma adiponectin in patients with type 2 diabetes mellitus. *Circ J* 2009; 73: 343–351.
 58. Nissen SE, Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 2007; 356: 2457–2471.
 59. Singh S, Loke YK, Furberg CD. Long-term risk of cardiovascular events with rosiglitazone: a meta-analysis. *JAMA* 2007; 298: 1189–1195.
 60. Goldberg RB, Kendall DM, Deeg MA, *et al.* A comparison of lipid and glycemic effects of pioglitazone and rosiglitazone in patients with type 2 diabetes and dyslipidemia. *Diabetes Care* 2005; 28: 1547–1554.
 61. Duan SZ, Ivashchenko CY, Russell MW, *et al.* Cardiomyocyte-specific knockout and agonist of peroxisome proliferator-activated receptor- γ both induce cardiac hypertrophy in mice. *Circ Res* 2005; 97: 372–379.
 62. Jackson SM, Parhami F, Xi XP, *et al.* Peroxisome proliferator-activated receptor activators target human endothelial cells to inhibit leukocyte-endothelial cell interaction. *Arterioscler Thromb Vasc Biol* 1999; 19: 2094–2104.
 63. Pasceri V, Wu HD, Willerson JT, *et al.* Modulation of vascular inflammation *in vitro* and *in vivo* by peroxisome proliferator-activated receptor- γ activators. *Circulation* 2000; 101: 235–238.
 64. Lee H, Shi W, Tontonoz P, *et al.* Role of peroxisome proliferator-activated receptor α in oxidized phospholipid-induced synthesis of monocyte chemotactic protein-1 and interleukin-8 by endothelial cells. *Circ Res* 2000; 87: 516–521.
 65. Calnek DS, Mazzella L, Roser S, *et al.* Peroxisome proliferator-activated receptor γ ligands increase release of nitric oxide from endothelial cells. *Arterioscler Thromb Vasc Biol* 2003; 23: 52–57.
 66. Wong WT, Tian XY, Xu A, *et al.* Adiponectin is required for PPAR γ -mediated improvement of endothelial function in diabetic mice. *Cell Metab* 2011; 14: 104–115.

67. Wakino S, Kintscher U, Kim S, *et al.* Peroxisome proliferator-activated receptor γ ligands inhibit retinoblastoma phosphorylation and G1 \rightarrow S transition in vascular smooth muscle cells. *J Biol Chem* 2000; 275: 22435–22441.
68. Law RE, Goetze S, Xi XP, *et al.* Expression and function of PPAR γ in rat and human vascular smooth muscle cells. *Circulation* 2000; 101: 1311–1318.
69. Law RE, Meehan WP, Xi XP, *et al.* Troglitazone inhibits vascular smooth muscle cell growth and intimal hyperplasia. *J Clin Invest* 1996; 98: 1897–1905.
70. Okura T, Nakamura M, Takata Y, *et al.* Troglitazone induces apoptosis via the p53 and Gadd45 pathway in vascular smooth muscle cells. *Eur J Pharmacol* 2000; 407: 227–235.
71. Aizawa Y, Kawabe J, Hasebe N, *et al.* Pioglitazone enhances cytokine-induced apoptosis in vascular smooth muscle cells and reduces intimal hyperplasia. *Circulation* 2001; 104: 455–460.
72. Takeda K, Ichiki T, Tokunou T, *et al.* Peroxisome proliferator-activated receptor gamma activators downregulate angiotensin II type 1 receptor in vascular smooth muscle cells. *Circulation* 2000; 102: 1834–1839.
73. Ikeda U, Shimpo M, Murakami Y, *et al.* Peroxisome proliferator-activated receptor- γ ligands inhibit nitric oxide synthesis in vascular smooth muscle cells. *Hypertension* 2000; 35: 1232–1236.
74. Sugawara A, Takeuchi K, Uruno A, *et al.* Transcriptional suppression of type 1 angiotensin II receptor gene expression by peroxisome proliferator-activated receptor- γ in vascular smooth muscle cells. *Endocrinology* 2001; 142: 3125–3134.
75. Tham DM, Martin-McNulty B, Wang YX, *et al.* Angiotensin II is associated with activation of NF- κ B-mediated genes and downregulation of PPARs. *Physiol Genomics* 2002; 11: 21–30.
76. Tontonoz P, Nagy L, Alvarez J, *et al.* PPAR γ promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* 1998; 93: 241–252.
77. Feng J, Han J, Pearce AFA, *et al.* Induction of CD36 expression by oxidized LDL and IL-4 by a common signaling pathway dependent on protein kinase C and PPAR γ . *J Lipid Res* 2000; 41: 688–696.
78. Nakata A, Nakagawa Y, Nishida M, *et al.* CD36, a novel receptor for oxidized low-density lipoproteins, is highly expressed on lipid-laden macrophages in human atherosclerotic aorta. *Arterioscler Thromb Vasc Biol* 1999; 19: 1333–1339.
79. Chawla A, Boisvert WA, Lee CH, *et al.* A PPAR γ –LXR–ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell* 2001; 7: 161–171.
80. Chinetti G, Lestavel S, Bocher V, *et al.* PPAR- α and PPAR- γ activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nat Med* 2001; 7: 53–58.
81. Lawn RM, Wade DP, Garvin MR, *et al.* The Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated lipid removal pathway. *J Clin Invest* 1999; 104: R25–R31.
82. Oram JF, Vaughan AM. ABCA1-mediated transport of cellular cholesterol and phospholipids to HDL apolipoproteins. *Curr Opin Lipidol* 2000; 11: 253–260.
83. Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, *et al.* Macrophage-specific PPAR γ controls alternative activation and improves insulin resistance. *Nature* 2007; 447: 1116–1120.
84. Bouhlel MA, Derudas B, Rigamonti E, *et al.* PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab* 2007; 6: 137–143.
85. Jiang C, Ting AT, Seed B. PPAR- γ agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998; 391: 82–86.
86. Shu H, Wong B, Zhou G, *et al.* Activation of PPAR α or γ reduces secretion of matrix metalloproteinase 9 but not interleukin 8 from human monocytic THP-1 cells. *Biochem Biophys Res Commun* 2000; 267: 345–349.
87. Kintscher U, Goetze S, Wakino S, *et al.* Peroxisome proliferator-activated receptor and retinoid X receptor ligands inhibit monocyte chemotactic protein-1-directed migration of monocytes. *Eur J Pharmacol* 2000; 401: 259–270.
88. Gordon D, Reidy MA, Benditt EP, *et al.* Cell proliferation in human coronary arteries. *Proc Natl Acad Sci USA* 1990; 87: 4600–4604.
89. Rosenfeld ME, Ross R. Macrophage and smooth muscle cell proliferation in atherosclerotic lesions of WHHL and comparably hypercholesterolemic fat-fed rabbits. *Arteriosclerosis* 1990; 10: 680–687.
90. Spagnoli LG, Orlandi A, Santeusano G. Foam cells of the rabbit atherosclerotic plaque arrested in metaphase by colchicine show a macrophage phenotype. *Atherosclerosis* 1991; 88: 87–92.
91. Sakai M, Miyazaki A, Hakamata H, *et al.* Lysophosphatidylcholine plays an essential role in the mitogenic effect of oxidized low density lipoprotein on murine macrophages. *J Biol Chem* 1994; 269: 31430–31435.
92. Matsumura T, Sakai M, Kobori S, *et al.* Two intracellular signaling pathways for activation of protein kinase C are involved in oxidized low-density lipoprotein-induced macrophage growth. *Arterioscler Thromb Vasc Biol* 1997; 17: 3013–3020.
93. Martens JS, Reiner NE, Herrera-Velitz P, *et al.* Phosphatidylinositol 3-kinase is involved in the induction of macrophage growth by oxidized low density lipoprotein. *J Biol Chem* 1998; 273: 4915–4920.
94. Hamilton JA, Myers D, Jessup W, *et al.* Oxidized LDL can induce macrophage survival, DNA synthesis, and enhanced proliferative response to CSF-1 and GM-CSF. *Arterioscler Thromb Vasc Biol* 1999; 19: 98–105.
95. Yano M, Matsumura T, Senokuchi T, *et al.* Troglitazone inhibits oxidized low-density lipoprotein-induced macrophage proliferation: impact of the suppression of

- nuclear translocation of ERK1/2. *Atherosclerosis* 2007; 191: 22–32.
96. Pascual G, Fong AL, Ogawa S, *et al.* A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR- γ . *Nature* 2005; 437: 759–763.
 97. Ogawa S, Lozach J, Benner C, *et al.* Molecular determinants of crosstalk between nuclear receptors and toll-like receptors. *Cell* 2005; 122: 707–721.
 98. Saydah SH, Fradkin J, Cowie CC. Poor control of risk factors for vascular disease among adults with previously diagnosed diabetes. *JAMA* 2004; 291: 335–342.
 99. Brown BG, Zhao XQ, Sacco DE, *et al.* Lipid lowering and plaque regression: new insights into prevention of plaque disruption and clinical events in coronary disease. *Circulation* 1993; 87: 1781–1791.
 100. Levine GN, Keaney JFJ, Vita JA. Cholesterol reduction in cardiovascular disease: clinical benefits and possible mechanisms. *N Engl J Med* 1995; 333: 512–521.
 101. Libby P, Aikawa M. New insights in plaque stabilization by lipid lowering. *Drugs* 1998; 56(suppl 1): 9–13.
 102. Vaughan CJ, Gotto AMJ, Basson CT. The evolving role of statins in the management of atherosclerosis. *J Am Coll Cardiol* 2000; 35: 1–10.
 103. Goff DC Jr, Gerstein HC, Ginsberg HN, *et al.* Prevention of cardiovascular disease in persons with type 2 diabetes mellitus: current knowledge and rationale for the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Am J Cardiol* 2007; 99: 4i–20i.
 104. American Diabetes Association. Standards of medical care in diabetes-2010. *Diabetes Care* 2010; 33(Suppl 1): S11–S61.
 105. Rosenson RS, Tangney CC. Antiatherothrombotic properties of statins: implications for cardiovascular event reduction. *JAMA* 1998; 279: 1643–1650.
 106. Davignon J, Laaksonen R. Low-density lipoprotein-independent effects of statins. *Curr Opin Lipidol* 1999; 10: 543–559.
 107. Corsini A, Bellosta S, Baetta R, *et al.* New insights into the pharmacodynamic and pharmacokinetic properties of statins. *Pharmacol Ther* 1999; 84: 413–428.
 108. Palinski W. New evidence for beneficial effects of statins unrelated to lipid lowering. *Arterioscler Thromb Vasc Biol* 2001; 21: 3–5.
 109. Kwak BR, Mach F. Statins inhibit leukocyte recruitment: new evidence for their anti-inflammatory properties. *Arterioscler Thromb Vasc Biol* 2001; 21: 1256–1258.
 110. Takemoto M, Liao JK. Pleiotropic effects of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Arterioscler Thromb Vasc Biol* 2001; 21: 1712–1719.
 111. Koh KK. Effects of statins on vascular wall: vasomotor function, inflammation, and plaque stability. *Cardiovasc Res* 2000; 47: 648–657.
 112. Grip O, Janciauskiene S, Lindgren S. Atorvastatin activates PPAR- γ and attenuates the inflammatory response in human monocytes. *Inflamm Res* 2002; 51: 58–62.
 113. Zelvyte I, Dominaitiene R, Crisby M, *et al.* Modulation of inflammatory mediators and PPAR γ and NF κ B expression by pravastatin in response to lipoproteins in human monocytes *in vitro*. *Pharmacol Res* 2002; 45: 147–154.
 114. Yano M, Matsumura T, Senokuchi T, *et al.* Statins activate peroxisome proliferator-activated receptor γ through extracellular signal-regulated kinase 1/2 and p38 mitogen-activated protein kinase-dependent cyclooxygenase-2 expression in macrophages. *Circ Res* 2007; 100: 1442–1451.
 115. Degraeve F, Bolla M, Blaie S, *et al.* Modulation of COX-2 expression by statins in human aortic smooth muscle cells. Involvement of geranylgeranylated proteins. *J Biol Chem* 2001; 276: 46849–46855.
 116. Chen JC, Huang KC, Wingerd B, *et al.* HMG-CoA reductase inhibitors induce COX-2 gene expression in murine macrophages: role of MAPK cascades and promoter elements for CREB and C/EBP β . *Exp Cell Res* 2004; 301: 305–319.
 117. Pahan K, Sheikh FG, Nambodiri AM, *et al.* Lovastatin and phenylacetate inhibit the induction of nitric oxide synthase and cytokines in rat primary astrocytes, microglia, and macrophages. *J Clin Invest* 1997; 100: 2671–2679.
 118. Veillard NR, Braunersreuther V, Arnaud C, *et al.* Simvastatin modulates chemokine and chemokine receptor expression by geranylgeranyl isoprenoid pathway in human endothelial cells and macrophages. *Atherosclerosis* 2006; 188: 51–58.
 119. Argmann CA, Edwards JY, Sawyez CG, *et al.* Regulation of macrophage cholesterol efflux through hydroxymethylglutaryl-CoA reductase inhibition: a role for RhoA in ABCA1-mediated cholesterol efflux. *J Biol Chem* 2005; 280: 22212–22221.
 120. Strawn WB, Chappell MC, Dean RH, *et al.* Inhibition of early atherogenesis by losartan in monkeys with diet-induced hypercholesterolemia. *Circulation* 2000; 101: 1586–1593.
 121. Hayek T, Attias J, Coleman R, *et al.* The angiotensin-converting enzyme inhibitor, fosinopril, and the angiotensin II receptor antagonist, losartan, inhibit LDL oxidation and attenuate atherosclerosis independent of lowering blood pressure in apolipoprotein E-deficient mice. *Cardiovasc Res* 1999; 44: 579–587.
 122. Takai S, Kim S, Sakonjo H, *et al.* Mechanisms of angiotensin II type 1 receptor blocker for anti-atherosclerotic effect in monkeys fed a high-cholesterol diet. *J Hypertens* 2003; 21: 361–369.
 123. Dol F, Martin G, Staels B, *et al.* Angiotensin AT1 receptor antagonist irbesartan decreases lesion size, chemokine expression, and macrophage accumulation in apolipoprotein E-deficient mice. *J Cardiovasc Pharmacol* 2001; 38: 395–405.
 124. Benson SC, Pershadsingh HA, Ho CI, *et al.* Identification of telmisartan as a unique angiotensin II receptor antagonist with selective PPAR γ -modulating activity. *Hypertension* 2004; 43: 993–1002.
 125. Schupp M, Janke J, Clasen R, *et al.* Angiotensin type 1 receptor blockers induce peroxisome proliferator-activated receptor-gamma activity. *Circulation* 2004; 109: 2054–2057.

126. Matsumura T, Kinoshita H, Ishii N, *et al.* Telmisartan exerts anti-atherosclerotic effects by activating PPAR γ in macrophages. *Arterioscler Thromb Vasc Biol* 2011; 31: 1268–1275.
127. Takaya T, Kawashima S, Shinohara M, *et al.* Angiotensin II type 1 receptor blocker telmisartan suppresses superoxide production and reduces atherosclerotic lesion formation in apolipoprotein E-deficient mice. *Atherosclerosis* 2006; 186: 402–410.
128. Blessing E, Preusch M, Kranzhofer R, *et al.* Anti-atherosclerotic properties of telmisartan in advanced atherosclerotic lesions in apolipoprotein E deficient mice. *Atherosclerosis* 2008; 199: 295–303.
129. Nagy N, Melchior-Becker A, Fischer JW. Long-term treatment with the AT1-receptor antagonist telmisartan inhibits biglycan accumulation in murine atherosclerosis. *Basic Res Cardiol* 2010; 105: 29–38.
130. Fukuda D, Enomoto S, Hirata Y, *et al.* The angiotensin receptor blocker, telmisartan, reduces and stabilizes atherosclerosis in ApoE and AT1aR double deficient mice. *Biomed Pharmacother* 2010; 64: 712–717.
131. Ikejima H, Imanishi T, Tsujioka H, *et al.* Effects of telmisartan, a unique angiotensin receptor blocker with selective peroxisome proliferator-activated receptor- γ -modulating activity, on nitric oxide bioavailability and atherosclerotic change. *J Hypertens* 2008; 26: 964–972.
132. Haraguchi T, Takasarki K, Naito T, *et al.* Cerebroprotective action of telmisartan by inhibition of macrophages/microglia expressing HMGB1 via a peroxisome proliferator-activated receptor γ -dependent mechanism. *Neurosci Lett* 2009; 464: 151–155.
133. Iwanami J, Mogi M, Tsukuda K, *et al.* Low dose of telmisartan prevents ischemic brain damage with peroxisome proliferator-activated receptor-gamma activation in diabetic mice. *J Hypertens* 2010; 28: 1730–1737.
134. Cianchetti S, Del Fiorentino A, Colognato R, *et al.* Anti-inflammatory and anti-oxidant properties of telmisartan in cultured human umbilical vein endothelial cells. *Atherosclerosis* 2008; 198: 22–28.
135. Baden T, Yamawaki H, Saito K, *et al.* Telmisartan inhibits methylglyoxal-mediated cell death in human vascular endothelium. *Biochem Biophys Res Commun* 2008; 373: 253–257.
136. Bian C, Wu Y, Chen P. Telmisartan increases the permeability of endothelial cells through zonula occludens-1. *Biol Pharm Bull* 2009; 32: 416–420.
137. Honda A, Matsuura K, Fukushima N, *et al.* Telmisartan induces proliferation of human endothelial progenitor cells via PPAR γ -dependent PI3K/Akt pathway. *Atherosclerosis* 2009; 205: 376–384.
138. Yamamoto K, Ohishi M, Ho C, *et al.* Telmisartan-induced inhibition of vascular cell proliferation beyond angiotensin receptor blockade and peroxisome proliferator-activated receptor- γ activation. *Hypertension* 2009; 54: 1353–1359.
139. Imayama I, Ichiki T, Inanaga K, *et al.* Telmisartan downregulates angiotensin II type 1 receptor through activation of peroxisome proliferator-activated receptor γ . *Cardiovasc Res* 2006; 72: 184–190.
140. Nakaya K, Ayaori M, Hisada T, *et al.* Telmisartan enhances cholesterol efflux from THP-1 macrophages by activating PPAR γ . *J Atheroscler Thromb* 2007; 14: 133–141.
141. Pitt B, Byington RP, Furberg CD, *et al.* Effect of amlodipine on the progression of atherosclerosis and the occurrence of clinical events. PREVENT Investigators. *Circulation* 2000; 102: 1503–1510.
142. Zanchetti A, Bond MG, Hennig M, *et al.* Calcium antagonist lacidipine slows down progression of asymptomatic carotid atherosclerosis: principal results of the European Lacidipine Study on Atherosclerosis (ELSA), a randomized, double-blind, long-term trial. *Circulation* 2002; 106: 2422–2427.
143. Henry PD, Bentley KI. Suppression of atherogenesis in cholesterol-fed rabbit treated with nifedipine. *J Clin Invest* 1981; 68: 1366–1369.
144. Atkinson JB, Swift LL. Nifedipine reduces atherogenesis in cholesterol-fed heterozygous WHHL rabbits. *Atherosclerosis* 1990; 84: 195–201.
145. Willis AL, Nagel B, Churchill V, *et al.* Antiatherosclerotic effects of nifedipine and nifedipine in cholesterol-fed rabbits. *Atherosclerosis* 1985; 5: 250–255.
146. Lichtlen PR, Hugenholtz PG, Rafflenbeul W, *et al.* Retardation of angiographic progression of coronary artery disease by nifedipine. Results of the International Nifedipine Trial on Antiatherosclerotic Therapy (INTACT). INTACT Group Investigators. *Lancet* 1990; 335: 1109–1113.
147. Brown MJ, Palmer CR, Castaigne A, *et al.* Morbidity and mortality in patients randomised to double-blind treatment with a long-acting calciumchannel blocker or diuretic in the International Nifedipine GITS study: intervention as a Goal in Hypertension Treatment (INSIGHT). *Lancet* 2000; 356: 366–372.
148. Motro M, Shemesh J. Calcium channel blocker nifedipine slows down progression of coronary calcification in hypertensive patients compared with diuretics. *Hypertension* 2001; 37: 1410–1413.
149. Simon A, Levenson J. Effects of calcium channel blockers on atherosclerosis: new insights. *Acta Cardiol* 2002; 57: 249–255.
150. Shinoda E, Yui Y, Kodama K, *et al.* Quantitative coronary angiogram analysis: nifedipine retard versus angiotensin-converting enzyme inhibitors (JMIC-B side arm study). *Hypertension* 2005; 45: 1153–1158.
151. Ishii N, Matsumura T, Kinoshita H, *et al.* Nifedipine induces peroxisome proliferator-activated receptor- γ activation in macrophages and suppresses the progression of atherosclerosis in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 2010; 30: 1598–1605.
152. Mulvaney JM, Zhang T, Fewtrell C, *et al.* Calcium influx through L-type channels is required for selective activation

- of extracellular signal-regulated kinase by gonadotropin-releasing hormone. *J Biol Chem* 1999; 274: 29796–29804.
153. Hirata A, Igarashi M, Yamaguchi H, *et al.* Nifedipine suppresses neointimal thickening by its inhibitory effect on vascular smooth muscle cell growth via a MEK-ERK pathway coupling with Pyk2. *Br J Pharmacol* 2000; 131: 1521–1530.
 154. Szabó C, Mitchell JA, Gross SS, *et al.* Nifedipine inhibits the induction of nitric oxide synthase by bacterial lipopolysaccharide. *J Pharmacol Exp Ther* 1993; 265: 674–680.
 155. Matsumori A, Nishio R, Nose Y. Calcium channel blockers differentially modulate cytokine production by peripheral blood mononuclear cells. *Circ J* 2010; 74: 567–571.
 156. Matsumoria A, Nunokawa Y, Sasayama S. Nifedipine inhibits activation of transcription factor NF- κ B. *Life Sci* 2000; 67: 2655–2661.
 157. Wu L, Iwai M, Li Z, *et al.* Nifedipine inhibited angiotensin II-induced monocyte chemoattractant protein 1 expression: involvement of inhibitor of nuclear factor κ B kinase and nuclear factor κ B-inducing kinase. *J Hypertens* 2006; 24: 123–130.
 158. Suzuki S, Nishimaki-Mogami T, Tamehiro N, *et al.* Verapamil increases the apolipoprotein-mediated release of cellular cholesterol by induction of ABCA1 expression via liver X receptor-independent mechanism. *Arterioscler Thromb Vasc Biol* 2004; 24: 519–525.
 159. Hashimoto R, Umemoto S, Guo F, *et al.* Nifedipine activates PPAR γ and exerts antioxidative action through Cu/ZnSOD independent of blood-pressure lowering in SHRSP. *J Atheroscler Thromb* 2010; 17: 785–795.
 160. Matsui T, Yamagishi S, Takeuchi M, *et al.* Nifedipine, a calcium channel blocker, inhibits advanced glycation end product (AGE)-elicited mesangial cell damage by suppressing AGE receptor (RAGE) expression via peroxisome proliferator-activated receptor- γ activation. *Biochem Biophys Res Commun* 2009; 385: 269–272.
 161. Matsui T, Yamagishi S, Takeuchi M, *et al.* Nifedipine inhibits advanced glycation end products (AGEs) and their receptor (RAGE) interaction-mediated proximal tubular cell injury via peroxisome proliferator-activated receptor- γ activation. *Biochem Biophys Res Commun* 2010; 398: 326–330.
 162. Lewis JD, Ferrara A, Peng T, *et al.* Risk of bladder cancer among diabetic patients treated with pioglitazone: interim report of a longitudinal cohort study. *Diabetes Care* 2011; 34: 916–922.