

EDGE ARTICLE

Cite this: *Chem. Sci.*, 2022, 13, 798

All publication charges for this article have been paid for by the Royal Society of Chemistry

Ultramacrocyclization in water *via* external templation†Qiong Chen,^a Ye Lei,^a Guangcheng Wu,^a Qing Li,^{*b} Yuanjiang Pan ^a and Hao Li ^{*ac}

Condensing a dihydrazide and each of a series of cationic bisaldehyde compounds bearing polymethylene chains in weakly acidic water produces either a macrocycle in a [1 + 1] manner or its dimer namely a [2] catenane, or their mixture. The product distribution is determined by the length of the bisaldehydes. Addition of cucurbit[8]uril (CB[8]) drives the catenane/macrocycle equilibria to the side of macrocycles, by forming ring-in-ring complexes with the latter. When the polymethylene unit of the bisaldehyde is replaced with a more rigid *p*-xylene linker, its self-assembly with the dihydrazide leads to quantitative formation of a [2]catenane. Upon addition of CB[8], the [2]catenane is transformed into an ultra-large macrocycle condensed in a [2 + 2] manner, which is encircled by two CB[8] rings. The framework of this macrocycle contains one hundred and two atoms, whose synthesis would be a formidable task without the external template CB[8]. Removal of CB[8] with a competitive guest leads to recovery of the [2] catenane.

Received 10th November 2021

Accepted 19th December 2021

DOI: 10.1039/d1sc06236k

rsc.li/chemical-science

Introduction

Synthesis of macrocyclic compounds represents one of the major focuses in the field of supramolecular chemistry.¹ When the sizes of the rings become larger, their production becomes more challenging, given that macrocyclization of rings containing more atoms implies more entropy loss. High dilution conditions² and/or addition of feasible guest templates³ have proven to be efficient approaches to suppress or avoid oligomeric or polymeric byproducts. However, these approaches become less useful in the case that the sizes of macrocyclic products are ultra-large, especially for those rings containing more than one hundred atoms. In addition, finding a feasible internal guest template to fit within the cavity of an ultra-large macrocycle becomes more difficult, despite a few exceptions.⁴

Our group⁵ and others⁶ discovered that hydrazone condensation represents an ideal dynamic covalent reaction that is amenable to use in water. A variety of [2]catenanes,^{5a,d-f} rings,^{5c,d,6d,e} and cages^{5b,g} were self-assembled in aqueous media, thanks to the reversible nature of hydrazone that allows the systems to perform error checking. In addition, cucurbit[8]uril

(CB[8])^{7c,e,g,i,k} was observed as a host that is able to accommodate a two π -electron guest, driven by the hydrophobic effect and dipole–cation interactions in the case of cationic guests such as pyridinium derivatives. We thus envision that by taking advantage of the marriage of dynamic covalent chemistry^{5,6,8} and external templation based on CB[8] rings,⁷ we might be able to self-assemble an ultra-large macrocycle. Here, by condensing a dihydrazide and a dicationic bisaldehyde containing a polymethylene chain in water, rings with medium sizes containing fifty to sixty atoms in their framework were self-assembled in a [1 + 1] manner as the major products. When the polymethylene chain becomes longer, the macrocycles start to undergo equilibration with their dimerized form namely [2] catenanes, driven by the hydrophobic effect. In the presence of CB[8], the macrocycle-[2]catenane equilibria shifted to the side of macrocycles, each of which was accommodated within the cavity of a CB[8] ring, forming a set of [2]pseudorotaxanes. The formation of these ring-in-ring complexes⁹ is favored by the CB[8]–pyridinium interactions between the host and the guest. When the poly-methylene chain in the bisaldehyde was changed to a more rigid *p*-xylene unit, its self-assembly with the dihydrazide linker resulted in the exclusive formation of a [2]catenane in close to a quantitative yield, thanks to the dynamic nature of hydrazone bonds that allows error checking. Addition of CB[8] led to decomposition of this [2]catenane. However, the putative [2]pseudorotaxane was not obtained as occurred in the case of the aforementioned polymethylene counterparts. This is probably on account of the *p*-xylene linker, which renders the [1 + 1] macrocycle too large to fit within the CB[8] cavity. Instead, the bisaldehyde and bishydrazide underwent condensation in

^aDepartment of Chemistry Institution, Zhejiang University, Hangzhou 310027, China. E-mail: lihao2015@zju.edu.cn

^bKey Laboratory of Macrocyclic and Supramolecular Chemistry, Guizhou University, Guiyang 550025, China. E-mail: nyliqing2012@163.com

^cZJU-Hangzhou Global Scientific and Technological Innovation Center, Hangzhou 310027, China

† Electronic supplementary information (ESI) available. CCDC 2014868, 2095808, 2096625 and 2096626. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1sc06236k



a [2 + 2] manner, forming an ultra-large macrocycle that was encircled by two CB[8] rings. This macrocycle contains more than one hundred atoms in its framework, whose formation would be rather challenging without the external template CB[8] rings. Upon removal of the CB[8] by adding a competitive guest, the ultra-large macrocycle underwent decomposition, regaining the [2]catenane.

Results and discussion

Each of the biscationic dialdehyde compounds, including $1x^{2+} \cdot 2Br^-$ ($x = a, b, c, d$ and e), contains two cationic pyridinium-benzaldehydes that are bridged by a polymethylene chain containing three to seven CH_2 units. A dihydrazine linker **2**, contains two phenylacetylhydrazide units that are connected by a glycol chain. The detailed synthetic procedures are described in the ESI.†

Self-assembly was performed by combining the dihydrazine linker **2** (2.5 mM) with each of biscationic dialdehydes $1x^{2+} \cdot 2Br^-$ ($x = a, b, c, d$ and e ; 2.5 mM) in D_2O . A catalytic amount of DCl was used to catalyze hydrazone condensation (Fig. 1 and Schemes S5–S7†). After heating the corresponding solutions at 60 °C for no less than 8 h, the corresponding 1H NMR spectra remained unchanged, showing that all the reactants were almost completely consumed. In the case of $1a^{2+}$ and $1b^{2+}$, two macrocycles ($1a^{2+} \cdot 2$) and ($1b^{2+} \cdot 2$), whose counteranions could be either Cl^- or Br^- , were self-assembled in a [1 + 1] manner as the major products, as indicated by both 1H NMR spectroscopy (Fig. S2 and S11†) and mass spectrometry (Fig. S1 and S10†). The yields of ($1a^{2+} \cdot 2$) and ($1b^{2+} \cdot 2$) were determined to be 76% and 65%, respectively, by using DMSO as an internal standard in the 1H NMR samples (Fig. S62B and S63B†). In the case of $1c^{2+}$ and $1d^{2+}$, the self-assembly products included both the macrocycles namely ($1c^{2+} \cdot 2$) and ($1d^{2+} \cdot 2$), as well as their

corresponding dimerized forms namely the [2]catenanes ($1c^{2+} \cdot 2$)₂ and ($1d^{2+} \cdot 2$)₂, whose counteranions could be either Cl^- or Br^- . The NMR yields of the [2]catenanes ($1c^{2+} \cdot 2$)₂ and ($1d^{2+} \cdot 2$)₂, as well as the rings ($1c^{2+} \cdot 2$) and ($1d^{2+} \cdot 2$) were calculated to be 61%, 71%, 33% and 13%, respectively (Fig. S64B and S65B†). In the case of the biscationic dialdehyde $1e^{2+}$ containing the longest polymethylene namely a $(CH_2)_7$ chain, a [2]catenane ($1e^{2+} \cdot 2$)₂ (counteranions could be either Cl^- or Br^-) was produced as the only observable product in the 1H NMR spectrum, whose yield is 88% (Fig. S66B†). Obviously, a dialdehyde with a longer polymethylene chain favors the formation of a [2]catenane to a more significant extent compared to its shorter counterparts.

Upon addition of an equimolar amount of CB[8] to a D_2O solution of a 1 : 1 mixture of **2** (2.5 mM) and each of $1x^{2+} \cdot 2Br^-$ ($x = a, b, c, d$ and e ; 2.5 mM) in the presence of DCl (Fig. 1 and Scheme S8†), a set of [2]pseudorotaxanes ($1x^{2+} \cdot 2$)₂CB[8] ($x = a, b, c, d$ and e) were observed to form as the predominant products. The yields of ($1x^{2+} \cdot 2$)₂CB[8] ($x = a, b, c, d$ and e) (counteranions could be either Cl^- or Br^-) were determined to be 80%, 74%, 88%, 90% and 85% (Fig. S62A, S63A, S64A, S65A and S66A†), respectively, calculated by using an internal standard in the corresponding 1H NMR samples. The structures of these [2]pseudorotaxanes were fully characterized by both 1H NMR spectroscopy and mass spectrometry (Fig. S6, S7, S15, S16, S25, S26, S33, S34, S42 and S43†). The 1H NMR spectra of both ($1a^{2+} \cdot 2$) and ($1a^{2+} \cdot 2$)₂CB[8] are shown in Fig. 2D. The resonances corresponding to the protons b and c in the guest undergo remarkable upfield shifts, indicating that the CB[8] host encircles the phenylene units in the $1a^{2+}$ residue, driven by the hydrophobic effect. As a comparison, the phenylene units in the **2** residue are located outside the pocket of CB[8], as indicated by the downfield shifts of the corresponding protons. CB[8] chooses to reside on the phenylene units in the $1a^{2+}$ residue




Fig. 1 Structural formulae of a series of dicationic bisaldehydes $1x^{2+}$ ($x = a, b, c, d$ and e) and a bishydrazide **2**. A series of dicationic macrocycles ($1x^{2+} \cdot 2$) ($x = a, b, c$ and d), as well as a set of tetracationic [2]catenanes ($1x^{2+} \cdot 2$)₂ ($x = c, d$ and e) were self-assembled by combining each of the bisaldehydes and the bishydrazide in water. Addition of CB[8] into each of these self-assembly products led to formation of a series of [2]pseudorotaxanes ($1x^{2+} \cdot 2$)₂CB[8] ($x = a, b, c, d$ and e). Charges are balanced by Cl^- or Br^- counteranions, which are omitted here for the sake of clarity. The yields shown are determined by 1H NMR spectroscopy. It seems that the counterions have little impact on the self-assembly.




Fig. 2 (A) The formation of $(1a^{2+} \cdot 2) \subset CB[8]$ by combining $(1a^{2+} \cdot 2)$ and $CB[8]$ in water. (B) Partial UV/Vis absorption spectra of $(1a^{2+} \cdot 2) \cdot 2Cl^-$ after adding different amounts of $CB[8]$ in H_2O at 298 K. $[(1a^{2+} \cdot 2)] = 3.0 \times 10^{-6} M$ for all spectra. (C) Plot of the changes of the absorbance intensity at $\lambda = 400$ nm in (B) versus the host/guest ratio namely $[CB[8]]/[(1a^{2+} \cdot 2)]$. (D) Partial 1H NMR spectra (500 MHz, D_2O , 298 K) of (i) $(1a^{2+} \cdot 2)$ and (ii) $(1a^{2+} \cdot 2) \subset CB[8]$, counteranions could be either Cl^- or Br^- .

instead of those in **2**, because of the cation–dipole interactions between the $CB[8]$ host and the pyridinium cations in the $1a^{2+}$ residue of the macrocyclic guest. The host/guest association and dissociation occurred at a relatively slow rate on the time-scale of 1H NMR spectroscopy, as indicated by the observation that the two protons of $CB[8]$ become chemically inequivalent in the 1H NMR spectrum of $(1a^{2+} \cdot 2) \subset CB[8]$ (Fig. 2D). Addition of an equimolar amount of $CB[8]$ to the corresponding pre-self-assembled [2]catenanes also produced the same [2]pseudorotaxane products, indicating that the corresponding [2]pseudorotaxanes $(1x^{2+} \cdot 2) \subset CB[8]$ ($x = a, b, c, d$ and e) are thermodynamic products, instead of kinetically trapped ones. Apparently, the formation of [2]pseudorotaxanes is more favored by dipole–cation interactions resulting from $CB[8]$, compared to the homo [2]catenanes.

In order to shed more light on the formation mechanism of the ring-in-ring pseudorotaxane, we measured the binding constants (K_a) of $(1a^{2+} \cdot 2) \subset CB[8]$ by using UV–Vis titration spectroscopic experiments (Fig. 2B). A pure sample of $(1a^{2+} \cdot 2) \cdot 2Cl^-$ was isolated *via* counterion exchange without chromatographic purification, which is water soluble. Addition of $CB[8]$ led to a decrease of the absorption peak of $(1a^{2+} \cdot 2)$ centered at 349 nm, while an increase of the absorption peak centered at 357 nm that corresponds to $(1a^{2+} \cdot 2) \subset CB[8]$. K_a of $(1a^{2+} \cdot 2) \subset CB[8]$ was determined (Fig. 2C) to be $5.6(\pm 1.1) \times 10^6 M^{-1}$, by fitting a 1 : 1 bind curve. Due to the fact that $CB[8]$ is barely soluble in water, titration experiments were performed at ultra-low

concentrations namely $[(1a^{2+} \cdot 2)] = 3.0 \times 10^{-3} mM$. Under such conditions, the concentrations of the compounds namely either $(1a^{2+} \cdot 2)$ or $CB[8]$ used in titration might be pseudo-accurate, which might be responsible for the absence of a clear isosbestic point in Fig. 2B. We thus performed competitive binding experiments at higher concentrations. That is, when $(1a^{2+} \cdot 2)$, paraquat and $CB[8]$, the concentration of each of which is 2.5 mM, were combined in D_2O , the 1H NMR spectrum (Fig. S67†) indicated that the $(1a^{2+} \cdot 2) \subset CB[8]$ was formed exclusively. Such an experiment implies that $(1a^{2+} \cdot 2)$ has a stronger binding affinity, compared to paraquat whose K_a is $1.1 \times 10^5 M^{-1}$.¹⁰ A similar competitive binding experiment (Fig. S68†) also clearly demonstrated that $(1a^{2+} \cdot 2) \subset CB[8]$ is a stronger complex compared to dodecane-1,12-diammonium $\subset CB[8]$, whose K_a is $1.1 \times 10^6 M^{-1}$.¹¹ We thus propose that the value of K_a of $(1a^{2+} \cdot 2) \subset CB[8]$ namely $5.6(\pm 1.1) \times 10^6 M^{-1}$ might still be reliable, even though not accurate.

$1f^{2+} \cdot 2Cl^-$, an analogue of $1x^{2+} \cdot 2Br^-$ ($x = a, b, c, d$ and e) whose polymethylene chains were replaced by a *p*-xylene unit, was synthesized. $1f^{2+} \cdot 2Cl^-$ (2.5 mM) and **2** (2.5 mM) were combined in D_2O in the presence of a catalytic amount of DCl (Fig. 4 and Scheme S9†). The solution was heated at 60 °C for 8 h. Both 1H NMR spectroscopy and mass spectrometry (Fig. 4D, S46 and S47†) demonstrated that a [2]catenane ($1f^{2+} \cdot 2$)₂·4Cl⁻ was produced as the predominant product. The NMR yield was calculated to be 94% (Fig. S60†). This yield is higher than those of the catenane containing polymethylene chains including $(1c^{2+} \cdot 2)_2$ and $(1d^{2+} \cdot 2)_2$, probably because the ring ($1f^{2+} \cdot 2$) contains a more rigid *p*-xylene unit. This linker affords the ring a more preorganized cavity that favours catenation to a more significant extent.

Upon the addition of 2 equiv. of $CB[8]$ to a D_2O solution of $(1f^{2+} \cdot 2)_2$ (Fig. 4 and Scheme S10†), the resonances corresponding to the catenane completely disappeared, while a new set of resonances was observed after heating the solution at 60 °C for 24 h (Fig. S57D†). Surprisingly, both the 1H NMR spectrum (Fig. 4B) and mass spectrum (Fig. S52†) indicated that a [3]pseudorotaxane namely $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ was self-assembled, containing two $CB[8]$ rings that encircle an ultra-large macrocycle $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$, which condensed in a [2 + 2] manner. The yield of $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ was determined to be 68%, by using an internal standard in the 1H NMR sample (Fig. S61†). It is noteworthy that a putative [2]pseudorotaxane $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset CB[8]$ containing only one $CB[8]$ was not observed during the self-assembly. The binding of two $CB[8]$ rings on the $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$ ring is positively cooperative. This proposition was supported by the observation that, addition of one equiv. of $CB[8]$ to a D_2O solution of $(1f^{2+} \cdot 2)_2$ led to a 1 : 1 mixture of the unreacted [2]catenane and the [3]pseudorotaxane (Fig. S57C†). It is likely that upon recognition of the first $CB[8]$ ring, the $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$ guest became more preorganized to accommodate the second ring.

Adamantan-1-ol,¹² a competitive guest binding with $CB[8]$, was added into the aqueous solution of $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ (Fig. 4 and S58†). After heating the corresponding solution at 60 °C for 48 h, the 1H NMR spectrum (Fig. S58F†) demonstrated that $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ was almost completely transformed into adamantan-1-ol $\subset CB[8]$, leading to recovery of the




Fig. 3 Solid-state structures of (A) $(1f^{2+} \cdot 2)_2$, (B) $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ (side view), and (C) $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ (rear view), obtained from single-crystal X-ray diffraction analysis. Oxygen atoms, red; nitrogen, blue; carbon, gray. Hydrogen atoms, counteranions, and disordered solvent molecules are omitted for clarity. Some of the close contacts are labeled with dashed lines, indicating the occurrence of intramolecular interactions such as π - π interactions.

[2]catenane $(1f^{2+} \cdot 2)_2$. This observation indicated that (i) adamantan-1-ol represents a better guest compared to $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$, with respect to the $CB[8]$ host; (ii) without the external templating of the $CB[8]$ ring, the ultra-large ring $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$ is no longer a thermodynamically stable compound in water compared to its [2]catenane isomer namely $(1f^{2+} \cdot 2)_2$.

Because $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2)$ can not be isolated, accurate determination of K_{a1} and K_{a2} of $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ was difficult or impossible. However, we could qualitatively evaluate K_{a2} by using competitive experiments. Upon heating a 2 : 1 : 2 mixture of $CB[8]$, the [2]catenane $(1f^{2+} \cdot 2)_2$ and the ring $(1a^{2+} \cdot 2)$ (Fig. S69[†]) for 24 h at 60 °C, the [2]pseudorotaxane $(1a^{2+} \cdot 2) \subset CB[8]$ was self-assembled selectively, leaving $(1f^{2+} \cdot 2)_2$ unreacted in water. As a comparison, heating a 1 : 2 : 2 mixture of the [2]catenane $(1f^{2+} \cdot 2)_2$, paraquat, and $CB[8]$ yielded $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ as the major product, accompanied by a small amount of paraquat $\subset CB[8]$ as the minor product (Fig. S70[†]). Such an experiment indicated that K_{a2} might be comparable to or a little larger than the K_a of paraquat $\subset CB[8]$, namely $1.1 \times 10^5 M^{-1}$, while smaller than the K_a of $(1a^{2+} \cdot 2) \subset CB[8]$.

Single crystals of the [2]catenane $(1f^{2+} \cdot 2)_2$ and the [3]pseudorotaxane $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ were obtained by either slow vapor diffusion of THF into the aqueous solution of $(1f^{2+} \cdot 2)_2$ (Fig. 3A), or slow evaporation of the aqueous solution of $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$ (Fig. 3B and C), which provided




Fig. 4 Structural formulae of a [2]catenane $(1f^{2+} \cdot 2)_2$ and a [3]pseudorotaxane $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$, by condensing a dicationic bisaldehyde $1f^{2+}$ and a bishydrazide **2** in water, in the absence and presence of $CB[8]$ respectively. Partial 1H NMR spectra (500 MHz, D_2O , 298 K) of (A) $1f^{2+}$, (B) $(1f^{2+} \cdot 2 \cdot 1f^{2+} \cdot 2) \subset 2CB[8]$, (C) **2** and (D) $(1f^{2+} \cdot 2)_2$. The assignment of each resonance was made using the corresponding two-dimensional NMR spectra shown in the ESI.[†] Charges are balanced by Cl^- counteranions, which are omitted here for the sake of clarity.

unambiguous evidence to convince their architectures. In the case of [2]catenane ($1\mathbf{f}^{2+}\cdot 2$)₂, the cavity of each of the two mutually mechanically interlocked rings is almost fully occupied by another ring, leading to a variety of intercomponent close contacts. For example, each of the pyridinium building blocks in one ring undergoes π - π interactions with an adjacent phenylene unit in another ring, as inferred from their short interplane distances, *i.e.*, around 3.4 Å (Fig. 3A). Close contacts also indicate the occurrence of CH \cdots O hydrogen bonding and CH- π interactions. In the case of the ring-in-rings complex namely the [3]pseudorotaxane ($1\mathbf{f}^{2+}\cdot 2\cdot 1\mathbf{f}^{2+}\cdot 2$) \subset 2CB[8] (Fig. 3B and C), each of the $1\mathbf{f}^{2+}$ residues adopts a different conformation compared to that in the [2]catenane ($1\mathbf{f}^{2+}\cdot 2$)₂, *i.e.*, in the former system, the two 4-phenylpyridinium units orient in different directions with respect to the central *p*-xylene unit. The two methylene units in the $1\mathbf{f}^{2+}$ residue are separated by 5.9 Å (Fig. 3A). This distance implies that the putative [1 + 1] ring, namely ($1\mathbf{f}^{2+}\cdot 2$), would be too large to fit within the cavity of CB[8]. As a consequence, the putative [2]pseudorotaxane ($1\mathbf{f}^{2+}\cdot 2$) \subset CB[8] can not form. Each of the CB[8] rings encircles two 4-phenylpyridinium units belonging to two different $1\mathbf{f}^{2+}$ residues. Such behavior favors the formation of the ultra-large macrocycle in a [2 + 2] manner. Within the cavity of the CB[8] ring, two phenylene units undergo π - π interactions, as inferred from the close contact namely 3.7 Å (Fig. 3C). The two pyridinium units, as a comparison, exhibit a displaced parallel orientation, which helps to avoid pyridinium-pyridinium electrostatic repulsion while strengthening ion-dipole and CH-O interactions between the cationic pyridinium units and the carbonyl groups in CB[8].

Conclusion

In summary, by combining the corresponding bishydrazide and a set of cationic bisaldehydes bearing polymethylene chains in aqueous media, a series of [1 + 1] macrocycles were self-assembled, accompanied with the corresponding dimers namely [2]catenanes. The product distribution was determined by the length of the polymethylene chains in the bisaldehyde precursors. That is, the longer chains favor the formation of [2]catenanes, while the shorter ones favor the macrocycles more. Upon addition of CB[8], the macrocycles form a set of [2]pseudorotaxanes, driving the catenane/macrocycle equilibria to the side of macrocycles. When the polymethylene units in the bisaldehyde compounds are replaced by a *p*-xylene unit, the putative [1 + 1] condensed ring is too large to fit within the ring of CB[8]. Instead, an ultra-large macrocycle was self-assembled in a [2 + 2] manner, which was encircled by two CB[8] rings that act as the external templates. The framework of the ultra-large ring contains more than one hundred atoms, whose synthesis would be thermodynamically disfavored in the absence of the external template, namely CB[8]. We envision that the success in formation of a ring-in-rings complex would be taken advantage of in the synthesis of more complex architectures, such as Borromean rings. Such trials are ongoing in our laboratory.

Data availability

Data for this paper, including the synthesis, structural characterization are available at ESI.†

Author contributions

Conceptualization: Hao Li, Qiong Chen. **Data curation:** Qiong Chen, Ye Lei. **Formal analysis:** Qiong Chen, Hao Li. **Investigation:** Qiong Chen, Ye Lei. **Methodology:** Qiong Chen, Guangcheng Wu, Yuanjiang Pan. **Project administration:** Qiong Chen, Qing Li. **Resources:** Hao Li. **Software:** Qiong Chen. **Supervision:** Qiong Chen, Hao Li. **Validation:** Qiong Chen, Ye Lei. **Roles/Writing – original draft:** Qiong Chen. **Writing – review & editing:** Hao Li.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

H. L. thanks the National Natural Science Foundation of China (No. 91856116, 21772173 and 21922108), the Natural Science Foundation of Zhejiang Province (No. LR18B020001), as well as the Fundamental Research Funds for the Central Universities (No. 2019FZA3007) for financial support. We thank Ms Ling He, Dr Yaqin Liu, and Prof. Mingxin Yu from the Analysis and Test Platform, Department of Chemistry, Zhejiang University for efficient NMR measurements.

Notes and references

- (a) D. J. Cram and J. M. Cram, *Science*, 1974, **183**, 803–809; (b) D. J. Cram, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 1009–1020.
- B. Odell, M. V. Reddington, A. M. Z. Slawin, N. Spencer, J. F. Stoddart and D. J. Williams, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 1547–1550.
- (a) M. Asakawa, W. Dehaen, G. L'abbé, S. Menzer, J. Nouwen, F. M. Raymo, J. F. Stoddart and D. J. Williams, *J. Org. Chem.*, 1996, **61**, 9591–9595; (b) C. D. Meyer, C. S. Joiner and J. F. Stoddart, *Chem. Soc. Rev.*, 2007, **36**, 1705–1723; (c) Y. Shen, L. Zou and Q. Wang, *New J. Chem.*, 2017, **41**, 7857–7860; (d) G. Wu, C. Y. Wang, T. Jiao, H. Zhu, F. Huang and H. Li, *J. Am. Chem. Soc.*, 2018, **140**, 5955–5961; (e) P. S. Bols, M. Rickhaus, L. Tejerina, H. Gotfredsen, K. Eriksen, M. Jirasek and H. L. Anderson, *J. Am. Chem. Soc.*, 2020, **142**, 13219–13226.
- (a) M. C. O'Sullivan, J. K. Sprafke, D. V. Kondratuk, C. Rinfray, T. D. W. Claridge, A. Saywell, M. O. Blunt, J. N. O'Shea, P. H. Beton, M. Malfois and H. L. Anderson, *Nature*, 2011, **469**, 72–75; (b) L. Favereau, A. Cossen, J. B. Kelber, J. Q. Gong, R. M. Oetterli, J. Cremers, L. M. Herz and H. L. Anderson, *J. Am. Chem. Soc.*, 2015, **137**, 14256–14259; (c) S. Wang, Y. Shen, B. Zhu, J. Wu and S. Li, *Chem. Commun.*, 2016, **52**, 10205–10216; (d) M. D. Peeks, T. D. W. Claridge and H. L. Anderson, *Nature*, 2017, **541**, 200–203; (e) J. Cremers, R. Haver, M. Rickhaus,

- J. Q. Gong, L. Favereau, M. D. Peeks, T. D. W. Claridge, L. M. Herz and H. L. Anderson, *J. Am. Chem. Soc.*, 2018, **140**, 5352–5355; (f) P. S. Bols and H. L. Anderson, *Acc. Chem. Res.*, 2018, **51**, 2083–2092.
- 5 (a) H. Li, H. Zhang, A. D. Lammer, M. Wang, X. Li, V. M. Lynch and J. L. Sessler, *Nat. Chem.*, 2015, **7**, 1003–1008; (b) X. Zheng, Y. Zhang, G. Wu, J. R. Liu, N. Cao, L. Wang, Y. Wang, X. Li, X. Hong, C. Yang and H. Li, *Chem. Commun.*, 2018, **54**, 3138–3141; (c) Y. Zhang, X. Zheng, N. Cao, C. Yang and H. Li, *Org. Lett.*, 2018, **20**, 2356–2359; (d) G. Wu, C.-Y. Wang, T. Jiao, H. Zhu, F. Huang and H. Li, *J. Am. Chem. Soc.*, 2018, **140**, 5955–5961; (e) C.-Y. Wang, G. Wu, T. Jiao, L. Shen, G. Ma, Y. Pan and H. Li, *Chem. Commun.*, 2018, **54**, 5106–5109; (f) Q. Chen, L. Chen, C.-Y. Wang, T. Jiao, Y. Pan and H. Li, *Chem. Commun.*, 2019, **55**, 13108–13111; (g) H. Wang, S. Fang, G. Wu, Y. Lei, Q. Chen, H. Wang, Y. Wu, C. Lin, X. Hong, S. K. Kim, J. L. Sessler and H. Li, *J. Am. Chem. Soc.*, 2020, **142**, 20182–20190.
- 6 (a) F. B. L. Cougnon, K. Caprice, M. Pupier, A. Bauza and A. Frontera, *J. Am. Chem. Soc.*, 2018, **140**, 12442–12450; (b) K. Caprice, M. Pupier, A. Bauza, A. Frontera and F. B. L. Cougnon, *Angew. Chem., Int. Ed.*, 2019, **58**, 8053–8057; (c) I. Neira, A. Blanco-Gomez, J. M. Quintela, C. Peinador and M. D. Garcia, *Org. Lett.*, 2019, **21**, 8976–8980; (d) A. Blanco-Gómez, Á. Fernández-Blanco, V. Blanco, J. Rodríguez, C. Peinador and M. D. García, *J. Am. Chem. Soc.*, 2019, **141**, 3959–3964; (e) A. Blanco-Gómez, I. Neira, J. L. Barriada, M. Melle-Franco, C. Peinador and M. D. Garcia, *Chem. Sci.*, 2019, **10**, 10680–10686; (f) K. Caprice, A. Aster, F. B. L. Cougnon and T. Kumpulainen, *Chem.–Eur. J.*, 2020, **26**, 1576–1587.
- 7 (a) S.-Y. Kim, I.-S. Jung, E. Lee, J. Kim, S. Sakamoto, K. Yamaguchi and K. Kim, *Angew. Chem., Int. Ed.*, 2001, **40**, 2119–2121; (b) H.-J. Kim, J. Heo, W. S. Jeon, E. Lee, J. Kim, S. Sakamoto, K. Yamaguchi and K. Kim, *Angew. Chem., Int. Ed.*, 2001, **40**, 1526–1529; (c) W. S. Jeon, A. Y. Ziganshina, J. W. Lee, Y. H. Ko, J.-K. Kang, C. Lee and K. Kim, *Angew. Chem., Int. Ed.*, 2003, **42**, 4097–4100; (d) Y. H. Ko, H. Kim, Y. Kim and K. Kim, *Angew. Chem., Int. Ed.*, 2008, **47**, 4106–4109; (e) L. Cera and C. A. Schalley, *Chem. Sci.*, 2014, **5**, 2560–2567; (f) S. J. Barrow, S. Kasera, M. J. Rowland, J. Barrio and O. A. Scherman, *Chem. Rev.*, 2015, **115**, 12320–12406; (g) Y. Shen, Q. Zhou, L. Zou and Q. Wang, *ChemistrySelect*, 2017, **2**, 11977; (h) N. A. Thompson, H. Barbero and E. Masson, *Chem. Commun.*, 2019, **55**, 12160–12163; (i) G. Wu, I. Szabó, E. Rosta and O. A. Scherman, *Chem. Commun.*, 2019, **55**, 13227–13230; (j) H. Wu, Y. Wang, L. O. Jones, W. Liu, B. Song, Y. Cui, K. Cai, L. Zhang, D. Shen, X.-Y. Chen, Y. Jiao, C. L. Stern, X. Li, G. C. Schatz and J. F. Stoddart, *J. Am. Chem. Soc.*, 2020, **142**, 16849–16860; (k) X.-K. Ma, W. Zhang, Z. Liu, H. Zhang, B. Zhang and Y. Liu, *Adv. Mater.*, 2021, 2007476; (l) H. Barbero and E. Masson, *Chem. Sci.*, 2021, **12**, 9962–9968.
- 8 (a) S. J. Rowan, S. J. Cantrill, G. R. L. Cousins, J. K. M. Sanders and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2002, **41**, 898–952; (b) L. Wang, M. O. Vysotsky, A. Bogdan, M. Bolte and V. Böhmer, *Science*, 2004, **304**, 1312–1314; (c) H. Y. Au-Yeung, G. D. Pantoş and J. K. M. Sanders, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 10466–10470; (d) Y. Jin, C. Yu, R. J. Denman and W. Zhang, *Chem. Soc. Rev.*, 2013, **42**, 6634–6654; (e) G. Zhang, O. Presly, F. White, I. M. Oppel and M. Mastalerz, *Angew. Chem., Int. Ed.*, 2014, **53**, 1516–1520; (f) S. Klotzbach and F. Beuerle, *Angew. Chem., Int. Ed.*, 2015, **54**, 10356–10360; (g) Q. Wang, C. Yu, H. Long, Y. Du, Y. Jin and W. Zhang, *Angew. Chem., Int. Ed.*, 2015, **54**, 7550–7554; (h) Q. Wang, C. Yu, C. Zhang, H. Long, S. Azarnoush, Y. Jin and W. Zhang, *Chem. Sci.*, 2016, **7**, 3370–3376; (i) S. Lee, A. Yang, T. P. Moneypenny and J. S. Moore, *J. Am. Chem. Soc.*, 2016, **138**, 2182–2185; (j) H. Löw, E. Mena-Osteritz and M. von Delius, *Chem. Commun.*, 2019, **55**, 11434–11437; (k) J. Hu, S. K. Gupta, J. Ozdemir and M. H. Beyzavi, *ACS Appl. Nano Mater.*, 2020, **3**, 6239–6269; (l) T. Jiao, G. Wu, Y. Zhang, L. Shen, Y. Lei, C.-Y. Wang, A. C. Fahrenbach and H. Li, *Angew. Chem., Int. Ed.*, 2020, **59**, 18350–18367; (m) F. V. Lijsebetten, J. O. Holloway, J. M. Winne and F. E. D. Prez, *Chem. Soc. Rev.*, 2020, **49**, 8425–8438; (n) A. J. Greenice, C. I. Wendell, M. M. Cencer, S. D. Laffoon and J. S. Moore, *Trends Chem.*, 2020, **2**, 1043–1051; (o) Y. Chen, G. Wu, B. Chen, H. Qu, T. Jiao, Y. Li, C. Ge, C. Zhang, L. Liang, X. Zeng, X. Cao, Q. Wang and H. Li, *Angew. Chem., Int. Ed.*, 2021, **60**, 2–8.
- 9 (a) S.-Y. Kim, I.-S. Jung, E. Lee, J. Kim, S. Sakamoto, K. Yamaguchi and K. Kim, *Angew. Chem., Int. Ed.*, 2001, **40**, 2119–2121; (b) A. I. Day, R. J. Blanch, A. P. Arnold, S. Lorenzo, G. R. Lewis and I. Dance, *Angew. Chem., Int. Ed.*, 2002, **41**, 275–277; (c) S.-H. Chiu, A. R. Pease, J. F. Stoddart, A. J. P. White and D. J. Williams, *Angew. Chem., Int. Ed.*, 2002, **41**, 270–274; (d) J. C. Loren, M. Yoshizawa, R. F. Haldimann, A. Linden and J. S. Siegel, *Angew. Chem., Int. Ed.*, 2003, **42**, 5702–5705; (e) T. Iwanaga, R. Nakamoto, M. Yasutake, H. Takemura, K. Sako and T. Shinmyozu, *Angew. Chem., Int. Ed.*, 2006, **45**, 3643–3647; (f) R. S. Forgan, C. Wang, D. C. Friedman, J. M. Spruell, C. L. Stern, A. A. Sarjeant, D. Cao and J. F. Stoddart, *Chem.–Eur. J.*, 2012, **18**, 202–212; (g) B. Sun, M. Wang, Z. Lou, M. Huang, C. Xu, X. Li, L. J. Chen, Y. Yu, G. L. Davis, B. Xu, H. B. Yang and X. Li, *J. Am. Chem. Soc.*, 2015, **137**, 1556–1564; (h) J. K. Klosterman, J. Veliks, D. K. Frantz, Y. Yasui, M. Loepfe, E. Zysman-Colman, A. Linden and J. S. Siegel, *Org. Chem. Front.*, 2016, **3**, 661–666; (i) Y. Lu, H. N. Zhang and G. X. Jin, *Acc. Chem. Res.*, 2018, **51**, 2148–2158; (j) M. C. Lipke, Y. Wu, I. Roy, Y. Wang, M. R. Wasielewski and J. F. Stoddart, *ACS Cent. Sci.*, 2018, **4**, 362–371; (k) K. Zhu, G. Baggi and S. J. Loeb, *Nat. Chem.*, 2018, **10**, 625–630; (l) H. Wu, Y. Wang, L. O. Jones, W. Liu, B. Song, Y. Cui, K. Cai, L. Zhang, D. Shen, X.-Y. Chen, Y. Jiao, C. L. Stern, X. Li, G. C. Schatz and J. F. Stoddart, *J. Am. Chem. Soc.*, 2020, **142**, 16849–16860.
- 10 W. S. Jeon, H.-J. Kim, C. Lee and K. Kim, *Chem. Commun.*, 2002, **17**, 1828–1829.
- 11 K. Baek, Y. Kim, H. Kim, M. Yoon, I. Hwang, Y. H. Ko and K. Kim, *Chem. Commun.*, 2010, **46**, 4091–4093.
- 12 S. Moghaddam, C. Yang, M. Rekharsky, Y. H. Ko, K. Kim, Y. Inoue and M. K. Gilson, *J. Am. Chem. Soc.*, 2011, **133**, 3570–3581.