




# Children born during the hunger season are at a higher risk of severe acute malnutrition: Findings from a Guinea Sahelian ecological zone in Northern Ghana

Engelbert A. Nonterah<sup>1,2</sup>  | Paul Welaga<sup>1,3</sup> | Samuel T. Chatio<sup>1</sup> | Sarah H. Kehoe<sup>4</sup>  | Winfred Ofofu<sup>5</sup> | Kate A. Ward<sup>4,6,8</sup> | Keith M. Godfrey<sup>4</sup>  | Abraham R. Oduro<sup>1</sup> | Marie-Louise Newell<sup>7,8</sup> | as members of the INPreP study group

<sup>1</sup>Navrongo Health Research Centre, Ghana Health Service, Navrongo, Ghana

<sup>2</sup>Julius Global Health, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

<sup>3</sup>School of Medicine and Dentistry, C K Tedam University of Technology and Applied Sciences, Navrongo, Upper East Region, Ghana

<sup>4</sup>MRC Lifecourse Epidemiology Unit and NIHR Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust, University of Southampton, Southampton, UK

<sup>5</sup>Upper East Regional Health Directorate, PMB, Bolgatanga, Ghana

<sup>6</sup>Global Health Research Institute, School of Health and Human Development, University of Southampton, Southampton, UK

<sup>7</sup>Department of Human Development, University of Southampton, Southampton, UK

<sup>8</sup>School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

## Correspondence

Engelbert A. Nonterah, Navrongo Health Research Centre, Ghana Health Service, Navrongo, Ghana.  
Email: [drenanonterah@gmail.com](mailto:drenanonterah@gmail.com)

## Funding information

National Institute for Health Research, Grant/Award Number: 17\63\154

## Abstract

Heightened food insecurity in the hunger season increases the risk of severe acute malnutrition (SAM) in childhood. This study examined the association of season of birth with SAM in a Guinean Sahelian ecological zone. We analyzed routine health and sociodemographic surveillance data from the Navrongo Health and Socio-demographic Surveillance System collected between 2011 and 2018. January–June, the period of highest food insecurity, was defined as the hunger season. We defined moderate acute malnutrition as child mid-upper arm circumference (MUAC) between 115 mm and 135 mm and SAM as MAUC  $\leq$  115 mm. We used adjusted logistic regression to quantify the association between the season of birth and SAM in children aged 6–35 months. From the 29,452 children studied, 24% had moderate acute malnutrition. Overall, 1.4% had SAM, with a higher prevalence (1.8%) in the hunger season of birth. Compared with those born October–December, adjusted odds ratios (aOR) and 95% confidence interval (95% CI) for SAM were increased for children born in the hunger season: January–March (1.77 [1.31–2.39]) and April–June (1.92 [1.44–2.56]). Low birth weight, age at an assessment of nutritional status, and ethno-linguistic group were also significantly associated with SAM in adjusted analyses. Our study established that being born in the hunger season is associated with a higher risk of severe acute malnutrition. The result implies improvement in the food supply to pregnant and lactating mothers through sustainable agriculture or food system change targeting the hunger season may reduce the burden of severe acute malnutrition.

## KEYWORDS

food insecurity, Ghana, malnutrition, severe acute malnutrition, seasonal variation

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Maternal & Child Nutrition* published by John Wiley & Sons Ltd.

## 1 | INTRODUCTION

After decades of undernutrition, Africa is now seeing a surge in overnutrition (Bain et al., 2013; Davies et al., 2018; UNICEF World Health Organization & World Bank, 2020). Despite the surge in overnutrition and the double burden, undernutrition remains an important public health problem in many African countries and requires a multidisciplinary approach to address it (Doku & Neupane, 2015). Like many countries in sub-Saharan Africa (SSA), Ghana has a high burden of malnutrition among children aged 0–59 months with stunting, underweight and wasting all being highly prevalent (19%, 11%, and 5%, respectively in 2014) (Black et al., 2008; Ghana Statistical Service Ghana Health Service & ICF International, 2014), with highest rates in the northern part of Ghana (Ghana Statistical Service Ghana Health Service & ICF International, 2014; Yawson et al., 2017). In sub-Saharan African countries most communities rely on rain-enabled agriculture due to minimal access to irrigation facilities (Abizari et al., 2017; Chikungu & Madise, 2014; Fentahun et al., 2018; Hagos et al., 2014). This creates a seasonal pattern of access to food, where the “hunger season” exposes families to food insecurity and its profound effect on nutritional status, particularly in children under age 5 years (Belayneh et al., 2020; Fentahun et al., 2018). The hunger season, a period between planting and harvest when food runs out is usually the worst hit period. This comes with profound adverse health effects on mothers and children. The hunger season is a challenging period for subsistence farmers and their families, who solely rely on what they grow.

The established underlying causes of malnutrition across all age groups likely follow a seasonal pattern, with consequences for morbidity and mortality across the life cycle (Bryson et al., 2021). Hence establishing the seasonal patterns in food availability and its effect on nutritional health or morbidity have policy implications in planning the distribution of scarce resources. While Demographic and Health Surveys do not measure the impact of seasonal variations in food supply and nutritional status, several epidemiological studies (though not at the country level) have provided insights (Sorgho et al., 2016). Most of these studies were not, however, conducted during the first 1000daysPlus (preconception through pregnancy, delivery and post-delivery and infancy up to 5 years), which is considered by the World Health Organization as a critical period for optimizing growth. Further evidence accruing from the developmental origins of health and diseases concept further highlights the importance of the 1000day-Plus period as a highly vulnerable period where adverse exposures such as malnutrition can have lasting effects on health in later life (Davies et al., 2018).

Previous studies from SSA have reported inconsistent findings regarding any link between seasonal fluctuations in access to food and the nutritional status of young children (Chikungu & Madise, 2014). Some studies from Gambia (Tomkins et al., 1986) and Malawi (Chikungu & Madise, 2014) have reported seasonal trends in child growth and demonstrated weight and height deficits during the rainy season (period for planting with minimal food availability). A study from northern Ghana had also observed a seasonal effect of

### Key Points

- Children born in the hunger season are at a higher risk of severe acute malnutrition in this Guinea-Saharan ecological zone.
- Children born with low birth weight were at a higher risk of severe acute malnutrition. Being female and of the Nankam ethno-linguistic group increased ones odds of severe acute malnutrition.
- Our observations imply sustainable food supply to pregnant and lactating mothers through improved agriculture or food system change targeting the hunger season may reduce the burden of severe acute malnutrition.

low dietary diversity and hence low micronutrient intake during the hunger season for both mothers and children (Abizari et al., 2017). In contrast, a study in Kenya observed increased nutrient intake in the rainy season. The preharvest season is often seen as a vulnerable period for acute malnutrition among exposed children. The effect is especially felt in drought-prone areas which rely mostly on rain-enabled agriculture or subsistence farming.

These current analyses of children aged 6–35 months from the Navrongo Health and Demographic Surveillance System (HDSS) (Oduro et al., 2012) was conducted as part of a broader INPreP - Improved nutrition preconception, during pregnancy and post-delivery study (<https://www.southampton.ac.uk/global-health/research/lifecourse-epidemiology/inprep/about-us.page>). We set out to determine the seasonal variation in the burden of severe acute malnutrition. We further determined the association of season of birth with SAM focusing especially on the effect of being born during the hunger season and the risk of SAM. We also reported on the unique factors in the Guinean Sahelian ecological zone and established factors that influence the nutritional status of children (Bain et al., 2013; Madan et al., 2018).

## 2 | METHODS

We used data from the Navrongo Health and Demographic Surveillance system (HDSS) in Ghana (Oduro et al., 2012) for this study. As part of data collection in the HDSS catchment area, trained field workers visited all households 2–3 times a year to collect relevant data. The HDSS currently monitors 165,000 individuals; data collected includes pregnancies, births, deaths (through verbal autopsies), hospitalization, immunization, and migration. In addition, the field workers assess the nutritional status of children between 6 and 35 months by measuring their mid-upper arm circumferences (MUAC). Other data collected include sociodemographic characteristics of the mothers and children including sex, age, maternal education, and household socioeconomic status (in wealth quintiles). Data on birth

weight were extracted from the child health records of the children or the maternal health record book. For this study, we extracted and utilized HDSS data from 2011 to 2018.

## 2.1 | Study setting

The Navrongo Health and Demographic Surveillance System (HDSS) (Oduro et al., 2012) operational area lie in an ecological belt that is food insecure due to climatic conditions. The area is highly dependent on rain-enabled seasonal subsistence agriculture for livelihoods. We conducted this study in the Kasena–Nankana West district and Kasena–Nankana municipality of Northern Ghana. The study area lies within the Guinea-Saharan ecological zone in Northern Ghana, which shares a border with Burkina Faso in the north-eastern corner. It covers a land area of 1675 km<sup>2</sup> with an estimated population of 165,000 in over 32,000 households under continuous demographic surveillance. There are two main ethnic groups, the Kasena and Nankani, who together constitute about 96% of the population.

The main occupation of the people is rain-enabled subsistence agriculture, and many parts (about 80%) are rural (Oduro et al., 2012). The study area faces cyclic food insecurity that alternates between pre- and postharvest seasons, making the population vulnerable to seasonal food shortages. The single rainy season in rural Northern Ghana often begins in May and ends around October. The area usually experiences heavy rainfall in August and September. Before some health interventions were introduced in the study area by the Navrongo Health Research Center (NHRC) both childhood and adult mortality were among the highest in the country. Under-five mortality rates, as well as family sizes, have since declined considerably in the study area. The NHRC manages the Navrongo HDSS which monitors morbidity and mortality of the entire population in the study area. The HDSS collects and updates data on births, deaths, pregnancies, marriages, in-migrations, out-migrations, and other vital events by visiting households 2–3 times annually (Oduro et al., 2012). Through key informants in the community, new houses are identified for which field workers visit and then determine and register the number of households as well as resident individuals. These are then subsequently visited every 6 months to update the records and collect new information when needed.

## 2.2 | Data management and statistical analyses

Routine data collected by the field workers are doubly entered and verified by a data manager for inconsistent entries using Visual FoxPro. We extracted the verified data and used STATA 16.0 for data cleaning and analysis. This analysis was limited to children with at least one MUAC measurement between January 2011 and December 2018. During each household visit, the mid-upper arm circumference (MUAC) of each child below 3 years was measured. We used the first MUAC score of the child if the child had more than one. MUAC of <115 mm was categorized as severe acute malnutrition (SAM),

MUAC between 115 mm and 135 mm as moderate acute malnutrition, and a MUAC of >135 mm as well-nourished (World Health Organization and UNICEF, 2009).

Birth weight was categorized as low (birth weight <2.5 kg) or normal (birth weight ≥2.5 kg) and according to 11th edition of the World Health organizations' (WHO) international classification of diseases (WHO, 2019). Season of birth was grouped into four categories to reflect the weather patterns and food availability within a year, that is, January–March (postharvest season), April–June (immediate pre-planting), July–September (planting season), and October–December (immediate harvest season). The hunger season was thus defined as the period between January and June, corresponding to the period of severely reduced availability of food.

We computed household wealth index using principal component analysis (PCA) from 30 separate household items, from large assets (e.g., land and car ownership) to smaller household items (e.g., phone, fan ownership). The principal component analysis of the household assets or items was done to predict factor scores. These scores were then categorized into wealth quintiles: Q1 = poorest, Q2 = poorer, Q3 = poor, Q4 = less poor, and Q5 = least poor.

We used logistic regression to examine the association between seasonality and severe acute malnutrition. Unadjusted and adjusted odds ratios (aOR) with 95% confidence intervals (95% CIs) were computed. We included all potential confounding variables in our data in our multivariable logistic regression model. These variables include season of birth, sex of child, birth weight, maternal education, wealth quintile, age of child, maternal age, mother's marital status, place of residence, and ethnicity. We did not find any statistically significant interaction between variables. The test of statistical significance was set at 5% ( $p < 0.05$ ).

## 3 | RESULTS

A total of 29,452 children aged 6–35 months and born between 2008 and 2018 were included in the analysis. Their median age was 11 months (interquartile range: 8–18). About 9% of the children had low birth weight, 50.2% were male and over half (55%) were age <12 months. The majority of the children (68.2%) were born to mothers aged 20–34 years, and 87% were residents in rural parts of the two districts. Almost 23% of the children were born to mothers with no formal education and 17.6% were born to mothers with secondary or tertiary education. Children born to unmarried women constituted 4% of the study population (Table 1)

Figure 1 shows the malnutrition rates by season of birth. Overall 24% of children were malnourished, with 1.4% having SAM and 22.5% moderate acute malnutrition; 76% were well-nourished. Children born in the second half of the hunger season, that is, April to June, had the highest rate of SAM (1.8%), while those born in the season after the harvest period (October–December) had the lowest SAM rate (1.0%). Low birth weight children had a SAM rate of 2.4% compared with 1.3% for normal birth weight children. Additionally, the SAM rate was 1.5% among females and 1.2% among males. Children aged below 12 months had a SAM rate of 1.8%.

**TABLE 1** Background characteristics by severe acute malnutrition rate of children age 6–35 months included in the analysis in the Navrongo Health and Demographic Surveillance System from 2011 to 2018

| Variable               | Number (%)    | Number with SAM | Percent with SAM | p-value          |
|------------------------|---------------|-----------------|------------------|------------------|
| Month of birth         |               |                 |                  |                  |
| January–March          | 6479 (22)     | 107             | 1.7              | <b>&lt;0.001</b> |
| April–June             | 7871 (26.7)   | 141             | 1.8              |                  |
| July–September         | 7656 (26)     | 88              | 1.1              |                  |
| October–December       | 7446 (25.3)   | 72              | 1.0              |                  |
| Sex                    |               |                 |                  |                  |
| Male                   | 14,778 (50.2) | 182             | 1.2              | <b>0.023</b>     |
| Female                 | 14,674 (49.8) | 226             | 1.5              |                  |
| Birth weight           |               |                 |                  |                  |
| Low birth weight       | 2615 (8.9)    | 62              | 2.4              | <b>&lt;0.001</b> |
| Normal birth weight    | 20,402 (69.3) | 271             | 1.3              |                  |
| Don't know             | 6435 (21.8)   | 75              | 1.2              |                  |
| Maternal education     |               |                 |                  |                  |
| No education           | 6683 (22.7)   | 93              | 1.4              | 0.82             |
| Primary-JSS            | 17,591 (59.7) | 248             | 1.4              |                  |
| Secondary-tertiary     | 5178 (17.6)   | 67              | 1.3              |                  |
| Household wealth       |               |                 |                  |                  |
| Poorest                | 8345 (28.3)   | 106             | 1.3              | <b>0.011</b>     |
| Poorer                 | 6802 (23.1)   | 109             | 1.6              |                  |
| Poor                   | 5500 (18.7)   | 94              | 1.7              |                  |
| Less poor              | 5584 (19)     | 69              | 1.2              |                  |
| Least poor             | 3221 (10.9)   | 30              | 0.9              |                  |
| Age at measurement     |               |                 |                  |                  |
| Under 1 year           | 16,240 (55.1) | 287             | 1.8              | <b>&lt;0.001</b> |
| 1–2 years              | 8846 (30.0)   | 102             | 1.2              |                  |
| 2–3 years              | 4366 (14.8)   | 19              | 0.4              |                  |
| Maternal age           |               |                 |                  |                  |
| <20 years              | 3291 (11.2)   | 35              | 1.1              | 0.108            |
| 20–34                  | 20,075 (68.2) | 276             | 1.4              |                  |
| 35–49                  | 6086 (20.7)   | 97              | 1.6              |                  |
| Marital status         |               |                 |                  |                  |
| Married                | 28,317 (96.1) | 388             | 1.4              | 0.268            |
| Not married            | 1135 (3.9)    | 20              | 1.8              |                  |
| Residence              |               |                 |                  |                  |
| Urban                  | 3740 (12.7)   | 44              | 1.2              | 0.242            |
| Rural                  | 25,711 (87.3) | 364             | 1.4              |                  |
| Ethno-linguistic group |               |                 |                  |                  |
| Kasem                  | 14,427 (49)   | 170             | 1.2              | <b>0.006</b>     |
| Nankam                 | 13,686 (46.5) | 221             | 1.6              |                  |

TABLE 1 (Continued)

| Variable | Number (%) | Number with SAM | Percent with SAM | p-value |
|----------|------------|-----------------|------------------|---------|
| Buli     | 608 (2.1)  | 11              | 1.8              |         |
| Other    | 731 (2.5)  | 6               | 0.8              |         |
| Total    | 29,452     | 408             | 1.4              |         |

Note: p-value calculated using Pearson's  $\chi^2$  test to show differences between the groups of the variables in the row data.

Abbreviations: JSS, junior high school; SAM, severe acute malnutrition.

**FIGURE 1** Proportion of children aged 6–35 months with severe acute malnutrition according to seasons in the Navrongo Health and Demographic Surveillance Site from 2011 to 2018




Figure 2 shows the yearly proportion of SAM cases among children 6–35 months in the NHDSS from 2011 to 2018. 2011 and 2015 recorded the lowest proportions of deaths while 2018 had the highest proportion. Mortality was fairly similar for the other years.

In Table 2, we show crude (OR) and aOR of sociodemographic characteristics, season of birth, economic characteristics of the mother and child characteristics in relation to SAM. In crude analyses, being born between January and June (hunger season), a female child, low birth weight, less than 2 years of age at the time of the survey, and coming from the Nankam ethno-linguistic group all increased odds of SAM. In adjusted multivariable analyses, being born between April and June was associated with nearly two-fold increased odds for SAM (aOR = 1.92, 95% CI [1.44–2.56]) compared to being born in October–December. Similarly, being born in the second half of the hunger season, January–April was associated with 1.8-fold increased odds of SAM compared with being born in October–December (aOR = 1.77, 95% CI [1.31–2.39]).


Children with low birth weight had an independent 90% increased odds of SAM compared to those with normal birth weight, (aOR = 1.88, 95% CI [1.42–2.50]). Children aged below 12 months at the time of the survey had four times greater odds of SAM compared to those aged 2 years or more, aOR = 4.05, 95% CI [(2.52–6.51)]. Being aged 12–23 months at the time of survey was associated greater than 160% increased odds for SAM compared to those aged two years or more (aOR = 2.64, 95% CI [1.61–4.33]). Females had 1.2-fold increased odds for SAM compared to males (aOR = 1.23, 95% CI [1.01–1.50]). Children from the Nankani ethno-linguistic

group had a higher odds of SAM compared with the Kasem ethno-linguistic group (Table 2).

## 4 | DISCUSSION

We showed an overall SAM rate of 1.4% and a moderate acute malnutrition rate of 22.5%. The highest SAM rate by a season of birth was observed in children born between April and June, the period considered to be the severe hunger season. This is the immediate pre-planting period and several months postharvest season. During this period most farmers are usually actively cultivating their farms and preparing to plant their crops. The INPreP study, a global health research group, had previously reported in a qualitative study that communities in the Navrongo HDSS see food insecurity as a major challenge to optimal nutrition due to lack of irrigated agriculture and improved agriculture (Debpuur et al., 2020; Watson et al., 2020). The highest rate of SAM was observed in 2018 and this may be due to the worse climatic conditions leading to poor yield and harvest. Fagariba et al. (2018) in study conducted in a neighboring Talensi district observed that 2018 experienced highest draught and temperature hence affecting food production despite climate change adaptation practices such as a change in time of planting, improved crop varieties and land rotation. Low birth weight children were most likely to be severely malnourished compared children with birth weight greater than 2.5 kg.

In multivariable analysis, we found that being born between April and June, being female, having low birth weight (birth weight < 2.5 kg),



**FIGURE 2** Yearly proportions of severe acute malnutrition cases in children aged 6–35 months in the Navrongo Health and Demographic Surveillance Site from 2011 to 2018

born into a poor household, being less than one year old and being from the Nankam ethnic group in the study area conferred the highest odds for SAM. The season in which a child is born is important to the child's growth and health. Agriculture serves as the primary livelihood of the people with a majority of the inhabitants being subsistence farmers who live in small scattered settlements. In northern Ghana, the agricultural cycle is driven by agro-ecological conditions and by annual cycles in the weather. Our study showed that children born between April and June, which is considered to be the lean season, had increased odds for SAM compared to children born during the harvest period of July to September and the immediate postharvest period of October to December. It is possible that during the hunger season with food insecurity, the diet of a lactating mother may not be optimal and could therefore affect the child's adequate lactation which could acutely affect nutritional status of the child. Previous studies in sub-Saharan Africa have also observed the association of seasonal variation with incidence of low birth weight and severe acute malnutrition (Rogawski McQuade et al., 2019).

Previous studies have shown that the mother's diet in pregnancy and lactation may program her offspring where the fetus adapts to undernutrition by changing its metabolism, altering its production of hormones and sensitivity of tissues to them, redistributing its blood flow, and slowing its growth rate (Morrison & Regnault, 2016). Children born in the harvest period of October–December had the lowest SAM rate, perhaps due to the good supply of food and hence higher nutritional status of the lactating mother.

Our results showed a strong association between birth weight and SAM among children in the population. The odds ratios of SAM in low birth weight children was nearly twice the odds of severe acute malnutrition in normal birth weight children. This observation is consistent with similar studies conducted in Tanzania (Rogawski McQuade et al., 2019), Malawi (Chikhungu & Madise, 2014), Ethiopia (Fentahun et al., 2018), India (Choudhary et al., 2019), and Bangladesh (Hillbruner & Egan, 2008). One of the major reasons for the association of low birth weight with SAM is suboptimal maternal nutrient intake. In the hunger season, there is greater likelihood that milk production among lactating mothers is reduced (Roba et al., 2016). Children born in the lean season also have a greater susceptibility to weight loss in the first

two weeks of life and this may also contribute to the higher risk of SAM. This may be explained by low quantity and/or quality of breast milk during this time. Some studies observed that these disparities were persistent and associated with smaller but sustained deficits at the age of 2 years. This suggests incomplete catch-up growth among children born in the high food insecurity season. Low birth weight children may also be prone to infection (such as diarrheal diseases and respiratory infections), loss of appetite and fatigue, jaundice, and anemia (Jones & Berkley, 2014). These competing morbidities may further hinder catch-up growth among these children and set them on a trajectory of sustained malnutrition and poor physical growth till after the age of five years.

In the study area, the hunger season is also associated with harsh climatic conditions including the seasonal north-east harmattan winds blowing from the Sahara and extremely high temperatures, which may affect dry season gardening due to lack of irrigated agriculture. These may act together to increase respiratory infections and further worsen nutrient intake in the said period. The Ghana Nutrition profile shows that younger children are at a higher risk of SAM but data on age-categorized sub-analysis was not provided (<https://www.usaid.gov/sites/default/files/documents/1864/Ghana-Nutrition-Profile>). Largely consistent with our findings, a study from northern Ghana had previously established that malnutrition was higher among children aged 6–8 and 12–23 months (Saaka et al., 2015). The authors observed that the lower age groups (<12 months) were less likely to recover from severe acute malnutrition (Akparibo et al., 2017).

We observed that lower household socioeconomic status was associated with higher risk of severe acute malnutrition while relatively wealthier households having lower risk of SAM. Similar findings have been previously reported in studies from western Nigeria (Imam et al., 2020), Pakistan (Ahmad et al., 2020), and Jamaica (Thompson et al., 2017). A study from Nepal also observed that families with higher socioeconomic status were likely to present with lower SAM (Hossain et al., 2020). These findings were further corroborated by a study in Ghana that observed that lower family income was associated with high levels of SAM (Tette et al., 2016). A plausible reason is that higher economic status is associated with a higher purchasing

**TABLE 2** Crude and adjusted odds ratios of factors associated with severe acute malnutrition in children aged 6–35 months in the Navrongo Health and Demographic Surveillance System from 2011 to 2018

| Variable               | Number (%)    | Crude OR (95% CI) | p-value          | Adjusted OR (95% CI) | p-value          |
|------------------------|---------------|-------------------|------------------|----------------------|------------------|
| Month of birth         |               |                   |                  |                      |                  |
| January–March          | 6479 (22)     | 1.72 (1.27–2.32)  | <b>&lt;0.001</b> | 1.77 (1.31–2.39)     | <b>&lt;0.001</b> |
| April–June             | 7871 (26.7)   | 1.87 (1.40–2.49)  |                  | 1.92 (1.44–2.56)     |                  |
| July–September         | 7656 (26)     | 1.19 (0.87–1.63)  |                  | 1.19 (0.87–1.63)     |                  |
| October–December       | 7446 (25.3)   | Ref               |                  | Ref                  |                  |
| Sex                    |               |                   |                  |                      |                  |
| Male                   | 14,778 (50.2) | Ref               | <b>0.024</b>     | Ref                  | <b>0.041</b>     |
| Female                 | 14,674 (49.8) | 1.25 (1.03–1.53)  |                  | 1.23 (1.01–1.50)     |                  |
| Birth Weight           |               |                   |                  |                      |                  |
| Low birth weight       | 2615 (8.9)    | 1.80 (1.36–2.38)  | <b>&lt;0.001</b> | 1.88 (1.42–2.50)     | <b>&lt;0.001</b> |
| Normal birth weight    | 20,402 (69.3) | Ref               |                  | Ref                  |                  |
| Don't know             | 6435 (21.8)   | 0.88 (0.68–1.13)  |                  | 1.06 (0.81–1.38)     |                  |
| Maternal education     |               |                   |                  |                      |                  |
| No education           | 6683 (22.7)   | Ref               | 0.821            | Ref                  | 0.636            |
| Primary/JSS            | 17,591 (59.7) | 1.01 (0.8–1.29)   |                  | 1.10 (0.86–1.42)     |                  |
| Secondary/Tertiary     | 5178 (17.6)   | 0.93 (0.68–1.27)  |                  | 1.18 (0.83–1.67)     |                  |
| Wealth quintile        |               |                   |                  |                      |                  |
| Poorest                | 8345 (28.3)   | Ref               | <b>0.012</b>     | Ref                  | <b>0.035</b>     |
| Poorer                 | 6802 (23.1)   | 1.27 (0.97–1.66)  |                  | 1.25 (0.96–1.64)     |                  |
| Poor                   | 5500 (18.7)   | 1.35 (1.02–1.79)  |                  | 1.36 (1.03–1.81)     |                  |
| Less poor              | 5584 (19)     | 0.97 (0.72–1.32)  |                  | 0.99 (0.72–1.36)     |                  |
| Least poor             | 3221 (10.9)   | 0.73 (0.49–1.10)  |                  | 0.73 (0.44–1.20)     |                  |
| Age at measurement     |               |                   |                  |                      |                  |
| Under 1 year           | 16,240 (55.1) | 4.12 (2.58–6.56)  | <b>&lt;0.001</b> | 4.05 (2.52–6.51)     | <b>&lt;0.001</b> |
| 1–2 years              | 8846 (30.0)   | 2.67 (1.63–4.36)  |                  | 2.64 (1.61–4.33)     |                  |
| 2–3 years              | 4366 (14.8)   | Ref               |                  | Ref                  |                  |
| Maternal age           |               |                   |                  |                      |                  |
| <20 years              | 3291 (11.2)   | Ref               | 0.11             | Ref                  | 0.068            |
| 20–34                  | 20,075 (68.2) | 1.30 (0.91–1.85)  |                  | 1.40 (0.98–2.00)     |                  |
| 35–49                  | 6086 (20.7)   | 1.51 (1.02–2.22)  |                  | 1.62 (1.08–2.43)     |                  |
| Marital status         |               |                   |                  |                      |                  |
| Married                | 28,317 (96.1) | Ref               | 0.269            | Ref                  | 0.449            |
| Not married            | 1135 (3.9)    | 1.29 (0.82–2.03)  |                  | 1.20 (0.75–1.90)     |                  |
| Residence              |               |                   |                  |                      |                  |
| Urban                  | 3740 (12.7)   | Ref               | 0.243            | Ref                  | 0.270            |
| Rural                  | 25,711 (87.3) | 1.21 (0.88–1.65)  |                  | 0.81 (0.55–1.18)     |                  |
| Ethno-linguistic group |               |                   |                  |                      |                  |
| Kasem                  | 14,427 (49)   | Ref               | <b>0.007</b>     | Ref                  | <b>0.025</b>     |

(Continues)



TABLE 2 (Continued)

| Variable | Number (%)    | Crude OR (95% CI) | p-value | Adjusted OR (95% CI) | p-value |
|----------|---------------|-------------------|---------|----------------------|---------|
| Nankam   | 13,686 (46.5) | 1.38 (1.13–1.68)  |         | 1.36 (1.10–1.69)     |         |
| Buli     | 608 (2.1)     | 1.55 (0.84–2.86)  |         | 1.55 (0.83–2.88)     |         |
| Other    | 731 (2.5)     | 0.69 (0.31–1.57)  |         | 0.84 (0.36–1.96)     |         |

Abbreviations: CI, confidence intervals; JSS, junior high school and OR, odds ratios.

power and hence access to food during the lean season. This may thus maintain a good or healthy dietary diversity across the seasons.

We further observed that maternal education was not associated with SAM. This has previously been observed in a study from Bangladesh, where in addition to maternal education, variations in household income did not influence levels of SAM (Rahman et al., 2016). The importance of the influence of maternal education with malnutrition is important because education is known to influence maternal food choices, practices of good sanitary conditions and prevention of infection. Though access to formal education to the girl child is still a burning issue in some countries in SSA, our findings may result from the absence of indirect factors that mediate the relation of education to SAM.

We observed female children were likely to be severely malnourished during the hunger season. Consistent with our findings was findings from a study from Nepal that previously reported that male children were less likely to have SAM compared with female children (Hossain et al., 2020). This was further observed by a study in Tanzania (Rogawski McQuade et al., 2019) but contrary to findings from 16 demographic health surveys reports from SSA (Wamani et al., 2007). A further systematic review also reported boys are more vulnerable to malnutrition and stunting compared with girls (Thurstans et al., 2020). These inconsistent findings may suggest the varying reasons currently provided for these differences are more speculative rather than informed by direct evidence. Further longitudinal studies may be required to explain the observed findings.

Overall our findings have important implications for policies and program implementers in the study area. These results suggest that children, particularly girls, born in the hunger season are vulnerable to SAM and hence poor health outcomes. The implications may extend into adulthood as severe malnutrition and poor growth have been associated with reduced cognitive function and risk of chronic diseases later in life (Liu et al., 2003). Prenatal care interventions, including nutritional supplementation, could be targeted during high food insecurity months to reduce seasonal disparities. Implementing initiatives to promote food market development and improve economic prospects especially during the hunger season are likely to help prevent seasonal declines in nutritional status. Further studies are also needed to elucidate the possible reasons for sex differences in nutritional status.

## 4.1 | Strengths and limitation

Our study does have several strengths including a large representative sample size which makes our results applicable to the HDSS study area. Our study also highlights the need to tailor interventions to the most vulnerable periods in the year. We would like to highlight a few limitations of our study. Assessing the effect of seasonal variability severe acute malnutrition poses methodological challenges. These include exposure assessment, ecological fallacies and the complexity of relationships. Ecological fallacy in our study may be minimized by the minimal local variations in seasons. However, residual confounding may still persist due to some unmeasured factors that may influence the findings of this study, and hence interpretations should be done with caution.

The use of crude MUAC measurements unadjusted for age may lead to misclassification of some participants and may also affect the rates of malnutrition reported. The World Health Organization growth and development classifications (Z-scores/norms) are usually criticized for their inappropriateness in some SSA contexts due to potential misclassifications. There are currently conflicting reports with several studies reporting agreement between waist-for-height (WHZ) scores and MUAC while other studies have reported disparities (Mehta et al., 2013). This implies the scores perform differently in different populations. However, MUAC measurements have been validated and found to be viable, low-cost, low-burden alternative for community-level nutritional status assessment.

## 5 | CONCLUSION

We established that being born in the hunger season is associated with a higher risk of SAM in this Guinea-Saharan ecological zone. The result implies improvement in the food supply to pregnant and lactating mothers through sustainable agriculture or food system change targeting the hunger may reduce the burden of SAM. Further work is also needed to understand the mechanisms underlying these observed associations, especially the role of the pregnancy/lactation period so as to guide the development of appropriate interventions.

## ACKNOWLEDGMENTS

We are grateful to the chiefs, elders and people from communities within the catchment of the Navrongo sociodemographic



surveillance site. We acknowledge all the field workers and researchers of the Navrongo Health Research Centre who did due diligence with data collection, entry, quality control and extraction. We appreciate the support of all INPreP study team members from NHRC including: Edith Dambayi, Esmond Nonterah, Doreen Ayi-Bisah, James Adoctor, Edna Kukuri and Paula Beeri. This work was supported by the National Institute for Health Research (NIHR) under grant 17\63\154 using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care. KMG is supported by the UK Medical Research Council (MC\_UU\_12011/4), the National Institute for Health Research (NIHR Senior Investigator (NF-SI-0515-10042), NIHR Southampton 1000DaysPlus Global Nutrition Research Group (17/63/154) and NIHR Southampton Biomedical Research Centre (IS-BRC-1215-20004)), the European Union (Erasmus+ Programme ImpENSA 598488-EPP-1-2018-1-DE-EPPKA2-CBHE-JP), the British Heart Foundation (RG/15/17/3174), and US National Institute On Aging of the National Institutes of Health (Award No. U24AG047867).

#### CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

#### ETHICS STATEMENT

Ethical approval for the operations of the Navrongo HDSS was obtained from the Navrongo Health Research Centre's Institutional Review Board (NHRC-IRB). An additional ethics approval for the analyses of the data through the INPreP project was obtained from the NHRC-IRB (approval number NHRCIRB325).

#### AUTHOR CONTRIBUTIONS

EAN, KMG, and M-LN designed the INPreP study. EAN and PW conceived the paper. PW analyzed the data and EAN interpreted the results and wrote the first draft. PW, STC, SHK, KAW, KMG, and M-LN contributed to writing the paper by commenting on successive drafts. ARO is head of the Navrongo Health and Socio-Demographic Surveillance System and WO is head of Upper East Regional Health Directorate and both are responsible for the data. All authors critically reviewed the paper and approved the final draft for submission.

#### DATA AVAILABILITY STATEMENT

Data used in this paper are available upon reasonable request from the corresponding author. The Navrongo Health and Demographic Surveillance System data can be obtained through a request to the Navrongo Health Research Centre's director (<https://www.navrongo-hrc.org>).

#### ORCID

Engelbert A. Nonterah  <http://orcid.org/0000-0001-8491-6478>

Sarah H. Kehoe  <http://orcid.org/0000-0003-2584-7999>

Keith M. Godfrey  <http://orcid.org/0000-0002-4643-0618>

#### REFERENCES

- Abizari, A. R., Azupogo, F., Nagasu, M., Creemers, N., & Brouwer, I. D. (2017). Seasonality affects dietary diversity of school-age children in northern Ghana. *PLoS One*, *12*(8), e0183206.
- Ahmad, D., Afzal, M., & Imtiaz, A. (2020). Effect of socioeconomic factors on malnutrition among children in Pakistan. *Future Business Journal*, *6*(1), 80.
- Akparibo, R., Harris, J., Blank, L., Campbell, M. J., & Holdsworth, M. (2017). Severe acute malnutrition in children aged under 5 years can be successfully managed in a non-emergency routine community healthcare setting in Ghana. *Maternal & Child Nutrition*, *13*(4), e12417.
- Bain, L. E., Awah, P. K., Geraldine, N., Kindong, N. P., Sigal, Y., Bernard, N., et al. (2013). Malnutrition in Sub-Saharan Africa: burden, causes and prospects. *The Pan African Medical Journal*, *15*, 120.
- Belayneh, M., Loha, E., & Lindtjorn, B. (2020). Seasonal variation of household food insecurity and household dietary diversity on wasting and stunting among young children in a drought prone area in South Ethiopia: A cohort study. *Ecology of Food and Nutrition*, *1*–26.
- Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., de Onis, M., Ezzati, M., et al. (2008). Maternal and child undernutrition: Global and regional exposures and health consequences. *The Lancet*, *371*(9608), 243–260.
- Bryson, J. M., Patterson, K., Berrang-Ford, L., Lwasa, S., Namanya, D. B., & Twesigomwe, S. (2021). Seasonality, climate change, and food security during pregnancy among indigenous and non-indigenous women in rural Uganda: Implications for maternal-infant health. *PLoS One*, *16*(3), e0247198
- Chikhungu, L. C., & Madise, N. J. (2014). Seasonal variation of child under nutrition in Malawi: Is seasonal food availability an important factor? Findings from a national level cross-sectional study. *BMC Public Health*, *14*, 1146.
- Choudhary, T. S., Srivastava, A., Chowdhury, R., Taneja, S., Bahl, R., Martinez, J., Bhan, M. K., & Bhandari, N. (2019). Severe wasting among Indian infants <6 months: Findings from the National Family Health Survey 4. *Maternal & Child Nutrition*, *15*, e12866.
- Davies, J. I., Macnab, A. J., Byass, P., Norris, S. A., Nyirenda, M., Singhal, A., Sobngwi, E., & Daar, A. S. (2018). Developmental origins of health and disease in Africa—influencing early life. *The Lancet Global Health*, *6*(3), e244–e245.
- Debpuur, C., Nonterah, E. A., Chatio, S. T., Adoctor, J. K., Dambayi, E., Beeri, P., Nonterah, E. W., Ayi-Bisah, D., Watson, D., Kehoe, S. H., Dalaba, M. A., Ofosu, W., Aborigo, R., Welaga, P., Oduro, A. R., Newell, M. L., Barker, M., & INPreP Study, G. (2020). Supporting maternal and child nutrition: Views from community members in rural Northern Ghana. *Public Health Nutrition*, *24*, 1–8.
- Doku, D. T., & Neupane, S. (2015). Double burden of malnutrition: increasing overweight and obesity and stall underweight trends among Ghanaian women. *BMC Public Health*, *15*, 670.
- Fagariba, C. J., Song, S., & Baoro, S. K. G. S. (2018). Climate change in Upper East Region of Ghana; challenges existing in farming practices and new mitigation policies. *Open. Agriculture (London)*, *3*(1), 524–536.
- Fentahun, N., Belachew, T., Coates, J., & Lachat, C. (2018). Seasonality and determinants of child growth velocity and growth deficit in rural southwest Ethiopia. *BMC Pediatrics*, *18*(1), 20.
- Ghana Statistical Service Ghana Health Service & ICF International. (2014). Ghana demographic and health survey. GSS, GHS and ICF International. <https://dhsprogram.com/pubs/pdf/FR307/FR307.pdf>
- Hagos, S., Lunde, T., Mariam, D. H., Woldehanna, T., & Lindtjorn, B. (2014). Climate change, crop production and child under nutrition in Ethiopia: A longitudinal panel study. *BMC Public Health*, *14*, 14.

- Hillbruner, C., & Egan, R. (2008). Seasonality, household food security, and nutritional status in Dinajpur, Bangladesh. *Food and Nutrition Bulletin*, 29(3), 221–231.
- Hossain, A., Niroula, B., Duwal, S., Ahmed, S., & Kibria, M. G. (2020). Maternal profiles and social determinants of severe acute malnutrition among children under-five years of age: A case-control study in Nepal. *Heliyon*, 6(5), e03849.
- Imam, A., Hassan-Hanga, F., Sallahdeen, A., & Farouk, Z. L. (2020). Socio-demographic and household-level risk factors for severe acute malnutrition in pre-school children in north-western Nigeria. *Journal of Tropical Pediatrics*, 66(6), 589–97.
- Jones, K. D., & Berkley, J. A. (2014). Severe acute malnutrition and infection. *Paediatr Int Child Health*, 34, Suppl 1, S1–S29.
- Liu, J., Raine, A., Venables, P. H., Dalais, C., & Mednick, S. A. (2003). Malnutrition at age 3 years and lower cognitive ability at age 11 years: Independence from psychosocial adversity. *Archives of Pediatrics and Adolescent Medicine*, 157, 593–600.
- Madan, E. M., Haas, J. D., Menon, P., & Gillespie, S. (2018). Seasonal variation in the proximal determinants of undernutrition during the first 1000 days of life in rural South Asia: A comprehensive review. *Global Food Security*, 19, 11–23.
- Mehta, N. M., Corkins, M. R., Lyman, B., Malone, A., Goday, P. S., Carney, L. N., et al. (2013). Defining pediatric malnutrition: a paradigm shift toward etiology-related definitions. *JPEN. Journal of Parenteral and Enteral Nutrition*, 37(4), 460–481.
- Morrison, J. L., & Regnault, T. R. (2016). Nutrition in pregnancy: Optimising maternal diet and fetal adaptations to altered nutrient supply. *Nutrients*, 8(6), 342.
- Oduro, A. R., Wak, G., Azongo, D., Debpuur, C., Wontuo, P., Kondayire, F., et al. (2012). Profile of the Navrongo health and demographic surveillance system. *International Journal of Epidemiology*, 41(4), 968–976.
- Rahman, M. S., Howlader, T., Masud, M. S., & Rahman, M. L. (2016). Association of low-birth weight with malnutrition in children under five years in Bangladesh: Do mother's education, socio-economic status, and birth interval matter? *PLoS One*, 11(6), e0157814.
- Roba, K. T., O'Connor, T. P., Belachew, T., & O'Brien, N. M. (2016). Variations between post- and pre-harvest seasons in stunting, wasting, and Infant and Young Child Feeding (IYCF) practices among children 6–23 months of age in lowland and midland agro-ecological zones of rural Ethiopia. *The Pan African Medical Journal*, 24, 163.
- Rogawski McQuade, E. T., Clark, S., Bayo, E., Scharf, R. J., DeBoer, M. D., Patil, C. L., et al. (2019). Seasonal food insecurity in Haydom, Tanzania, is associated with low birthweight and acute malnutrition: Results from the MAL-ED Study. *Am J Trop Med Hyg*, 100(3), 681–687.
- Saaka, M., Larbi, A., Hoeschle-Zeledon, I., & Appiah, B., Child malnutrition in Northern Ghana. (2015). Evidence, factors and recommendations from a new study. IITA report. IITA, Ibadan, Nigeria. 22.
- Sorgho, R., Franke, J., Simboro, S., Phalkey, R., & Saeuborn, R. (2016). NUTRition and CLIMate (NUTRICLIM): Investigating the relationship between climate variables and childhood malnutrition through agriculture, an exploratory study in Burkina Faso. *Public Health Reviews*, 37, 16.
- Tette, E. M. A., Sifah, E. K., Nartey, E. T., Nuro-Ameyaw, P., Tete-Donkor, P., & Biritwum, R. B. (2016). Maternal profiles and social determinants of malnutrition and the MDGs: What have we learnt? *BMC Public Health*, 16, 214.
- Thompson, D. S., Younger-Coleman, N., Lyew-Ayee, P., Greene, L. G., Boyne, M. S., & Forrester, T. E. (2017). Socioeconomic factors associated with severe acute malnutrition in Jamaica. *PLoS One*, 12(3), e0173101.
- Thurstans, S., Opondo, C., Seal, A., Wells, J., Khara, T., Dolan, C., et al. (2020). Boys are more likely to be undernourished than girls: a systematic review and meta-analysis of sex differences in undernutrition. *BMJ Global Health*, 5(12), e004030.
- Tomkins, A. M., Dunn, D. T., Hayes, R. J., & Bradley, A. K. (1986). Seasonal variations in the nutritional status of urban Gambian children. *British Journal of Nutrition*, 56(3), 533–543.
- UNICEF World Health Organization & World Bank. (2020). UNICEF-WHO-the world bank: Joint child malnutrition estimates. New York/Geneva/Washington: UNICEF, WHO, and World Bank.
- Wamani, H., Astrom, A. N., Peterson, S., Tumwine, J. K., & Tylleskar, T. (2007). Boys are more stunted than girls in sub-Saharan Africa: A meta-analysis of 16 demographic and health surveys. *BMC Pediatrics*, 7, 17.
- Watson, D., Kehoe, S. H., Erzse, A., Compaoré, A., Debpuur, C., Nonterah, E. A., et al. (2020). Community perspectives on maternal and child health during nutrition and economic transition in sub-Saharan Africa. *Public Health Nutrition*, 1–9.
- WHO. (2019). International Classification of Diseases, 11th Revision (ICD-11). <https://www.who.int/classifications/icd/en/>
- World Health Organization and UNICEF. (2009). WHO child growth standards and the identification of severe acute malnutrition in infants and children. A Joint Statement by the World Health Organization and the United Nations Children's Fund. WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
- Yawson, A. E., Amoafu, E. O., Senaya, L. K., Yawson, A. O., Aboagye, P. K., & Mahama, A. B., et al (2017). The Lancet series nutritional interventions in Ghana: A determinants analysis approach to inform nutrition strategic planning. *BMC. Nutrition*, 3(1), 1–8.

**How to cite this article:** Nonterah, E. A., Welaga, P., Chatio, S. T., Kehoe, S. H., Ofori, W., Ward, K. A., Godfrey, K. M., Oduro, A. R., Newell, M.-L., & as members of the INPreP study group (2022). Children born during the hunger season are at a higher risk of severe acute malnutrition: Findings from a Guinea Sahelian ecological zone in Northern Ghana. *Maternal & Child Nutrition*, 18, e13313.

<https://doi.org/10.1111/mcn.13313>