

1-(2-Methylbenzoyl)-3-*m*-tolylthioureaB. M. Yamin,^a S. Yousuf,^{b*} M. S. M. Yusof^c and T. N. D. T. Zakaria^c

^aSchool of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43500 Bangi Selangor, Malaysia, ^bHEJ Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan, and ^cDepartment of Chemistry, Universiti Malaysia Terengganu, Mannagabang Telipot, Terengganu, Malaysia
Correspondence e-mail: sammer_yousuf@yahoo.com


Received 10 April 2008; accepted 28 April 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.043; wR factor = 0.125; data-to-parameter ratio = 14.9.

The molecule of the title compound, $\text{C}_{16}\text{H}_{16}\text{N}_2\text{OS}$, is not planar; the two aromatic rings are inclined to one another by 37.59 (9)°. There are intramolecular hydrogen bonds between the benzoyl O atom and the H atom of the thioamide N atom, and between the thiourea S atom and the H atom of the tolyl group. These hydrogen bonds stabilize the molecule in such a way that the thiourea group adopts a *trans-cis* geometry. In the crystal structure, molecules are linked by $\text{N}-\text{H}\cdots\text{S}$ intermolecular hydrogen bonds, forming centrosymmetric dimers.

Related literature

For the crystal structure of 1-(2,3-dimethylphenyl)-3-(2-methylbenzoyl)thiourea, see: Khawar Rauf *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{16}\text{N}_2\text{OS}$
 $M_r = 284.37$
 Triclinic, $P\bar{1}$
 $a = 6.440$ (3) Å
 $b = 10.201$ (5) Å
 $c = 11.415$ (5) Å
 $\alpha = 77.310$ (7)°
 $\beta = 89.896$ (8)°
 $\gamma = 86.468$ (8)°
 $V = 730.1$ (6) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.22$ mm⁻¹
 $T = 298$ (2) K
 $0.35 \times 0.34 \times 0.22$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\min} = 0.927$, $T_{\max} = 0.954$
 7240 measured reflections
 2703 independent reflections
 2202 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.125$
 $S = 1.02$
 2703 reflections
 181 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{S1}^i$	0.86	2.74	3.407 (2)	136
$\text{N2}-\text{H2A}\cdots\text{O1}$	0.86	1.97	2.658 (2)	136
$\text{C7}-\text{H7C}\cdots\text{O1}$	0.96	2.52	2.933 (3)	106
$\text{C15}-\text{H15A}\cdots\text{S1}$	0.93	2.54	3.168 (3)	125

Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL, PARST (Nardelli, 1995) and PLATON (Spek, 2003).

The authors thank the Ministry of Higher Education of Malaysia for the Fundamental Research Grants UKM-OUP-BTT-28-2007 and UMT-FRGS-59001, and University Kebangsaan Malaysia, University Malaysia Terengganu and the HEJ Research Institute of Chemistry, University of Karachi, for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2052).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bruker (2000). SADABS, SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Khawar Rauf, M., Badshah, A. & Bolte, M. (2007). *Acta Cryst. E* **63**, o1256–o1257.
 Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o1227 [doi:10.1107/S1600536808012300]

1-(2-Methylbenzoyl)-3-*m*-tolylthiourea

B. M. Yamin, S. Yousuf, M. S. M. Yusof and T. N. D. T. Zakaria

Comment

The title compound, (I), is analogous to 1-(2,3-Dimethylphenyl)-3-(2-methylbenzoyl)thiourea (II) (Khawar Rauf *et al.*, 2007), but with the 2,3-dimethyl phenyl group replaced by a 2-methyl phenyl (*m*-tolyl) group (Fig. 1). The molecule maintains the *trans-cis* configuration with respect to the position of the methyl benzoyl and 3-methyl benzene groups, respectively, relative to the thiono S1 atom. The bond lengths and angles are in normal ranges (Allen *et al.*, 1987). The central thiourea moiety, S1/N1/N2/C9, the 2-methylbenzoyl ring, (C1—C8), and the *m*-tolyl group (C10—C15,C16) are all relatively planar, with a maximum deviation from any best mean plane of 0.015 (2) Å for atom C10. The central thiourea moiety makes dihedral angles with the 2-methylbenzoyl and *m*-tolyl fragments of 49.61 (7) and 17.87 (9)°, respectively. The *trans-cis* geometry of the thiourea moiety is stabilized by N2—H2A···O1, C7—H7C···O1 and C15—H15A···S1 intramolecular hydrogen bonds (Table 1).

In the crystal structure of (I), symmetry related molecules are linked by the N1—H1A···S1 intermolecular hydrogen bonds (Table 1) to form centrosymmetric dimers (Fig 2).

Experimental

2-methylbenzoyl chloride (9.720 g, 0.025mole) was mixed with an equimolar amount of ammonium thiocyanate (1.903 g, 0.025 mol) and 3-methyl aniline (2.701 g, 0.025 mol) in 45 ml dry acetone. The mixture was refluxed with stirring for 4 h. The solution was then filtered and left to evaporate at room temperature. Colourless crystals, suitable for X-ray analysis, were obtained after a few days (Yield 85%).

Refinement

NH and C-bound H atoms were positioned geometrically and constrained to ride on their parent atoms: N—H = 0.86 and C—H = 0.93 - 0.96 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{CH and NH})$, and $1.5U_{\text{eq}}(\text{CH}_3)$.

Figures




Fig. 1. The molecular structure of compound (1), with displacement ellipsoids drawn at 50% probability level (The dashed lines indicate the intramolecular hydrogen bonds).




Fig. 2. The crystal packing diagram of compound (1), showing the formation of the N-H...S hydrogen bonded dimers (Hydrogen bonds are shown by dashed lines).

1-(2-Methylbenzoyl)-3-*m*-tolylthiourea

Crystal data

$C_{16}H_{16}N_2OS$

$M_r = 284.37$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.440$ (3) Å

$b = 10.201$ (5) Å

$c = 11.415$ (5) Å

$\alpha = 77.310$ (7)°

$\beta = 89.896$ (8)°

$\gamma = 86.468$ (8)°

$V = 730.1$ (6) Å³

$Z = 2$

$F_{000} = 300$

$D_x = 1.293$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 2987 reflections

$\theta = 1.8$ – 25.5 °

$\mu = 0.22$ mm⁻¹

$T = 298$ (2) K

Block, colorless

$0.35 \times 0.34 \times 0.22$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 83.66 pixels mm⁻¹

$T = 298$ (2) K

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2000)

$T_{\min} = 0.927$, $T_{\max} = 0.954$

7240 measured reflections

2703 independent reflections

2202 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.023$

$\theta_{\max} = 25.5$ °

$\theta_{\min} = 1.8$ °

$h = -7 \rightarrow 7$

$k = -12 \rightarrow 12$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.125$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.077P)^2 + 0.124P]$

$S = 1.02$	where $P = (F_o^2 + 2F_c^2)/3$
2703 reflections	$(\Delta/\sigma)_{\max} < 0.000$
181 parameters	$\Delta\rho_{\max} = 0.23 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.32952 (9)	0.66913 (5)	0.02660 (4)	0.0657 (2)
O1	0.5770 (2)	0.43595 (14)	0.39009 (11)	0.0589 (4)
N1	0.5747 (2)	0.51255 (14)	0.18710 (13)	0.0474 (5)
N2	0.3091 (2)	0.62497 (14)	0.26753 (12)	0.0463 (4)
C1	0.9854 (3)	0.3839 (2)	0.19650 (18)	0.0549 (6)
C2	1.1512 (3)	0.2991 (2)	0.1779 (2)	0.0663 (8)
C3	1.1594 (3)	0.1659 (2)	0.2386 (2)	0.0708 (8)
C4	1.0059 (3)	0.1190 (2)	0.3173 (2)	0.0632 (7)
C5	0.8385 (3)	0.20259 (18)	0.33940 (17)	0.0502 (6)
C6	0.8294 (2)	0.33688 (17)	0.27600 (15)	0.0450 (5)
C7	0.6744 (3)	0.1451 (2)	0.4262 (2)	0.0664 (7)
C8	0.6511 (3)	0.43112 (17)	0.29297 (15)	0.0445 (5)
C9	0.4011 (3)	0.60340 (16)	0.16802 (15)	0.0445 (5)
C10	0.1164 (2)	0.69333 (16)	0.28323 (15)	0.0421 (5)
C11	0.0350 (3)	0.65998 (18)	0.39738 (16)	0.0508 (6)
C12	-0.1571 (3)	0.7154 (2)	0.42026 (19)	0.0611 (7)
C13	-0.2680 (3)	0.80384 (19)	0.33001 (19)	0.0568 (7)
C14	-0.1870 (3)	0.84061 (19)	0.21718 (18)	0.0560 (6)
C15	0.0074 (3)	0.78556 (19)	0.19379 (16)	0.0551 (6)
C16	-0.3049 (4)	0.9400 (3)	0.1191 (2)	0.0903 (10)
H1A	0.64410	0.50630	0.12390	0.0570*
H1B	0.97800	0.47350	0.15540	0.0660*
H2A	0.37820	0.59190	0.33290	0.0560*
H2B	1.25610	0.33120	0.12520	0.0800*
H3A	1.26980	0.10770	0.22600	0.0850*
H4A	1.01400	0.02880	0.35700	0.0760*
H7A	0.71130	0.05180	0.45960	0.1000*

supplementary materials

H7B	0.54280	0.15370	0.38500	0.1000*
H7C	0.66430	0.19330	0.48970	0.1000*
H11A	0.10970	0.60040	0.45840	0.0610*
H12A	-0.21230	0.69310	0.49690	0.0730*
H13A	-0.39920	0.83900	0.34580	0.0680*
H15A	0.06450	0.81070	0.11790	0.0660*
H16A	-0.43520	0.96820	0.14950	0.1350*
H16B	-0.33020	0.89820	0.05330	0.1350*
H16C	-0.22440	1.01680	0.09180	0.1350*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0808 (4)	0.0649 (3)	0.0427 (3)	0.0316 (3)	0.0035 (2)	-0.0031 (2)
O1	0.0592 (8)	0.0699 (8)	0.0445 (7)	0.0245 (6)	-0.0033 (6)	-0.0134 (6)
N1	0.0440 (8)	0.0508 (8)	0.0439 (8)	0.0123 (6)	0.0053 (6)	-0.0071 (6)
N2	0.0435 (8)	0.0519 (8)	0.0413 (7)	0.0144 (6)	-0.0033 (6)	-0.0100 (6)
C1	0.0418 (10)	0.0589 (11)	0.0643 (12)	0.0012 (8)	0.0004 (8)	-0.0155 (9)
C2	0.0393 (10)	0.0854 (15)	0.0775 (14)	0.0030 (10)	0.0073 (9)	-0.0269 (12)
C3	0.0463 (11)	0.0783 (15)	0.0936 (16)	0.0214 (10)	-0.0050 (11)	-0.0382 (13)
C4	0.0573 (12)	0.0524 (11)	0.0802 (14)	0.0146 (9)	-0.0122 (10)	-0.0202 (10)
C5	0.0450 (9)	0.0498 (10)	0.0570 (10)	0.0062 (8)	-0.0082 (8)	-0.0165 (8)
C6	0.0366 (9)	0.0512 (9)	0.0484 (9)	0.0063 (7)	-0.0059 (7)	-0.0159 (8)
C7	0.0646 (13)	0.0550 (11)	0.0743 (14)	0.0019 (10)	0.0050 (11)	-0.0044 (10)
C8	0.0399 (9)	0.0463 (9)	0.0469 (9)	0.0048 (7)	-0.0028 (7)	-0.0116 (7)
C9	0.0420 (9)	0.0400 (8)	0.0494 (10)	0.0053 (7)	0.0013 (7)	-0.0078 (7)
C10	0.0406 (9)	0.0409 (8)	0.0456 (9)	0.0055 (7)	-0.0024 (7)	-0.0134 (7)
C11	0.0534 (10)	0.0515 (10)	0.0448 (9)	0.0085 (8)	0.0006 (8)	-0.0082 (7)
C12	0.0589 (12)	0.0628 (12)	0.0598 (12)	0.0062 (9)	0.0156 (9)	-0.0122 (9)
C13	0.0421 (10)	0.0567 (11)	0.0730 (13)	0.0097 (8)	0.0028 (9)	-0.0211 (9)
C14	0.0534 (11)	0.0524 (10)	0.0614 (12)	0.0180 (8)	-0.0097 (9)	-0.0163 (9)
C15	0.0590 (11)	0.0556 (10)	0.0462 (10)	0.0176 (9)	0.0015 (8)	-0.0071 (8)
C16	0.0882 (17)	0.0929 (17)	0.0776 (16)	0.0491 (14)	-0.0173 (13)	-0.0073 (13)

Geometric parameters (\AA , $^\circ$)

S1—C9	1.6605 (19)	C12—C13	1.379 (3)
O1—C8	1.216 (2)	C13—C14	1.372 (3)
N1—C8	1.380 (2)	C14—C16	1.506 (3)
N1—C9	1.393 (2)	C14—C15	1.390 (3)
N2—C9	1.335 (2)	C1—H1B	0.9300
N2—C10	1.416 (2)	C2—H2B	0.9300
N1—H1A	0.8600	C3—H3A	0.9300
N2—H2A	0.8600	C4—H4A	0.9300
C1—C6	1.388 (3)	C7—H7A	0.9600
C1—C2	1.378 (3)	C7—H7B	0.9600
C2—C3	1.381 (3)	C7—H7C	0.9600
C3—C4	1.371 (3)	C11—H11A	0.9300
C4—C5	1.390 (3)	C12—H12A	0.9300

C5—C6	1.400 (3)	C13—H13A	0.9300
C5—C7	1.503 (3)	C15—H15A	0.9300
C6—C8	1.492 (3)	C16—H16A	0.9600
C10—C15	1.386 (3)	C16—H16B	0.9600
C10—C11	1.383 (3)	C16—H16C	0.9600
C11—C12	1.377 (3)		
C8—N1—C9	129.20 (15)	C13—C14—C15	119.23 (18)
C9—N2—C10	130.59 (14)	C10—C15—C14	120.21 (17)
C8—N1—H1A	115.00	C2—C1—H1B	120.00
C9—N1—H1A	115.00	C6—C1—H1B	120.00
C10—N2—H2A	115.00	C1—C2—H2B	120.00
C9—N2—H2A	115.00	C3—C2—H2B	120.00
C2—C1—C6	120.68 (19)	C2—C3—H3A	120.00
C1—C2—C3	119.10 (19)	C4—C3—H3A	120.00
C2—C3—C4	120.45 (19)	C3—C4—H4A	119.00
C3—C4—C5	121.78 (19)	C5—C4—H4A	119.00
C4—C5—C6	117.43 (17)	C5—C7—H7A	110.00
C4—C5—C7	119.31 (17)	C5—C7—H7B	110.00
C6—C5—C7	123.23 (17)	C5—C7—H7C	109.00
C1—C6—C8	119.24 (16)	H7A—C7—H7B	110.00
C5—C6—C8	120.22 (15)	H7A—C7—H7C	109.00
C1—C6—C5	120.54 (16)	H7B—C7—H7C	109.00
O1—C8—N1	122.48 (17)	C10—C11—H11A	120.00
O1—C8—C6	123.92 (16)	C12—C11—H11A	120.00
N1—C8—C6	113.60 (14)	C11—C12—H12A	120.00
N1—C9—N2	115.10 (15)	C13—C12—H12A	120.00
S1—C9—N1	117.23 (13)	C12—C13—H13A	120.00
S1—C9—N2	127.65 (14)	C14—C13—H13A	120.00
N2—C10—C15	124.98 (15)	C10—C15—H15A	120.00
C11—C10—C15	119.83 (15)	C14—C15—H15A	120.00
N2—C10—C11	115.18 (15)	C14—C16—H16A	109.00
C10—C11—C12	119.69 (17)	C14—C16—H16B	109.00
C11—C12—C13	120.27 (19)	C14—C16—H16C	109.00
C12—C13—C14	120.72 (18)	H16A—C16—H16B	109.00
C13—C14—C16	120.97 (18)	H16A—C16—H16C	110.00
C15—C14—C16	119.80 (18)	H16B—C16—H16C	109.00
C9—N1—C8—O1	-5.0 (3)	C7—C5—C6—C1	-179.65 (18)
C9—N1—C8—C6	174.26 (16)	C7—C5—C6—C8	0.2 (3)
C8—N1—C9—S1	-170.78 (15)	C1—C6—C8—O1	-137.6 (2)
C8—N1—C9—N2	7.9 (3)	C1—C6—C8—N1	43.1 (2)
C10—N2—C9—S1	8.9 (3)	C5—C6—C8—O1	42.6 (3)
C10—N2—C9—N1	-169.59 (16)	C5—C6—C8—N1	-136.71 (17)
C9—N2—C10—C11	159.19 (18)	N2—C10—C11—C12	-176.51 (17)
C9—N2—C10—C15	-19.4 (3)	C15—C10—C11—C12	2.1 (3)
C6—C1—C2—C3	0.6 (3)	N2—C10—C15—C14	175.98 (17)
C2—C1—C6—C5	0.5 (3)	C11—C10—C15—C14	-2.5 (3)
C2—C1—C6—C8	-179.37 (18)	C10—C11—C12—C13	-0.1 (3)
C1—C2—C3—C4	-0.7 (3)	C11—C12—C13—C14	-1.7 (3)

supplementary materials

C2—C3—C4—C5	-0.3 (3)	C12—C13—C14—C15	1.3 (3)
C3—C4—C5—C6	1.4 (3)	C12—C13—C14—C16	-178.4 (2)
C3—C4—C5—C7	179.67 (19)	C13—C14—C15—C10	0.8 (3)
C4—C5—C6—C1	-1.5 (3)	C16—C14—C15—C10	-179.6 (2)
C4—C5—C6—C8	178.38 (17)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1A...S1 ⁱ	0.86	2.74	3.407 (2)	136
N2—H2A...O1	0.86	1.97	2.658 (2)	136
C7—H7C...O1	0.96	2.52	2.933 (3)	106
C15—H15A...S1	0.93	2.54	3.168 (3)	125

Symmetry codes: (i) $-x+1, -y+1, -z$.

Fig. 1




Fig. 2

