

MEETING ABSTRACT

Open Access

# Zebrafish as model organism for cNMP research

Fanni Dittmar<sup>1\*</sup>, Salim Seyfried<sup>2</sup>, Volkhard Kaever<sup>3</sup>, Roland Seifert<sup>1</sup>

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications  
Trier, Germany. 19-21 June 2015

## Introduction

The zebrafish *Danio rerio* has become an important vertebrate model organism for a wide range of scientific research questions including development, genetics, and disease [1]. The zebrafish is particularly well suited for this research due to its small size, rapid development, short generation time, optical transparency of embryos and larvae as well as functional conservation of genes [1]. Furthermore, application of drugs is easy since zebrafish readily absorb compounds from their surrounding media [2].

The aim of our study was to determine the composition and to quantify the endogenous level of different cyclic nucleotides (cNMPs) in various developmental stages and organs of *Danio rerio*.

## Methods

Wild-type AB zebrafish were mated; for tissue preparation, we harvested embryos at 24 hours post fertilization (hpf) and larvae at 5 days post fertilization (dpf). Organs (eyes, brain, heart, entrails, eggs and testes) from adult female and male wild-type AB zebrafish were dissected according to the guidance of Gupta and Mullins [3]. All samples were then prepared for measurement of cNMP concentrations via a highly sensitive and specific method, namely high-performance liquid chromatography tandem mass spectrometry (HPLC-MS/MS) [4].

## Results

We could detect all studied cNMPs (namely cAMP, cCMP, cGMP, cIMP and cUMP) with specific variations as outlined below. (1) cGMP and cAMP could be detected in tissue samples of both developmental stages and within all harvested organs. Remarkably, they are the only cyclic nucleotides detected in brain and entrails.

The highest cGMP concentrations were present in eyes; the highest cAMP concentrations in larvae tissue. (2) cIMP was only detected in some organs with the highest concentration in heart. (3) cCMP could not be detected in embryos but in larvae and within several organs. Like cIMP, the highest cCMP concentrations were present in heart. (4) cUMP was detected in both developmental stages as well as in some organs. This cNMP was most abundant within the heart.

## Conclusion

The zebrafish *Danio rerio* constitutes a good model organism not only for studies focused on development, genetics and disease but also in the field of cyclic nucleotides, especially related to the role of cNMPs in cardiovascular biology. We observed specific cNMP patterns in development and in different organs, which is in support of the hypothesis of a distinct cNMP signaling code [5].

## Future studies

Zebrafish embryos will be treated with the NO-independent sGC activator cinaciguat (BAY 58-2667) and the NO-synergistic sGC stimulator 3-(4-amino-5-cyclopropylpyrimidine-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine (BAY 41-2272), from one day post fertilization to five days post fertilization. cNMP concentrations and morphological changes will be determined afterwards. Another research area will be the elucidation of the controversial role of cIMP as second messenger [6,7].

## Authors' details

<sup>1</sup>Institute of Pharmacology, Hannover Medical School (MHH), Hannover, Germany. <sup>2</sup>Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany. <sup>3</sup>Research Core Unit Metabolomics, Institute of Pharmacology, Hannover Medical School (MHH), Hannover, Germany.

Published: 2 September 2015

\* Correspondence: [dittmar.fanni@mh-hannover.de](mailto:dittmar.fanni@mh-hannover.de)

<sup>1</sup>Institute of Pharmacology, Hannover Medical School (MHH), Hannover, Germany

Full list of author information is available at the end of the article

**Table 1. cNMP concentrations of zebrafish embryos, larvae and various organs**

n=2	cGMP[pmol/mg protein]	cIMP[pmol/mg protein]	cAMP[pmol/mg protein]	cCMP[pmol/mg protein]	cUMP[pmol/mg protein]
<b>Embryos 24 hpf</b>	0.282 ± 0.009	not detected	2.29 ± 0.37	not detected	0.414 ± 0.067
<b>Larvae 5 dpf</b>	1.99 ± 0.32	not detected	23.66 ± 3.97	0.328 ± 0.119	0.890 ± 0.422
<b>Eyes</b>	2.69 ± 0.03	0.111 ± 0.000	2.73 ± 0.34	0.043 ± 0.004	0.087 ± 0.040
<b>Brain</b>	0.149 ± 0.073	not detected	10.02 ± 2.89	not detected	not detected
<b>Heart</b>	0.607 ± 0.124	0.452 ± 0.069	23.12 ± 17.40	0.504 ± 0.143	1.11 ± 0.10
<b>Entrails</b>	0.292 ± 0.031	not detected	16.52 ± 2.30	not detected	not detected
<b>Eggs</b>	0.002 ± 0.003	not detected	0.29 ± 0.02	0.004 ± 0.001	0.008 ± 0.001
<b>Testes</b>	0.802 ± 1.134	0.407 ± 0.037	12.15 ± 3.80	0.116 ± 0.164	0.617 ± 0.052

#### References

1. Lawrence C: **The husbandry of zebrafish (*Danio rerio*): A review.** *Aquaculture* 2007, **269**(1-4):1-20.
2. Langheinrich U: **Zebrafish: A new model on the pharmaceutical catwalk.** *BioEssays* 2003, **25**(9):904-912.
3. Gupta T, Mullins MC: **Dissection of organs from the adult zebrafish.** *J Vis Exp* 2010, **37**.
4. Bähre H, Danker KY, Stasch JP, Kaever V, Seifert R: **Nucleotidyl cyclase activity of soluble guanylyl cyclase in intact cells.** *Biochem Biophys Res Commun* 2014, **443**(4):1195-1199.
5. Seifert R, Schneider EH, Bähre H: **From canonical to non-canonical cyclic nucleotides as second messengers: Pharmacological implications.** *Pharmacol Ther* 2015, **148**:154-184.
6. Chen Z, Zhang X, Ying L, Dou D, Li Y, Bai Y, et al: **cIMP synthesized by sGC as a mediator of hypoxic contraction of coronary arteries.** *AJP Hear Circ Physiol* 2014, **307**(3):H328-H336.
7. Seifert R: **Is cIMP a second messenger with functions opposite to those of cGMP?** *Naunyn Schmiedebergs Arch Pharmacol* 2014, **387**(9):897-899.

doi:10.1186/2050-6511-16-S1-A45

Cite this article as: Dittmar et al.: Zebrafish as model organism for cNMP research. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A45.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

