



Signal Transduction for TNF α -Induced Type II SOCS Expression and Its Functional Implication in Growth Hormone Resistance in Carp Hepatocytes

OPEN ACCESS

Edited by:

Chun Peng,
York University, Canada

Reviewed by:

Wei Hu,
Institute of Hydrobiology (CAS), China
Elisabeth Eppler,
University of Zurich, Switzerland

*Correspondence:

Anderson O. L. Wong
olwong@hku.hk

† Present address:

Xue Jiang,
State Key Laboratory of
Pharmaceutical Biochemistry,
University of Hong Kong,
Hong Kong, China;
Joint Laboratory of Guangdong and
Hong Kong on Metabolic Diseases,
Guangdong Pharmaceutical
University, Guangdong, China

Specialty section:

This article was submitted to
Experimental Endocrinology,
a section of the journal
Frontiers in Endocrinology

Received: 28 November 2019

Accepted: 13 January 2020

Published: 30 January 2020

Citation:

Jiang X, He M, Bai J, Chan CB and
Wong AOL (2020) Signal Transduction
for TNF α -Induced Type II SOCS
Expression and Its Functional
Implication in Growth Hormone
Resistance in Carp Hepatocytes.
Front. Endocrinol. 11:20.
doi: 10.3389/fendo.2020.00020

Xue Jiang[†], Mulan He, Jin Bai, Chi Bun Chan and Anderson O. L. Wong*

School of Biological Sciences, The University of Hong Kong, Hong Kong, China

In mammals, local production of tumor necrosis factor α (TNF α) inhibits growth hormone (GH)-induced IGF-I expression at tissue level and contributes to GH resistance caused by sepsis/endotoxemia and inflammation. Although the loss of GH responsiveness can be mediated by a parallel rise in SOCS expression, the signaling mechanisms for TNF α -induced SOCS expression at the hepatic level have not been characterized and the comparative aspects of the phenomenon, especially in lower vertebrates, are still unknown. Recently, type II SOCS, including SOCS1-3 and CISH, have been cloned in grass carp and shown to act as the feedback repressors for GH signaling via JAK₂/STAT₅ pathway. To shed light on the mechanisms for TNF α -induced GH resistance in fish model, grass carp TNF α was cloned and confirmed to be a single-copy gene expressed in various tissues including the liver. In carp hepatocytes, incubation with the endotoxin LPS induced TNF α expression with parallel rises in SOCS1-3 and CISH mRNA levels. Similar to LPS, TNF α treatment could block GH-induced IGF-I/-II mRNA expression and elevate SOCS1, SOCS3, and CISH transcript levels. However, TNF α was not effective in altering SOCS2 expression. In parallel experiment, LPS blockade of IGF-I/-II signals caused by GH could be partially reverted by TNF α receptor antagonism. At hepatocyte level, TNF α induction also triggered rapid phosphorylation of I κ B α , MEK_{1/2}, ERK_{1/2}, MKK_{3/6}, P₃₈^{MAPK}, Akt, JAK₂, and STAT_{1,3,5}, and TNF α -induced SOCS1, SOCS3, and CISH mRNA expression could be negated by inhibiting the IKK/NF κ B, MAPK, PI3K/Akt, and JAK/STAT cascades. Our findings, as a whole, suggest that local production of TNF α may interfere with IGF-I/-II induction by GH in the carp liver by up-regulation of SOCS1, SOCS3, and CISH via IKK/NF κ B, MAPK, PI3K/Akt, and JAK/STAT-dependent mechanisms, which may contribute to GH resistance induced by endotoxin in carp species.

Keywords: TNF α , SOCS, CISH, GH, IGF, signal transduction, hepatocytes, grass carp

INTRODUCTION

Tumor necrosis factor α (TNF α) is a member of the proinflammatory cytokines and plays a key role in regulating immune cell activation/migration, cell proliferation/apoptosis, angiogenesis, and insulin resistance/growth impairment caused by chronic stress/infection (1). TNF α is also involved in diseases/ pathological conditions related to immune disorder, e.g., rheumatoid arthritis, ankylosing spondylitis, psoriasis, and inflammatory bowel disease (2). TNF α , together with other proinflammatory cytokines (e.g., IL-1 β and IL-6), can also act as local mediators for GH resistance induced by sepsis/endotoxemia or chronic inflammation (3–5). During the process, serum GH can be elevated with current drops in circulating IGF-I (6) and its production at tissue level (e.g., in muscle and liver) (4, 6). Meanwhile, a loss in hepatic responsiveness for GH-induced JAK₂/STAT₅ signaling and IGF-I gene expression has also been reported, e.g., in rat hepatocytes (7) and CWSV-1 hepatic cells (8, 9). GH resistance triggered by local production of cytokines can be attributed to a drop in GH receptor (GHR) expression via inhibition on GHR promoter activity (10–12) with parallel activation of SOCS expression (e.g., SOCS1, SOCS3, and CISH) (3, 12, 13), which can lead to failure in linear growth during childhood development (14) as well as muscle wasting, poor wound healing, cachexia, hepatic steatosis and insulin resistance in adult stage (14, 15). Although modulation of GH responsiveness by cytokines represents a major mechanism for the functional crosstalk of the immune system with somatotrophic axis (16), the studies in this area are restricted to mammalian models and the comparative aspects of the phenomenon, especially in lower vertebrates, are still unknown.

SOCS proteins are feedback inhibitors for cytokine signaling and their inhibitory effects are mediated through interference of the JAK/STAT pathway functionally coupled with cytokine receptors (17). They are widely expressed at the tissue level (18) and can be induced by microbial/viral infection (19, 20) via local production of cytokines (e.g., IL-4 and IFN γ) (21, 22). By limiting the duration/magnitude of cytokine signaling, SOCS expression can prevent hyper activation of immune responses and bring the system back to a homeostatic state (23), and dysregulation of SOCS can be associated with autoimmune diseases, chronic inflammation and carcinogenesis (24, 25). At present, at least 10 members of SOCS family, including SOCS1–9 and CISH, have been identified. They can be classified into type I (SOCS4–7 and SOCS5b–9) and type II subfamily (SOCS1–3 and CISH) with the type II SOCS closely related to the ancestral lineage found in invertebrates (26). Members of SOCS family all share a common structural organization with a SH2 domain in the central core followed by a SOCS

box in the C-terminal. The SH2 domain can bind to the phosphotyrosine residues in cytokine receptors or activated JAKs and is essential for target recognition, and presumably, can also interfere with receptor coupling with the JAK/STAT pathway (27). The SOCS box can trigger E3 ubiquitin ligase assembly to induce ubiquitination and proteosomal degradation of the signaling complex formed by JAK and cytokine receptor (28). An additional motif, namely the kinase inhibitory region, has been reported in SOCS1 and SOCS3 but not in other SOCS members (29), which can act as a pseudo-substrate domain to inhibit JAK activity (30). In different cell types (e.g., hepatocytes and macrophages), TNF α is involved in SOCS expression (e.g., SOCS3) induced by endotoxin/inflammation (31–33). SOCS expression, in agreement with its role as feedback signals for cytokines, has been reported to inhibit TNF α actions, e.g., SOCS1 over-expression can block TNF α -induced caspase activity and apoptosis in cardiomyocyte (34) and fibroblast culture prepared from rodents (35). Although TNF α receptors, namely TNFR1 and TNFR2, are known to mediate TNF α actions via activation of the IKK/NF κ B, JAK/STAT, MAPK, and TRADD/caspase pathways (36–38), there is a general lack of information on the mechanisms for TNF α -induced SOCS expression. Except for two reports, one in 3T3 cells (39) and the other in smooth muscle cells (40), showing the involvement of MAPK signaling in TNF α -induced SOCS3 expression, not much is known for the signal transduction for SOCS induction by TNF α . Of note, the post-receptor signaling for the corresponding responses occurred at the hepatic level, especially during GH resistance induced by endotoxin, has not been examined and represents an important topic for further investigation.

Recently, the key members of type II SOCS, SOCS1–3 and CISH, have been cloned in grass carp and confirmed to be inducible by GH and serve as the feedback repressors for GH signaling via the JAK₂/STAT₅ pathway with inhibitory effects on both basal and GH-induced IGF-I gene transcription at the hepatic level (41). In fish models, cytokine induction by microbial/viral infection [e.g., TNF α and interleukins in carp species (42, 43)] and IGF-I modulation by cytokines [e.g., TNF α] have been reported [e.g., in immune cells/tissues of trout and sea bass, see (44) for a recent review]. In turbot, bacterial infection is also effective in increasing SOCS3 mRNA levels in various tissues including the liver and SOCS3 over-expression can inhibit TNF α and IL-1 β expression in macrophages isolated from the head kidney (45). These findings not only suggest that the role of SOCS as feedback inhibitors to fine tune the immune responses is well-conserved from fish to mammals, but also raise the possibility that SOCS expression can form a functional link between the somatotrophic axis and immune system in fish models. Since the spreading of bacterial/viral infection is a major concern in fish culture, which will also adversely affect the growth performance of culture species, study has been initiated in our laboratory to examine the mechanisms for GH resistance in fish model induced by endotoxin of bacteria origin. Grass carp was chosen as the animal model as it is a representative of the carp family and by itself constitutes a major aquacultural output in Asian countries (~4.6 million tons per year and account for 15.6% of global finfish production)

Abbreviations: TNF α , Tumor necrosis factor α ; SOCS, Suppressor of cytokine signaling; CISH, Cytokine-inducible SH2-containing protein; GH, Growth hormone; IGF, Insulin-like growth factor; LPS, Lipopolysaccharide; IKK, I κ B kinase; NF κ B, Nuclear factor kappa beta; MEK_{1/2}, Mitogen-activated protein kinase kinase 1/2; ERK_{1/2}, Extracellular signal-regulated kinase 1/2; P₃₈^{MAPK}, P₃₈ Mitogen-activated protein kinase; PI3K, Phosphoinositide 3-kinase; Akt, Protein kinase B, JAK₂, Janus kinase 2; STAT, Signal transducer and activator of transcription.

(FAO yearbook of Fishery and Aquaculture Statistics 2011). In this study, the questions to be addressed are: (i) Can TNF α be induced by lipopolysaccharide (LPS, an endotoxin in gram-negative bacteria) and contribute to GH resistance in the carp liver along with the corresponding responses in SOCS? (ii) What are the subtypes of SOCS members induced by LPS at the hepatic level mediated by local production of TNF α signal? (iii) What are the signal transduction mechanisms involved in TNF α -induced SOCS expression which may contribute to GH resistance? As a first step, TNF α was cloned in grass carp and its tissue expression, especially in the liver, was confirmed by RT-PCR and LC/MS/MS. Using primary culture of carp hepatocytes, TNF α expression was examined with exposure to LPS and correlated to the corresponding changes in SOCS1-3 and CISH mRNA levels using real-time PCR. To evaluate if GH resistance could occur in the carp liver by TNF α signal induced by LPS, GH-induced IGF-I and -II mRNA expression in carp hepatocytes were tested with co-treatment of LPS or TNF α , respectively. Using a combination of pharmacological approach coupled with immunoblotting of the phosphorylation status of respective signaling targets/kinases, the signal transduction mechanisms for SOCS expression induced by TNF α were also elucidated at the hepatic level. Our studies, as a whole, have shed light on the mechanisms involved in GH resistance induced by endotoxin occurring in the liver of a fish model.

MATERIALS AND METHODS

Animals and Test Substances

One-year-old grass carp (*Ctenopharyngodon idellus*) were purchased from local wholesale markets and maintained in well-aerated 200 L aquaria at 18°C under at 12L:12D photoperiod. Since the carp at this stage was pre-pubertal and sexual dimorphism was not apparent, fish of mixed sexes were used for tissue sampling and hepatocyte preparation. During the process, the fish was killed by anesthesia in 0.05% MS222 (Sigma, St. Louis, MO) followed by spinosectomy according to the protocol approved by the Committee for Animal Use in Teaching and Research at the University of Hong Kong (Hong Kong). Human TNF α and bovine GH were obtained from R&D Systems (Minnesota, MN). The I κ B kinase (IKK) inhibitor N-(4-Pyrrolidin-1-yl-piperidin-1-yl)-[4-(4-benzo[b]thiophen-2-yl-pyrimidinylamino) phenyl]carboamide (IKK16) and NF- κ B activation blocker N-[(Phenyl methoxy)carbonyl]-L-leucyl-N-[(1S)-1-formyl-3-methylbutyl]-L-leucinamide (MG132) were acquired from TOCRIS (Eillsville, MO). The JAK₂ inhibitor 1,2,3,4,5,6-hexabromocyclohexane (Hex), STAT₁ inhibitor Fludarabine (FA), STAT₃ inhibitor Ethyl-1-(4-cyano-2,3,5,6-tetrafluorophenyl)-6,7,8-trifluoro-4-oxo-1,4-dihydroquinoline-3-carboxylate (ETDDC), STAT₅ inhibitor N1-(11H-Indolo[3,2-c] quinolin-6-yl)-N2,N2-dimethylethane-1,2-diamine (IQDMA), MEK_{1/2} inhibitor U0126, ERK_{1/2} inhibitor SCH 772984, P₃₈^{MAPK} inhibitor PD169316, PI3K inhibitor Ly294002, Akt inhibitor HIMOC and the endotoxin LPS were procured from Calbiochem (San Diego, CA). Test substances, except for TNF α and GH dissolved in double-distilled deionized water, were dissolved in DMSO

and stored frozen at -80°C in small aliquots. On the day of experimentation, test substances were thawed on ice and diluted to appropriate levels with prewarmed medium 15 min prior to drug administration. The final dilutions of DMSO were always $\leq 0.1\%$ and did not affect TNF α and SOCS mRNA expression in carp hepatocytes. In our studies, GH and TNF α of mammalian origin were used as the functional substitutes for the fish counterparts, as these mammalian proteins are commercially available and confirmed to be bioactive by previous reports in fish models [e.g., in carp (41) and rainbow trout (46, 47)].

Molecular Cloning, Gene Copy Number, and Tissue Expression of TNF α

For molecular cloning of grass carp TNF α , total RNA was isolated from the carp liver with TRIZOL and subjected to 5'/3' RACE using primers designed based on the conserved regions of TNF α reported in zebrafish. Sequence alignment, 3D protein modeling, and phylogenetic analysis were conducted using CLUSTAL-W (<https://www.genome.jp/tools-bin/clustalw>), SWISS-MODEL (<https://swissmodel.expasy.org/>) and MEGA 6.0 (<https://www.megasoftware.net/>), respectively. To deduce the gene copy number of TNF α , Southern blot was performed in genomic DNA isolated from the whole blood of grass carp as described previously (41). For tissue expression of TNF α , RT-PCR was conducted in selected tissues and brain areas using primers for carp TNF α (Forward primer: 5'-GCTTCACGC TCAACAAGT CTCA-3'; Reverse primer: 5'-AGCCTGGTCTGTTCACCTCT-3'). PCR for TNF α expression was conducted with denaturation at 94°C for 3 min followed by 37 cycles of denaturation at 94°C for 30 s, annealing at 56°C for 30 s, and extension at 72°C for 30 s, and concluded with a final extension step at 72°C for 5 min. The authenticity of the PCR products obtained was confirmed by Southern blot using a DIG-labeled cDNA probe for carp TNF α and parallel PCR for β actin was used as an internal control. Using LC/MS/MS, protein expression of TNF α in the carp liver was also evaluated using a SCIEX TripleTOF-5600 system (AB SCIEX, Concord, ON, Canada) according to the standard protocol in our laboratory (48).

TNF α and Type II SOCS mRNA Expression in Carp Hepatocytes

Primary culture of grass carp hepatocytes was prepared by collagenase digestion (41) and maintained in 24-well plates at a seeding density of $\sim 0.7 \times 10^6$ cells/ml/well. After drug treatment, total RNA was extracted from individual wells by TRIZOL, digested with DNase I, and reversely transcribed using Superscripts II (Invitrogen) with Oligo-dT as the primer. RT samples prepared were then subjected to real-time PCR for TNF α and type II SOCS mRNA measurement using a LightCycler 480 SYBR Green Master I Kit (Roche) with the RotorGene-Q qPCR System (Qiagen). In parallel experiments to evaluate GH resistance at the hepatic level, carp hepatocytes were exposed to GH with co-treatment of LPS or TNF α and the RT samples prepared were used for real-time PCR measurement of GHR and IGF-I/-II transcripts. Real-time PCR for respective gene targets, including TNF α , GHR, IGF-I/-II, and different members of type

II SOCS, will be conducted according to the conditions described in **Supplemental Table 1**. In our studies, serial dilutions of plasmids with the ORF of the target genes were used as the qPCR standards for data calibration and parallel measurement of 18S RNA expression was used as the internal control. After the assays, the authenticity of PCR products was routinely confirmed by melting curve analysis.


Western Blot of Signaling Targets and Kinases in Carp Hepatocytes

To test if TNF α can induce IKK/NF- κ B, JAK/STAT, MAPK, and PI3K/Akt activation in the carp liver, Western blot was performed in cell lysate prepared from carp hepatocytes with TNF α treatment using the antibodies for phosphorylated form (“p-” form) and total protein (“t-” form) of I κ B α (1:1,000; Santa Cruz), MEK $_{1/2}$ (1:1,000; Cell Signaling), Erk $_{1/2}$ (1:5,000; Sigma), P $_{38}^{MAPK}$ (1:1,000; Cell Signaling), Akt (1:1,000, Cell

Signaling), JAK $_2$ (1:1,000; Santa Cruz), STAT $_1$ (1:1,000; Cell Signaling), STAT $_3$ (1:1,000; Cell Signaling) and STAT $_5$ (1:1,000; Cell Signaling), respectively. In these studies, parallel blotting of β actin with an Actin Ab-1 Kit (Calbiochem) was used as the internal control.

Data Transformation and Statistical Analysis

For real-time PCR of TNF α , GHR, IGF-I/-II, SOCS1-3, and CISH mRNA levels, standard curves constructed with serial dilutions of plasmid DNA carrying the ORF of the respective gene targets with a dynamic range of $\geq 10^5$, amplification efficiency ≥ 0.98 and a correlation coefficient of ≥ 0.95 were used for data calibration with the RotorGene Q-Rex software (Qiagen). Given that the data for transcript expression of 18S RNA, the internal control for real-time PCR, did not exhibit significant changes in our studies, the raw data for target gene expression were simply






FIGURE 2 | Effects of LPS on TNF α and type II SOCS expression in primary culture of carp hepatocytes. Hepatocytes were treated with LPS (1 μ g/ml) for the duration as indicated. At the respective time points, total RNA was (Continued)

FIGURE 2 | extracted from the cell culture, reversely transcribed and subjected to real-time PCR for TNF α (A), SOCS1 (B), SOCS2 (C), SOCS3 (D), and CISH mRNA measurement (E). Data presented are expressed as mean \pm SEM ($N = 6$) and experimental groups denoted by different letters represent a significant difference at $p < 0.05$ (ANOVA followed by Newman-Keuls Test).

transformed as a percentage of the mean value in the control group without drug treatment (as “%Ctrl”). For Western blot, except for the data of I κ B α expressed as a ratio of I κ B α signals over β actin expression in the same sample (due to a drop of total I κ B α after treatment), the Western blot signals detected were quantitated by densitometry scanning in “arbitrary density unit” and expressed as the ratio of phosphorylated form over total protein of the same target. The normalized data were then expressed as fold increase compared to the control group without drug treatment (as “fold induction”). Data presented, expressed as mean \pm SEM ($N = 6$), are pooled results from six independent experiments and analyzed using one-way (for dose-dependence/co-treatment studies with signaling inhibitors)/two-way ANOVA (for time course) followed by Newman-Keuls test. Differences between groups were considered as significant at $p < 0.05$.

RESULTS

Molecular Cloning, Structural Characterization, and Tissue Expression of Grass Carp TNF α

To establish the structural identity of TNF α expressed in carp species, the full-length cDNA of grass carp TNF α (GenBank accession No. JQ040498) was cloned and found to be 1251 bp in size with a 720 bp ORF encoding a 239 a.a. protein (with deduced MW of \sim 26 kDa) flanking by a 126 bp 5’UTR and a 405 bp 3’UTR (Supplemental Figure 1). In the 3’UTR, five AU-rich elements (ARE, “atata”) were also located in the region overlapping with three polyadenylation signals (“aaaag” and “atata”) upstream of the poly(A) tail. Phylogenetic analysis of the nucleotide sequence obtained using the neighbor-joining method revealed that the newly cloned cDNA could be clustered within the clade of fish TNF α and closely related to the TNF α in carp species (Figure 1A). In the deduced a.a. sequence, the transmembrane domain and signature motif of the TNF family (IIIPDDGIYFVYSVSF) could be identified along with a TACE cleavage site (TL), three putative N-linked glycosylation sites (NXT/S) and two well-conserved Cys residues. Protein sequence alignment of grass carp TNF α with the corresponding sequences found in other species using CLUSTAL-W also confirms that the grass carp sequence is highly homologous to the TNF α reported in the carp family and to a lower extent when compared with the corresponding sequences in other fish species and tetrapods (Supplemental Figure 2). Of note, the twelve β sheets (namely β sheet 1–12) as a major structural characteristic of TNF α could also be identified in grass carp TNF α and the a.a. sequences of β sheet 5–6 (covering the signature motif of TNF α), β sheet 7–8, β sheet 9–10, and β sheet 12 were found to be highly conserved

among different species. *In silico* protein modeling using the crystal structure of human TNF α as the template also showed that the 3D structure of the secreted form of carp TNF α (covering β sheet 3–12) could fit into a highly packed structure with 10 anti-parallel β strands arranged in a β -jellyroll topography (Figure 1B). The 3D model deduced for carp TNF α , especially the spatial arrangement and orientation of β sheets, was found

to be highly comparable if not identical to that of the human counterpart except for the absence of two short helices in the linker between β sheet 6 and 7.

To determine the gene copy number of TNF α , Southern blot was performed in DNA sample isolated from the whole blood of grass carp using a DIG-labeled probe for carp TNF α . After digestion of DNA sample with restriction enzymes including




FIGURE 3 | LPS and TNF α treatment on IGF-I/II and GHR expression in carp hepatocytes. To examine the effects of LPS (A) and TNF α (B) on IGF-I/II expression, hepatocyte culture was exposed to GH (500 ng/ml) with/without the co-treatment of LPS (1 μ g/ml) or TNF α (100 ng/ml), respectively. In parallel experiments, hepatocytes were also treated with LPS (1 μ g/ml) or TNF α alone (100 ng/ml) to study their effects on GHR expression. To shed light on the role of TNF α in LPS-induced GH resistance at the hepatic level, the effects of LPS (1 μ g/ml) on IGF-I/II responses induced by GH (500 ng/ml) and basal level of GHR expression were also tested with co-treatment of the TNF α receptor R7050 (10 μ M) (C). In these studies, the duration of drug treatment was fixed at 24 h. After treatment, total RNA was extracted for real-time PCR measurement of IGF-I/II and GHR transcripts, respectively. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

BamHI, Bgl II, Cla I, EcoR V, and Pst I, respectively, a single band was consistently observed in individual lanes of the Southern blot (**Figure 1C**), implying that TNF α is a single-copy gene in the carp genome. For tissue expression profiling of TNF α expression, RT-PCR was also conducted in selected tissues and brain areas in grass carp. As shown in **Figure 1D**, PCR signals for TNF α were found to be expressed at high levels in the pituitary, thymus

and gonad, to a lower extent in the brain, spleen, and liver, and not detectable in the gills, heart, muscle, blood, kidney, intestine, and head kidney. In selected brain areas, TNF α signals were also detected in the hypothalamus, olfactory bulbs, optic tectum, medulla oblongata, and spinal cord, but not in the telencephalon and cerebellum. In these experiments, the PCR signals for β actin were consistently detected in all the samples examined,




FIGURE 4 | Effects of TNF α on type II SOCS expression in carp hepatocytes. **(A)** Time course and **(B)** dose dependence of TNF α treatment on SOCS1, SOCS2, SOCS3, and CISH expression. For time course study, hepatocytes were incubated with TNF α (100 ng/ml) for the duration as indicated up to 24 h. For dose dependence, cell culture were treated for 6 h with increasing concentrations of TNF α (0.1–100 ng/ml). After drug treatment, total RNA was extracted, reversely transcribed and subjected to real-time PCR for SOCS1, SOCS2, SOCS3, and CISH mRNA measurement, respectively. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

suggesting that the lack of TNF α signals due to RNA degradation was unlikely.

TNF α Induction by LPS and Its Effects on IGF-I/-II and SOCS/CISH Expression in Carp Hepatocytes

Using LC/MS/MS, peptide fragments originated from TNF α were also detected in protein sample prepared from the carp liver with trypsin digestion (protein coverage by peptides identified: 92.7% for peptides with $\geq 75\%$ confidence and 85.4% for peptides with 99% confidence), suggesting that the transcript signals of TNF α detected by RT-PCR can be properly translated into target protein in the carp liver (**Supplemental Figure 3**). To test for the presence of a functional TNF α /SOCS system at the hepatic level, primary culture of grass carp hepatocytes was challenged with the endotoxin LPS (1 $\mu\text{g/ml}$). In this case, a transient rise in TNF α mRNA levels with a peak response at 6 h was noted and

the elevation in TNF α signals reduced gradually and returned to basal by the end of 24 h (**Figure 2A**). Meanwhile, LPS induction was also effective in increasing SOCS1-3 and CISH transcript expression with peak responses at 3 h (**Figures 2B–E**). Of note, a significant rise in TNF α signal (at 1.5 h) could be observed prior to the corresponding changes in SOCS1-3 and CISH (at 3 h) and the peak responses of these type II SOCS also occurred before the “recovery phase” of TNF α expression (from 6 to 24 h).

In carp hepatocytes, GH induction (500 ng/ml) consistently induced IGF-I and -II mRNA expression and these stimulatory effects could be blocked by co-treatment with LPS (1 $\mu\text{g/ml}$, **Figure 3A**) or TNF α (100 ng/ml, **Figure 3B**), respectively. The blockade on GH-induced IGF-I and -II signals by LPS, however, was found to be partially reverted by the TNF α receptor antagonist R7050 (10 μM , **Figure 3C**). In these studies, LPS but not TNF α was also effective in reducing basal levels of IGF-I and -II as well as GHR mRNA expression (**Figures 3A,B**) but the inhibitory action on GHR by LPS was not affected by




FIGURE 5 | Functional role of IKK/NF κ B pathway in TNF α -induced SOCS1, SOCS3, and CISH expression at the hepatic level. **(A)** TNF α treatment on I κ B α phosphorylation in carp hepatocytes. Hepatocytes were treated with TNF α (100 ng/ml) for the duration as indicated. After that, cell lysate was prepared and subjected to Western blot with antibodies for the phosphorylated form (p-form) and total protein (t-form) of I κ B α . In this study, parallel blotting of β actin was used as the internal control. The signals for the p-form and t-form of I κ B α were quantified by densitometry scanning and normalized as a ratio of β actin expressed in the same sample. **(B)** Blocking IKK/NF κ B pathway on TNF α -induced SOCS1, SOCS3, and CISH mRNA expression. Hepatocytes were treated for 6 h with TNF α (100 ng/ml) in the presence/absence of the IKK inhibitor IKK16 (10 μM) or NF κ B inhibitor MG132 (10 μM). After that, total RNA was prepared and used for real-time PCR measurement of the respective transcripts. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

co-treatment with R7050 (Figure 3C). To test if the effects of LPS on SOCS expression could be mimicked by TNF α , time course and dose dependence studies were also conducted in carp hepatocytes with TNF α treatment. Interestingly, TNF α induction was found to up-regulate SOCS1, SOCS3, and CISH but not SOCS2 mRNA in a time- (Figure 4A) and dose-dependent manner (Figure 4B). Similar to the kinetics of LPS-induced TNF α mRNA expression, the gene expression of SOCS1, SOCS3, and CISH induced by TNF α reached their respective peaks at 6 h and gradually returned to basal by the end of 24 h.

Signal Transduction for TNF α -Induced SOCS1, SOCS3, and CISH mRNA Expression at Hepatic Level

To shed light on the post-receptor signaling mediating TNF α -induced SOCS1, SOCS3, and CISH expression at

the hepatic level, Western blot was conducted to monitor the effects of TNF α on protein phosphorylation of various signaling targets in carp hepatocytes. In this case, short-term treatment with TNF α (100 ng/ml, up to 30 min) was found to be effective in triggering rapid phosphorylation of I κ B (Figure 5A), MEK $_{1/2}$ (Figure 6A), ERK $_{1/2}$ (Figure 6A), MKK $_{3/6}$ (Figure 7A), P $_{38}^{MAPK}$ (Figure 7A), JAK $_2$ (Figure 8A), STAT $_1$ (Figure 8A), STAT $_3$ (Figure 9A), STAT $_5$ (Figure 9A), and Akt (Figure 10A), respectively. Except for the drop in total content of I κ B observed (Figure 5A), the total protein for other signaling targets examined did not show noticeable changes after TNF α treatment (Figures 6A–10A). In parallel studies, SOCS1, SOCS3, and CISH mRNA expression in carp hepatocytes were consistently up-regulated by TNF α (100 ng/ml) and these stimulatory effects could be reduced/abolished by co-treatment with the IKK inhibitor IKK16 (Figure 5B), NF κ B inhibitor MG132 (Figure 5B), MEK $_{1/2}$ inhibitor U0126




FIGURE 6 | Functional role of MEK/ERK pathway in TNF α -induced SOCS1, SOCS3, and CISH expression at the hepatic level. **(A)** TNF α treatment on MEK $_{1/2}$ and ERK $_{1/2}$ phosphorylation in carp hepatocytes. Hepatocytes were treated with TNF α (100 ng/ml) for various durations up to 30 min. After that, cell lysate was prepared and used in Western blot with antibodies for the phosphorylate form (p-form) and total protein (t-form) of MEK $_{1/2}$ and ERK $_{1/2}$, respectively, and parallel blotting of β actin was used as the internal control. The signals for the two forms of protein targets were quantified by densitometry scanning and expressed as a ratio of the p-form/t-form in the same sample. **(B)** Blocking MEK/ERK pathway on TNF α -induced SOCS1, SOCS3, and CISH mRNA expression. Hepatocytes were treated for 6 h with TNF α (100 ng/ml) in the presence/absence of the MEK $_{1/2}$ inhibitor U0126 (10 μ M) or ERK $_{1/2}$ inhibitor SCH772984 (10 μ M). After that, total RNA was prepared and used for real-time PCR measurement of the respective transcripts. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.




FIGURE 7 | Functional role of MKK_{3/6}/P₃₈^{MAPK} cascade in TNF α -induced SOCS1, SOCS3, and CISH expression at the hepatic level. **(A)** TNF α treatment on MKK_{3/6} and P₃₈^{MAPK} phosphorylation in carp hepatocytes. Hepatocytes were treated with TNF α (100 ng/ml) for the duration as indicated. After that, cell lysate was prepared and used in Western blot with antibodies for the phosphorylated form (p-form) and total protein (t-form) of MKK_{3/6} and P₃₈^{MAPK}, respectively, and parallel blotting of β actin was used as the internal control. **(B)** Blocking MKK_{3/6}/P₃₈^{MAPK} pathway on TNF α -induced SOCS1, SOCS3, and CISH mRNA expression. Hepatocytes were treated for 6 h with TNF α (100 ng/ml) in the presence/absence of the P₃₈^{MAPK} inhibitor PD169316 (10 μ M). After that, total RNA was isolated and used for real-time PCR measurement of the respective transcripts. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

(Figure 6B), ERK_{1/2} inhibitor SCH 772984 (Figure 6B), P₃₈^{MAPK} inhibitor PD169316 (Figure 7B), JAK₂ inhibitor HEX (Figure 8B), STAT₁ inhibitor FA (Figure 8B), STAT₃ inhibitor ETDDC (Figure 9B), STAT₅ inhibitor IQDMA (Figure 9B), PI3K inhibitor Ly294002 (Figure 10B), and Akt inhibitor HIMOC (Figure 10B), respectively. Besides, basal levels of SOCS1, SOCS3, and CISH transcript expression were also suppressed by the NF κ B inhibitor MG132 (Figure 5B), MEK_{1/2} inhibitor U0126 (Figure 6B), P₃₈^{MAPK} inhibitor PD169316 (Figure 7B), JAK₂ inhibitor HEX (Figure 8B), STAT₅ inhibitor IQDMA (Figure 9B), and Akt inhibitor HIMOC (Figure 10B).

DISCUSSION

As a first step to examine TNF α -induced GH resistance in the carp liver, the structural identity of grass carp TNF α was established by 5'/3'RACE. Based on phylogenetic analysis, the full-length cDNA obtained could be clustered in the clade of fish TNF α and closely related to the corresponding sequences in silver carp and common carp, the other members of the carp family. In fish models, two isoforms of TNF α encoded in separate genes are commonly reported in tetraploid species, including rainbow trout (49), common carp (50), and goldfish (51), which is assumed to be the result of the "fish-specific"




FIGURE 8 | Functional role of JAK₂ and STAT₁ in TNF α -induced SOCS1, SOCS3, and CISH expression at the hepatic level. **(A)** TNF α treatment on JAK₂ and STAT₁ phosphorylation in carp hepatocytes. Hepatocytes were treated with TNF α (100 ng/ml) up to 30 min and cell lysate prepared at different time points was used in Western blot with antibodies for phosphorylated form (p-form) and total protein (t-form) of JAK₂ and STAT₁, respectively, and parallel blotting of β actin was used as the internal control. **(B)** Blocking JAK₂ and STAT₁ on TNF α -induced SOCS1, SOCS3, and CISH mRNA expression. Hepatocytes were treated for 6 h with TNF α (100 ng/ml) in the presence/absence of the JAK₂ inhibitor HEX (50 μ M) or STAT₁ inhibitor FA (50 μ M). After that, total RNA was isolated and used for real-time PCR measurement of the respective transcripts. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

(or 3R) whole genome duplication occurred prior to tetrapod evolution (52). In our study, a single form of TNF α was cloned and confirmed to be a single-copy gene in the carp genome, which is consistent with the fact that grass carp is a diploid fish, and presumably, had been branched off without the 3R genome duplication of the teleost lineage (53). The ORF of grass carp TNF α encodes a 239 a.a. protein with the 14 a.a. signature motif of the TNF family, a highly conserved transmembrane domain, and the typical feature of 12 antiparallel β sheets arranged in a “ β -jellyroll” topography. The size of grass carp TNF α is similar to that of mammals, e.g., 233 a.a. for human TNF α (54) and 235 a.a. for mouse TNF α (55), but notably smaller than that of the gilthead seabream (253 a.a.), flounder (256 a.a.), and rainbow trout (246 a.a.) (56). *In silico* protein modeling also reveals that,

except for the lack of 2 short helices in the linker between β sheet 6 and 7, the 3D structure of carp TNF α , especially for the spatial arrangement and orientation of the β sheets forming the compact jellyroll structure, is highly comparable to its human counterpart. Apparently, despite the a.a. substitutions occurred at protein level from fish to mammals, the 3D structure of TNF α is well conserved during vertebrate evolution, which may account for the bioactivity of human TNF α observed in immune cells of fish origin reported previously, e.g., inducing respiratory burst in macrophages (47) and cell proliferation in leucocytes in rainbow trout (46). It is also worth mentioning that five ARE elements could be located among the multiple polyadenylation signals within the 3’UTR of grass carp TNF α . These AU-rich cis-acting elements are commonly found in 3’UTR of “unstable





FIGURE 9 | Functional role of STAT₃ and STAT₅ in TNF α -induced SOCS1, SOCS3, and CISH expression at the hepatic level. **(A)** TNF α treatment on STAT₃ and STAT₅ phosphorylation in carp hepatocytes. Hepatocytes were treated with TNF α (100 ng/ml) up to 30 min and cell lysate prepared was used in Western blot with antibodies for phosphorylated form (p-form) and total protein (t-form) of STAT₃ and STAT₅, respectively, and parallel blotting of β actin was used as the internal control. **(B)** Blocking STAT₃ and STAT₅ on TNF α -induced SOCS1, SOCS3, and CISH mRNA expression. Hepatocytes were treated for 6 h with TNF α (100 ng/ml) in the presence/absence of the STAT₃ inhibitor ETDDC (300 nM) or STAT₅ inhibitor IQDMA (50 μ M). After that, total RNA was isolated and used for real-time PCR measurement of the respective transcripts. Experimental groups denoted by different letters represent a significant difference at $P < 0.05$.

transcripts” and serve as the binding sites for ARE-binding proteins, including TTP, KSRP, and AUF1, which are known to trigger mRNA deadenylation and exonucleolytic decay (57). Presence of multiple ARE elements in the 3’UTR of TNF α has also been reported in rainbow trout (49), catfish (58), goldfish (51), and flounder (59), implying that the TNF α transcript in fish models may not be very stable, and probably, has a fast turnover rate, which is in agreement with the transient nature of pro-inflammatory responses mediated by TNF α (60).

In mammals, TNF α is expressed in immune cells including macrophages, monocytes, lymphocytes and neutrophils, especially on exposure to the bacterial endotoxin LPS. Similar findings have also been reported in fish models, e.g., in carp species (61, 62). Of note, TNF α expression can also be detected

in non-immune cells/tissues, e.g., in astrocytes, microglia, fibroblast, and smooth muscle [for review, see (63)]. In rainbow trout (64), goldfish (51), tuna (65), and catfish (58), constitutive expression of TNF α has been reported in various tissues, especially in immune organs (e.g., thymus and head kidney) as well as in tissues with high chance of exposure to microbes (e.g., gills and intestine). In grass carp, as revealed by RT-PCR, high levels of TNF α signal were detected in the thymus, pituitary, and gonad, to a lower extent in the brain, spleen, and liver, but not in other tissues examined. Within the brain, low levels of TNF α signals were also located in the olfactory bulb, hypothalamus, optic tectum and medulla oblongata but not in other brain areas. In contrast to the wide spread/ubiquitous pattern of expression reported in other fish



species, grass carp tends to have a tissue-specific pattern of TNF α expression. Although TNF α signal could be detected in the carp thymus, it was not found in the gills and head kidney, which are the major sites of TNF α expression in other fish models. The cause of the discrepancy is unclear and may be related to species-specific variations/culture environment of the species concerned. Apparently, basal expression of TNF α in most of the tissues/brain areas examined in grass carp (i.e., without experimentally induced infection/inflammation) is rather low/undetectable. Of note, our study also demonstrated for the first time that high levels of TNF α signals could be identified in the carp pituitary and gonad. The relevance of these findings related to endocrine regulation and reproductive functions is still unclear and future investigations are clearly warranted.


In our study with grass carp, detectable levels of TNF α transcript (by RT-PCR) and protein signals (by LC/MS/MS) could be identified in the liver, which is consistent with the idea that the liver can serve as a major immune target (e.g., for hepatitis virus) (66) as well as an immunological organ in vertebrates (e.g., as a source of complement factors and

immune cells) (67). In primary culture of carp hepatocytes, LPS induction could trigger a transient rise in TNF α gene expression with parallel elevations in SOCS1, SOCS2, SOCS3, and CISH mRNA levels. A significant rise in TNF α signal was found to occur prior to those of type II SOCS and a gradual drop in TNF α response was also noted after the peak responses of SOCS1, SOCS2, SOCS3, and CISH expression. These findings corroborate with the idea that SOCS expression induced by infection/inflammation can serve as feedback inhibitors for cytokine signaling (17, 23). Of note, TNF α treatment was effective in mimicking the stimulatory effects of LPS on SOCS1, SOCS3, and CISH but not SOCS2 mRNA expression. In parallel experiments, transcript expression of IGF-I and -II induced by GH could be blocked by co-treatment with LPS and these inhibitory actions were also mimicked by TNF α and partially recovered by TNF α receptor antagonism. These findings, as a whole, suggest that (i) a functional system composed of TNF α and type II SOCS signals is present in the carp liver and inducible by endotoxin, presumably forming an integral component of the innate immunity in carp species, and (ii) type II SOCS

expression induced by LPS at hepatic level, including SOCS1, SOCS3, and CISH, could be mediated by local production of TNF α , which may contribute to GH resistance for IGF-I/-II expression in the carp liver induced by endotoxin. In fish models including rainbow trout (64), catfish (58), and common carp (61, 62), TNF α expression induced by LPS has been reported, e.g., in macrophages and leucocytes, but the functional relevance of the phenomenon in GH resistance has not been examined. In mammals, endotoxin exposure can induce GH resistance at tissue level, e.g., in the liver, muscle and intestine (3, 4, 68). The effect is mediated by local production of cytokines including TNF α , IL-1 β , and IL-6 (5, 68), which are known to reduce GH responsiveness by reducing GHR expression (10, 11) or blocking GHR signaling via SOCS expression (3, 13). Apparently, different cytokines have their distinct role in GH resistance induced by endotoxin (33). In liver cells or hepatoma cell lines (e.g., Huh-7 cells), TNF α and IL-1 β induced by LPS only have minor effects on SOCS expression and their effects on GH resistance are mediated by reducing GHR gene transcription (32, 33), probably through inhibition of Sp1/3 binding to GHR promoter (10). In contrast, IL-6 signal induced by LPS has no effect on GHR expression but can serve as a potent stimulator for SOCS3 and CISH expression in these cell models (33, 69). In our recent study, type II SOCS including SOCS1-3 and CISH were up-regulated by GH in carp hepatocytes and over-expression of these SOCS members could inhibit JAK₂/STAT₅ signaling and block GH-induced IGF-I promoter activation, implying that type II SOCS can serve as feedback repressors for GH signaling in carp species (41). In our current study with carp hepatocytes, GH resistance induced by LPS occurred with parallel rises in TNF α and type II SOCS signals with a concurrent drop in GHR expression. Unlike mammals, TNF α treatment did not alter GHR expression but was effective in increasing SOCS1, SOCS3, and CISH signals at the hepatic level. These results suggest that TNF α may exert its effect on GH resistance by “cross-inhibition” on GH signaling via SOCS expression to down-regulate IGF-I/-II responses in the carp liver.

Although TNF α -induced SOCS expression has been reported in different cell models, including the fibroblasts, macrophages and hepatocytes in rodents (32, 39), and TNF α receptors are well-documented to be functionally coupled to the IKK/NF κ B, MAPK, PI3K/Akt, and JAK/STAT cascades (37, 38), the post-receptor signaling for TNF α -induced SOCS expression at the hepatic level is largely unknown. In carp hepatocytes, TNF α could induce a rapid phosphorylation of I κ B, MEK_{1/2}, ERK_{1/2}, MKK_{3/6}, P₃₈^{MAPK}, JAK₂, STAT_{1,3,5}, and Akt, respectively. Furthermore, pharmacological inhibition of IKK, NF κ B, MEK_{1/2}, ERK_{1/2}, P₃₈^{MAPK}, PI3K, Akt, JAK₂, and STAT_{1,3,5} were also effective in blocking SOCS1, SOCS3, and CISH mRNA expression induced by TNF α . These findings, as a whole, provide evidence that TNF α -induced type II SOCS expression in the carp liver (except for the lack of SOCS2 response) was mediated by the IKK/NF κ B, MAPK, PI3K/Akt, and JAK/STAT cascades and the subsequent rises of SOCS/CISH signals presumably can play a role in TNF α -induced GH resistance at the hepatic level. In our study, the cellular content of I κ B was reduced with concurrent rise in I κ B phosphorylation after TNF α treatment. These findings imply that, similar to the mechanisms for NF κ B activation in

mammals (70), TNF α activation of IKK/NF κ B pathway in carp liver may also involve IKK phosphorylation and proteosomal degradation of I κ B to allow for nuclear translocation of NF κ B and target gene transcription. In different cell models, TNF α activation of IKK/NF κ B [e.g., IKK2 in hepatocytes (71)], MAPK [e.g., ERK_{1/2} in trophoblasts (72) and P₃₈^{MAPK} in macrophages (39)], PI3K/Akt [e.g., Akt in myoblasts (73)], and JAK/STAT cascades [e.g., JAK₁ in lymphoma B cells (74) and STAT_{1,3,5} in 3T3-L1 adipocytes (75)] have been reported and these post-receptor signaling pathways can also exhibit functional crosstalk at cellular level. For examples, TNF α -induced MMP-9 promoter activity via NF κ B in HIPEC-65 cells is dependent on ERK_{1/2} activation (72). Previous studies on post-receptor signaling of TNFR1 in HEK293 cells reveal that MEKK1, the upstream activator of JNK_{1/2}, can activate IKK with subsequent induction of I κ B/NF κ B cascade. Meanwhile, NF κ B-inducing kinase (NIK) associated with TNFR1, an upstream activator of IKK, is also effective in cross-activation of JNK kinase, the upstream activator of P₃₈^{MAPK} and JNK_{1/2} (76). In MCF-7 cancer cells, TNF α -induced NF κ B activity is partly mediated by PI3K/Akt activation and NIK coupling to TNFR1 (77). Whether similar crosstalk in post-receptor signaling can also occur in carp hepatocytes



and contribute to TNF α -induced GH resistance is unclear and remains to be elucidated.

In summary, grass carp TNF α was cloned and confirmed to be a single copy gene in the carp genome. Its tissue expression, including the liver, has been confirmed and functional studies in carp hepatocytes also reveal that TNF α together with its induction of type II SOCS, including SOCS1, SOCS3, and CISH, may constitute a local immune response induced by LPS and exert negative effects on GH-induced IGF -I and -II expression at hepatic level (**Figure 11**). Apparently, the stimulatory actions of TNF α on SOCS1, SOCS3, and CISH expression in carp hepatocytes are mediated via activation of IKK/NF κ B, MAPK, PI3K/Akt, and JAK/STAT cascades. Our studies, as a whole, have shed light on the functional crosstalk between the immune system and somatotrophic axis at the hepatic level in a fish model. In this case, local production of TNF α with subsequent induction of different SOCS members probably can mediate GH resistance commonly observed in the carp liver after infection with microbes/exposure to endotoxin. In carp hepatocytes, LPS was shown to elevate SOCS2 but suppress GHR expression and TNF α treatment was not effective in these regards. The discrepancy observed raises the possibility that other cytokines can also be induced by endotoxin exposure and contribute to GH resistance by functional coupling with SOCS2 and GHR regulation. In trout hepatocytes, cytokine expression induced by LPS (e.g., IL-8) can be blocked by cortisol co-treatment and cortisol alone is known to trigger SOCS1/2 up-regulation in the same model (78). Local production of other cytokines induced by LPS and their functional interactions with signals from the hypothalamo-pituitary-adrenal axis in GH resistance for sure can be an interesting topic for our future study in carp model.

DATA AVAILABILITY STATEMENT

The datasets generated for this study can be found in GenBank, at this link: <https://www.ncbi.nlm.nih.gov/nucleotide/JQ040498>.

REFERENCES

- Aggarwal BB, Gupta SC, Kim JH. Historical perspectives on tumor necrosis factor and its superfamily: 25 years later, a golden journey. *Blood*. (2012) 119:651–65. doi: 10.1182/blood-2011-04-325225
- Holbrook J, Lara-Reyna S, Jarosz-Griffiths H, McDermott M. Tumour necrosis factor signalling in health and disease. *F1000Res*. (2019) 8:111–123. doi: 10.12688/f1000research.17023.1
- Chen Y, Sun D, Krishnamurthy VM, Rabkin R. Endotoxin attenuates growth hormone-induced hepatic insulin-like growth factor I expression by inhibiting JAK₂/STAT₅ signal transduction and STAT_{5b} DNA binding. *Am J Physiol Endocrinol Metab*. (2007) 292:E1856–62. doi: 10.1152/ajpendo.00581.2006
- Frost RA, Nystrom GJ, Lang CH. Tumor necrosis factor α decreases insulin-like growth factor-I messenger ribonucleic acid expression in C2C12 myoblasts via a Jun N-terminal kinase pathway. *Endocrinology*. (2003) 144:1770–9. doi: 10.1210/en.2002-220808
- Wang P, Li N, Li JS, Li WQ. The role of endotoxin, TNF α , and IL-6 in inducing the state of growth hormone insensitivity. *World J Gastroenterol*. (2002) 8:531–6. doi: 10.3748/wjg.v8.i3.531

ETHICS STATEMENT

The study was carried out according to the guidelines for the care and use of animals for research and teaching at the University of Hong Kong (Hong Kong). The protocol used in our study was approved by the Committee on the Use of Live Animal for Teaching and Research, University of Hong Kong.

AUTHOR CONTRIBUTIONS

AW was the PI and grant holder. AW and XJ were responsible for project planning and data analysis. XJ and JB were involved in molecular cloning of TNF α and functional studies in grass carp hepatocyte culture. MH was responsible for Western blot studies and LC/MS/MS detection of TNF α expression in the liver. Manuscript preparation was done by AW, CC, and XJ.

FUNDING

This project was supported by GRF Grants (17113918, 17117716, and 17128215) from the Research Grant Council (Hong Kong).

ACKNOWLEDGMENTS

Support from the School of Biological Sciences, University of Hong Kong (Hong Kong), in the form of postgraduate studentship (to JB and XJ) is acknowledged. We also thank Dr. Gerald Brown for his help in setting up the assay systems for GHR and IGF-I/-II transcript expression in grass carp.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fendo.2020.00020/full#supplementary-material>

- Lang CH, Hong-Brown L, Frost RA. Cytokine inhibition of JAK-STAT signaling: - A new mechanism of growth hormone resistance. *Pediatr Nephrol*. (2005) 20:306–12. doi: 10.1007/s00467-004-1607-9
- Thissen JP, Verniers J. Inhibition by interleukin-1 β and tumor necrosis factor α of the insulin-like growth factor I messenger ribonucleic acid response to growth hormone in rat hepatocyte primary culture. *Endocrinology*. (1997) 138:1078–84. doi: 10.1210/endo.138.3.4966
- Ahmed T, Yumet G, Shumate M, Lang CH, Rotwein P, Cooney RN. Tumor necrosis factor inhibits growth hormone-mediated gene expression in hepatocytes. *Am J Physiol Gastrointest Liver Physiol*. (2006) 291:G35–44. doi: 10.1152/ajpgi.00550.2005
- Yumet G, Shumate ML, Bryant P, Lin CM, Lang CH, Cooney RN. Tumor necrosis factor mediates hepatic growth hormone resistance during sepsis. *Am J Physiol Endocrinol Metab*. (2002) 283:E472–81. doi: 10.1152/ajpendo.00107.2002
- Denson LA, Menon RK, Shaull A, Bajwa HS, Williams CR, Karpen SJ. TNF α downregulates murine hepatic growth hormone receptor expression by inhibiting Sp1 and Sp3 binding. *J Clin Invest*. (2001) 107:1451–8. doi: 10.1172/JCI10994
- DiFedele LM, He J, Bonkowski EL, Han X, Held MA, Bohan A, et al. Tumor necrosis factor α blockade restores growth hormone

- signaling in murine colitis. *Gastroenterology*. (2005) 128:1278–91. doi: 10.1053/j.gastro.2005.02.003
12. Han X, Benight N, Osuntokun B, Loesch K, Frank SJ, Denson LA. Tumour necrosis factor α blockade induces an anti-inflammatory growth hormone signalling pathway in experimental colitis. *Gut*. (2007) 56:73–81. doi: 10.1136/gut.2006.094490
 13. Yumet G, Shumate ML, Bryant DP, Lang C, Cooney RN. Hepatic growth hormone resistance during sepsis is associated with increased suppressors of cytokine signaling expression and impaired growth hormone signaling. *Crit Care Med*. (2006) 34:1420–7. doi: 10.1097/01.CCM.0000215113.66070.E0
 14. Ahmed SF, Farquharson C. The effect of GH and IGF1 on linear growth and skeletal development and their modulation by SOCS proteins. *J Endocrinol*. (2010) 206:249–59. doi: 10.1677/JOE-10-0045
 15. Liu Z, Cordoba-Chacon J, Kineman RD, Cronstein BN, Muzumdar R, Gong Z, et al. Growth hormone control of hepatic lipid metabolism. *Diabetes*. (2016) 65:3598–609. doi: 10.2337/db16-0649
 16. Szalecki M, Malinowska A, Prokop-Piotrkowska M, Janas R. Interactions between the growth hormone and cytokines - A review. *Adv Med Sci*. (2018) 63:285–9. doi: 10.1016/j.advms.2018.03.001
 17. Yoshimura A, Ito M, Chikuma S, Akanuma T, Nakatsukasa H. Negative regulation of cytokine signaling in immunity. *Cold Spring Harb Perspect Biol*. (2018) 10:a028571. doi: 10.1101/cshperspect.a028571
 18. Delgado-Ortega M, Melo S, Meurens F. Expression of SOCS1-7 and CIS mRNA in porcine tissues. *Vet Immunol Immunopathol*. (2011) 144:493–8. doi: 10.1016/j.vetimm.2011.08.002
 19. Cheng SM, Li JC, Lin SS, Lee DC, Liu L, Chen Z, et al. HIV-1 transactivator protein induction of suppressor of cytokine signaling-2 contributes to dysregulation of IFN γ signaling. *Blood*. (2009) 113:5192–201. doi: 10.1182/blood-2008-10-183525
 20. Akhtar LN, Benveniste EN. Viral exploitation of host SOCS protein functions. *J Virol*. (2011) 85:1912–21. doi: 10.1128/JVI.01857-10
 21. Hebenstreit D, Luft P, Schmiedlechner A, Regl G, Frischauf A, Aberger F, et al. IL-4 and IL-13 induce SOCS-1 gene expression in A549 cells by three functional STAT $_6$ -binding motifs located upstream of the transcription initiation site. *J Immunol*. (2003) 171:5901–7. doi: 10.4049/jimmunol.171.11.5901
 22. Sato T, Saito R, Jinushi T, Tsuji T, Matsuzaki J, Koda T, et al. IFN γ -induced SOCS-1 regulates STAT $_6$ -dependent eotaxin production triggered by IL-4 and TNF α . *Biochem Biophys Res Commun*. (2004) 314:468–75. doi: 10.1016/j.bbrc.2003.12.124
 23. Linossi EM, Calleja DJ, Nicholson SE. Understanding SOCS protein specificity. *Growth Factors*. (2018) 36:104–17. doi: 10.1080/08977194.2018.1518324
 24. Jiang M, Zhang WW, Liu P, Yu W, Liu T, Yu J. Dysregulation of SOCS-mediated negative feedback of cytokine signaling in carcinogenesis and its significance in cancer treatment. *Front Immunol*. (2017) 8:70. doi: 10.3389/fimmu.2017.00070
 25. Yoshimura A, Suzuki M, Sakaguchi R, Hanada T, Yasukawa H. SOCS, inflammation, and autoimmunity. *Front Immunol*. (2012) 3:20. doi: 10.3389/fimmu.2012.00020
 26. Jin HJ, Shao JZ, Xiang LX, Wang H, Sun LL. Global identification and comparative analysis of SOCS genes in fish: - Insights into the molecular evolution of SOCS family. *Mol Immunol*. (2008) 45:1258–68. doi: 10.1016/j.molimm.2007.09.015
 27. Liao NP, Laktyushin A, Babon JJ. Purification of SOCS SH2 domains for structural and functional studies. In: Machida K, Liu BA, editors. *Methods in Molecular Biology*. Vol. 1555. Springer (2017). p. 173–82. doi: 10.1007/978-1-4939-6762-9_10
 28. Durham GA, Williams JJ, Nasim MT, Palmer TM. Targeting SOCS proteins to control JAK-STAT signalling in disease. *Trends Pharmacol Sci*. (2019) 40:298–308. doi: 10.1016/j.tips.2019.03.001
 29. Piessevaux J, Lavens D, Peelman F, Tavernier J. The many faces of the SOCS box. *Cytokine Growth Factor Rev*. (2008) 19:371–81. doi: 10.1016/j.cytogfr.2008.08.006
 30. Sasaki A, Yasukawa H, Suzuki A, Kamizono S, Syoda T, Kinjyo I, et al. Cytokine-inducible SH2 protein-3 (CIS3/SOCS3) inhibits Janus tyrosine kinase by binding through the N-terminal kinase inhibitory region as well as SH2 domain. *Genes Cells*. (1999) 4:339–51. doi: 10.1046/j.1365-2443.1999.00263.x
 31. Bode JG, Nimmegern A, Schmitz J, Schaper F, Schmitt M, Frisch W, et al. LPS and TNF α induce SOCS3 mRNA and inhibit IL-6-induced activation of STAT $_3$ in macrophages. *FEBS Lett*. (1999) 463:365–70. doi: 10.1016/S0014-5793(99)01662-2
 32. Colson A, Le Cam A, Maiter D, Edery M, Thissen JP. Potentiation of growth hormone-induced liver suppressors of cytokine signaling messenger ribonucleic acid by cytokines. *Endocrinology*. (2000) 141:3687–95. doi: 10.1210/endo.141.10.7724
 33. Zhao Y, Xiao X, Frank SJ, Lin HY, Xia Y. Distinct mechanisms of induction of hepatic growth hormone resistance by endogenous IL-6, TNF α , and IL-1 β . *Am J Physiol Endocrinol Metab*. (2014) 307:E186–98. doi: 10.1152/ajpendo.00652.2013
 34. Yan L, Tang Q, Shen D, Peng S, Zheng Q, Guo H, et al. SOCS-1 inhibits TNF α -induced cardiomyocyte apoptosis via ERK $_{1/2}$ pathway activation. *Inflammation*. (2008) 31:180–8. doi: 10.1007/s10753-008-9063-5
 35. Kimura A, Naka T, Nagata S, Kawase I, Kishimoto T. SOCS-1 suppresses TNF α -induced apoptosis through the regulation of JAK activation. *Int Immunol*. (2004) 16:991–9. doi: 10.1093/intimm/dxh102
 36. Baud V, Karin M. Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol*. (2001) 11:372–7. doi: 10.1016/S0962-8924(01)02064-5
 37. Li J, Yin Q, Wu H. Structural basis of signal transduction in the TNF receptor superfamily. *Adv Immunol*. (2013) 119:135–53. doi: 10.1016/B978-0-12-407707-2.00005-9
 38. Wullaert A, van Loo G, Heynincx K, Beyaert R. Hepatic tumor necrosis factor signaling and nuclear factor-kappa B: effects on liver homeostasis and beyond. *Endocr Rev*. (2007) 28:365–86. doi: 10.1210/er.2006-0031
 39. Ehrling C, Lai WS, Schaper F, Brenndorfer ED, Matthes RJ, Heinrich PC, et al. Regulation of SOCS3 mRNA stability by TNF α involves activation of the MKK $_6$ /P $_{38}$ ^{MAPK}/MK2 cascade. *J Immunol*. (2007) 178:2813–26. doi: 10.4049/jimmunol.178.5.2813
 40. Hsu CK, Lee IT, Lin CC, Hsiao LD, Yang CM. Nox2/ROS-dependent human antigen R translocation contributes to TNF α -induced SOCS-3 expression in human tracheal smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol*. (2014) 306:L521–33. doi: 10.1152/ajplung.00274.2013
 41. Jiang X, Xiao J, He M, Ma A, Wong AO. Type II SOCS as a feedback repressor for GH-induced IGF-I expression in carp hepatocytes. *J Endocrinol*. (2016) 229:171–86. doi: 10.1530/JOE-15-0423
 42. Basu M, Swain B, Maiti NK, Routray P, Samanta M. Inductive expression of toll-like receptor 5 (TLR5) and associated downstream signaling molecules following ligand exposure and bacterial infection in the Indian major carp, mrigal (*Cirrhinus mrigala*). *Fish Shellfish Immunol*. (2012) 32:121–31. doi: 10.1016/j.fsi.2011.10.031
 43. Samanta M, Basu M, Swain B, Panda P, Jayasankar P. Molecular cloning and characterization of Toll-like receptor 3, and inductive expression analysis of type I IFN, Mx and pro-inflammatory cytokines in the Indian carp, rohu (*Labeo rohita*). *Mol Biol Rep*. (2013) 40:225–35. doi: 10.1007/s11033-012-2053-6
 44. Franz AC, Faass O, Kollner B, Shved N, Link K, Casanova A, et al. Endocrine and local IGF-I in the bony fish immune system. *Biology*. (2016) 5:9. doi: 10.3390/biology5010009
 45. Zhang M, Xiao ZZ, Sun L. Suppressor of cytokine signaling 3 inhibits head kidney macrophage activation and cytokine expression in *Scophthalmus maximus*. *Dev Comp Immunol*. (2011) 35: 174–81. doi: 10.1016/j.dci.2010.09.006
 46. Hardie LJ, Chappell LH, Secombes CJ. Human tumor necrosis factor α influences rainbow trout *Oncorhynchus mykiss* leucocyte responses. *Vet Immunol Immunopathol*. (1994) 40:73–84. doi: 10.1016/0165-2427(94)90016-7
 47. Nova F, Figueras A, Ashton I, Secombes CJ. *In vitro* studies on the regulation of rainbow trout (*Oncorhynchus mykiss*) macrophage respiratory burst activity. *Dev Comp Immunol*. (1996) 20:207–16. doi: 10.1016/0145-305X(96)00011-0
 48. Wong MK, Sze KH, Chen T, Cho CK, Law HC, Chu IK, et al. Goldfish spexin: Solution structure and novel function as a satiety factor in

- feeding control. *Am J Physiol Endocrinol Metab.* (2013) 305:E348–66. doi: 10.1152/ajpendo.00141.2013
49. Zou J, Wang T, Hirono I, Aoki T, Inagawa H, Honda T, et al. Differential expression of two tumor necrosis factor genes in rainbow trout, *Oncorhynchus mykiss*. *Dev Comp Immunol.* (2002) 26:161–72. doi: 10.1016/S0145-305X(01)00058-1
50. Saeji JP, Stet RJ, de Vries BJ, van Muiswinkel WB, Wiegertjes GF. Molecular and functional characterization of carp TNF: a link between TNF polymorphism and trypanotolerance? *Dev Comp Immunol.* (2003) 27:29–41. doi: 10.1016/S0145-305X(02)00064-2
51. Grayfer L, Walsh JG, Belosevic M. Characterization and functional analysis of goldfish (*Carassius auratus L.*) tumor necrosis factor α . *Dev Comp Immunol.* (2008) 32:532–43. doi: 10.1016/j.dci.2007.09.009
52. Meyer A, Van de Peer Y, From 2R to 3R: evidence for a fish-specific genome duplication (FSGD). *Bioessays.* (2005) 27:937–45. doi: 10.1002/bies.20293
53. Zhu C, Tong J, Yu X, Guo W. Comparative mapping for bighead carp (*Aristichthys nobilis*) against model and non-model fishes provides insights into the genomic evolution of cyprinids. *Mol Genet Genomics.* (2015) 290:1313–26. doi: 10.1007/s00438-015-0992-z
54. Iris FJ, Bougueleret L, Prieur S, Caterina D, Primas G, Perrot V, et al. Dense Alu clustering and a potential new member of the NF κ B family within a 90 kilobase HLA class III segment. *Nat Genet.* (1993) 3:137–45. doi: 10.1038/ng0293-137
55. Cheng J, Turksen K, Yu QC, Schreiber H, Teng M, Fuchs E. Cachexia and graft-vs.-host-disease-type skin changes in keratin promoter-driven TNF α transgenic mice. *Genes Dev.* (1992) 6:1444–56. doi: 10.1101/gad.6.8.1444
56. Garcia-Castillo J, Pelegrin P, Mulero V, Meseguer J. Molecular cloning and expression analysis of tumor necrosis factor α from a marine fish reveal its constitutive expression and ubiquitous nature. *Immunogenetics.* (2002) 54:200–7. doi: 10.1007/s00251-002-0451-y
57. Bolognani F, Perrone-Bizzozero NI. RNA-protein interactions and control of mRNA stability in neurons. *J Neurosci Res.* (2008) 86:481–9. doi: 10.1002/jnr.21473
58. Zou J, Secombes CJ, Long S, Miller N, Clem LW, Chinchar VG. Molecular identification and expression analysis of tumor necrosis factor in channel catfish (*Ictalurus punctatus*). *Dev Comp Immunol.* (2003) 27:845–58. doi: 10.1016/S0145-305X(03)00085-5
59. Hirono I, Nam BH, Kurobe T, Aoki T. Molecular cloning, characterization, and expression of TNF cDNA and gene from Japanese flounder *Paralichthys olivaceus*. *J Immunol.* (2000) 165:4423–7. doi: 10.4049/jimmunol.165.8.4423
60. Bradley JR. TNF-mediated inflammatory disease. *J Pathol.* (2008) 214:149–60. doi: 10.1002/path.2287
61. Duan D, Sun Z, Jia S, Chen Y, Feng X, Lu Q. Characterization and expression analysis of common carp *Cyprinus carpio* TLR5M. *DNA Cell Biol.* (2013) 32:611–20. doi: 10.1089/dna.2013.2051
62. Yuan C, Pan X, Gong Y, Xia A, Wu G, Tang J, et al. Effects of astragalus polysaccharides (APS) on the expression of immune response genes in head kidney, gill and spleen of the common carp, *Cyprinus carpio L.* *Int Immunopharmacol.* (2008) 8:51–8. doi: 10.1016/j.intimp.2007.10.009
63. Goetz FW, Planas JV, MacKenzie S. Tumor necrosis factors. *Dev Comp Immunol.* (2004) 28:487–97. doi: 10.1016/j.dci.2003.09.008
64. Laing KJ, Wang T, Zou J, Holland J, Hong S, Bols N, et al. Cloning and expression analysis of rainbow trout *Oncorhynchus mykiss* tumour necrosis factor α . *Eur J Biochem.* (2001) 268:1315–22. doi: 10.1046/j.1432-1327.2001.01996.x
65. Kadowaki T, Harada H, Sawada Y, Kohchi C, Soma G, Takahashi Y, et al. Two types of tumor necrosis factor α in bluefin tuna genes: molecular cloning and expression profile in response to several immunological stimulants. *Fish Shellfish Immunol.* (2009) 27:585–94. doi: 10.1016/j.fsi.2008.12.006
66. Eksteen B. Infections and the liver. *Dig Dis.* (2011) 29:184–90. doi: 10.1159/000323884
67. Racanelli V, Rehermann B. The liver as an immunological organ. *Hepatology.* (2006) 43:S54–62. doi: 10.1002/hep.21060
68. Soendergaard C, Kvist PH, Thygesen P, Reslow M, Nielsen OH, Kopchick JJ, et al. Characterization of growth hormone resistance in experimental and ulcerative colitis. *Int J Mol Sci.* (2017) 18:E2046. doi: 10.3390/ijms18102046
69. Denson LA, Held MA, Menon RK, Frank SJ, Parlow AF, Arnold DL. Interleukin-6 inhibits hepatic growth hormone signaling via upregulation of CIS and SOCS-3. *Am J Physiol Gastrointest Liver Physiol.* (2003) 284:G646–54. doi: 10.1152/ajpgi.00178.2002
70. Schuster M, Annemann M, Plaza-Sirvent C, Schmitz I. Atypical I κ B proteins - nuclear modulators of NF κ B signaling. *Cell Commun Signal.* (2013) 11:23. doi: 10.1186/1478-811X-11-23
71. Luedde T, Assmus U, Wustefeld T, Meyer zu Vilsendorf A, Roskams T, Schmidt-Supprian M, et al. Deletion of IKK $_2$ in hepatocytes does not sensitize these cells to TNF-induced apoptosis but protects from ischemia/reperfusion injury. *J Clin Invest.* (2005) 115:849–59. doi: 10.1172/JCI23493
72. Cohen M, Meisser A, Haeggeli L, Bischof P. Involvement of MAPK pathway in TNF α -induced MMP-9 expression in human trophoblastic cells. *Mol Hum Reprod.* (2006) 12:225–32. doi: 10.1093/molehr/gal023
73. Plaisance I, Morandi C, Murigande C, Brink M. TNF α increases protein content in C2C12 and primary myotubes by enhancing protein translation via the TNF-R1, PI3K, and MEK. *Am J Physiol Endocrinol Metab.* (2008) 294:E241–50. doi: 10.1152/ajpendo.00129.2007
74. Miscia S, Marchisio M, Grilli A, Di Valerio V, Centurione L, Sabatino G, et al. Tumor necrosis factor α activates JAK $_1$ /STAT $_3$ -STAT $_5B$ signaling through TNFR-1 in human B cells. *Cell Growth Differ.* (2002) 13:13–8.
75. Guo D, Dunbar JD, Yang CH, Pfeffer LM, Donner DB. Induction of JAK/STAT signaling by activation of the type 1 TNF receptor. *J Immunol.* (1998) 160:2742–50.
76. Karin M, Delhase M. JNK or IKK, AP-1 or NF κ B, which are the targets for MEK kinase 1 action? *Proc Natl Acad Sci USA.* (1998) 95:9067–9. doi: 10.1073/pnas.95.16.9067
77. Burow ME, Weldon CB, Melnik LI, Duong BN, Collins-Burow BM, Beckman BS, et al. PI3K/Akt regulation of NF κ B signaling events in suppression of TNF-induced apoptosis. *Biochem Biophys Res Commun.* (2000) 271:342–5. doi: 10.1006/bbrc.2000.2626
78. Philip AM, Kim DS, Vijayan MM. Cortisol modulates the expression of cytokines and suppressors of cytokine signaling (SOCS) in rainbow trout hepatocytes. *Dev Comp Immunol.* (2012) 38:360–7. doi: 10.1016/j.dci.2012.07.005

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Jiang, He, Bai, Chan and Wong. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.