



Importance of Lung Epithelial Injury in COVID-19-associated Acute Respiratory Distress Syndrome: Value of Plasma Soluble Receptor for Advanced Glycation End-Products

To the Editor:

The respiratory form of coronavirus disease (COVID-19) has led to an unprecedented number of hospitalizations for acute respiratory distress syndrome (ARDS). To date, the pathophysiology of COVID-19-associated ARDS (CARDS) remains poorly understood. This has led to discussion about a different presentation from non-COVID-19 ARDS, regarding lung mechanics abnormalities and hypoxemia mechanisms (1, 2).

However, little attention has been paid to the value of biomarkers of lung injury. The soluble form of the receptor for advanced glycation end-products (sRAGE) is a well-characterized marker of lung alveolar epithelial injury (3) and has been associated with both prognostic and pathogenic values in patients with ARDS (4).

This study aims to investigate the value of baseline plasma sRAGE in CARDS and how it could differ between COVID-19 and non-COVID-19 ARDS.

Patients and Methods

We prospectively enrolled all consecutive adult patients admitted to the medical ICU of the Saint-Louis hospital, Paris, France, between March 1 and June 1, 2020, for CARDS according to the Berlin definition (5). This study was approved by the Ethics Committee of the Société de Réanimation de Langue Française (CE SRLF no. 20-32).

Management of patients included protective volume-controlled ventilation, neuromuscular blockers, and prone position if needed. All measurements were performed within 24 hours after intubation. Ventilator settings and respiratory mechanics measures were collected, together with dead space fraction, ventilatory ratio, and shunt fraction. When available, measurements of the recruitment-to-inflation ratio were collected (6). A value ≤ 0.5 was considered as a potential for lung recruitment.

The severity of lung edema was assessed using the Radiographic Assessment of Lung Edema (RALE) score, evaluated by two independent physicians on the chest radiography of the day of mechanical ventilation (MV) initiation.

This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern (dgern@thoracic.org).

Supported by the Université de Paris.

Author Contributions: G.D., J.-M.C., E.Y., and N.K. designed and performed research. G.D. and J.-M.C. analyzed the data. G.D., N.K., and J.-M.C. wrote the manuscript. G.D., J.-M.C., E.Y., N.K., G.M., V.L., M.D., C.d.M.-M., M.J., and E.A. collected the data. All authors approved the final manuscript.

Originally Published in Press as DOI: 10.1164/rccm.202104-1070LE on May 25, 2021

Levels of plasma sRAGE were measured in duplicate from thawed samples collected within 24 hours after MV initiation. A commercially available sandwich enzyme immunoassay kit (Human sRAGE Quantikine ELISA Kit; R&D Systems) was used following recommendations from the manufacturer.

Patients with CARDS were then compared with a historical multicentric prospective cohort of patients with ARDS in whom plasma sRAGE had been measured (7) and with control patients (e.g., mechanically ventilated patients without COVID-19 infection or ARDS, $n = 15$).

Continuous variables are described as medians (interquartile ranges) and compared using the Wilcoxon's rank sum test or the Kruskal-Wallis test; categorical variables are summarized by counts (percentages) and compared using the Fisher exact test. Correlations were assessed with the Rho Spearman's correlation test. Prognosis value of sRAGE on Day-90 mortality was assessed using Cox model adjusted on potential confounders (e.g., ARDS etiology, cardiovascular risk factors, body mass index, driving pressure, and $\text{PaO}_2/\text{FiO}_2$).

All tests were two-sided and P values $< 5\%$ were considered to indicate significant associations. Analyses were performed using R statistical platform, version 3.0.2.

Results

Characteristics of patients with COVID-19 ARDS. Overall, 50 patients with CARDS (median [interquartile range], 62.0 [54.0–68.7] yr of age; 68% male) were included. Median time from symptoms onset to invasive MV initiation was 9.0 (7.0–14.0) days. For further details, see Table 1.

Baseline plasma sRAGE correlates with lung injury severity and outcome in COVID-19. At baseline, plasma sRAGE was 4,044.0 (1,763.0–4,768.0) pg/ml and significantly differed from control (525.0 [411.0–638.5] pg/ml; $P < 0.001$; Figure 1).

Baseline plasma sRAGE correlated with $\text{PaO}_2/\text{FiO}_2$ (Spearman's $\rho = -0.49$; $P = 0.001$), ventilatory ratio ($\rho = 0.36$; $P = 0.019$), shunt ($\rho = 0.39$; $P = 0.01$), and RALE score (median score, 28 [18–36]; $\rho = 0.64$; $P < 0.01$).

The recruitment-to-inflation ratio was measured in 16 patients (32%) and high potential for recruitability was observed in 6 (37.5%). Plasma sRAGE levels were higher in patients with high potential for recruitability (4,245.0 [3,795.0–4,854.0] pg/ml vs. 2,890.0 [2,312.0–3,566.0] pg/ml; $P = 0.02$).

Of note, baseline plasma sRAGE was significantly higher in Day-90 decedents than in survivors (4,403.1 [2,564.0–4,990.2] pg/ml vs. 2,708.0 [1,965.9–4,304.5] pg/ml; $P = 0.04$).

Comparison between patients with CARDS and those with non-COVID-19 ARDS. Compared with patients with non-COVID-19 ARDS, patients with CARDS were significantly different with regard to body mass index, cardiovascular risk factors, and incidence of ARDS severity at Day 1 (Table 1). Median static compliance of respiratory system was similar between patients with ARDS with or without COVID-19 (29.5 [26.2–35.0] vs. 28.6 [21.9–34.5] ml/cm H_2O , respectively; $P = 0.17$).

Baseline sRAGE levels were significantly higher in CARDS compared with non-COVID-19 ARDS (4,044.0 [1,763.0–4,768.0] pg/ml vs. 2,230.0 [1,156.0–3,954.0] pg/ml; $P = 0.005$; Figure 1). Overall, Day-90 mortality rate was 54% in CARDS and 36% in




Figure 1. Value of plasma soluble receptor for advanced glycation end-products (sRAGE) levels at baseline in patients with coronavirus disease (COVID-19) or non-COVID-19 acute respiratory distress syndrome (ARDS). (A) Comparison of sRAGE levels between survivors and nonsurvivors in patients with COVID-19 ARDS. (B) Correlations between baseline plasma sRAGE and RALE score in patients with COVID-19 ARDS. (C) Plasma sRAGE levels in a subset of 16 patients with measurement of R/I ratio available on the day of mechanical ventilation initiation. (D) Comparison of plasma sRAGE levels in patients with COVID-19, patients with non-COVID-19 ARDS, and control patients. Correlations have been tested with the calculation of Spearman's rank correlation coefficient R (ρ). RALE = Radiographic Assessment of Lung Edema; R/I = recruitment/inflation.

non-COVID-19 ARDS ($P = 0.045$). Adjusted on potential confounders, baseline plasma sRAGE levels were significantly associated with mortality (adjusted hazard ratio, 1.51 [1.05–2.16] per one log increment; $P = 0.02$).

Discussion

Whether CARDS-related lung injury is similar to that from other causes of ARDS is an important question. The answer may guide the ventilatory strategy and carry some prognostic information. Using a well-characterized marker of lung epithelial injury, this study suggests that CARDS includes a component of pulmonary alveolar damage higher than other causes of ARDS. Moreover, as in non-COVID-19 ARDS, plasma sRAGE is associated with CARDS severity and outcome, especially lung edema, assessed by baseline RALE score and oxygenation impairment.

Since the onset of the pandemic, CARDS has been suggested to be an atypical subset of ARDS (2, 8). This assertion has been recently challenged, mainly through comparisons of lung mechanics parameters (9). Although sRAGE production could

have several sources, numerous works have provided evidence that alveolar type I cells are the main source of plasma sRAGE, and that sRAGE is a reliable marker of diffuse lung alveolar injury and impaired fluid clearance in both clinical and experimental models of ARDS (3). In this study, we found a marked elevation in sRAGE levels among patients with CARDS, which argues for intense lung epithelial injury. This is consistent with recent pathological reports from postmortem lung biopsies, in which diffuse alveolar damage was the most common histological finding (10).

This study has some limitations. First, the limited number of patients from a single center requires additional data to confirm this hypothesis. Second, plasma sRAGE was only measured at baseline and the value of changes over time is unknown.

In summary, our findings suggest that lung epithelial injury, as reflected by plasma sRAGE, may be a key pathophysiological feature with prognostic information in CARDS. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Table 1. Comparisons between Patients with COVID-19–related ARDS and Patients with ARDS from Other Causes

	COVID-19 ARDS (N = 50)	Non-COVID-19 ARDS (N = 117)	P Value
Demographic			
Age, yr	62.0 (54.0–68.7)	60.0 (45.0–70.0)	0.38
Sex, M	34 (68)	80 (68)	1.00
BMI, kg/m ²	27.7 (24.3–30.7)	25.9 (22.2–28.5)	0.008
Hypertension	27 (54)	38 (32)	0.015
Diabetes	18 (36)	22 (19)	0.029
Dyslipidemia	24 (48)	17 (15)	<0.0001
At least one cardiovascular risk factor	32 (64)	54 (46)	0.052
SAPS II	45.0 (36.0–56.0)	49.0 (39.0–64.0)	0.10
ARDS cause			<0.05
Pulmonary	50 (100)	85 (73)	
Lung infection	50 (100)	85 (100)	
Extrapulmonary	—	27 (27)	
Intraabdominal infection	—	22 (81)	
Acute pancreatitis	—	5 (19)	
ARDS severity			0.001
Mild	13 (26)	8 (7)	
Moderate	24 (48)	56 (48)	
Severe	13 (26)	52 (45)	
Respiratory parameters, Day 1			
V _T , ml/kg PBW	6.0 (6.0–6.17)	6.6 (6.0–7.3)	<0.0001
Pplat, cm H ₂ O	23.0 (21.0–25.0)	28.0 (24.0–30.0)	<0.0001
PEEP, cm H ₂ O	10.0 (8.0–12.0)	10.0 (8.0–13.0)	0.57
Crs, ml/cm H ₂ O	29.5 (26.2–35.0)	28.6 (21.9–34.5)	0.17
Ventilatory ratio	1.6 (1.4–1.9)	2.0 (1.7–2.4)	<0.0001
Biological data			
Pa _{O₂} /Fi _{O₂} , mm Hg	126 (99.25–199.8)	107.2 (71.11–147.2)	0.005
pH	7.38 (7.34–7.42)	7.35 (7.27–7.40)	0.008
Pa _{CO₂} , mm Hg	40.5 (36.9–45.8)	44.0 (37.7–50.0)	0.070
Baseline plasma sRAGE, pg/ml	4,044.0 (1,763.0–4,768.0)	2,230.0 (1,156.0–3,954.0)	0.005
Treatments			
Prone position use	28 (56)	24 (21)	<0.0001
NO therapy	4 (8)	33 (28)	0.007
VV-ECMO	4 (8)	2 (2)	0.11
Outcomes			
Duration of MV, d	12.0 (4.0–17.0)	11.0 (6.0–20.0)	0.31
ICU LOS, d	14.0 (10.0–22.0)	18.0 (10.0–34.2)	0.063
In-ICU mortality	27 (54)	38 (33)	0.016
Day-90 mortality	27 (54)	42 (36)	0.045

Definition of abbreviations: ARDS = acute respiratory distress syndrome; BMI = body mass index; COVID-19 = coronavirus disease; Crs = static compliance of respiratory system; LOS = length of stay; MV = mechanical ventilation; NO = nitric oxide; PBW = predicted body weight; PEEP = positive end-expiratory pressure; Pplat = inspiratory plateau pressure; SAPS II = simplified acute physiology score; sRAGE = soluble receptor for advanced glycation end-products; VV-ECMO = venovenous extracorporeal membrane oxygenation. Results are presented as *n* (%) or median (interquartile range).

Natacha Kapandji, M.D.
Hôpital Saint-Louis
Paris, France

and
Université de Paris
Paris, France

Elise Yvin, M.D.
Hôpital Saint-Louis
Paris, France

Constance de Margerie-Mellon, M.D., Ph.D.
Hôpital Saint-Louis
Paris, France

and
Université de Paris
Paris, France

and
Université de Paris
Paris, France

Magali Devriese, M.D.
Hôpital Saint-Louis
Paris, France

Giulia Moratelli, M.D.
Virginie Lemiale, M.D.
Hôpital Saint-Louis
Paris, France



IgG Levels and Mortality in Chronic Obstructive Pulmonary Disease

To the Editor:

Chronic obstructive pulmonary disease (COPD) is a major cause of morbidity and mortality worldwide (1). Most of the poor outcomes of COPD occur during acute exacerbations (AECOPD) (2), which are frequently triggered by a respiratory tract infection (3). IgG deficiency has long been identified as a cause of recurrent respiratory tract infections and is thought to be a risk factor for developing chronic respiratory diseases such as idiopathic bronchiectasis (4). Recently, we showed that serum IgG deficiency is a significant risk factor for AECOPD and related hospitalizations (1). Here, we examined the relationship between serum IgG levels and 1-year mortality in patients with COPD. Some of the results of this study have been previously reported in an abstract (5).

Methods

All patients provided written informed consent. The study was approved by the University of British Columbia Providence Health Care Research Ethics Board (certificate number H11-00786) for patients enrolled at St. Paul's Hospital, Vancouver, Canada, and the University of British Columbia Clinical Research Ethics Board (certificate number H13-00790) for patients enrolled at Vancouver General Hospital, Vancouver, Canada. Written informed consent was provided by each participant in accordance with the Ethics Board. We collected blood samples from the Rapid Transition Program (clinicaltrials.gov identifier NCT02050022). Details of the study, including its inclusion and exclusion criteria, have been published previously (3). Briefly, the Rapid Transition Program included patients hospitalized with AECOPD ($n = 489$) and clinically stable patients, who were recruited from a COPD clinic in the same hospital ($n = 132$). None had significant bronchiectasis either by history or by thoracic computed tomography scan. Samples were collected and were processed per standardized protocol and stored at -80°C . Serum IgG levels were measured via liquid chromatography–tandem mass spectrometry as previously described (6). IgG measurements were processed in the clinical laboratory at St. Paul's Hospital, Vancouver, British Columbia, Canada. After enrollment, patients were followed for 1 year, during which their vital status was ascertained through hospital records, which were validated by death certificates. Cox regression modeling, which adjusted

Supported by Genome Canada, Genome British Columbia, Genome Quebec, Canadian Institutes of Health Research, Providence Health Care, and St. Paul's Foundation.

Author Contributions: N.M.A. had full access to all the data in the study, takes responsibility for the integrity of the data and the accuracy of the data analysis, and contributed to drafting the manuscript. D.D.S. contributed substantially to the study design, data analysis, and interpretation and had authority over manuscript preparation and the decision to submit the manuscript for publication. All authors contributed to the critical revision of the manuscript for important intellectual content and gave final approval of the version to be published.

Originally Published in Press as DOI: 10.1164/rccm.202102-0382LE on May 4, 2021

Matthieu Jabaudon, M.D., Ph.D.
CHU ClermontFerrand and Université Clermont Auvergne
ClermontFerrand, France

Elie Azoulay, M.D., Ph.D.
Hôpital Saint-Louis
Paris, France

and
Université de Paris
Paris, France

Jean-Michel Constantin, M.D., Ph.D.
Sorbonne University and PitiéSalpêtrière Hospital
Paris, France

Guillaume Dumas, M.D., PhD.*
Hôpital Saint-Louis
Paris, France

and
Université de Paris
Paris, France

ORCID ID: 0000-0002-8121-1680 (M.J.).

*Corresponding author (e-mail: dumas.guillaume1@gmail.com).

References

- Gattinoni L, Chiumello D, Caironi P, Busana M, Romitti F, Brazzi L, *et al*. COVID-19 pneumonia: different respiratory treatments for different phenotypes? *Intensive Care Med* 2020;46:1099–1102.
- Chiumello D, Busana M, Coppola S, Romitti F, Formenti P, Bonifazi M, *et al*. Physiological and quantitative CT-scan characterization of COVID-19 and typical ARDS: a matched cohort study. *Intensive Care Med* 2020;46:2187–2196.
- Jabaudon M, Blondonnet R, Roszyk L, Bouvier D, Audard J, Clairefond G, *et al*. Soluble receptor for advanced glycation end-products predicts impaired alveolar fluid clearance in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2015;192:191–199.
- Jabaudon M, Blondonnet R, Pereira B, Cartin-Ceba R, Lichtenstern C, Mauri T, *et al*. Plasma sRAGE is independently associated with increased mortality in ARDS: a meta-analysis of individual patient data. *Intensive Care Med* 2018;44:1388–1399.
- ARDS Definition Task Force; Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, *et al*. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526–2533.
- Chen L, Del Sorbo L, Grieco DL, Junhasavasdikul D, Rittayamai N, Soliman I, *et al*. Potential for lung recruitment estimated by the recruitment-to-inflation ratio in acute respiratory distress syndrome: a clinical trial. *Am J Respir Crit Care Med* 2020;201:178–187.
- Mrozek S, Jabaudon M, Jaber S, Paugam-Burtz C, Lefrant J-Y, Rouby J-J, *et al*; Azorea network. Elevated plasma levels of sRAGE are associated with nonfocal CT-based lung imaging in patients with ARDS: a prospective multicenter study. *Chest* 2016;150:998–1007.
- Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S, Chiumello D. COVID-19 does not lead to a "typical" acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2020;201:1299–1300.
- Grasselli G, Tonetti T, Protti A, Langer T, Girardis M, Bellani G, *et al*. Pathophysiology of COVID-19-associated acute respiratory distress syndrome: a multicentre prospective observational study. *Lancet Respir Med* 2020;8:1201–1208.
- Bradley BT, Maioli H, Johnston R, Chaudhry I, Fink SL, Xu H, *et al*. Histopathology and ultrastructural findings of fatal COVID-19 infections in Washington State: a case series. *Lancet* 2020;396:320–332.

Copyright © 2021 by the American Thoracic Society