


 Cite this: *RSC Adv.*, 2021, 11, 19310

 Received 24th April 2021  
 Accepted 16th May 2021

DOI: 10.1039/d1ra03209g

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

# Me<sub>3</sub>SI-promoted chemoselective deacetylation: a general and mild protocol†

 Aakanksha Gurawa,‡ Manoj Kumar‡ and Sudhir Kashyap \*


A Me<sub>3</sub>SI-mediated simple and efficient protocol for the chemoselective deprotection of acetyl groups has been developed *via* employing KMnO<sub>4</sub> as an additive. This chemoselective deacetylation is amenable to a wide range of substrates, tolerating diverse and sensitive functional groups in carbohydrates, amino acids, natural products, heterocycles, and general scaffolds. The protocol is attractive because it uses an environmentally benign reagent system to perform quantitative and clean transformations under ambient conditions.

The divergent protection–deprotection approach and extensive functional group manipulation continues to serve as an important chemical tool to access biologically potent molecules and complex natural products.<sup>1</sup> In particular, the hydroxyl moieties and their derivatives are ubiquitous in natural products and recognized as renowned scaffolds because of the overwhelming number of chemical and biological applications that they can be used in.<sup>2,3</sup> The selective and orthogonal protection–deprotection of free hydroxyl group(s) are significant chemical transformations and frequently employed in target-oriented synthesis (TOS).<sup>4</sup> Due to the widespread use and relative ease of protection–deprotection, introduction of an *O*-Ac group to mask the hydroxyl-moiety has remained as a highly reliable and convenient strategy, especially in synthetic carbohydrate chemistry.<sup>1–3</sup> However, the chemoselective deprotection of *O*-Ac in the presence of analogous and sensitive *O*-protective groups such as benzoyl (Bz) or pivaloyl (Pv) is a notoriously challenging yet important task.<sup>1–3</sup>

Thus, considerable effort has been devoted to developing robust and selective methods for the deprotection of the acetate ester. As outlined in Scheme 1, the cleavage of *O*-Ac is conventionally performed under a homogeneous reagent system including: (a) Brønsted acids/Lewis acids such as HCl/MeOH,<sup>5</sup> HBF<sub>4</sub>·Et<sub>2</sub>O,<sup>6</sup> BF<sub>3</sub>·Et<sub>2</sub>O,<sup>7</sup> *p*-TsOH or CSA,<sup>8</sup> (b) inorganic/organic basic conditions employing Zemplén hydrolysis (NaOMe/MeOH),<sup>1</sup> ammonia solution,<sup>9</sup> hydrazine/AcOH/pyridine,<sup>10</sup> DBU/PhH,<sup>11</sup> guanidine/EtOH/DCM,<sup>12</sup> Mg–metal or Mg(OMe)<sub>2</sub>,<sup>13</sup> KCN/EtOH,<sup>14</sup> K<sub>2</sub>CO<sub>3</sub>/MeOH/H<sub>2</sub>O,<sup>15</sup> NaHCO<sub>3</sub>/H<sub>2</sub>O,<sup>16</sup> (c) metallic compounds or oxidants such as MoO<sub>2</sub>Cl<sub>2</sub>,<sup>17</sup> molecular iodine/

MeOH,<sup>18a</sup> Sm/I<sub>2</sub>,<sup>18b</sup> Bu<sub>2</sub>SnO/heating,<sup>19a</sup> Bu<sub>3</sub>SnOMe/DCE.<sup>19b</sup> Although, the use of heterogeneous catalysts such as CuFe<sub>2</sub>O<sub>4</sub> nanoparticles<sup>20</sup> and enzymes<sup>21</sup> have also been demonstrated for deacetylation with limited substrate scope. Recently, a tetranuclear zinc cluster, Zn<sub>4</sub>(OCOFC<sub>3</sub>)<sub>6</sub>O has been investigated for use in trans-esterification and deacylation with discriminate selectivity.<sup>22</sup>

Despite the synthetic challenges and demands placed on modern chemistry, advancing green chemical syntheses employing milder and environmentally benign reagent systems remains a constant motivation and is actively pursued. In this context and our continued interest<sup>23</sup> developing versatile and selective protocols inspired the development of a simple and chemoselective deacetylation method with remarkable and distinctive synthetic applicability. Recently the dual-reactivity of KMnO<sub>4</sub> for selective deacetylation and one-pot deacetylation–oxidation of benzyl-*O*-acetates under controlled reaction conditions was successfully investigated.<sup>23a</sup>



**Scheme 1** Previous advances, and the Me<sub>3</sub>SI-catalyzed chemoselective deacetylation developed in this work.

Carbohydrate Chemistry Research Laboratory (CCRL), Department of Chemistry, Malaviya National Institute of Technology Jaipur, (MNIT), Jaipur-302017, India. E-mail: [skashyap.chy@mnit.ac.in](mailto:skashyap.chy@mnit.ac.in); [skr.kashyap@gmail.com](mailto:skr.kashyap@gmail.com)

† Electronic supplementary information (ESI) available: General procedures, characterization data, and spectral charts. See DOI: 10.1039/d1ra03209g

‡ Aakanksha Gurawa and Manoj Kumar contributed equally.



Emerging from these precedents, the Me<sub>3</sub>SI-catalyzed, simple and chemoselective removal of *O*-acetate, an elegant and promising alternative approach for the deprotection of the acetate ester (Scheme 1d) is presented herein. Inspired by recent research,<sup>23f</sup> an initial experiment employing Me<sub>3</sub>SI(OAc)<sub>2</sub>, generated *in situ* oxidative transfer of the acetyl groups from PhI(OAc)<sub>2</sub> to Me<sub>3</sub>SI, which gave the regioselective 2-iodoglycosylation of enol ether functionality in 3,4,6-tri-*O*-acetyl-*D*-glucal (**1a**) with methanol together with traces of the deacetylated product **1** (Table 1, entry 1). The preliminary results led to the exploration of the possibility of using sulfonium iodate salts in the selective removal of the *O*-acetyl protection group. As summarized in Table 1, subsequent optimization of the reaction was performed by altering the reagents and solvent systems. Accordingly, the reactions of TOAc-glucal **1a** using varying amounts of Me<sub>3</sub>SI and NaIO<sub>4</sub> in the presence of MeOH facilitated the smooth deacetylation providing the desired product **1** with excellent efficiency under ambient conditions (Table 1, entries 2–5). It is worth noting, that the use of Me<sub>3</sub>SI in combination with NaIO<sub>4</sub> in MeOH as the solvent, promoted the deprotection of acetates in a chemoselective manner, however the direct 1,2-addition product, 2-deoxy-glycoside was not

detected.<sup>23f,g</sup> Meanwhile, the reaction using 0.2 equiv. of both Me<sub>3</sub>SI and NaIO<sub>4</sub> was found to suffice for obtaining the complete deacetylation of **1a** after 40 min (Table 1, entry 4).


Consequently, TOAc-glucal **1a** was subjected to various complementary halide salts employing NaIO<sub>4</sub> (0.1 equiv.) in MeOH under ambient reaction conditions (Table 1, entries 6–12). Use of the catalytic *n*-Bu<sub>4</sub>NI and KI provided the desired product with good to modest transformation, 90% and 50% yields, respectively (Table 1, entries 6 and 7 vs. 5). It was found that NaI can efficiently promote the deacetylation in improved yields (90%) after 16 h under the present conditions (Table 1, entry 8). In contrast, Me<sub>3</sub>SOI and bromide salts such as Me<sub>3</sub>SBr, *n*-Bu<sub>4</sub>NBr, and KBr failed to deliver the deacetylated product even after increasing the amount of the salts and additive NaIO<sub>4</sub> to 1.0 equiv. and a prolonged reaction time (Table 1, entries 9–12). Remarkably, switching to KMnO<sub>4</sub> (0.1 equiv.) as the additive in a Me<sub>3</sub>SI-catalyzed reaction led to simple deacetylation of **1a** with almost quantitative yields (100%), within 5 min (Table 1, entry 13). A rapid screening of common organic solvents including CH<sub>3</sub>CN, toluene, THF or CH<sub>2</sub>Cl<sub>2</sub> were unsuccessful and did not give any desired transformations (Table 1, entries 14–17). Further optimization employing analogous additives such as K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, oxone, KBrO<sub>3</sub>, NaBO<sub>3</sub>·H<sub>2</sub>O, and Na<sub>3</sub>BO<sub>3</sub> were found to be too sluggish or resulted in poor conversions (Table 1, entries 18–22).

Having improved and optimized the reagent system, the generality and limitation of Me<sub>3</sub>SI-catalyzed selective deacetylation protocol was investigated next, and the results are summarized in Scheme 2. Of the particular note, diverse and commonly used *O*-protecting groups in *D*-glucal comprising benzoyl (**1b**), benzyl (**1c**), methyl (**1d**), and *tert*-butyldimethylsilyl ether (**1e**) were fully tolerated, thus unambiguously establishing the potential synthetic applicability of the representative chemoselective procedure. Subsequently, the selective deacetylation of several per-*O*-acetylated glycals: *D*-galactal (**2a**), *D*-rhamnal (**3a**), *L*-rhamnal (**4a**), *D*-xylal (**5a**), *D*-arabinal (**6a**) and *L*-arabinal (**7a**), also disaccharide derived glycals bearing 1,4-glycosidic linkages, for example *D*-lactal (**8a**) and *D*-maltal (**9a**), were performed successfully to obtain the corresponding products **2–9** in excellent yields.

Furthermore, the substrates bearing *O*-TIPS (**10a**), *O*-Bn (**11a–13a**), and 4,6-*O*-benzylidene (**14a**) in carbohydrate substrates were uniformly sustained under selective deacetylation conditions giving the hydroxyl products **10–14**, respectively, (up to 98% yields). Indeed, the glycosides comprising sensitive *O*-isopropylidene linkages in *D*-galactopyranose (**15a**), *D*-glucofuranose (**16a**), *D*-ribose (**17a**), and unprotected uridine (**18a**) were selectively deacetylated to generate the corresponding free-hydroxyl compounds **15–18** in good yields. Importantly, the amino acid derivatives **19a–23a** including relatively subtle or labile substituents such as *tert*-butylcarbamate (Boc), fluorenylmethoxycarbonyl (Fmoc) carbobenzoxy (Cbz), and methyl ester (–CO<sub>2</sub>Me), underwent efficient chemoselective deacetylation giving the desired products **19–23** in good to excellent yields.

Encouraged by these results, the scope of the selective protocol was extended by performing the systematic


**Table 1** The screening and optimization of the deacetylation reagent system<sup>a</sup>



Entry	Catalyst (equiv.)	Additive (equiv.)	Solvent	Time	Yield <sup>b</sup>
1	Me <sub>3</sub> SI (1.0)	PhI(OAc) <sub>2</sub> (1.0)	MeOH	24 h	Trace
2	Me <sub>3</sub> SI (1.0)	NaIO <sub>4</sub> (1.0)	MeOH	10 min	100%
3	Me <sub>3</sub> SI (0.4)	NaIO <sub>4</sub> (0.4)	MeOH	30 min	100%
4	Me <sub>3</sub> SI (0.2)	NaIO <sub>4</sub> (0.2)	MeOH	40 min	100%
5	Me <sub>3</sub> SI (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	95%
6	<i>n</i> -Bu <sub>4</sub> NI (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	90%
7	KI (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	50%
8	NaI (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	16 h	90%
9 <sup>c</sup>	Me <sub>3</sub> SOI (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	NR
10 <sup>c</sup>	Me <sub>3</sub> SBr (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	NR
11 <sup>c</sup>	<i>n</i> -Bu <sub>4</sub> NBr (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	NR
12 <sup>c</sup>	KBr (0.1)	NaIO <sub>4</sub> (0.1)	MeOH	12 h	NR
13	<b>Me<sub>3</sub>SI (0.1)</b>	<b>KMnO<sub>4</sub> (0.1)</b>	<b>MeOH</b>	<b>5 min</b>	<b>100%</b>
14	Me <sub>3</sub> SI (0.1)	KMnO <sub>4</sub> (0.1)	MeCN	24 h	NR
15	Me <sub>3</sub> SI (0.1)	KMnO <sub>4</sub> (0.1)	Toluene	24 h	NR
16	Me <sub>3</sub> SI (0.1)	KMnO <sub>4</sub> (0.1)	THF	24 h	NR
17	Me <sub>3</sub> SI (0.1)	KMnO <sub>4</sub> (0.1)	DCM	24 h	NR
18	Me <sub>3</sub> SI (0.1)	K <sub>2</sub> S <sub>2</sub> O <sub>8</sub> (0.1)	MeOH	12 h	Trace
19	Me <sub>3</sub> SI (0.1)	Oxone (0.1)	MeOH	12 h	Trace
20	Me <sub>3</sub> SI (0.1)	KBrO <sub>3</sub> (0.1)	MeOH	12 h	Trace
21	Me <sub>3</sub> SI (0.1)	NaBO <sub>3</sub> ·H <sub>2</sub> O (0.1)	MeOH	12 h	40%
22	Me <sub>3</sub> SI (0.1)	Na <sub>3</sub> BO <sub>3</sub> (0.1)	MeOH	12 h	Trace

<sup>a</sup> Reaction conditions: **1a** (1.0 equiv.), salt (0.1 to 1.0 equiv.), additive (0.1 to 1.0 equiv.), solvent (1 mL) in open-air at room temperature.


<sup>b</sup> The isolated and unoptimized yields, based on the starting material **1a**. <sup>c</sup> Increasing the amount of both salt and additive up to 1.0 equiv., 24 h. NR = no reaction.



**Scheme 2** The study of the substrate scope and a functional group compatibility investigation for the chemoselective deacetylation. Reaction conditions: **1a**–**23a** (1.0 equiv.), Me<sub>3</sub>SI (0.1 equiv.), additive (0.1 equiv.), MeOH (1 mL), 25 °C under atmospheric pressure, 1–12 h. The isolated and unoptimized yields are shown.

deacetylation of substrates involving benzylic, alkylic, allylic, phenolic, naphthyl, heterocyclic, alicyclic, spirocyclic, natural-products, long chain aliphatic, and aminoxy moieties. As illustrated in Scheme 3, a series of substrates enclosing comparatively susceptible and electronically diverse *O*-protecting groups (*O*-Ac; **24a** vs. *O*-Piv; **24b**, *O*-Boc; **24c**, *O*-Ms; **24d**, *O*-THP; **24e**, *O*-Tr; **24f**) were well protected under present conditions, further confirming the effectiveness of the chemoselective method. The Me<sub>3</sub>SI-catalyzed simple removal of acetyl groups, in menthyl (**25a**), geranyl (**26a**), cholesteryl (**27a**), and diversely substituted aryl substrates **28a**–**35a**, was conducted to efficiently obtain the desired products **25**–**35** in good-to-excellent yields (90–99%). It is worth noting, that the selective deacetylation for a substrate consisting of two electronically different acetates was accomplished successfully (cleavage of *O*-acetate vs. *N*-acetate in **29a**).

Next, the diversely substituted phenolic or naphthyl *O*-acetates **36a**–**42a**, eugenyl **43a**, vanillyl **44a**, and estrone acetate **45a** were subjected, under ambient conditions, to access the corresponding free-phenols **36**–**45** in acceptable yields (72–96%). Likewise, the reaction employing heterocyclic furfuryl acetate (**46a**), 2-amino-(Cbz)-ethyl acetate (**47a**), and alicyclic substrates with different ring sizes (five, six, or seven) **48a**–**50a** proceeded smoothly providing the corresponding products **46**–**50** (80–99% yields). Additionally, long chain aliphatic acetates **51a** and **52a**, a spirocycle bearing sensitive dioxo moiety **53a**,



**Scheme 3** Extended scope of the Me<sub>3</sub>SI-catalyzed selective deacetylation. Reaction conditions: **24a**–**55a** (1.0 equiv.), Me<sub>3</sub>SI (0.1 equiv.), additive (0.1 equiv.), MeOH (1 mL), 25 °C under atmospheric pressure, 1–12 h. The isolated and unoptimized yields are shown.

sterically hindered adamantyl **54a**, and aminoxy-*O*-acetate **55a** were competent, when used in the applied protocol, of producing their corresponding hydroxyls **51**–**55** in satisfactory yields (90–99%).

To gain further insight into the Me<sub>3</sub>SI-promoted deacetylation, the reaction of TOAc-glucal **1a** was performed with equimolar amounts of trimethylsulfonium iodide and an additive (0.1 equiv. each) employing deuterated methanol (CD<sub>3</sub>OD) as the solvent (Fig. 1). The <sup>1</sup>H-NMR analyses of the crude reaction mixture revealed the presence of a distinctive resonance which was due to an anomeric proton ( $\delta$  6.42 dd,  $J_{1-2}$  = 6.10, 1.4 Hz, H-1) and other characteristic chemical shifts which conformed with that of the desired product **1'** (tri-*O*-<sup>2</sup>H-D-glucal).

With no variance, the acetyl protons of methyl acetate (CH<sub>3</sub>CO-OCD<sub>3</sub>) were found at  $\delta$  2.10 (s, 9H) which potentially resulted from the methanolysis of the acetate groups with CD<sub>3</sub>OD. In addition, substantial amounts of trimethylsulfonium hydroxide species were observed consistently at  $\delta$  3.05 as a singlet (vs. Me<sub>3</sub>SI at  $\delta$  1.56),<sup>24</sup> and this established the role of trimethylsulfonium iodide in selective and simple cleavage of acetates. A set of control experiments were performed to




Fig. 1 Experiments with deuterated methanol and  $^1\text{H}$ -NMR analyses, used to provide mechanistic rationalization for  $\text{Me}_3\text{SI}$ -promoted deacetylation.

identify the possible intermediate iodine species in this transformation. The studies revealed the absence of molecular iodine in the present reaction conditions as confirmed by a negative starch solution test (the color did not change to an intense blue color).<sup>24</sup> Indeed, the standard  $\text{Me}_3\text{SI}$ -mediated deacetylation reaction employing a common radical scavenger such as TEMPO (a stable aminoxy radical) or 2,6-di-*tert*-butyl-4-methylphenol (BHT) preceded smoothly, and further ruled out a radical pathway in the process. Although, attempting a reaction of TOAc-glucal **1a** using a catalytic amount of iodine in methanol solvent was inconsistent (50–60% conversion) resulting in the mono-deacetylation of the primary acetate (6-OAc) even after a prolonged reaction time (48 h).<sup>18a</sup> It is worth noting that the characteristic Lewis acidity of molecular iodine ( $\text{I}_2$ ) has been exploited in the selective *O*-Ac as well *N*-Boc protection of diverse substrates.<sup>25</sup>

Based on the previously mentioned experimental observations and with reference to precedents in the literature, a plausible mechanistic rationalization for a  $\text{Me}_3\text{SI}$ -catalyzed reaction is postulated in Scheme 4. Initially,  $\text{KMnO}_4$  showed rapid disproportionation, resulting in permanganic acid ( $\text{HMnO}_4 \leftrightarrow \text{H}^+ + \text{MnO}_4^-$ ) and potassium methoxide ( $\text{KOMe} \leftrightarrow \text{K}^+ + \text{MeO}^-$ ) ions in the presence of methanol, which was attributed to the reducing behavior of methanol in the present system.<sup>26</sup> The main oxidizing reagent  $\text{Mn}(\text{VII})\text{O}_4^-$  was capable of inverting the

polarity of the reactivity of the halide salt  $\text{Me}_3\text{SI}$  in slightly acidic conditions which resulted in the iodate ( $\text{I}^+$ ) species [A], and was disproportionate to  $\text{Mn}(\text{IV})\text{O}_2$  and  $\text{H}_2\text{O}$ . The reduced  $\text{Mn}(\text{IV})$  species would then readily oxidize back to the higher  $\text{Mn}(\text{VI})\text{O}_4^-$  in a slightly basic medium ( $\text{KOMe}/\text{H}_2\text{O}$ ) under the normal atmosphere<sup>26c</sup> and this was likely to facilitate the release of trimethylsulfonium hydroxide. Meanwhile, the incipient electrophilic intermediate [A] initiated from the oxidation of  $\text{Me}_3\text{SI}$  coordinated with the oxygen atoms of the carbonyl functionality ( $\text{CH}_3\text{CO}_2^-$ ) in the acetate compound [1a] to generate a transient species [B].<sup>25b,25c,27</sup> This significantly induced the polarization of the  $\text{C}=\text{O}$  double bond and increased the electrophilicity of the carbonyl carbon as well contributing towards the bond lengthening of the carbonyl  $\text{C}=\text{O}$  moiety in [B].

Subsequent coordination of the  $\text{K}^+$  ion with alkyl-oxygen ( $\text{RO}^-$ ) and attack of methoxide ( $\text{MeO}^-$ ) as the nucleophile on the electrophilic carbon in [B], would then eventually produce the tetrahedral intermediate [C]. Following the elimination of the penultimate transient species [D] and simultaneously trapping of the respective alkoxide ion ( $\text{RO}^-$ ) in a hydrolytic environment would unambiguously lead to the formation of the respective alkanol ( $\text{ROH}$ ) [1] with the liberation of potassium methoxide. The proposed intermediate [D] finally displaced the corresponding methyl acetate ( $\text{CH}_3\text{CO}_2\text{Me}$ ) by releasing the required iodate species [A] which presumably participated in another catalytic cycle *en route* to the selective deacetylation process.

In summary, a catalytic and practical procedure for chemoselective deacetylation with a general substrate scope employing environmentally benign reagents under ambient reaction conditions is reported. It is worth noting that the catalytic protocol is broadly applicable to numerous substrates, including carbohydrates, amino acids, and natural products, tolerating orthogonal and sensitive groups (esters, ethers, silyl ethers, carbamates, carboxybenzyl, mesyl, 2-tetrahydropyranyl, trityl, aldehydes, and also alkenes). Furthermore, the method is advantageous as it involves a safe and convenient reaction, ensuring smooth and quantitative conversion as well as preventing trans-esterification.

## Experimental

### Representative procedure for chemoselective deacetylation

To a previously prepared solution of acetate substrate (50 mg, 1.0 equiv.) in MeOH (1 mL) was added  $\text{Me}_3\text{SI}$  (0.1 equiv.) and  $\text{KMnO}_4$  (0.1 equiv.). The mixture was stirred at room temperature in an open-air environment and the reaction progress was monitored by TLC. After the complete consumption of the starting material had occurred, typically 5 min to 12 h, the reaction mixture was filtered and washed with EtOAc (10 mL). The filtrate was treated with saturated aqueous  $\text{NaHCO}_3$  (5 mL), and the aqueous layer was extracted with EtOAc ( $3 \times 30$  mL). The combined organic layers were then washed with water and brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , and concentrated *in vacuo* to obtain the analytically pure products.

## Conflicts of interest

There are no conflicts to declare.

Scheme 4 A plausible mechanism for  $\text{Me}_3\text{SI}$ -promoted selective deacetylation.

## Acknowledgements

SK gratefully acknowledges the Department of Science and Technology, India for the Early Career Research Award and Research Grant for Young Scientists (SERB-ECR/2017/001477). SK thanks TEQIP-III for their generous support and MRC MNIT (Advanced Analytical and Characterization Centre) for providing the NMR spectroscopic and analytical data. The authors are also grateful to the Director of MNIT for providing the necessary infrastructure for this research. AG acknowledges a UGC Fellowship.

## Notes and references

- (a) T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Wiley, New York, 4th edn, 2006; (b) P. J. Kocienski, *Protecting Groups*, Thieme, New York, 1994.
- (a) K.-T. Huang, B.-C. Wu, C.-C. Lin, S.-C. Luo, C. Chen, C.-H. Wong and C.-C. Lin, *Carbohydr. Res.*, 2006, **341**, 2151–2155; (b) J. B. Schwarz, S. D. Kuduk, X.-T. Chen, D. Sames, P. W. Glunz and S. J. Danishefsky, *J. Am. Chem. Soc.*, 1999, **121**, 2662–2673; (c) X. Mao, P. Li, T. Li, M. Zhao, C. Chen, J. Liu, Z. Wang and L. Yu, *Chin. Chem. Lett.*, 2020, **31**, 3276–3278; (d) X. Xiao, Z. Shao and L. Yu, *Chin. Chem. Lett.*, 2021, DOI: 10.1016/j.cclet.2021.03.047.
- (a) J. Žemlička, J. Beránek and J. Smrť, *Collect. Czech. Chem. Commun.*, 1962, **27**, 2784–2795; (b) C. B. Reese and J. C. M. Stewart, *Tetrahedron Lett.*, 1968, **9**, 4273–4276; (c) V. Schwarz, *Collect. Czech. Chem. Commun.*, 1962, **27**, 2567–2574; (d) T. B. Winholdz and D. B. R. Johnston, *Tetrahedron Lett.*, 1967, **8**, 2555–2557; (e) B. Srinivas, T. R. Reddy and S. Kashyap, *Carbohydr. Res.*, 2015, **406**, 86–92; (f) T. R. Reddy, S. K. Battina and S. Kashyap, *J. Carbohydr. Chem.*, 2015, **34**, 133–144.
- (a) J. K. Baskin, *Chem. Rev.*, 2000, **100**, 4265–4266; (b) R. W. Dugger, J. A. Ragan and D. H. B. Ripin, *Org. Process Res. Dev.*, 2005, **9**, 253–258; (c) J. S. Carey, D. Laffan, C. Thomson and M. T. Williams, *Org. Biomol. Chem.*, 2006, **4**, 2337–2347.
- (a) V. Pozsgay, *J. Am. Chem. Soc.*, 1995, **117**, 6673–6681; (b) N. Yamamoto, T. Nishikawa and M. Isobe, *Synlett*, 1995, 505–506; (c) E. J. Corey, D. A. Clark, G. Goto, A. Marfat, C. Mioskowski, B. Samuelsson and C. Hammarström, *J. Am. Chem. Soc.*, 1980, **102**, 1436–1439; (d) N. E. Byramova, M. V. Ovchinnikov, L. Backinowsky and N. K. Kochetkov, *Carbohydr. Res.*, 1983, **124**, C8–C11.
- V. Pozsgay and B. Coxon, *Carbohydr. Res.*, 1993, **257**, 189–215.
- D. Askin, C. Angst and S. Danishefsky, *J. Org. Chem.*, 1987, **52**, 622–635.
- A. G. González, I. Brouard, F. León, J. I. Padrón and B. Jaime, *Tetrahedron Lett.*, 2001, **42**, 3187–3188.
- T. Neilson and E. S. Werstiuk, *Can. J. Chem.*, 1971, **49**, 493–499.
- Y. Ishido, N. Nakazaki and N. Sakairi, *J. Chem. Soc., Perkin Trans. 1*, 1979, 2088–2098.
- L. H. B. Baptistella, J. F. dos Santos, K. C. Ballabio and A. J. Marsaioli, *Synthesis*, 1989, 436–439.
- (a) N. Kunesch, C. Meit and J. Poisson, *Tetrahedron Lett.*, 1987, **28**, 3569–3572; (b) U. Ellervik and G. Magnusson, *Tetrahedron Lett.*, 1997, **38**, 1627–1628.
- (a) Y.-C. Xu, A. Bizuneh and C. Walker, *J. Org. Chem.*, 1996, **61**, 9086–9089; (b) Y.-C. Xu, E. Lebeau and C. Walker, *Tetrahedron Lett.*, 1994, **35**, 6207–6210; (c) Y.-C. Xu, A. Bizuneh and C. Walker, *Tetrahedron Lett.*, 1996, **37**, 455–458.
- J. Herzig, A. Nudelman, H. E. Gottlieb and B. Fischer, *J. Org. Chem.*, 1986, **51**, 727–730.
- (a) N. C. Barua and R. R. Schmidt, *Synthesis*, 1986, **11**, 891–893; (b) J. J. Plattner, R. D. Gless and H. Rapoport, *J. Am. Chem. Soc.*, 1972, **94**, 8613–8615.
- Q. Y. Zheng, L. G. Darbie, X. Cheng and C. K. Murray, *Tetrahedron Lett.*, 1995, **36**, 2001.
- C.-Y. Liu, H.-L. Chen, C.-M. Ko and C.-T. Chen, *Tetrahedron*, 2011, **67**, 872–876.
- (a) B. Ren, L. Cai, L.-R. Zhang, Z.-J. Yang and L.-H. Zhang, *Tetrahedron Lett.*, 2005, **46**, 8083–8086; (b) R. Yanada, N. Negoro, K. Bessho and K. Yanada, *Synlett*, 1995, 1261–1263.
- (a) M. G. Perez and M. S. Maier, *Tetrahedron Lett.*, 1995, **36**, 3311–3314; (b) A. Nudelman, J. Herzig, H. E. Gottlieb, E. Keinan and J. Sterling, *Carbohydr. Res.*, 1987, **162**, 145–152.
- J. E. Tasca, A. Ponzinibbio, G. Diaz, R. D. Bravo, A. Lavat and M. G. Gonzalez, *Top. Catal.*, 2010, **53**, 1087–1090.
- (a) K. Naemura, N. Takahashi and H. Chikamatsu, *Chem. Lett.*, 1988, **30**, 1717–1720; (b) C. R. Johnson and C. H. Senanayake, *J. Org. Chem.*, 1989, **54**, 735–736; (c) O. Houille, T. Schmittberger and D. Uguen, *Tetrahedron Lett.*, 1996, **37**, 625–628; (d) R. López, E. Montero, F. Sánchez, J. Cañada and A. Fernández-Mayoralas, *J. Org. Chem.*, 1994, **59**, 7027–7032; (e) E. W. Holla, V. Sinnwell and W. Klaffke, *Synlett*, 1992, 413–414; (f) T. Itoh, A. Uzu, N. Kanda and Y. Takagi, *Tetrahedron Lett.*, 1996, **37**, 91–92; (g) K. Takabe, N. Mase, T. Hisano and H. Yoda, *Tetrahedron Lett.*, 2003, **44**, 3267–3269; (h) T. Hisano, K. Onodera, Y. Toyabe, N. Mase, H. Yoda and K. Takabe, *Tetrahedron Lett.*, 2005, **46**, 6293–6295.
- T. W. Lin, A. K. Adak, H. J. Lin, A. Das, W. C. Hsiao, T. C. Kuan and C. C. Lin, *RSC Adv.*, 2016, **6**, 58749–58754.
- (a) A. Gurawa, M. Kumar, D. S. Rao and S. Kashyap, *New J. Chem.*, 2020, **44**, 16702–16707; (b) D. S. Rao, T. R. Reddy, A. Gurawa, M. Kumar and S. Kashyap, *Org. Lett.*, 2019, **21**, 9990–9994; (c) T. R. Reddy, D. S. Rao and S. Kashyap, *Chem. Commun.*, 2019, **55**, 2833–2836; (d) D. S. Rao, T. R. Reddy and S. Kashyap, *Org. Biomol. Chem.*, 2018, **16**, 1508–1518; (e) D. S. Rao, T. R. Reddy, K. Babachary and S. Kashyap, *Org. Biomol. Chem.*, 2016, **14**, 7529–7543; (f) T. R. Reddy, D. S. Rao, K. Babachary and S. Kashyap, *Eur. J. Org. Chem.*, 2016, 291–311; (g) G. Kundoor, D. S. Rao and S. Kashyap, *Asian J. Org. Chem.*, 2016, **5**, 264–270; (h) S. Chittela, T. R. Reddy, P. R. Krishna and S. Kashyap, *J. Org. Chem.*, 2015, **80**, 7108–7116; (i) T. R. Reddy, D. S. Rao

- and S. Kashyap, *RSC Adv.*, 2015, **5**, 28338–28343; (f) S. Chittela, T. R. Reddy, P. R. Krishna and S. Kashyap, *RSC Adv.*, 2014, **4**, 46327–46331.
- 24 See the ESI.†
- 25 (a) G. Basumatary and G. Bez, *Tetrahedron Lett.*, 2017, **58**, 4312–4315; (b) R. Varala, S. Nuvula and S. R. Adapa, *J. Org. Chem.*, 2006, **71**, 8283–8286; (c) K. P. R. Kartha and R. A. Field, *Tetrahedron*, 1997, **53**, 11753–11766.
- 26 (a) P. K. Sen, P. R. Samaddar and K. Das, *Transition Met. Chem.*, 2005, **30**, 261–267; (b) S. Dash, S. Patel and B. K. Mishra, *Tetrahedron*, 2009, **65**, 707–739, and references cited therein; ; (c) N. Singh and D. G. Lee, *Org. Process Res. Dev.*, 2000, **5**, 599–603.
- 27 The electrophilic iodine ( $I^+$ ) species generated in the proposed pathway likely to coordinate with oxygen atoms of carbonyl group under basic conditions. See: (a) W. A. Szarek, A. Zamojski, K. N. Tiwari and E. R. Ison, *Tetrahedron Lett.*, 1986, **27**, 3827–3830; (b) J. L. Wahlstrom and R. C. Ronald, *J. Org. Chem.*, 1998, **63**, 6021–6022.