

Research



Cite this article: van Lieshout SHJ, Badás EP, Mason MWT, Newman C, Buesching CD, Macdonald DW, Dugdale HL. 2020 Social effects on age-related and sex-specific immune cell profiles in a wild mammal. *Biol. Lett.* **16**: 20200234.

<http://dx.doi.org/10.1098/rsbl.2020.0234>

Received: 8 April 2020

Accepted: 25 June 2020

Subject Areas:

ecology, evolution, molecular biology

Keywords:

European badger, social effects, innate immunity, adaptive immunity, neutrophil, lymphocyte

Author for correspondence:

Sil H. J. van Lieshout

e-mail: sil.vanlieshout@gmail.com

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5050354>.

Social effects on age-related and sex-specific immune cell profiles in a wild mammal

Sil H. J. van Lieshout¹, Elisa P. Badás^{1,2}, Michael W. T. Mason¹, Chris Newman³, Christina D. Buesching³, David W. Macdonald³ and Hannah L. Dugdale^{1,2}

¹School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK

²Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9747 AG Groningen, The Netherlands

³Wildlife Conservation Research Unit, Department of Zoology, University of Oxford, The Recanati-Kaplan Centre, Abingdon, Oxfordshire OX13 5QL, UK

SHJvL, 0000-0003-4136-265X; EPB, 0000-0001-9398-5440; MWTM, 0000-0003-3264-1569; CN, 0000-0002-9284-6526; CDB, 0000-0002-4207-5196; DWM, 0000-0003-0607-9373; HLD, 0000-0001-8769-0099

Evidence for age-related changes in innate and adaptive immune responses is increasing in wild populations. Such changes have been linked to fitness, and knowledge of the factors driving immune response variation is important for understanding the evolution of immunity. Age-related changes in immune profiles may be owing to factors such as immune system development, sex-specific behaviour and responses to environmental conditions. Social environments may also contribute to variation in immunological responses, for example, through transmission of pathogens and stress arising from resource and mate competition. Yet, the impact of the social environment on age-related changes in immune cell profiles is currently understudied in the wild. Here, we tested the relationship between leukocyte cell composition (proportion of neutrophils and lymphocytes [innate and adaptive immunity, respectively] that were lymphocytes) and age, sex and group size in a wild population of European badgers (*Meles meles*). We found that the proportion of lymphocytes in early life was greater in males in smaller groups compared to larger groups, but with a faster age-related decline in smaller groups. By contrast, the proportion of lymphocytes in females was not significantly related to age or group size. Our results provide evidence of sex-specific age-related changes in immune cell profiles in a wild mammal, which are influenced by the social environment.

1. Introduction

The immune system involves multiple mechanisms that protect the host against pathogens [1]. The functioning of the immune system is related to sex [2,3], changes throughout life [4–9] and has been linked to mortality in the wild [9]. Investigation of how such factors drive variation in immune responses is important for understanding the evolution of immunity.

The immune system principally comprises two components: innate and adaptive immunity [1]. The innate immune response is the first defence against pathogens, involving phagocytic cells (e.g. neutrophils, macrophages and dendritic cells) that detect antigens and produce cytokines, which trigger other

parts of the immune system [10–14]. The activation of adaptive immunity includes the cell-mediated immune response, with the stimulation of T lymphocytes and humoral immunity, and activated B lymphocytes that differentiate to produce immunoglobulins against specific antigens [13,15]. The relative components of innate and adaptive immunity are therefore often reflected in the neutrophil–lymphocyte ratio, respectively [16–19].

The adaptive immune system generally undergoes an age-related decline in performance, i.e. immunosenescence, and evidence for this process has been emerging in wild populations [4–9]. By contrast, the innate immune response is usually maintained, or even enhanced with age [4–9]. This enhanced innate immune response can be a consequence of overstimulation of the immune system, owing to a reduced T cell repertoire and bias towards CD8+ effector memory cells, leading to chronic inflammation and accelerated immunosenescence, as seen in humans [20,21].

The innate and adaptive immune responses, mediated by genes and hormones, are sex-specific [2,3]. For example, in the human innate immune response, males typically have higher neutrophil and macrophage phagocytic activity than females [22,23], whereas in the adaptive immune response, females typically have stronger antibody responses, higher basal immunoglobulin levels and more B cells than males [22,24]. Such sex differences in immune responses may become exacerbated with age [3,25]. For example, male Soay sheep (*Ovis aries*) exhibit steeper sex-specific changes in leukocyte cell composition with age [26]. However, such changes may be species-specific since no sex differences in leukocyte cell composition with age were detected in roe deer (*Capreolus capreolus*; [5]).

Social stress is also emerging as a potential driver of variation in immune responses in the wild [27–29], with stress being reflected in the neutrophil–lymphocyte ratio [30]. Gregarious individuals often experience greater stress owing to more social interactions or increased mate competition [28,31,32]. Testosterone can have a suppressive effect on the immune system ([33,34], but see [35]), and polygynous males have more circulating testosterone than conspecific females or monogamous males. Thus, the social system and the environment can have a sex-specific effect on immune cell profiles. Social individuals may also experience greater costs of pathogen exposure owing to group-living, compared with solitary individuals [29]. For example, greater early-life exposure to pathogen variety and intensity within social groups could prime the immune system and result in enhanced later-life immunity but with the risk of later-life auto-immunity [36,37]. However, to date, there has been no clear evidence for the effects of the social environment on sex-specific immune cell profiles and their age-related changes in the wild.

Here, we use blood samples collected from a wild population of European badgers (*Meles meles*; hereafter ‘badger’) to explore longitudinal changes in sex-specific immune cell profiles in relation to social conditions. We quantify the relative components in the immune system through the proportion of neutrophils and lymphocytes that are lymphocytes (henceforth ‘proportion of lymphocytes’), which reflects the relative balance between innate and adaptive immunity [16–19]. Specifically, we test whether the proportion of lymphocytes: (i) changes with age, (ii) exhibits sex differences and (iii) is linked to group size.

2. Methods

(a) Study species and data collection

We conducted this study in Wytham Woods, Oxfordshire, UK (51°46′24″ N, 1°20′04″ W), a 424 ha semi-natural woodland surrounded by mixed arable pasture [38]. The resident high-density badger population (mean \pm s.e. = 36 ± 3 badgers/km²; [39]) consists of large mixed-sex social groups (mean group size = 11, range = 2–29; [40]). Badgers have a polygynandrous mating system with high rates of extra-group paternity [41,42], where males exhibit seasonal peaks in testosterone levels [43,44]. Badgers are exposed to pathogens, such as coccidia, that negatively impact development and cause juvenile mortality [45–47].

Trapping was undertaken three times per year, for three consecutive days per social group in 2017 and 2018. Trapped badgers were anaesthetized using an intra-muscular injection of 0.2 ml ketamine hydrochloride per kg body weight [48]. Individuals were identified by a unique tattoo number on the left inguinal region, with capture date, social group affiliation and sex recorded. Age was determined as the difference between capture date and the 14th of February in the birth year, since implantation and parturition dates are highly synchronous in badgers [49–51]. Badgers first caught as adults were aged through tooth wear (scale 1–5), where a score of 2 typically indicates a 1-year-old adult [52]. Blood was collected through jugular venipuncture into vacutainers with EDTA anticoagulant. Badgers were released at their setts, after full recovery from anaesthesia. Additionally, bait-marking was conducted periodically to delimit social groups [53] and calculate group sizes using dispersal rules (see electronic supplementary material).

Immediately after blood collection, one drop of blood was smeared on a microscope slide. Slides were air-dried for 1 h then stained using Kwik-Diff (Thermo Scientific, Manchester, UK) according to the manufacturer’s protocol. Leukocyte cell counts were conducted by the same observer (blind to group size and sex) by counting 100 cells per slide (4 repeats per slide, not consecutively to avoid bias; $n = 82$ slides, 23 individuals; 9 females, 14 males), at 40 \times magnification using the battlement technique [54]. Cells were identified as neutrophils, eosinophils and basophils (i.e. granulocytes) or lymphocytes and monocytes (i.e. agranulocytes; [55]). Basophils (<0.1%), eosinophils (1.4%) and monocytes (3.4%) were rarely observed, thus we only used neutrophils and lymphocytes to calculate the lymphocyte proportion from these data [56].

(b) Statistical analyses

Statistical analyses were conducted in R. 3.3.1 [57], using parametric bootstrapping ($n = 5000$) as a robust method to determine significance of predictors and 95% confidence intervals in *lme4* 1.1–14 [58–60]. The mixed model had a binomial error distribution (link = logit), as recommended with a proportional response variable [61] (proportion of lymphocytes), with an offset to account for the number of cells counted per slide ($n = 7$ repeats, 5 slides, where a total of 100 neutrophils and lymphocytes were counted on a slide). Models were run separately for males and females to test for a sex-specific effect with both age [3,26] and group size [62]. To ensure that separating our models by sex did not alter out conclusions (e.g. owing to reduced statistical power), we also ran a model with both sexes included.

We first compared the fit of the relationship between age versus logarithmic age and the proportion of lymphocytes using AICc values; a negative logarithmic pattern was best supported in the full dataset (Δ AICc = -3.8) and males (Δ AICc = -2.3), but with little difference in females (Δ AICc = 0.2). Logarithmic age was therefore included in the mixed model analysing the full dataset and in the separate models for

Table 1. Parameter estimates and 95% confidence intervals of fixed effects from a mixed model and subsequent parametric bootstrapping testing age and group size effects on the proportion of neutrophils and lymphocytes that were lymphocytes in male European badgers. β = direction and magnitude of effect, s.e. = standard error, 95% CI = 95% confidence intervals; reference terms in brackets = reference level for factors; \times = interaction. Significant parameters (95% CI does not overlap zero) are in italics. Random effect estimates (variance): individual ID ($<1.000 \times 10^{-12}$), slide nested in individual ID (1.378×10^{-1}), social group (1.979×10^{-2}), cohort ($<1.000 \times 10^{-12}$), observation (1.080×10^{-1}).

parameter (reference level)	β	s.e.	95% CI
<i>intercept</i>	-2.325	0.127	-2.570 to -2.073
<i>log age</i>	-0.211	0.095	-0.403 to -0.015
<i>group size</i>	0.220	0.087	0.050 to 0.388
year (2017)			
2018	0.421	0.137	0.148 to 0.693
season (spring)			
summer	-0.046	0.131	-0.310 to 0.215
autumn	0.617	0.224	0.156 to 1.069
<i>body condition index</i>	-0.255	0.097	-0.446 to -0.065
<i>log age \times group size</i>	0.202	0.052	0.101 to 0.304

males and females, but the female models were also checked with linear age. We then used AICc to determine *a priori* whether interactions between age, group size and sex (full dataset) and between age and group size (sex-specific datasets) should be included (electronic supplementary material, table S1). When multiple models were plausible ($\Delta\text{AICc} < 7$; [63]) and the interaction was non-significant, we re-ran the model without the interaction to also accurately test the first-order effects. We also included season, year and body condition index ($\log_{10}\text{weight}/\log_{10}\text{body length}$; [44,64]) as fixed effects since these affect immune cell concentrations [65–67]. Cohort, social group, slide nested within individual ID and observation (for each unique measure to account for overdispersion [68]) were included as random effects.

3. Results

In males, we found an interaction between age and group size on the proportion of lymphocytes (table 1). Males living in smaller groups had a higher proportion of lymphocytes in early life, which declined more steeply with age than in males in larger groups, such that the proportion of lymphocytes decreased with age by 50% for males in larger groups compared to 80% for males in smaller groups (figure 1 and table 1). By contrast, for females, the proportion of lymphocytes did not differ significantly according to group size or age (table 2 and electronic supplementary material, table S2), or when using linear age (electronic supplementary material, tables S3 and S4). The full dataset showed an interaction between age, group size and sex on the proportion of lymphocytes (electronic supplementary material, table S5), indicating an interaction between age and group size that differs between males and females, thus providing similar results to the models analysing the sexes separately.

4. Discussion

We found that social conditions (i.e. group size) have sex-specific effects on individual immune cell profiles with age. In male badgers in larger groups, early life exposure to a greater diversity, or higher intensity, of pathogens or greater

stress associated with resource or mate competition could have possibly led to a stronger bias toward innate over adaptive immune cell profiles with age. Male badgers grow to maturity faster than females, resulting in a slight sexual dimorphism, and male growth is predominantly affected by social factors, whereas weather conditions predominantly affect female development [69]. According to the ‘hygiene-hypothesis’ [37,70], early life exposure to pathogens could alleviate the detrimental consequences of increased pathogen pressure in later life and thus slow age-related changes in immune cell profiles. In smaller groups, lower exposure to pathogens in early life could have the opposite effect [71,72], accelerating changes in immune cell profiles with age. Moreover, if fewer conspecifics share the pathogen burden, this could lead to a stronger pressure on the immune response and rapid changes in the proportion of lymphocytes. Indeed, we found that the proportion of lymphocytes in early life was greater in male badgers living in smaller social groups, but with a steeper age-related decline. There was no significant effect in females. This is supported by a previous study in this same population showing that coccidiosis caused by *Eimeria melis* has a more severe effect on male badger cub development [45]. Thus, the greater proportion of neutrophils to lymphocytes that we observed in males in early life, compared to females, could reflect their greater immune response to juvenile coccidiosis.

We also found a relative decrease in the proportion of lymphocytes with age in males but not females. Possibly, female badgers develop a stronger immune response against pathogens in early life, as observed in Soay sheep, where males had a steeper decline in lymphocyte proportion with age than did females [26]. Male badgers, given the polygynandrous mating system, have high testosterone levels [43], particularly compared to monogamous species [44], which may lead to immunosuppression and stronger decreases in adaptive immunity (i.e. lymphocytes) with age ([33,34], but see [35]). The potentially immuno-suppressive effect of testosterone in male badgers accords with sex-specific responses to environmental conditions and associated sex differences in immune responses seen in other species [2,3].




Figure 1. The interplay between age and group size on the proportion of neutrophils and lymphocytes that were lymphocytes for males. Raw data points are shown. Group size was modelled as a continuous variable in the mixed model, but for visualization is shown in small (range = 1–9; $n = 99$ repeats; 25 slides; 9 individuals; brown triangles and dashed line) and large (range = 10–16; $n = 96$ repeats; 24 slides; 8 individuals; blue circles and solid line) groups. Three males were part of a large group at one time point and a small group at another time point, hence total sample size differs from the methods section (§2). Fitted lines represent the model prediction for age interacting with group size, with associated 95% confidence intervals as shaded areas.

Table 2. Parameter estimates and 95% confidence intervals of fixed effects from a mixed model and subsequent parametric bootstrapping testing age and group size effects on the proportion of neutrophils and lymphocytes that were lymphocytes in female European badgers. β = direction and magnitude of effect, s.e. = standard error, 95% CI = 95% confidence intervals; reference terms in brackets = reference level for factors; \times = interaction. Significant parameters (95% CI does not overlap zero) are in *italics*. Random effect estimates (variance): individual ID (4.310×10^{-2}), slide nested in individual ID (1.879×10^{-1}), social group ($<1.000 \times 10^{-12}$), cohort ($<1.000 \times 10^{-12}$), observation (1.206×10^{-1}).

parameter (reference level)	β	s.e.	95% CI
<i>intercept</i>	–2.284	0.188	–2.661 to –1.911
<i>log age</i>	–0.078	0.151	–0.386 to 0.214
<i>group size</i>	–0.107	0.115	–0.344 to 0.122
<i>year (2017)</i>			
2018	–0.017	0.211	–0.447 to 0.405
<i>season (spring)</i>			
summer	0.137	0.194	–0.228 to 0.527
autumn	0.558	0.317	–0.038 to 1.203
<i>body condition index</i>	–0.262	0.143	–0.551 to 0.015
<i>log age \times group size</i>	–0.015	0.117	–0.245 to 0.210

The greatest changes in immune cell profiles in males occurred in early life, when the immune response is developing. Early-life changes may have arisen owing to there being quantitatively fewer acquired immunity cells, or more innate cells being produced. The later-life decrease in the proportion of lymphocytes with age seen in this study has been associated with age-related reduction in thymus size in humans [73,74], accompanied by lower numbers of naïve T cells [75] and CD4⁺ T and CD8⁺ subpopulations with age, which has detrimental implications for effective immune responses

to new antigens [10,76–80]. Alternatively, innate immune mechanisms may become more active with age through increased production of pro-inflammatory cytokines [81]. Such low-grade chronic inflammation in older individuals has detrimental effects on health and contributes to senescence and the development of age-related pathologies [21].

While we cannot provide direct evidence of immunosenescence, as we measured the proportion of lymphocytes rather than the absolute number of leukocytes per unit volume of blood, the relative decrease in adaptive immune

cells and increase in innate immune cells that we detected with age accords with previous studies in the wild [4–6]. Furthermore, understanding the changes in immune cell profiles with age in mammals is important for the interpretation of leukocyte telomere dynamics [52]. Since granulocytes (e.g. neutrophils) have longer telomeres than agranulocytes (e.g. lymphocytes) in humans and baboons [82,83], apparent changes in telomere length with age could be owing to a changing leukocyte cell composition, or selective loss of leukocytes, with age, leading to spurious inferences on telomere shortening.

We were unable to sample individuals until at least three months of age owing to welfare legislation (Protection of Badgers Act, 1992), and thus we cannot rule out the possibility of selective disappearance of individuals with poor innate immune responses, potentially linked to coccidiosis mortality [45]. Additionally, while we provide evidence of age-related changes in one immune parameter (i.e. leukocyte cell composition), immunity is complex and future studies should analyse multiple immune markers (e.g. specific antibodies, inflammatory parameters) together to understand trade-offs and drivers of variation in immune responses. Nonetheless, our results indicate that age-related changes in immune

profiles are associated with the social environment and these effects differ between the sexes.

Data accessibility. Data are available on Dryad (<https://dx.doi.org/10.5061/dryad.2547d7wmx>) [56] and the R code is archived at GitHub (<https://github.com/DugdaleResearchGroup/Leukocyte>).

Authors' contributions. The study was conceived by S.H.J.v.L. and H.L.D. and developed by E.P.B., M.W.T.M., C.N., C.D.B. and D.W.M.; slides were prepared by S.H.J.v.L., and analysed by M.W.T.M.; statistical analyses were conducted by S.H.J.v.L. with input from E.P.B. and H.L.D.; the paper was written by S.H.J.v.L., E.P.B. and H.L.D.; all authors revised the paper for important intellectual content. All authors gave approval for publication and agree to be held accountable for the content.

Competing interests. We declare we have no competing interests.

Funding. S.H.J.v.L. was funded by a Leeds Anniversary Research Scholarship from the University of Leeds with support from a Heredity Fieldwork Grant from the Genetics Society and a Priestley Centre Climate Bursary from the University of Leeds. H.L.D. was supported by a Royal Society Research Grant (RG170425).

Acknowledgements. We thank all members of the Wytham badger team, past and present, for collecting data. We also thank Alexandra Sparks for comments on an earlier draft of this manuscript, and three anonymous reviewers for their comments which greatly improved the manuscript.

References

1. Hoebe K, Janssen E, Beutler B. 2004 The interface between innate and adaptive immunity. *Nat. Immunol.* **5**, 971–974. (doi:10.1038/ni1004-971)
2. Restif O, Amos W. 2010 The evolution of sex-specific immune defences. *Proc. R. Soc. B* **277**, 2247–2255. (doi:10.1098/rspb.2010.0188)
3. Klein SL, Flanagan KL. 2016 Sex differences in immune responses. *Nat. Rev. Immunol.* **16**, 626–638. (doi:10.1038/nri.2016.90)
4. Nussey DH, Watt K, Pilkington JG, Zamojska R, McNeilly TN. 2012 Age-related variation in immunity in a wild mammal population. *Aging Cell* **11**, 178–180. (doi:10.1111/j.1474-9726.2011.00771.x)
5. Cheynel L *et al.* 2017 Immunosenescence patterns differ between populations but not between sexes in a long-lived mammal. *Sci. Rep.* **7**, 13700. (doi:10.1038/s41598-017-13686-5)
6. Peters A, Delhey K, Nakagawa S, Aulsebrook A, Verhulst S. 2019 Immunosenescence in wild animals: meta-analysis and outlook. *Ecol. Lett.* **22**, 1709–1722. (doi:10.1111/ele.13343)
7. Palacios MG, Winkler DW, Klasing KC, Hasselquist D, Vleck CM. 2011 Consequences of immune system aging in nature: a study of immunosenescence costs in free-living Tree Swallows. *Ecology* **92**, 952–966. (doi:10.1890/10-0662.1)
8. Schneeberger K, Courtiol A, Czirjak GA, Voigt CC. 2014 Immune profile predicts survival and reflects senescence in a small, long-lived mammal, the Greater Sac-Winged Bat (*Saccopteryx bilineata*). *PLoS ONE* **9**, e108268. (doi:10.1371/journal.pone.0108268)
9. Froy H, Sparks AM, Watt K, Sinclair R, Bach F, Pilkington JG, Pemberton JM, McNeilly TN, Nussey DH. 2019 Senescence in immunity against helminth parasites predicts adult mortality in a wild mammal. *Science* **365**, 1296–1298. (doi:10.1126/science.aaw5822)
10. Weiskopf D, Weinberger B, Grubeck-Loebenstein B. 2009 The aging of the immune system. *Transpl. Int.* **22**, 1041–1050. (doi:10.1111/j.1432-2277.2009.00927.x)
11. Akira S, Uematsu S, Takeuchi O. 2006 Pathogen recognition and innate immunity. *Cell* **124**, 783–801. (doi:10.1016/j.cell.2006.02.015)
12. Nathan C. 2006 Neutrophils and immunity: challenges and opportunities. *Nat. Rev. Immunol.* **6**, 173–182. (doi:10.1038/nri1785)
13. Mantovani A, Cassatella MA, Costantini C, Jaillon S. 2011 Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat. Rev. Immunol.* **11**, 519–531. (doi:10.1038/nri3024)
14. Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, Yokoyama WM, Ugolini S. 2011 Innate or adaptive immunity? The example of natural killer cells. *Science* **331**, 44–49. (doi:10.1126/science.1198687)
15. Iwasaki A, Medzhitov R. 2010 Regulation of adaptive immunity by the innate immune system. *Science* **327**, 291–295. (doi:10.1126/science.1183021)
16. Fest J, Ruiter R, Ikram MA, Voortman T, van Eijck CHJ, Stricker BH. 2018 Reference values for white blood-cell-based inflammatory markers in the Rotterdam Study: a population-based prospective cohort study. *Sci. Rep.* **8**, 10566. (doi:10.1038/s41598-018-28646-w)
17. Hu B *et al.* 2014 Systemic immune-inflammation index predicts prognosis of patients after curative resection for hepatocellular carcinoma. *Clin. Cancer Res.* **20**, 6212–6222. (doi:10.1158/1078-0432.Ccr-14-0442)
18. Templeton AJ *et al.* 2014 Prognostic role of neutrophil-to-lymphocyte ratio in solid tumors: a systematic review and meta-analysis. *J. Natl Cancer Inst.* **106**, dju124. (doi:10.1093/jnci/dju124)
19. van der Willik KD, Fani L, Rizopoulos D, Licher S, Fest J, Schagen SB, Ikram MK, Ikram MA. 2019 Balance between innate versus adaptive immune system and the risk of dementia: a population-based cohort study. *J. Neuroinflamm.* **16**, 68. (doi:10.1186/s12974-019-1454-z)
20. Sansoni P *et al.* 2014 New advances in CMV and immunosenescence. *Exp. Gerontol.* **55**, 54–62. (doi:10.1016/j.exger.2014.03.020)
21. Franceschi C, Garagnani P, Parini P, Giuliani C, Santoro A. 2018 Inflammaging: a new immunometabolic viewpoint for age-related diseases. *Nat. Rev. Endocrinol.* **14**, 576–590. (doi:10.1038/s41574-018-0059-4)
22. Abdullah M, Chai PS, Chong MY, Tohit ERM, Ramasamy R, Pei CP, Vidyadaran S. 2012 Gender effect on *in vitro* lymphocyte subset levels of healthy individuals. *Cell. Immunol.* **272**, 214–219. (doi:10.1016/j.cellimm.2011.10.009)
23. Spitzer JA. 1999 Gender differences in some host defense mechanisms. *Lupus* **8**, 380–383. (doi:10.1177/096120339900800510)
24. Furman D, Hejblum BP, Simon N, Jovic V, Dekker CL, Thiebaut R, Tibshirani RJ, Davis MM. 2014 Systems analysis of sex differences reveals an immunosuppressive role for testosterone in the response to influenza vaccination. *Proc. Natl Acad.*

- Sci. USA* **111**, 869–874. (doi:10.1073/pnas.1321060111)
25. Campo JL, Davila SG. 2002 Estimation of heritability for heterophil:lymphocyte ratio in chickens by restricted maximum likelihood. Effects of age, sex, and crossing. *Poult. Sci.* **81**, 1448–1453. (doi:10.1093/ps/81.10.1448)
 26. Watson RL *et al.* 2017 Sex differences in leukocyte telomere length in a free-living mammal. *Mol. Ecol.* **26**, 3230–3240. (doi:10.1111/mec.13992)
 27. Côté IM, Poulin R. 1995 Parasitism and group-size in social animals – a meta-analysis. *Behav. Ecol.* **6**, 159–165. (doi:10.1093/beheco/6.2.159)
 28. Creel S, Dantzer B, Goymann W, Rubenstein DR. 2013 The ecology of stress: effects of the social environment. *Funct. Ecol.* **27**, 66–80. (doi:10.1111/j.1365-2435.2012.02029.x)
 29. Altizer S *et al.* 2003 Social organization and parasite risk in mammals: integrating theory and empirical studies. *Annu. Rev. Ecol. Evol. Syst.* **34**, 517–547. (doi:10.1146/annurev.ecolsys.34.030102.151725)
 30. Davis AK, Maney DL, Maerz JC. 2008 The use of leukocyte profiles to measure stress in vertebrates: a review for ecologists. *Funct. Ecol.* **22**, 760–772. (doi:10.1111/j.1365-2435.2008.01467.x)
 31. Blumstein DT, Williams DM, Lim AN, Kroeger S, Martin JGA. 2018 Strong social relationships are associated with decreased longevity in a facultatively social mammal. *Proc. R. Soc. B* **285**, 20171934. (doi:10.1098/rspb.2017.1934)
 32. Martin LB. 2009 Stress and immunity in wild vertebrates: timing is everything. *Gen. Comp. Endocr.* **163**, 70–76. (doi:10.1016/j.ygcen.2009.03.008)
 33. Klein SL. 2000 Hormones and mating system affect sex and species differences in immune function among vertebrates. *Behav. Process.* **51**, 149–166. (doi:10.1016/S0376-6357(00)00125-X)
 34. Hillgarth N, Wingfield JC. 1997 Testosterone and immunosuppression in vertebrates: implications for parasite-mediated sexual selection. In *Parasites and pathogens: effects on host hormones and behaviour* (ed. NE Beckage), pp. 143–155. New York, NY: Chapman & Hall.
 35. Roberts ML, Buchanan KL, Evans MR. 2004 Testing the immunocompetence handicap hypothesis: a review of the evidence. *Anim. Behav.* **68**, 227–239. (doi:10.1016/j.anbehav.2004.05.001)
 36. Olszak T *et al.* 2012 Microbial exposure during early life has persistent effects on natural killer T cell function. *Science* **336**, 489–493. (doi:10.1126/science.1219328)
 37. von Mutius E. 2007 Allergies, infections and the hygiene hypothesis – The epidemiological evidence. *Immunobiology* **212**, 433–439. (doi:10.1016/j.imbio.2007.03.002)
 38. Macdonald DW, Newman C. 2002 Population dynamics of badgers (*Meles meles*) in Oxfordshire, UK: numbers, density and cohort life histories, and a possible role of climate change in population growth. *J. Zool.* **256**, 121–138. (doi:10.1017/S0952836902000158)
 39. Macdonald DW, Newman C, Nouvellet PM, Buesching CD. 2009 An analysis of Eurasian badger (*Meles meles*) population dynamics: implications for regulatory mechanisms. *J. Mammal.* **90**, 1392–1403. (doi:10.1644/08-MAMM-A-356R1.1)
 40. da Silva J, Macdonald DW, Evans PGH. 1994 Net costs of group living in a solitary forager, the Eurasian badger (*Meles meles*). *Behav. Ecol.* **5**, 151–158. (doi:10.1093/beheco/5.2.151)
 41. Dugdale HL, Macdonald DW, Pope LC, Burke T. 2007 Polygyny, extra-group paternity and multiple-paternity litters in European badger (*Meles meles*) social groups. *Mol. Ecol.* **16**, 5294–5306. (doi:10.1111/j.1365-294X.2007.03571.x)
 42. Annavi G, Newman C, Dugdale HL, Buesching CD, Sin YW, Burke T, Macdonald DW. 2014 Neighbouring-group composition and within-group relatedness drive extra-group paternity rate in the European badger (*Meles meles*). *J. Evol. Biol.* **27**, 2191–2203. (doi:10.1111/jeb.12473)
 43. Buesching CD, Heistermann M, Macdonald DW. 2009 Seasonal and inter-individual variation in testosterone levels in badgers *Meles meles*: evidence for the existence of two endocrinological phenotypes. *J. Comp. Physiol. A* **195**, 865–871. (doi:10.1007/s00359-009-0465-0)
 44. Sugianto NA, Newman C, Macdonald DW, Buesching CD. 2019 Heterochrony of puberty in the European badger (*Meles meles*) can be explained by growth rate and group-size: Evidence for two endocrinological phenotypes. *PLoS ONE* **14**, e0203910. (doi:10.1371/journal.pone.0203910)
 45. Newman C, Macdonald DW, Anwar MA. 2001 Coccidiosis in the European badger, *Meles meles* in Wytham Woods: infection and consequences for growth and survival. *Parasitology* **123**, 133–142. (doi:10.1017/S0031182001008265)
 46. Anwar MA, Newman C, MacDonald DW, Woolhouse MEJ, Kelly DW. 2000 Coccidiosis in the European badger (*Meles meles*) from England, an epidemiological study. *Parasitology* **120**, 255–260. (doi:10.1017/S0031182099005491)
 47. Sin YW, Annavi G, Dugdale HL, Newman C, Burke T, Macdonald DW. 2014 Pathogen burden, co-infection and Major Histocompatibility Complex variability in the European badger (*Meles meles*). *Mol. Ecol.* **23**, 5072–5088. (doi:10.1111/mec.12917)
 48. McLaren GW, Thornton PD, Newman C, Buesching CD, Baker SE, Mathews F, Macdonald DW. 2005 The use and assessment of ketamine-medetomidine-butorphanol combinations for field anaesthesia in wild European badgers (*Meles meles*). *Vet. Anaesth. Analg.* **32**, 367–372. (doi:10.1111/j.1467-2995.2005.00206.x)
 49. Canivenc R, Bonnin M. 1979 Delayed implantation is under environmental control in the badger (*Meles meles* L.). *Nature* **278**, 849–850. (doi:10.1038/278849a0)
 50. Dugdale HL, Ellwood SA, Macdonald DW. 2010 Alloparental behaviour and long-term costs of mothers tolerating other members of the group in a plurally breeding mammal. *Anim. Behav.* **80**, 721–735. (doi:10.1016/j.anbehav.2010.07.011)
 51. Woodroffe R. 1995 Body condition affects implantation date in the European badger, *Meles meles*. *J. Zool.* **236**, 183–188. (doi:10.1111/j.1469-7998.1995.tb04486.x)
 52. van Lieshout SHJ, Bretman A, Newman C, Buesching CD, Macdonald DW, Dugdale HL. 2019 Individual variation in early life telomere length and survival in a wild mammal. *Mol. Ecol.* **28**, 4152–4165. (doi:10.1111/mec.15212)
 53. Delahay RJ, Brown JA, Mallinson PJ, Spyvee PD, Handoll D, Rogers LM, Cheeseman CL. 2000 The use of marked bait in studies of the territorial organization of the European badger (*Meles meles*). *Mammal Rev.* **30**, 73–87. (doi:10.1046/j.1365-2907.2000.00058.x)
 54. Schalm OW. 1965 *Veterinary hematology*. Philadelphia, PA: Lea & Febiger.
 55. Bain BJ. 2015 *Blood cells: a practical guide*. Oxford, UK: John Wiley & Sons.
 56. van Lieshout SHJ, Badás EP, Mason MWT, Newman C, Buesching CD, Macdonald DW, Dugdale HL. 2020 Data from: Social effects on age-related and sex-specific immune cell profiles in a wild mammal. *Dryad Digital Repository*. (doi:10.5061/dryad.2547d7wmw)
 57. R Development Core Team. 2019 *R: a language and environment for statistical computing*. Vienna, Austria: R foundation for statistical computing.
 58. Bates D, Machler M, Bolker BM, Walker SC. 2015 Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
 59. Nakagawa S, Schielzeth H. 2013 A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol. Evol.* **4**, 133–142. (doi:10.1111/j.2041-210x.2012.00261.x)
 60. Nakagawa S, Schielzeth H. 2010 Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists. *Biol. Rev.* **85**, 935–956. (doi:10.1111/j.1469-185X.2010.00141.x)
 61. Crawley MJ. 2012 *The R book*. Chichester, UK: John Wiley & Sons.
 62. Leech T, Evison SEF, Armitage SAO, Sait SM, Bretman A. 2019 Interactive effects of social environment, age and sex on immune responses in *Drosophila melanogaster*. *J. Evol. Biol.* **32**, 1082–1092. (doi:10.1111/jeb.13509)
 63. Burnham KP, Anderson DR, Huyvaert KP. 2011 AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behav. Ecol. Sociobiol.* **65**, 23–35. (doi:10.1007/s00265-010-1029-6)
 64. Noonan MJ, Markham A, Newman C, Trigoni N, Buesching CD, Ellwood SA, Macdonald DW. 2014 Climate and the individual: inter-annual variation in the autumnal activity of the European badger (*Meles meles*). *PLoS ONE* **9**, e83156. (doi:10.1371/journal.pone.0083156)
 65. Downs JD, Dochtermann NA, Ball R, Klasing KC, Martin LB. 2019 The effects of body mass on immune cell concentrations of mammals. *Am. Nat.* **195**, 107–114. (doi:10.1086/706235)
 66. Moller AP, Erritzoe J, Saino N. 2003 Seasonal changes in immune response and parasite impact

- on hosts. *Am. Nat.* **161**, 657–671. (doi:10.1086/367879)
67. Beaulieu M, Benoit L, Abaga S, Kappeler PM, Charpentier MJE. 2017 Mind the cell: seasonal variation in telomere length mirrors changes in leucocyte profile. *Mol. Ecol.* **26**, 5603–5613. (doi:10.1111/mec.14329)
68. Harrison XA. 2014 Using observation-level random effects to model overdispersion in count data in ecology and evolution. *PeerJ* **2**, e616. (doi:10.7717/peerj.616)
69. Sugianto NA, Newman C, Macdonald DW, Buesching CD. 2019 Extrinsic factors affecting cub development contribute to sexual size dimorphism in the European badgers (*Meles meles*). *Zoology* **135**, 125688. (doi:10.1016/j.zool.2019.04.005)
70. Strachan DP. 1989 Hay-fever, hygiene, and household size. *Brit. Med. J.* **299**, 1259–1260. (doi:10.1136/bmj.299.6710.1259)
71. Shaw SY, Blanchard JF, Bernstein CN. 2010 Association between the use of antibiotics in the first year of life and pediatric inflammatory bowel disease. *Am. J. Gastroenterol.* **105**, 2687–2692. (doi:10.1038/ajg.2010.398)
72. Goksor E, Alm B, Thengilsdottir H, Pettersson R, Aberg N, Wennergren G. 2011 Preschool wheeze – impact of early fish introduction and neonatal antibiotics. *Acta Paediatr.* **100**, 1561–1566. (doi:10.1111/j.1651-2227.2011.02411.x)
73. Flores KG, Li J, Sempowski GD, Haynes BF, Hale LP. 1999 Analysis of the human thymic perivascular space during aging. *J. Clin. Invest.* **104**, 1031–1039. (doi:10.1172/Jci7558)
74. George AJT, Ritter MA. 1996 Thymic involution with ageing: obsolescence or good housekeeping? *Immunol. Today* **17**, 267–272. (doi:10.1016/0167-5699(96)80543-3)
75. Fagnoni FF *et al.* 2000 Shortage of circulating naive CD8⁺ T cells provides new insights on immunodeficiency in aging. *Blood* **95**, 2860–2868. (doi:10.1182/blood.V95.9.2860.009k35_2860_2868)
76. Goronzy JJ, Weyand CM. 2005 T cell development and receptor diversity during aging. *Curr. Opin. Immunol.* **17**, 468–475. (doi:10.1016/j.coi.2005.07.020)
77. Effros RB, Cai ZL, Linton PJ. 2003 CD8 T cells and aging. *Crit. Rev. Immunol.* **23**, 45–64. (doi:10.1615/CritRevImmunol.v23.i12.30)
78. Pfister G, Weiskopf D, Lazuardi L, Kovaiov RD, Cioca DP, Keller M, Lorbeg B, Parson W, Grubeck-Loebenstein B. 2006 Naive T cells in the elderly: are they still there? *Ann. NY Acad. Sci.* **1067**, 152–157. (doi:10.1196/annals.1354.018)
79. Kohler S, Wagner U, Pierer M, Kimmig S, Oppmann B, Mowes B, Julke K, Romagnani C, Thiel A. 2005 Post-thymic *in vivo* proliferation of naive CD4⁺ T cells constrains the TCR repertoire in healthy human adults. *Eur. J. Immunol.* **35**, 1987–1994. (doi:10.1002/eji.200526181)
80. Haynes L, Eaton SM, Burns EM, Randall TD, Swain SL. 2003 CD4 T cell memory derived from young naive cells functions well into old age, but memory generated from aged naive cells functions poorly. *Proc. Natl Acad. Sci. USA* **100**, 15 053–15 058. (doi:10.1073/pnas.2433717100)
81. Fagiolo U, Cossarizza A, Scala E, Fanalesbelasio E, Ortolani C, Cozzi E, Monti D, Franceschi C, Paganelli R. 1993 Increased cytokine production in mononuclear cells of healthy elderly people. *Eur. J. Immunol.* **23**, 2375–2378. (doi:10.1002/eji.1830230950)
82. Baerlocher GM, Rice K, Vulto I, Lansdorp PM. 2007 Longitudinal data on telomere length in leukocytes from newborn baboons support a marked drop in stem cell turnover around 1 year of age. *Aging Cell* **6**, 121–123. (doi:10.1111/j.1474-9726.2006.00254.x)
83. Kimura M, Gazitt Y, Cao XJ, Zhao XY, Lansdorp PM, Aviv A. 2010 Synchrony of telomere length among hematopoietic cells. *Exp. Hematol.* **38**, 854–859. (doi:10.1016/j.exphem.2010.06.010)