

Research Article

Ethanol Extract of the Flower *Chrysanthemum morifolium* Augments Pentobarbital-Induced Sleep Behaviors: Involvement of Cl⁻ Channel Activation

Jae-Wook Kim,¹ Jin-Yi Han,² Jin Tae Hong,¹ Rihua Li,³ Jae Soon Eun,³ and Ki-Wan Oh¹

¹ College of Pharmacy, Chungbuk National University, Cheongju 361-763, Republic of Korea

² Research Institute of Veterinary Medicine, Chungbuk National University, Cheongju 361-763, Republic of Korea

³ College of Pharmacy, Woosuk University, Samrye 565-701, Republic of Korea

Correspondence should be addressed to Ki-Wan Oh, kiwan@chungbuk.ac.kr

Received 10 November 2010; Accepted 11 January 2011

Copyright © 2011 Jae-Wook Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dried *Chrysanthemum morifolium* flowers have traditionally been used in Korea for the treatment of insomnia. This study was performed to investigate whether the ethanol extract of *Chrysanthemum morifolium* flowers (EFC) enhances pentobarbital-induced sleep behaviors. EFC prolonged sleep time induced by pentobarbital similar to muscimol, a GABA_A receptors agonist. EFC also increased sleep rate and sleep time when administrated with pentobarbital at a subhypnotic dosage. Both EFC and pentobarbital increased chloride (Cl⁻) influx in primary cultured cerebellar granule cells. EFC increased glutamic acid decarboxylase (GAD) expression levels, but had no effect on the expression of α 1-, β 2-, and γ 2-subunits of the GABA_A receptor in the hippocampus of a mouse brain. This is in contrast to treatment with pentobarbital, which showed decreased α 1-subunit expression and no change in GAD expression. In conclusion, EFC augments pentobarbital-induced sleep behaviors; these effects may result from Cl⁻ channel activation.

1. Introduction

The flower of *Chrysanthemum morifolium* Ramat. (FC) has been used in oriental countries for hundreds of years and is widely consumed as a medicinal herbal tea [1, 2]. FC is reported to have various biological features including antioxidation [3], cardiovascular protection [4], antitumorigenesis [5], and anti-inflammation [6]. *Chrysanthemum* species have been shown to contain a wide variety of flavonoids, phenols, and phenolic acids [7]. Significant amounts of flavonoids and hydroxycinnamoylquinic acids are considered to be the biologically active components [4, 5, 8], and the health benefits of FC have been shown to be associated with the flavonoids [9]. In Korea, the dried FC in herbal tea has traditionally been used for the treatment of insomnia [3].

Insomnia symptoms are some of the most frequent sleep complaints in the general population, with an estimated prevalence varying from 10 to nearly 60%, depending in part on the use of varying definitions and data-collection methodologies [10]. Individuals reporting disturbed sleep

are more likely to report emotional distress and recurrent health problems [11]. This is not surprising since it has been shown that sleep deprivation has a great impact on the everyday life of healthy subjects, affecting alertness, attention, concentration, cognitive abilities, memory, mood, and pain.

A variety of modulators of GABA-transmission, including neurosteroids, benzodiazepines, barbiturates, and GABA agonists, have been investigated in both *in vitro* and *in vivo* models. It is now well known that their effects are related to their binding to specific GABA_A-receptor subtypes. Muscimol and other GABA_A receptor agonists that potentiate Cl⁻ influx also cause potentiation of Cl⁻ influx when administered with pentobarbital or other agonists [12]. The sedative and anticonvulsant effects of diazepam and zolpidem are primarily mediated by their interaction with the α 1-subunit [13]. Benzodiazepines and benzodiazepine-like compounds are the most widely used hypnotics; they primarily act to shorten sleep latency and enhance and consolidate sleep. It was postulated that the hypnotic properties of benzodiazepines, that is, their capacity to shorten

sleep latency, to reduce waking after sleep onset, and to consolidate sleep are associated with the $\alpha 1$ -subunit [14, 15]. The various side effects associated with benzodiazepines stimulated the search for alternative compounds for the treatment of insomnia. Therefore, the goal of this study was to evaluate the hypnotic effects of the ethanol extract of the flower of *C. morifolium* (EFC) on pentobarbital-induced sleep behaviors and to investigate its possible mechanisms.

2. Materials and Methods

2.1. Animals. Male ICR mice (Samtako, Korea) weighing 20–25 g, in groups of 10–12, were used for behavioral experiments. Animals were housed in acrylic cages (45 × 60 × 25 cm) with water and food available *ad libitum* under an artificial 12-h light/dark cycle (lights on at 7:00) and at a constant temperature (22 ± 2°C). Mice were housed in the departmental room for 1 week before testing to ensure adaptation to the new environment. All of the behavioral experiments were performed between 10:00 and 17:00. All of the experiments involving animals were carried out in accordance with the National Institutes of Health Guide for Care and Use of Laboratory Animals (NIH publication No. 85-23, revised 1985), and the Institutional Animal Care and Use Committee of Chungbuk National University approved the protocol.

2.2. Cell Culture. Primary cultures of cerebellar neurons enriched in granule cells were prepared from cerebella of 8-day-old Sprague-Dawley rats as previously described [16]. After 8 days in culture, these cells express functional GABA_A receptors, with an expression pattern similar to that of the cerebellum during postnatal development, but different from the pattern observed in the adult rat cerebellum [17]. Briefly, cells were plated (1 × 10⁵ cells per well) in 96-well microplates that had been coated with poly-L-lysine (50 µg/mL; Sigma, St. Louis, MO, USA) and were cultured in Dulbecco's modified Eagle's medium nutrient and Ham's F12 mixture media (Life Technologies, Gaithersburg, MD, USA) supplemented with 10% heat-inactivated fetal bovine serum (Life Technologies), glutamine (2 mM), gentamicin (100 µg/mL), antibiotic-antimycotic solution (10 mL/L; Sigma) and Potassium chloride (25 mM); a high concentration of potassium was necessary to induce persistent depolarization, which promotes the survival of granule cells. Cytosine arabinofuranoside (final concentration, 10 µM; Sigma) was added to cultures 18–24 h after plating, to inhibit the proliferation of nonneuronal cells.

2.3. Ethanol Extraction of *C. morifolium*. The plant materials were collected during October 2009 at Jiri Mt., Jeonbuk, Korea. The shade-dried *C. morifolium* flower (100 g) was extracted three times with 70% EtOH at 50°C. The extracts were filtered and concentrated using a rotary vacuum evaporator, followed by lyophilization. The yield of *C. morifolium* flower extracts (EFC) was approximately 2.5%.

2.4. Pentobarbital-Induced Sleep. Pentobarbital sodium (Hanlim Pharm. Co., Ltd., Korea) was diluted in physiological saline and administered to each mouse intraperitoneally (i.p.) to induce sleep. EFC was suspended in physiological saline and was administered orally (p.o.) to animals. Muscimol (Sigma, USA) was administered as a reference drug 15 min prior to administration of pentobarbital. All experiments were carried out between 13:00 and 17:00. Animals were food-deprived for 24 h prior to the experiment. Thirty minutes after the oral administration of the test samples, pentobarbital was given to animals placed in a box. Animals that stopped moving around the box, whereas animals that failed to fall asleep within 15 min after pentobarbital administration were excluded from the study [18, 19]. Mice that remained immobile for more than 3 min were judged to be asleep. The time that elapsed from receiving pentobarbital until each animal lost its righting reflex when positioned delicately on its back represented the latency to onset of sleep. The animals were observed constantly, and the time of awakening, characterized by righting of the animal, was noted. Sleep time was defined as the time required for the animal to regain spontaneous movements after having been transferred to the second box.

2.5. Measurement of Intracellular Cl⁻ Influx. The intracellular Cl⁻ concentration ([Cl⁻]_i) of cerebellar granule cells was estimated using Cl⁻ sensitive fluorescence probe N-(ethoxycarbonylmethyl)-6-methoxyquionolinium bromide (MQAE) according to the method of West and Molloy, with a slight modification [20]. The buffer (pH 7.4) contained the following components: 2.4 mM HPO₄²⁻, 0.6 mM H₂PO₄⁻, 10 mM HEPES, 10 mM D-glucose, and 1 mM MgSO₄. A variety of MQAE-loading conditions were assessed. The cells were incubated overnight in medium containing 10 mM MQAE (Dojindo, Japan). After loading, the cells were washed three times in the appropriate Cl⁻ containing buffer or Cl⁻-free buffer. The buffer was replaced with buffer with or without the compounds or Cl⁻-free buffer. Repetitive fluorescence measurements were initiated immediately using a FLUOstar plate reader (Excitation wavelength: 320 nm, emission wavelength: 460 nm; BMG LabTechnology, Germany). The data is presented as the relative fluorescence F_0/F , where F_0 is the fluorescence without Cl⁻ ions and F is the fluorescence as a function of time. The F_0/F values were directly proportional to [Cl⁻]_i.

2.6. GAD and GABA_A Receptors Subunits Expression. Mice were administered EFC or pentobarbital for 3 days and sacrificed. The mice were decapitated, their brains removed and the hippocampus dissected on ice according to the methods described by Glowinski and Iversen [21] and Segal and Kuczenski [22]. Mouse hippocampus was homogenized with lysis buffer. The extracts were centrifuged at 20,000 × g for 20 min. Equal amounts of proteins were separated on a 10% SDS/polyacrylamide gel and transferred to a nitrocellulose membrane (Hyboud ECL, Amersham Pharmacia Biotech Inc., Piscataway, NJ, USA). The blots were blocked

TABLE 1: Effects of EFC on sleep onset of mice treated by subhypnotic doses of pentobarbital (28 mg/kg, i.p.).

Group	Dose (mg/kg)	No. falling asleep/total	Sleep time (min)
Control	0	6/15	23.3 ± 1.8
Muscimol	0.2	13/14**	41.2 ± 4.2***
EFC	20	7/14	28.0 ± 4.0
	50	7/14	29.6 ± 3.2
	100	11/13*	35.9 ± 3.6*

Each value represents the mean (±S.E.M.) of 13–15 observations. * $P < .05$, ** $P < .01$, *** $P < .005$ versus control.

for 2 h at room temperature with 5% (w/v) nonfat dried milk in Tris-buffered saline solution (10 mM Tris, pH 8.0 and 150 mM NaCl) containing 0.1% Tween-20. The membrane was incubated with specific rabbit polyclonal antibodies against GABA_A receptor subunits (1 : 1000; Abcam Inc.) for 2 h at room temperature. The blot was then incubated with the corresponding antirabbit IgG-conjugated to horseradish peroxidase (Santa Cruz Biotechnology Inc.). The immunoreactive proteins were detected using the ECL western blotting detection system [23].

2.7. Statistical Analysis. The results are presented as the mean ± S.E.M. The significance of the effects of the compounds was assessed using analysis of variance (ANOVA). Where there was significant variability, the individual values were compared using Dunnett's test. For the subhypnotic pentobarbital dosage experiment, Chi-square test was used to compare the proportion of sleep onset between the group treated with a subhypnotic dose of pentobarbital alone and each of the groups that received pentobarbital in combination with another drug.

3. Results

3.1. Effects of EFC on the Onset and Duration of Sleep in Pentobarbital-Treated Mice. The administration of EFC increased sleep time. EFC produced a dose-dependent prolongation of pentobarbital-induced sleep time at dose of 50 mg/kg and 100 mg/kg; however, EFC did not affect the latency of sleep. Pretreatment of mice with muscimol (0.2 mg/kg, i.p.) as a positive control, 15 min before the administration of pentobarbital (40 mg/kg), produced an increase in total sleep time and a decrease in the latency of sleep (Figure 1).

3.2. Effects of EFC on Sleep Onset in Mice Treated with a Subhypnotic Dosage of Pentobarbital. Administration of EFC increased the rate of sleep onset and the duration of sleep time induced by a subhypnotic dosage of pentobarbital (28 mg/kg, i.p.). Pretreatment with muscimol also increased the rate of sleep onset and prolonged the duration of sleep time when given in combination with a subhypnotic dosage




FIGURE 1: Effects of EFC on onset and duration of sleep in pentobarbital-treated mice. Mice were food-deprived for 24 h before the experiment. Pentobarbital (40 mg/kg, i.p.) was administered to mice following administration of muscimol or EFC. The sleep time (a) and sleep latency (b) were recorded. Each column represents the mean with S.E.M. The significance of the effects of the compounds was assessed using analysis of variance (ANOVA). ** $P < .01$, *** $P < .005$, compared to control.

of pentobarbital (Table 1). EFC showed similar effects to muscimol at 100 mg/kg.

3.3. Effects of EFC on Cl⁻ Influx in Primary Cultured Cerebellar Granule Cells. Intracellular chloride ion influx in primary cultured cerebellar granule cells was measured. The measured data is presented as the relative fluorescence F_0/F , where F_0 is the fluorescence without chloride ions and F is the fluorescence as a function of each sample. The F_0/F values were directly proportional to intracellular chloride ion concentration. Treatment of granule cells with EFC (1–4 μ g/mL) produced a significant increase in chloride ion influx. Pentobarbital 10 μ M also increased the influx of Cl⁻ in primary cultured cerebellar granule cells (Figure 2).




FIGURE 2: Effects of EFC on chloride influx in primary cultured cerebellar granule cells. After the culture of cerebellar granule cells for 8 days, the cells were incubated with MQAE overnight, and then EFC (1–4 μg/mL) and pentobarbital (PENT, 10 μM) were added 5 min prior to measurement. PENT: pentobarbital. Each column represents the mean with S.E.M. The significance of the effects of the compounds was assessed using analysis of variance (ANOVA). * $P < .05$, *** $P < .005$, compared to control.

3.4. *Effects of EFC on Expression of Glutamic Acid Decarboxylase (GAD) and Subunit of GABA_A Receptor.* Mice were administered 100 mg/kg EFC or 40 mg/kg pentobarbital for 3 days, and they were sacrificed to examine the effect of these drugs on the abundance of glutamic acid decarboxylase (GAD) and GABA_A receptor subunits in the hippocampus. EFC treatment increased expression of GAD₆₅ (Figure 3) but did not influence the amounts of α1-, β2-, and γ2-subunits in the GABA_A receptor (Figure 4); however, pentobarbital significantly decreased amounts of the α1-subunit, but did not affect the abundance of β2- and γ2-subunit. Protein concentrations of GAD₆₅ following pentobarbital treatment also were not changed.

4. Discussion

The results demonstrate that EFC potentiates pentobarbital-induced sleep behaviors in mice. The increase and decrease of pentobarbital-induced sleep time can be a useful tool for examining the stimulatory or inhibitory effects on central nervous system (CNS), especially for investigating drug influences on GABAergic systems [24, 25]. In addition, pentobarbital is well known to potentiate the effects of GABA by acting at its own binding sites on the GABA/benzodiazepine receptor ionophore complex [25]. Many hypnotic, anti-anxiety and anti-epilepsy drugs have been shown to cause prolongation of pentobarbital-induced sleep time [26–28]. We were interested in whether EFC prolongs pentobarbital-induced sleep behaviors and interacts with pentobarbital in the CNS via the GABAergic systems.

We investigated the effects of different doses of EFC and muscimol in rodents with pentobarbital treatment. We found that EFC could potentiate pentobarbital-induced sleep




FIGURE 3: Effects of EFC on glutamic acid decarboxylase (GAD). Immunoblots of lysed mouse hippocampus 3 days following administration of EFC or pentobarbital are shown. GAPDH levels were measured to confirm equal protein loading. PENT: pentobarbital. Each column represents the mean with S.E.M. The significance of the effects of the compounds was assessed using analysis of variance (ANOVA). * $P < .05$, compared to control.

at 100 mg/kg. Additionally, EFC increased the rate of sleep onset and prolonged sleep time at subhypnotic dosages of pentobarbital (28 mg/kg). These results are similar to those of the GABA_A receptor agonist muscimol. This indicates that the hypnotic effect of EFC may be due to interaction with GABAergic systems. GABA_A receptors possess various binding sites, including binding sites for GABA, benzodiazepine, and barbiturates. GABA_A receptors form heteromeric GABA-gated Cl⁻ channels, which are assembled from a large family of subunit genes. GABA_A receptor channels open after binding GABA to give a net inward flux of negative Cl⁻ ions (outward current), hyperpolarizing the membrane and reducing neuronal firing [29]. Muscimol and other GABA_A receptor agonists that potentiate Cl⁻ influx also cause potentiation of Cl⁻ influx when administered with pentobarbital or other agonists [12]. EFC produced a significant increase in Cl⁻ influx; this increase was similar to that of pentobarbital. This suggests that EFC may act to induce Cl⁻ channel opening of GABA_A receptors.

Researchers have demonstrated that the pharmacological profile and different drug-induced behaviors of GABA_A receptors depend upon its subunit composition [30]. Glutamic acid decarboxylase (GAD), the rate-limiting enzyme in GABA biosynthesis, also plays an important role in maintaining GABA levels in the brain [31]. Hence, alteration of expression levels of this enzyme may change GABA transmission in the brain. We sought to determine GAD protein and GABA_A receptor subunit expression levels at the effective dosage of EFC and pentobarbital to determine the possible




FIGURE 4: Effects of EFC on GABA_A receptor subunits. Immunoblots of lysed mouse hippocampus 3 days following administration of EFC or pentobarbital are shown. GAPDH levels were measured to confirm equal protein loading. PENT: pentobarbital. Each column represents the mean with S.E.M. The significance of the effects of the compounds was assessed using analysis of variance (ANOVA). * $P < .05$, compared to control.

site of action by which EFC exerts its sleep-potentiating effects. GAD has two molecular forms (GAD₆₅ and GAD₆₇ with molecular weights of 65 kDa and 67 kDa, respectively), and we investigated expression levels of GAD₆₅. GAD₆₅ is responsible for vesicular GABA production; therefore this isoform is directly involved in GABA transmission at the synapse [32]. We also investigated the expression levels of GABA_A receptor $\alpha 1$ -, $\beta 2$ -, and $\gamma 2$ -subunits. The most abundant GABA_A receptor subunit composition, $\alpha 1\beta 2\gamma 2$, is present in most brain regions, including the hippocampus, and these subunits are related to the hypnotic/sedative effect of GABA_A receptor ligands [30]. Our results showed that neither EFC nor pentobarbital treatment influenced expression of GABA_A receptor $\beta 2$ - and $\gamma 2$ -subunits; however pentobarbital decreased abundance of $\alpha 1$ -subunits, and EFC increased levels of GAD.

Many herbal preparations and a diversity of drugs used to promote sleep are known to act on GABA_A receptors [33]. Drugs acting on GABA_A receptors mainly act to increase synaptic inhibition either by directly activating GABA_A receptors or, more usually, by enhancing the action of other ligands on GABA_A receptors. Our results suggest that EFC




FIGURE 5: A proposed mechanism of EFC in augmenting pentobarbital-induced sleep. EFC administration will increase expression levels of GAD, the rate-limiting enzyme in GABA biosynthesis, thereby increasing GABA levels. This will promote Cl⁻ channel opening, causing hyperpolarization of the post-synaptic neuron. Hyperpolarization may result in potentiation of pentobarbital-induced sleep.

has sleep-potentiating effects, which may be mediated by Cl⁻ channel opening (Figure 5).

The search for novel plant-derived pharmacotherapies for psychiatric illness has progressed significantly in the past decade. A considerable number of herbal constituents whose behavioral effects and pharmacological actions have been well characterized may be good candidates for further investigations that may ultimately lead to clinical use of these constituents. The potential benefits of herbal remedies such as St. John's wort and Kava-kava in psychiatric practice have been addressed previously [34]. EFC may be another good candidate for the treatment of psychiatric illnesses, such as sleep disorders.

5. Conclusions

EFC enhanced hypnotic effects in pentobarbital-treated mice. This enhancement may result from Cl⁻ channel activation. Further investigation is needed to determine the effects of other EFC derivatives with strong pharmacological action.

Acknowledgment

This work was supported by a grant from the Korean Ministry of Education, Science and Technology (The Regional Core Research Program/Center for Healthcare Technology Development).

References

- [1] J. P. Lai, Y. H. Lim, J. Su, H. M. Shen, and C. N. Ong, "Identification and characterization of major flavonoids and caffeoylquinic acids in three Compositae plants by LC/DAD-APCI/MS," *Journal of Chromatography B*, vol. 848, no. 2, pp. 215–225, 2007.
- [2] Q. Chu, L. Fu, Y. Guan, and J. Ye, "Determination and differentiation of Flos Chrysanthemum based on characteristic electrochemical profiles by capillary electrophoresis with electrochemical detection," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 26, pp. 7828–7833, 2004.
- [3] H. Kim and Y. S. Lee, "Identification of new dicaffeoylquinic acids from *Chrysanthemum morifolium* and their antioxidant activities," *Planta Medica*, vol. 71, no. 9, pp. 871–876, 2005.
- [4] H. Jiang, Q. Xia, W. Xu, and M. Zheng, "*Chrysanthemum morifolium* attenuated the reduction of contraction of isolated rat heart and cardiomyocytes induced by ischemia/reperfusion," *Pharmazie*, vol. 59, no. 7, pp. 565–567, 2004.
- [5] M. Miyazawa and M. Hisama, "Antimutagenic activity of flavonoids from *Chrysanthemum morifolium*," *Bioscience, Biotechnology and Biochemistry*, vol. 67, no. 10, pp. 2091–2099, 2003.
- [6] M. Ukiya, T. Akihisa, K. Yasukawa et al., "Constituents of Compositae plants. 2. Triterpene diols, triols, and their 3-O-fatty acid esters from edible *Chrysanthemum* flower extract and their anti-inflammatory effects," *Journal of Agricultural and Food Chemistry*, vol. 49, no. 7, pp. 3187–3197, 2001.
- [7] L. Z. Lin and J. M. Harnly, "Identification of the phenolic components of chrysanthemum flower (*Chrysanthemum morifolium* Ramat)," *Food Chemistry*, vol. 120, no. 1, pp. 319–326, 2010.
- [8] C. W. Beninger, M. M. Abou-Zaid, A. L. E. Kistner et al., "A flavanone and two phenolic acids from *Chrysanthemum morifolium* with phytotoxic and insect growth regulating activity," *Journal of Chemical Ecology*, vol. 30, no. 3, pp. 589–606, 2004.
- [9] M. G. L. Hertog, E. J. M. Feskens, P. C. H. Hollman, M. B. Katan, and D. Kromhout, "Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study," *The Lancet*, vol. 342, no. 8878, pp. 1007–1011, 1993.
- [10] M. M. Ohayon, "Methodology of a study on insomnia in the general population," *Encephale*, vol. 28, no. 3 I, pp. 217–226, 2002.
- [11] C. M. Morin and S. E. Gramling, "Sleep patterns and aging: comparison of older adults with and without insomnia complaints," *Psychology and aging*, vol. 4, no. 3, pp. 290–294, 1989.
- [12] A. Chistina Grobin, J. R. Inglefield, R. D. Schwartz-Bloom, L. L. Devaud, and A. L. Morrow, "Fluorescence imaging of GABA_A receptor-mediated intracellular [Cl⁻] in P19-N cells reveals unique pharmacological properties," *Brain Research*, vol. 827, no. 1-2, pp. 1–11, 1999.
- [13] H. Möhler, J. M. Fritschy, and U. Rudolph, "A new benzodiazepine pharmacology," *Journal of Pharmacology and Experimental Therapeutics*, vol. 300, no. 1, pp. 2–8, 2002.
- [14] I. Tobler, C. Kopp, T. Deboer, and U. Rudolph, "Diazepam-induced changes in sleep: role of the $\alpha 1$ GABA(A) receptor subtype," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 11, pp. 6464–6469, 2001.
- [15] U. Rudolph, F. Crestani, and H. Möhler, "GABA(A) receptor subtypes: dissecting their pharmacological functions," *Trends in Pharmacological Sciences*, vol. 22, no. 4, pp. 188–194, 2001.
- [16] S. Zhu and R. C. Baker, "Effects of inhalation anesthetics on kainate-induced glutamate release from cerebellar granule cells," *Life Sciences*, vol. 58, no. 16, pp. 1359–1366, 1996.
- [17] P. Follès, P. Porcu, C. Sogliano et al., "Changes in GABA_A receptor γ subunit gene expression induced by long-term administration of oral contraceptives in rats," *Neuropharmacology*, vol. 42, no. 3, pp. 325–336, 2002.
- [18] V. Darias, S. Abdala, D. Martin-Herrera, M. Luisa Tello, and S. Vega, "CNS effects of a series of 1,2,4-triazolyl heterocarboxylic derivatives," *Pharmazie*, vol. 53, no. 7, pp. 477–481, 1998.
- [19] C. Wolfman, H. Viola, M. Marder et al., "Anxiolytic properties of 6,3'-dinitroflavone, a high-affinity benzodiazepine receptor ligand," *European Journal of Pharmacology*, vol. 318, no. 1, pp. 23–30, 1996.
- [20] M. R. West and C. R. Molloy, "A microplate assay measuring chloride ion channel activity," *Analytical Biochemistry*, vol. 241, no. 1, pp. 51–58, 1996.
- [21] J. Glowinski and L. L. Iversen, "Regional studies of catecholamines in the rat brain. I. The disposition of [3H]norepinephrine, [3H]dopamine and [3H]dopa in various regions of the brain," *Journal of Neurochemistry*, vol. 13, no. 8, pp. 655–669, 1966.
- [22] D. S. Segal and R. Kuczenski, "Tyrosine hydroxylase activity: regional and subcellular distribution in brain," *Brain Research*, vol. 68, no. 2, pp. 261–266, 1974.
- [23] H. Han, Y. Ma, J. S. Eun et al., "Anxiolytic-like effects of sanjoinine A isolated from *Zizyphi spinosi* Semen: possible involvement of GABAergic transmission," *Pharmacology Biochemistry and Behavior*, vol. 92, no. 2, pp. 206–213, 2009.
- [24] F. C. F. De Sousa, B. A. Pereira, V. T. M. Lima et al., "Central nervous system activity of yangambin from *Ocotea duckei* Vattimo (Lauraceae) in mice," *Phytotherapy Research*, vol. 19, no. 4, pp. 282–286, 2005.
- [25] Y. Ma, H. Han, J. S. Eun, H. C. Kim, J. T. Hong, and K. I. W. Oh, "Sanjoinine A isolated from *Zizyphi spinosi* Semen augments pentobarbital-induced sleeping behaviors through the modification of GABA-ergic systems," *Biological and Pharmaceutical Bulletin*, vol. 30, no. 9, pp. 1748–1753, 2007.
- [26] A. L. Martínez, F. Domínguez, S. Orozco et al., "Neuropharmacological effects of an ethanol extract of the *Magnolia dealbata* Zucc. leaves in mice," *Journal of Ethnopharmacology*, vol. 106, no. 2, pp. 250–255, 2006.
- [27] Y. Ma, H. Ma, Y. J. Jo et al., "Honokiol potentiates pentobarbital-induced sleeping behaviors through GABA_A receptor Cl channel activation," *Biomolecules and Therapeutics*, vol. 16, no. 4, pp. 328–335, 2008.
- [28] H. Han, Y. Ma, J. S. Eun, J. T. Hong, and K. I. W. Oh, "Anxiolytic-like effects of cyclopeptide fraction alkaloids of *Zizyphi spinosi* semen: possible involvement of GABA_A receptors," *Biomolecules and Therapeutics*, vol. 16, no. 3, pp. 261–269, 2008.
- [29] R. L. Macdonald and R. W. Olsen, "GABA_A receptor channels," *Annual Review of Neuroscience*, vol. 17, pp. 569–602, 1994.
- [30] U. Rudolph and H. Möhler, "GABA-based therapeutic approaches: GABA_A receptor subtype functions," *Current Opinion in Pharmacology*, vol. 6, no. 1, pp. 18–23, 2006.
- [31] N. J. K. Tillakaratne, L. Medina-Kauwe, and K. M. Gibson, "Gamma-aminobutyric acid (GABA) metabolism in mammalian neural and nonneural tissues," *Comparative Biochemistry and Physiology A: Physiology*, vol. 112, no. 2, pp. 247–263, 1995.

- [32] C. Buddhala, C. C. Hsu, and J. Y. Wu, "A novel mechanism for GABA synthesis and packaging into synaptic vesicles," *Neurochemistry International*, vol. 55, no. 1–3, pp. 9–12, 2009.
- [33] M. Chebib and G. A. R. Johnston, "GABA-activated ligand gated ion channels: medicinal chemistry and molecular biology," *Journal of Medicinal Chemistry*, vol. 43, no. 8, pp. 1427–1447, 2000.
- [34] Z. J. Zhang, "Therapeutic effects of herbal extracts and constituents in animal models of psychiatric disorders," *Life Sciences*, vol. 75, no. 14, pp. 1659–1699, 2004.