

RESEARCH

Open Access

Evidence of multiple mechanisms providing carbamate and organophosphate resistance in field *An. gambiae* population from Atacora in Benin

Rock Aïkpon^{1,2*}, Michel Sèzonlin², Razaki Ossé^{1,2} and Martin Akogbéto^{1,2}

Abstract

Background: Insecticide resistance in *Anopheles gambiae* s.l is a major concern to malaria vector control programmes. In West Africa, resistance is mainly due to target-site insensitivity arising from a single point mutation. Metabolic-based resistance mechanisms have also been implicated and are currently being investigated in west Africa. The aim of this study is to better understand the origins of carbamate and organophosphate resistance in *An. gambiae* population from Atacora, Benin in West Africa.

Methods: Anopheles mosquitoes were reared from larvae collected in two districts (Kouandé and Tanguiéta) of the Atacora department. Mosquitoes were then exposed to WHO impregnated papers. Four impregnated papers were used: carbamates (0.1% bendiocarb, 0.1% propoxur) and organophosphates (0.25% pirimiphos methyl, 1% fenitrothion). PCR assays were run to determine the members of the *An. gambiae* complex, as well as phenotypes for insensitive acetylcholinesterase (AChE1). Biochemical assays were also carried out to detect any increase in the activity of enzyme typically involved in insecticide metabolism (oxidase, esterase and glutathion-S-transferase).

Results: 769 female of *An. gambiae* mosquitoes from Kouandé and Tanguiéta were exposed to bendiocarb, propoxur, pirimiphos methyl and fenitrothion. Bioassays showed resistance with low mortality to bendiocarb (78.57% to 80.17%), propoxur (77.21% to 89.77%), and fenitrothion (89.74% to 92.02%). On the other hand, the same populations of *An. gambiae* from Kouandé and Tanguiéta showed high susceptibility to pirimiphos methyl with recorded mortality of 99.02% and 100% respectively. The low rate of ace-1R allele frequency (3.75% among survivors and 0.48% among dead) added to the high proportion of homozygous susceptible specimens which survived the WHO bioassays (8/28), suggest that the ace-1 mutation could not entirely explain *Anopheles gambiae* resistance to carbamate and organophosphate. Biochemical assays suggest that resistance in this population is mediated by metabolic resistance with elevated level of GST, MFO and NSE compared to a susceptible strain *An. gambiae* Kisumu.

Conclusions: *Anopheles gambiae* populations resistance from Atacora is multifactorial and includes target-site mutation and metabolic mechanism. The co-implication of both resistance mechanisms in *An. gambiae* s.l may be a serious obstacle for the future success of malaria control operations based on LLINs and IRS.

Keywords: Resistance mechanisms, Carbamate, Organophosphate, *Anopheles gambiae*, Benin

* Correspondence: rockypremier@yahoo.fr

¹Centre de Recherche Entomologique de Cotonou (CREC), 06 BP 2604, Cotonou, Bénin

²Faculté des Sciences et Techniques, Université d'Abomey Calavi, Calavi, Bénin

Background

Malaria is a main cause of mortality and morbidity and *Anopheles gambiae* is one of the major vectors of this disease in sub-Saharan Africa [1]. The current effective vector control tools include the use of Long Lasting Insecticide Nets (LLIN) and Indoor Residual Spraying (IRS) [2]. The success of such interventions requires a good knowledge of vector populations particularly their susceptibility status to the main insecticides used for such control programmes in order to detect and monitor resistance to these insecticides.

Pyrethroids are the only group of insecticides currently recommended for net treatment and the other classes of insecticides (organochlorine, carbamate and organophosphate) are applied for IRS [3,4]. The main problem with LLINs and IRS is the development of insecticide resistance, particularly pyrethroid-resistance by several populations of *An. gambiae* [5-8]. Recently, the emergence of resistance in populations of *An. gambiae* to common classes of insecticides used in public health has been reported in many countries in Africa, including Côte d'Ivoire [5,9], Kenya [10], Benin [11,12], Niger [13], Burkina Faso [14], Mali [15], Nigeria [16,17], South Africa [18] and Cameroon [19]. More recently in Benin, several studies pointed out *An. gambiae* resistance to bendiocarb [20,21] and showed that the mean *Ace-1* mutation frequency had increased significantly from 2010 to 2012 after two years of IRS campaign using bendiocarb in Atacora region in the northern part of the country [22,23]. However, these studies clearly showed the evidence of bendiocarb resistance in *An. gambiae* populations but mentioned the *ace-1R* mutation could not entirely explain the resistance to bendiocarb observed.

The aim of this study is to better understand the origins of carbamate and organophosphate resistance in *An. gambiae* population from Atacora. We conducted bioassays, molecular and biochemical analysis to update resistance status in *An. gambiae* and detect other potential mechanisms of resistance that might have evolved.

Methods

Study area

The study was carried out in Atacora region located in the North-west of Benin and includes two districts: Kouandé and Tanguiéta (Figure 1). The two districts covered 5,128 km² and had an estimated population of 198,956 in 2012. Atacora region has a sub-equatorial type climate with one dry season (December-May) and only one rainy season (June to November). The annual mean rainfall is 1,300 mm and the mean monthly temperature ranges between 22 and 33°C. The region is irrigated by three major rivers: the Mekrou, the Pendjari and the Alibori. The major economic activity is agriculture and it is characterized by the production of cotton

and millet where various classes of pesticides are used for pest control. Since 2011 onwards the department has conducted a large scale Indoor Residual Spraying (IRS) campaign.

Mosquito collections

Anopheles gambiae s.l. larvae were collected in 2 districts and in each district, two villages were selected randomly. At each locality chosen, Anopheline larvae were collected from various natural breeding sites including ground pools, gutters, puddles and abandoned potholes, at the beginning of the rainy season during June 2014. Water was scooped using a plastic scoop and poured into small transparent plastic bowls. A strainer was used to sieve and pool together the third and fourth instar larvae in order to have sufficient adult emergence of the same physiological age. The mosquito larvae collected were transported in well labelled plastic bottles to the laboratory of the Centre de Recherche Entomologique de Cotonou, Benin (CREC) where they were maintained at 28 ± 2°C and 72 ± 5% relative humidity. A laboratory susceptible strain of *An. gambiae* Kisumu was used as a reference strain to compare the susceptibility levels of the field populations.

Insecticide susceptibility tests

Mosquitoes collected were assayed using WHO discriminating dosages with four insecticides of technical grade quality: two carbamates (0.1% propoxur, 0.1% bendiocarb) and two organophosphates (1% fenitrothion, 0.25% pirimiphos méthyl). Four batches of 25 unfed females, aged 2–5 days, were exposed to the diagnostic doses of insecticide treated papers for 60 min at 27 ± 1°C and 80% relative humidity. The twenty-five females of *An. gambiae* were introduced into each tube and monitored at different time intervals (10, 15, 20, 30, 45, 60 minutes), the number "knocked-down" were recorded. After one hour exposure, mosquitoes were transferred into holding tubes and provided with cotton wool soaked with a 10% honey solution. Batches exposed to untreated papers were used as control. Mortalities were recorded after 24 hours and the susceptibility status of the population was graded according to the WHO protocol [24]. Dead and surviving mosquitoes from this bioassay were kept separately in eppendorf tubes containing silica gel and stored at -20°C for further molecular analysis.

Species identification and PCR detection of *Ace-1R* mutation

Live and dead specimens of *An. gambiae* from the bioassay tests were subjected to the *An. gambiae* species specific PCR assays for species identification [25,26]. The PCR-RFLP diagnostic test was used to detect the presence of G119S mutation (*ace-1R* gene). Mosquito genomic DNA was amplified using the primers




Figure 1 Map of the department of Atacora showing the localities where Anopheline mosquitoes were collected.

Ex3AGdir 5' GATCGTGGACACCGTGTTCG3' and Ex3AGrev 5'AGGATGGCCCGCTGGAACAG3' according to [27]. One microlitre of total DNA extracted from a single mosquito was used as a template in a 25 ml PCR reaction containing Taq DNA polymerase buffer, 0.2 mM dNTP and 10 pmol of each primer. The PCR conditions were 94°C for 5 min and then 35 cycles of (94°C for 30 s, 54°C for 30 s and 72°C for 30 s) with a final 5 min extension at 72°C. Fifteen microlitres of PCR product were digested with 5U of AluI restriction enzyme (Promega) in a final volume of 25 ml. The PCR fragments were fractionated on a 2% agarose gel stained with ethidium bromide and visualized under UV light.

Biochemical analysis

Biochemical assays were performed to compare the levels of activity of mixed function oxidases (MFO),

non-specific esterases (NSE) using α -naphthyl acetate as a substrate and glutathione S-transferases (GST) [28] in the *An. gambiae* s.s. susceptible Kisumu and the field population from Tanguiéta and Kouandé districts. Mosquitoes used for the biochemical analysis have not been exposed to any insecticides prior to the assay.

Detoxifying enzyme activities were measured on single mosquitoes ($N = 50$) from each test locality, which were stored at 280uC within 24 h from emergence (above). Each mosquito was ground on ice in 200 μ l of distilled water and the homogenate was centrifuged at 1,4000 rpm for 2 mins. Two 10 ml replicates of supernatant were transferred into two adjacent wells of a microtitre plate for non-specific esterases (NSEs), glutathione S-transferases (GSTs). Monooxygenases assays were performed with two 20 ml replicates of supernatant.




Figure 2 Mortality rates in *Anopheles gambiae* 24-hours post exposure to propoxur 0.1%, bendiocarb 0.1%, fenitrothion 1% and pirimiphos methyl 0.25%. S = susceptible; R = resistant; RS = resistance suspected.

Non-specific esterases (NSEs)

Non-specific esterase activity was measured using a-naphthol acetate (aN_a) and b-naphthol acetate (bN_a). In each replicate well, 90 ml of phosphate buffer (PBS, pH = 6.5) and 100 ml of 0.6 M aN_a (or bN_a) were added to 10 ml of centrifuged mosquito homogenate. After 30 min incubation, 100 ml of Fast Garnett BC solution (8 g Fast Garnett Salt +10 ml distilled water) was added to stop the reaction. The concentration of the final product was determined at 550 nm as an endpoint calculated from standard curves of a- and b-Naphthol, respectively.

Glutathione-S-transferases (GST)

To measure glutathione- S-transferase (GST) activity in mosquitoes, 200 ml of GSH/CDNB working solution (100 ml of an extemporaneous solution of 0.6% weight/volume reduced glutathione in 0.1 M sodium phosphate buffer pH = 6.5 + 0.013 g of 1-chloro-2, 4-dinitrobenzene diluted in 1 ml of 70% methanol) were added to each replicate of mosquito homogenate. The reaction was read at 340 nm immediately as a kinetic assay for 5 minutes. An extinction coefficient of 5.76 mM⁻¹ (corrected for a path length of 0.6 cm) was used to convert absorbance values to moles of product. GST specific activity was reported as the rate of formation of GSH produced in mmol·min⁻¹·mg⁻¹ protein.

Oxydases (Cytochrome P450)

Cytochrome P450 activity was determined using the heme-peroxidase assay according to Brogdon et al. [29]. The assay detects the elevation in the amount of heme, which is then converted into equivalent units of cytochrome P450. Eighty ml of 0.625 M potassium phosphate buffer (pH = 7.2) were added to 20 ml of mosquito homogenate together with 200 ml Tetramethyl Benzidine solution (0.011 g 3,3',5,5' Tetramethyl Benzidine in 5 ml of

70% methanol +15 ml sodium acetate buffer 0.25 M pH = 5.0); 25 ml of 3% hydrogen peroxide were then added and the mixture was incubated for 30 min at room temperature. Absorbance was read at 630 nm and values calculated from a standard curve of cytochrome C.

Data analysis

The resistant status of mosquito samples was determined according to the WHO criteria [23]:

- Mortality rate is > 98%: the population was considered fully susceptible
- Mortality rates ranged between 90 - 98%: resistance suspected in the population
- Mortality rates < 90%, the population was considered resistant to the tested insecticides

Allelic frequencies of G119S mutation were analysed using the version 1.2 of Genepop [30].

To assess if the mutation frequencies were identical across populations, the test of genotypic differentiation was performed [31]. Biochemical assay data (enzymatic activity per mg of protein) of wild specimens of *An. gambiae* s.l were compared to the Kisumu susceptible strain using Mann-Whitney non-parametric *U*-test (Statistica software).

Results

Bendiocarb and organophosphate susceptibility status of *Anopheles gambiae* s.l

A total of 769 female of *An. gambiae* mosquitoes (3–5 days old) from Kouandé and Tanguiéta were exposed to bendiocarb (n = 214), propoxur (n = 167), pirimiphos methyl (n = 209) and fenitrothion (n = 179). Mortality rates of the Kisumu reference strain to all insecticides was 100% (Figure 2). In contrast, wild mosquitoes were

high resistant with mortality rates less than 90% for bendiocarb (80.17% and 78.57%) and propoxur (77.21% and 89.77%) in Kouandé and Tanguiéta respectively (Figure 2). As for the organophosphates, susceptibility to pirimiphos-methyl was assessed on all populations. Mortality rate were higher than 98% as per WHO criteria with 99.02% recorded in Kouandé and 100% in Tanguiéta. However, fenitrothion resistance was detected in Kouandé and Tanguiéta, with mortality rates of 92.07% and 89.74% respectively.

PCR- species identification

One hundred mosquitoes were identified to *Anopheles gambiae* species. All PCR analysis identifying *An. gambiae* s.l species in this study confirmed that all mosquitoes belonging to *An. gambiae* s.l were *An. gambiae*.

Involvement of ace-1R mutation in providing carbamate and organophosphate resistance

To assess the role of the ace-1R allele in conferring carbamate and organophosphate resistance in *An. gambiae* population, the ace-1R genotype was determined for dead and alive mosquitoes detected in the WHO bioassay using carbamate and organophosphate (Figure 3). A total of 59 mosquitoes were tested of which 31 dead and 28 alive. Among both bioassay survivors and non-survivors, all ace-1R genotypes (RR, RS and SS) were found. However, the homozygous resistant genotype RR was only found among the bioassay survivors, and the heterozygote genotype RS was the most prevalent genotype among the bioassay survivors. Although there was a significant ace-1R genotype differentiation between bioassay survivors and non-survivors ($p < 0.0001$), homozygous susceptible mosquitoes were found among bioassay survivors.

Biochemical assays

Biochemical assays were successfully performed on the two *An. gambiae* populations (Tanguiéta and Kouandé) from Atacora. Figure 4 shows the mean level of enzymatic

activity (MFO, NSE and GST) in field-collected mosquitoes compared to the reference susceptible strain Kisumu. All sampled populations showed a significantly higher MFOs activity (expressed in cytochrome P450 units) than the susceptible reference strain Kisumu ($p < 0.005$). The level of esterase activity (using alpha and beta naphthyl acetate as a substrate) did not reveal any significant differences between Kisumu and Kouandé *An. gambiae* population ($p = 0.65$). Overall, the mean level of alpha and beta esterase activity in Tanguiéta was significantly higher than that of the susceptible strain ($p < 0.05$). Besides, all sampled populations showed a significantly higher GST activity than the susceptible reference strain Kisumu ($p < 0.05$).

Discussion

This study highlighted the diversity of carbamate and organophosphate resistance in *Anopheles gambiae* s.l. populations from Atacora in Benin. The results presented here show that carbamate and organophosphate resistance in sampled *Anopheles gambiae* populations is multifactorial and includes target-site mutation and metabolic mechanism in this region.

An. gambiae displayed large variations in resistance levels to carbamates and organophosphates. Although the wild populations were all resistant to bendiocarb, resistance was less marked to propoxur and fenitrothion, at WHO diagnostic concentrations. However, all these populations were very susceptible to pirimiphos-methyl. This resistance of the mosquito population to carbamate and fenitrothion would be due to the strong selective pressure that represents the use of insecticides in households for public health purposes, notably IRS using bendiocarb [21] and massive quantities of carbamates and organophosphates in agricultural settings in the department of Atacora. Indeed, in the cotton growing areas in Atacora, farmers use huge amounts of insecticides to avoid substantial yield reduction of their crops. Several studies showed that agricultural practices seem to have




Figure 3 Ace-1 genotypes distribution among live and dead *An. gambiae* s.s individuals from WHO susceptibility test to carbamate and organophosphate.




Figure 4 Detoxifying enzyme activities in *An. gambiae* mosquitoes collected from two districts in Atacora department.

contributed to the emergence of insecticide resistance in *Anopheles* populations [8,12,32].

Cross-resistance to organophosphates and carbamates suggests the involvement of their common target-site: AChE-1 [33]. The high proportion of homozygous susceptible specimens, which survived the WHO bioassays, added to the low rate of ace-1R allele frequency suggest that the ace-1 mutation could not entirely explain *Anopheles gambiae* resistance to carbamate and organophosphate.

In the present study, the low ace-1 mutation frequency in sampled *An. gambiae* populations associated with the resistance to carbamate and organophosphate strongly supports the involvement of metabolic resistance based on the high activities of typical enzymes NSE, GST and MFO. Similar findings have been reported in *Culex quinquefasciatus* and *Anopheles gambiae* in four other sites in Benin, where greater oxidase and esterase activities were observed in *Cx. quinquefasciatus* and *An. gambiae*, but where ace-1 was absent [34]. The same findings were also reported in the field experimental station of Pito (Cameroun) [35,36].

Considering the multiple insecticide resistance mechanisms in *An. gambiae* s.l population from Atacora, resistance management becomes a particularly important issue. Insecticide rotation or synergistic use of different classes of insecticides to enhance the killing effects are being considered as alternative methods of vector control to slow down the spread of resistance genes.

Conclusions

The present study revealed the simultaneous presence of multiple resistance mechanisms in malaria vector *An. gambiae* s.l population from Atacora. The co-implication of both metabolic and ace-1 resistance mechanisms in *An. gambiae* s.l may be a serious obstacle for the future success of malaria control operations based on LLINs and IRS. Alternative innovative vector control tools are therefore urgently needed to manage these complex mechanisms and to complement insecticide-based strategies.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RA, MS, RO, and MA designed the study. RA and MA carried out the field activities. RA drafted the manuscript and analyzed the data. RO and MS critically revised the manuscript. RA conceived and designed the study and revised the manuscript for intellectual content. All authors read and approved the final manuscript.

Acknowledgements

This work was financially supported by PMI (President's Malaria Initiative) through USAID. We thank the team of CREC for their technical assistance during field work. We also thank the people of Atacora for their collaboration.

Received: 28 September 2014 Accepted: 24 November 2014

Published online: 02 December 2014

References

1. Gillies MT, Coetzee M: A supplement to the anophelinae Africa South of the Sahara (Afrotropical region), Johannesburg, South Africa. *S Afr Inst Med Res* 1987, 55:1–143.
2. Beier JC, Keating J, Githure JI, Macdonald MB, Impoinvil DE, Novak RJ: Integrated vector management for malaria control. *Malaria J* 2008, 7(suppl):54.
3. Guillet P, Chandre F, Mouchet J: L'utilisation des insecticides en santé publique: état et perspectives. *Med Mal Infect* 1997, 27:552–557.
4. Devine GJ, Oogusuku E: Adaptability is key when monitoring insecticide resistance. *Bull World Health Organ* 2009, 87:887.
5. Elissa N, Mouchet J, Riviere F, Meunier JY, Yao K: Resistance of *Anopheles gambiae* s.s. to pyrethroids in Côte-d'Ivoire. *Ann Soc Belge Med Trop* 1993, 73:291–294.
6. Akogbeto M, Yakoubou S: Résistance des vecteurs du paludisme vis-à-vis des pyréthrinoïdes utilisés pour l'imprégnation des moustiquaires au Bénin, Afrique de l'Ouest. *Bull Soc Pathol Exot* 1999, 92:123–130.
7. Chandre F, Darriet F, Duchon S, Finot L, Manguin S, Carnevale P, Guillet P: Modifications of pyrethroid effects associated with kdr mutation in *Anopheles gambiae*. *Med Vet Entomol* 2000, 14:81–88.
8. Akogbeto M, Djouaka RF, Kinde-Gazard DA: Screening of pesticide residues in soil water samples from agricultural settings. *Malaria J* 2006, 5:22.
9. Ahoua-Alou LP, Koffi AA, Adja MA, Tla E, Kouassi PK, Kone M, Chandre F: Distribution of ace-1R and resistance to carbamates and organophosphates in *Anopheles gambiae* s.s. populations from Côte d'Ivoire. *Malaria J* 2010, 9:167.
10. Vulule JM, Beach RF, Atieli FK, McAllister JC, Brogdon WG, Roberts JM, Mwangi RW, Hawley WA: Elevated oxidase and esterase levels associated with permethrin tolerance in *Anopheles gambiae* from Kenyan villages using permethrin impregnated nets. *Med Vet Entomol* 1999, 13:239–244.
11. Aïzoun N, Aïkpon R, Padonou GG, Oussou O, Oké-Agbö F, Gnanguenon V, Ossè R, Akogbeto M: Mixed-function oxidases and esterases associated with permethrin, deltamethrin and bendiocarb resistance in *Anopheles gambiae* s.l. in the south-north transect Benin, West Africa. *Parasit Vectors* 2013, 6:223.
12. Yadouleton AW, Asidi A, Djouaka RF, Braima J, Agossou CD, Akogbeto MC: Development of vegetable farming: a cause of the emergence of insecticide resistance in populations of *Anopheles gambiae* in urban areas of Benin. *Malaria J* 2009, 8:103.
13. Czeher C, Labbo R, Arzika I, Duchemin JB: Evidence of increasing Leu-Phe knockdown resistance mutation in *Anopheles gambiae* from Niger following a nationwide longlasting insecticide-treated nets implementation. *Malaria J* 2008, 7:189.
14. Diabate A, Baldet T, Chandre F, Guiguemde RT, Brengues C, Guillet P, Hemingway J, Hougard JM: First report of the kdr mutation in *Anopheles gambiae* M form from Burkina Faso, West Africa. *Parassitologia* 2002, 44:157–158.
15. Fanello C, Petrarca V, Della Torre A, Santolamazza F, Dolo G, Coulibaly M, Allouche A, Curtis CG, Toure YT, Coluzzi M: The pyrethroid knock-down resistance gene in the *Anopheles gambiae* complex in Mali and further indication of incipient speciation within An. gambiae s.s. *Insect Mol Biol* 2003, 12:241–245.
16. Awolola TS, Brooke BD, Koekemoer LL, Coetzee M: Resistance of the malaria vector *Anopheles gambiae* s.s. to pyrethroid insecticides, in south-western Nigeria. *Ann Trop Med Parasitol* 2002, 96:849–852.
17. Oduola AO, Idowu ET, Oyebola MK, Adeogun AO, Olojede JB, Otubanjo OA, Awolola TS: Evidence of carbamate resistance in urban populations of *Anopheles gambiae* s.s. mosquitoes resistant to DDT and deltamethrin insecticides in Lagos, South-Western Nigeria. *Parasit Vectors* 2012, 5:116.
18. Hargreaves K, Koekemoer LL, Brooke B, Hunt RH, Mthembu J, Coetzee M: *Anopheles funestus* resistant to pyrethroid insecticides in South Africa. *Med Vet Entomol* 2000, 14:181–189.
19. Etang J, Manga L, Chandre F, Guillet P, Fondjo E, Mimpfoundi R, Toto JC, Fontenille D: Insecticide susceptibility status of *Anopheles gambiae* s.l. (Diptera: Culicidae) in the Republic of Cameroon. *J Med Entomol* 2003, 40:491–497.
20. Aïkpon R, Agossa F, Ossè R, Oussou O, Aïzoun N, Oké-Agbö F, Akogbeto M: Bendiocarb resistance in *Anopheles gambiae* s.l. populations from Atacora department in Benin, West Africa: a threat for malaria vector control. *Parasit Vectors* 2013, 6:192.
21. Aïzoun N, Aïkpon R, Gnanguenon V, Oussou O, Agossa F, Padonou GG, Akogbeto M: Status of organophosphate and carbamate resistance in *Anopheles gambiae* sensu lato from the south and north Benin, West Africa. *Parasit Vectors* 2013, 6:274.
22. Aïkpon R, Aïzoun N, Sovi A, Oussou O, Govoetchan R, Gnanguenon V, Oké-Agbö R, Ossè R, Akogbeto M: Increase of Ace-1 resistance allele in the field population of *Anopheles gambiae* following a large scale indoor residual spraying (IRS) implementation using bendiocarb in Atacora region in Benin, West Africa. *J Cell Anim Biol* 2014, 8(1):15–22.
23. Aïkpon R, Aïzoun N, Ossè R, Oké-Agbö R, Oussou O, Govoetchan R, Sovi A, Akogbeto M: Seasonal variation of Ace-1R mutation in *Anopheles gambiae* s. l. populations from Atacora region in Benin, West Africa. *J Entomol Nematol* 2014, 6(1):14–18.
24. WHO: Guidelines for Testing Mosquito Adulticides Intended for Indoor Residual Spraying (IRS) and Insecticide Treated Nets (ITNs). Geneva: WHO/CDS/NTD/WHO/PDF/GCDDP/2006.3; 2006:2.
25. Scott J, Brogdon W, Collins F: Identification of single specimens of the *Anopheles gambiae* complex by PCR. *Am J Trop Med Hyg* 1993, 49:520–529.
26. Favia G, Lanfrancotti A, Spanos L, Siden-Kiamos I, Louis C: Molecular characterization of ribosomal DNA polymorphisms discriminating among chromosomal forms of *Anopheles gambiae* s.s. *Insect Mol Biol* 2001, 10:19–23.
27. Weill M, Lutfalla G, Mogensen K, Chandre F, Berthomieu A, Berticat C, Pasteur N, Philips A, Fort P, Raymond M: Insecticide resistance in mosquito vectors. *Nature* 2003, 423:136–137.
28. Hemingway J: Techniques to Detect Insecticide Resistance Mechanisms (Field and Laboratory Manual). Document WHO/CDS/CPC/MAL/98.6. Geneva, Switzerland: World Health Organ; 1998.
29. Brogdon WG, McAllister JC: Insecticide resistance and vector control. *Emerg Infect Dis* 1998, 4:605–613.
30. Raymond M, Rousset F: Genepop (version 1.2), population genetics software for exact tests and ecumenicism. *J Heredity* 1995, 86:248–249.
31. Goudet J, Raymond M, De Meeùs T, Rousset F: Testing differentiation in diploid populations. *Genetics* 1996, 144:1933–1940.
32. Diabaté A, Baldet T, Chandre F, Akogbeto M, Guiguemde TR, Darriet F, Brengues C, Guillet P, Hemingway J, Small GJ, Hougard JM: The role of agricultural use of insecticides in resistance to pyrethroids in *Anopheles gambiae* s.l. in Burkina Faso. *Am J Trop Med Hyg* 2002, 67:617–622.
33. Weill M, Malcolm C, Chandre F, Mogensen K, Berthomieu A, Marquine M, Raymond M: The unique mutation in Ace-1 giving high insecticide resistance is easily detectable in mosquito vectors. *Insect Mol Biol* 2004, 13:1–7.
34. Corbel V, N'Guessan R, Brengues C, Chandre F, Djogbénou L, Martin T, Akogbeto M, Hougard JM, Rowland M: Multiple insecticide resistance mechanisms in *Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta Trop* 2007, 101:207–216.
35. Etang J, Manga L, Toto JC, Guillet P, Fondjo E, Chandre F: Spectrum of metabolic-based resistance to DDT and pyrethroids in *Anopheles gambiae* s.l. populations from Cameroon. *J Vect Ecol* 2007, 32:123–133.
36. Chouaiou M, Simard F, Chandre F, Etang J, Darriet F, Hougard JM: Efficacy of bifenthrin-impregnated bednets against *Anopheles funestus* and pyrethroid-resistant *Anopheles gambiae* in North Cameroon. *Malar J* 2006, 5:77.

doi:10.1186/s13071-014-0568-5

Cite this article as: Aïkpon et al.: Evidence of multiple mechanisms providing carbamate and organophosphate resistance in field An. gambiae population from Atacora in Benin. *Parasites & Vectors* 2014 7:568.