

# Modification of neuropathic pain sensation through microglial ATP receptors

Kazuhide Inoue · Makoto Tsuda ·  
Hidetoshi Tozaki-Saitoh

Received: 31 May 2007 / Accepted: 2 August 2007 / Published online: 25 August 2007  
© Springer Science + Business Media B.V. 2007

**Abstract** Neuropathic pain that typically develops when peripheral nerves are damaged through surgery, bone compression in cancer, diabetes, or infection is a major factor causing impaired quality of life in millions of people worldwide. Recently, there has been a rapidly growing body of evidence indicating that spinal glia play a critical role in the pathogenesis of neuropathic pain. Accumulating findings also indicate that nucleotides play an important role in neuron-glia communication through P2 purinoceptors. Damaged neurons release or leak nucleotides including ATP and UTP to stimulate microglia through P2 purinoceptors expressing on microglia. It was shown in an animal model of neuropathic pain that microglial P2X<sub>4</sub> and P2X<sub>7</sub> receptors are crucial in pain signaling after peripheral nerve lesion. In this review, we describe the modification of neuropathic pain sensation through microglial P2X<sub>4</sub> and P2X<sub>7</sub>, with the possibility of P2Y<sub>6</sub> and P2Y<sub>12</sub> involvement.

**Keywords** Allodynia · ATP · Microglia · Neuropathic pain · P2X<sub>4</sub> · Spinal cord

## Abbreviations

ADP	adenosine 5'-diphosphate
ATP	adenosine 5'-triphosphate
BDNF	brain-derived neurotrophic factor
BzATP	2'- and 3'-O-(4-benzoylbenzoyl) adenosine 5'-triphosphate
[Ca <sup>2+</sup> ] <sub>i</sub>	intracellular Ca <sup>2+</sup> concentration
CNS	central nervous system

CR3	complement receptor 3
IL-1β	interleukin-1β
IL-6	interleukin-6
OATP	oxidized ATP
PK11195	[1-(2-chlorophenyl)-N-methyl-N-(1-methylpropyl)-3-isoquinolineiso-quinoline carboxamide]
PPADS	pyridoxalphosphate-6-azophenyl-2', 4'-disulphonic acid
TNP-ATP	2',3'-O-(2,4,6-trinitrophenyl)adenosine 5'-triphosphate
TNF-α	tumor necrosis factor-α
UDP	uridine 5'-diphosphate
IL-1β	interleukin-1β

## Introduction

There is a type of pain that does not go away even though the tissue has already healed. One type of this pain is called neuropathic pain that typically develops when peripheral nerves are damaged such as through surgery, bone compression in cancer, diabetes, or infection. Neuropathic pain is a major factor causing impaired quality of life in millions of people worldwide and is frequently resistant to all known analgesic drugs. Over the last decade, accumulating evidence concerning how peripheral nerve injury creates neuropathic pain has suggested that nerve injury produces molecular and cellular alterations that result in multiple forms of neuronal plasticity and anatomical reorganization in the dorsal horn of the spinal cord. These alterations have been proposed to be crucial in the pathogenesis of neuropathic pain [1, 2]. While the dominant theme in research on neuropathic pain has been to

K. Inoue (✉) · M. Tsuda · H. Tozaki-Saitoh  
Department of Molecular and System Pharmacology,  
Graduate School of Pharmaceutical Sciences, Kyushu University,  
3-1-1 Maidashi, Higashi,  
Fukuoka 812-8582, Japan  
e-mail: inoue@phar.kyushu-u.ac.jp

understand the roles of neurons in the peripheral nervous system and the dorsal horn, there is a rapidly growing body of evidence indicating that spinal glial cells play a critical role in the pathogenesis of neuropathic pain.

Recently, growing evidence has indicated that neuron-glia interaction is a key idea to understand functions of the central nervous system (CNS). Especially glia play important roles in pathophysiological situations of the CNS including psychiatric disorders, physical trauma, and infections [3]. Glia consist of three members: astrocytes, oligodendrocytes, and microglia. Accumulating findings also indicate that nucleotides play an important role in neuron-glia communication through P2 purinoceptors, even though ATP is recognized primarily to be a source of free energy and nucleotides are key molecules in cells. Microglia, which are thought to be residential macrophages in the CNS, express P2 purinoceptors, mainly P2X<sub>4</sub> and P2X<sub>7</sub> as well as P2Y<sub>2</sub>, P2Y<sub>6</sub>, and P2Y<sub>12</sub> [4]. Damaged neurons release or leak nucleotides including ATP and UTP to stimulate microglia [5, 6]. It is clear that these nucleotides trigger the release of various neurotoxic and neuroprotective cytokines and growth factors via different purine receptors [4, 7] or induce P2Y<sub>12</sub>-dependent chemotaxis in cultured cells [8, 9]. It was shown in an animal model of neuropathic pain that microglial P2X<sub>4</sub> and P2X<sub>7</sub> receptors are crucial in pain signaling after peripheral nerve lesion [10–12]. Other purine receptors are upregulated in microglia in response to neuronal injury, as was recently demonstrated for P2Y<sub>6</sub> [6]. The P2Y<sub>6</sub> receptor triggers microglial phagocytosis [6]. In this paper, we review the modification of pain sensation through microglial P2X<sub>4</sub> and P2X<sub>7</sub>, and also point out the possibility of P2Y<sub>6</sub> and P2Y<sub>12</sub> involvement in pain signaling.

### P2X<sub>4</sub> in neuropathic pain

We found that the marked tactile allodynia that develops following nerve injury was reversed by acutely administering TNP-ATP intrathecally but was unaffected by administering PPADS [10]. From the pharmacological profiles of TNP-ATP (blocking P2X<sub>4</sub> at high concentration) and PPADS (not blocking P2X<sub>4</sub>), it was suggested that tactile allodynia depends upon P2X<sub>4</sub> in the spinal cord. The expression of P2X<sub>4</sub> protein, normally low in the naïve spinal cord, progressively increased in the days following nerve injury with a time course parallel to that of the development of tactile allodynia. Double immunolabeling analysis demonstrated that not neurons or astrocytes but activated microglia in the dorsal horn were intensely positive for P2X<sub>4</sub> protein [10]. Moreover, intrathecally administered antisense oligodeoxynucleotide for P2X<sub>4</sub>R reduced the expression of P2X<sub>4</sub> protein in spinal microglia

and prevented the development of the nerve injury-induced tactile allodynia. In naïve rats, intrathecal administration of cultured microglia that were preincubated with ATP to activate P2X<sub>4</sub> produced tactile allodynia over the 3–5 h after the administration [10].

Since it was already reported that the nerve injury-induced tactile allodynia depends on a depolarizing shift in the  $E_{\text{anion}}$  of spinal lamina I (LI) neurons in the dorsal spinal cord, resulting in converting the GABA<sub>A</sub> receptor- and glycine receptor-mediated inhibition to excitation [13], it was considered that microglia may affect  $E_{\text{anion}}$  in LI neurons. To investigate this possibility, microglia were administered to the lumbar spinal level of naïve rats by an intrathecal catheter as described [11]. Administering microglia stimulated with ATP caused a progressive tactile allodynia over the 5 h after injection.  $E_{\text{anion}}$  in LI neurons from rats administered ATP-stimulated microglia was shifted to  $-61.6$  mV from  $-68.3$  mV of normal rats. In addition, we found that GABA response switched from hyperpolarizing in control rats to depolarizing in microglia-treated rats. Activated microglia secrete various biologically active molecules, one of which, BDNF, was implicated in the hypersensitivity of dorsal horn neurons that follows sensitization and inflammation [14–16] and in anion gradient shifts in the hippocampus [17]. Indeed, intrathecal administration of recombinant BDNF produced tactile allodynia comparable to that produced by ATP-stimulated microglia [11].  $E_{\text{anion}}$  of LI neurons in slices treated with BDNF (>90 min, *in vitro*) was significantly less negative than that of LI neurons from control slices. The rise in  $[\text{Ca}^{2+}]_i$  was prevented by the GABA<sub>A</sub> receptor blocker bicuculline, confirming that the effect was mediated by GABA<sub>A</sub> receptors. Thus, acute administration of BDNF in slices caused a depolarizing shift in  $E_{\text{anion}}$  and caused GABA to produce net excitation [11]. Moreover, a function-blocking antibody against the TrkB receptor (anti-TrkB) and a BDNF-sequestering fusion protein (TrkB-Fc) acutely inhibited the allodynia and the shift of  $E_{\text{anion}}$  of LI neurons [11]. The administration of ATP-stimulated microglia with either anti-TrkB or TrkB-Fc did not develop tactile allodynia. After pretreatment of microglia with double-stranded short interfering RNA directed against BDNF (BDNF siRNA), the ATP-stimulated microglia injected intrathecally into normal rats did not cause the allodynia [11]. Anti-TrkB and BDNF siRNA prevented the shift in  $E_{\text{anion}}$  induced by ATP-stimulated microglia. ATP stimulation caused release of BDNF from microglia in culture. This effect of ATP was blocked by treating the cultures with the P2X receptor blocker TNP-ATP. In addition, pretreatment of the microglia with BDNF siRNA prevented release of BDNF by ATP stimulation. By bath application of TNP-ATP to spinal slices taken from allodynic rats 2 weeks after nerve injury,  $E_{\text{anion}}$  of LI neurons was returned to normal value [11]. These findings


indicate that P2X<sub>4</sub>-dependent release of BDNF from microglia is necessary to sustain both the tactile allodynia and the depolarizing shift in E<sub>anion</sub> in LI neurons that result from nerve injury (Fig. 1).

### P2X<sub>7</sub> in neuropathic pain

P2X<sub>7</sub> purinoceptors are a ligand-gated nonselective cationic channel and are expressed predominantly on immune cells [18]. Stimulation of the P2X<sub>7</sub> receptors on microglia is associated with release of cytokines including tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) [5, 19], interleukin-6 (IL-6) [20], and interleukin-1 $\beta$  (IL-1 $\beta$ ) [21–23]. Several cytokines such as IL-1 $\beta$ , IL-6, and TNF- $\alpha$  in the dorsal horn are increased after nerve lesion [24–26] and have been implicated in contributing to neuropathic pain [24–27]. Recent evidence indicates the relationship between TNF- $\alpha$

and neuropathic pain [28–32], and TNF- $\alpha$  released after injury is proposed as an initiator of abnormal pain sensation. TNF- $\alpha$  is upregulated after nerve injury in both dorsal root ganglion (DRG) and spinal cord [33]. The inhibition of TNF- $\alpha$  reduces the hyperalgesia in neuropathic pain models [34]. Recent evidence indicates the relationship between inflammatory cytokines including IL-1 $\beta$  and neuropathic pain [25, 26, 28]. The expression of IL-1 $\beta$  is upregulated in the spinal cord of several rat neuropathy models [24–26]. These findings further support a role for central IL-1 $\beta$  in the development and maintenance of neuropathic pain through induction of a proinflammatory cytokine cascade (Fig. 1).

Recently, it was reported that in mice lacking P2X<sub>7</sub> inflammatory and neuropathic hypersensitivity is completely absent to both mechanical and thermal stimuli, whilst normal nociceptive processing is preserved [12]. Contribution of P2X<sub>7</sub> receptor to neuropathic pain is also demon-



**Fig. 1** Schematic illustration of potential mechanisms by which P2X/Y receptors in activated microglia modulate neuropathic pain signaling in the dorsal horn. Activated microglia in the spinal cord after nerve injury express ionotropic ATP receptors [e.g., P2X<sub>4</sub> receptor (P2X<sub>4</sub>R) or P2X<sub>7</sub>R]. P2X<sub>4</sub>R or P2X<sub>7</sub>R activation leads to the release of bioactive diffusible factors such as BDNF and other proinflammatory factors (cytokines and chemokines). BDNF causes a collapse of transmembrane anion gradient in dorsal horn lamina I neurons presumably through the downregulation of KCC2, which in turn renders GABA and glycine effects depolarizing, rather than hyper-

polarizing, in these neurons. Microglial factors may also interact with excitatory synapses of neighboring dorsal horn neurons and enhance the excitability in dorsal horn neurons. The net hyperexcitability in the dorsal horn pain network by these factors from activated microglia may be responsible for neuropathic pain. Microglia also express G protein-coupled ATP receptors [e.g., P2Y<sub>6</sub> receptor (P2Y<sub>6</sub>R) and P2Y<sub>12</sub>R]. Activating P2Y<sub>12</sub>R and P2Y<sub>6</sub>R leads to chemotaxis and phagocytosis, respectively, but their functional relevance to neuropathic pain remained to be determined

strated by using the recently developed selective antagonist for P2X<sub>7</sub> receptor A-740003 [35]. The knockout animals were unimpaired in their ability to produce mRNA for pro-IL-1 $\beta$ , and cytometric analysis of paw and systemic cytokines from knockout and wild-type animals following adjuvant insult suggested a selective effect of the gene deletion on release of IL-1 $\beta$  and IL-10, with systemic reductions in adjuvant-induced increases in IL-6 and MCP-1. In addition, P2X<sub>7</sub> receptors were upregulated in human dorsal root ganglia and injured nerves obtained from chronic neuropathic pain patients [12]. It was hypothesized that the P2X<sub>7</sub> receptor plays a common upstream transductional role in the development of pain of neuropathic and inflammatory origin via regulation of mature IL-1 $\beta$  production.

In addition, a recent study has shown that activation of P2X<sub>7</sub> receptors expressed on satellite glia that enwrap each DRG neuron leads to a release of TNF- $\alpha$ , which in turn increases the excitability of DRG neurons [36]. P2X<sub>7</sub> receptors are upregulated in surrounding satellite glial cells in the DRG in humans [12]. Implication of TNF- $\alpha$  has been reported in neuropathic pain [28–32]. Therefore, the contribution of P2X<sub>7</sub> receptor to neuropathic pain might be related not only to the activation of microglial P2X<sub>7</sub> receptors, but also to P2X<sub>7</sub> receptors on other cell types such as satellite glia.

### P2Y<sub>6</sub> and P2Y<sub>12</sub> in neuropathic pain?

When neurons are injured or dead, microglia are activated, resulting in their interaction with immune cells, active migration to the site of injury, release of proinflammatory substances, and the phagocytosis of damaged cells or debris. For such activation of microglial motilities, extracellular nucleotides have a central role. Extracellular ATP functions as a chemoattractant. Microglial chemotaxis by ATP via P2Y<sub>12</sub> receptors was originally found by Honda et al. [8] and has recently been confirmed in vivo in P2Y<sub>12</sub> receptor knockout animals [37]. Neuronal injury results in the release or leakage of ATP that appears to be a “find-me” signal from damaged neurons to microglia to cause chemotaxis. In addition to microglial migration by ATP, another nucleotide, UDP, an endogenous agonist of the P2Y<sub>6</sub> receptor, greatly activates the motility of microglia and orders microglia to eat damaged neurons. UDP does not cause chemotaxis, but instead causes phagocytosis by microglia [6]. Phagocytosis, a specialized form of endocytosis, is the uptake by the cell of relatively large particles (>1.0  $\mu$ m) into vacuoles and has a central role in tissue remodeling, inflammation, and the defense against infectious agents [38]. Phagocytosis is initiated by the activation

of various cell surface phagocytosis receptors, including Fc receptors, complement receptors, integrins, endotoxin receptors (CD18, CD14), mannose receptors, and scavenger receptors [39], which are activated by corresponding extracellular ligands. In the CNS, a full innate immune system, i.e., Fc receptors, complement system, scavenger receptors, and Toll-like receptors etc., has been described, and microglia reveal related roles as dedicated phagocytes. Since recognition is the first and the most important step for phagocytosis, extensive studies on phagocytosis receptors have been reported. It is well-known that dying cells express so-called eat-me signals such as phosphatidylserine on their surface membrane [39], by which microglia recognize the apoptotic cells to catch and remove them [39]. We first found that exogenously applied UDP caused microglial phagocytosis in a concentration-dependent manner, which was P2Y<sub>6</sub> receptor-dependent. We found that neuronal injury caused by kainic acid (KA) upregulated P2Y<sub>6</sub> receptors in microglia, the KA-evoked neuronal injury resulted in an increase in extracellular UTP, which was immediately metabolized into UDP in vivo and in vitro. Moreover, UDP leaked from injured neurons caused P2Y<sub>6</sub> receptor-dependent phagocytosis in vivo and in vitro. These results suggest that UDP could be a molecule that signals the crisis of damaged neurons to microglia, triggering phagocytosis. It should be noted that nucleotides could be both “find-me” and “eat-me” signals. Cells release ATP, and we also found that KA caused an increase in extracellular UTP/UDP. Therefore microglia might be attracted by ATP/ADP [8, 40, 41] and subsequently recognize UDP, leading to the removal of the dying cells and their debris.

Are P2Y<sub>6</sub> and P2Y<sub>12</sub> receptors involved in neuropathic pain? For this question, there is a report indicating that the UDP-sensitive P2Y<sub>6</sub> receptor produces inhibitory effects on spinal pain transmission in a neuropathic pain model [42]. In the neuropathic pain model, in which the sciatic nerves of rats were partially ligated, UDP (30 and 100 nmol/rat) produced significant antiallodynic effects. UDP (100 nmol/rat) caused no motor deficit in the inclined plane test [42]. The mechanism of this UDP-evoked inhibitory effect on neuropathic pain is unknown. There is no evidence to show the site (neuron or glia) of action of UDP. We are now investigating the involvement of P2Y<sub>12</sub> in pain signaling and have obtained evidence suggesting the role of P2Y<sub>12</sub> in the neuropathic pain state.

### Conclusion

Neuropathic pain is a major factor causing impaired quality of life in millions of people worldwide. We have to try to reveal the mechanism of this pain in order to develop effective drugs against the pain. Recently, there has been a

rapidly growing body of evidence indicating that spinal glia, especially microglia, play a critical role in the pathogenesis of neuropathic pain and that nucleotides play an important role in neuron-glia communication through P2 purinoceptors. We described the modification of neuropathic pain signaling through microglial P2X<sub>4</sub> and P2X<sub>7</sub>, with the possibility of P2Y<sub>6</sub> and P2Y<sub>12</sub> involvement (Fig. 1). It was shown in an animal model of neuropathic pain that microglial P2X<sub>4</sub> and P2X<sub>7</sub> receptors are crucial in pain signaling after peripheral nerve lesion. Microglial P2Y<sub>6</sub> and P2Y<sub>12</sub> play very interesting roles for phagocytosis and chemotaxis, respectively. Besides, there is a possibility that these receptors are involved in pain sensation. Additional experiments are needed to clarify the possibility.

## References

1. Woolf CJ, Mannion RJ (1999) Neuropathic pain: aetiology, symptoms, mechanisms, and management. *Lancet* 353:1959–1964
2. Scholz J, Woolf CJ (2002) Can we conquer pain? *Nat Neurosci* 5 Suppl:1062–1067
3. Miller G (2005) Neuroscience. The dark side of glia. *Science* 308:778–781
4. Inoue K (2006) The function of microglia through purinergic receptors: neuropathic pain and cytokine release. *Pharmacol Ther* 109:210–226
5. Wang X, Arcuino G, Takano T et al (2004) P2X<sub>7</sub> receptor inhibition improves recovery after spinal cord injury. *Nat Med* 10:821–827
6. Koizumi S, Shigemoto-Mogami Y, Nasu-Tada K et al (2007) UDP acting at P2Y<sub>6</sub> receptors is a mediator of microglial phagocytosis. *Nature* 446:1091–1095
7. Suzuki T, Hide I, Ido K et al (2004) Production and release of neuroprotective tumor necrosis factor by P2X<sub>7</sub> receptor-activated microglia. *J Neurosci* 24:1–7
8. Honda S, Sasaki Y, Ohsawa K et al (2001) Extracellular ATP or ADP induce chemotaxis of cultured microglia through Gi/o-coupled P2Y receptors. *J Neurosci* 21:1975–1982
9. Ohsawa K, Irino Y, Nakamura Y et al (2007) Involvement of P2X<sub>4</sub> (4) and P2Y<sub>12</sub> (12) receptors in ATP-induced microglial chemotaxis. *Glia* 55:604–616
10. Tsuda M, Shigemoto-Mogami Y, Koizumi S et al (2003) P2X<sub>4</sub> receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* 424:778–783
11. Coull JA, Beggs S, Boudreau D et al (2005) BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature* 438:1017–1021
12. Chessell IP, Hatcher JP, Bountra C et al (2005) Disruption of the P2X<sub>7</sub> purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain* 114:386–396
13. Coull JA, Boudreau D, Bachand K et al (2003) Trans-synaptic shift in anion gradient in spinal lamina I neurons as a mechanism of neuropathic pain. *Nature* 424:938–942
14. Mannion RJ, Costigan M, Decosterd I et al (1999) Neurotrophins: peripherally and centrally acting modulators of tactile stimulus-induced inflammatory pain hypersensitivity. *Proc Natl Acad Sci U S A* 96:9385–9390
15. Heppenstall PA, Lewin GR (2001) BDNF but not NT-4 is required for normal flexion reflex plasticity and function. *Proc Natl Acad Sci U S A* 98:8107–8112
16. Thompson SW, Bennett DL, Kerr BJ et al (1999) Brain-derived neurotrophic factor is an endogenous modulator of nociceptive responses in the spinal cord. *Proc Natl Acad Sci U S A* 96:7714–7718
17. Rivera C, Li H, Thomas-Crusells J et al (2002) BDNF-induced TrkB activation down-regulates the K<sup>+</sup>-Cl<sup>-</sup> cotransporter KCC2 and impairs neuronal Cl<sup>-</sup> extrusion. *J Cell Biol* 159:747–752
18. Collo G, Neidhart S, Kawashima E et al (1997) Tissue distribution of the P2X<sub>7</sub> receptor. *Neuropharmacology* 36:1277–1283
19. Hide I, Tanaka M, Inoue A et al (2000) Extracellular ATP triggers tumor necrosis factor- $\alpha$  release from rat microglia. *J Neurochem* 75:965–972
20. Shigemoto-Mogami Y, Koizumi S, Tsuda M et al (2001) Mechanisms underlying extracellular ATP-evoked interleukin-6 release in mouse microglial cell line, MG-5. *J Neurochem* 78:1339–1349
21. Ferrari D, Villalba M, Chiozzi P et al (1996) Mouse microglial cells express a plasma membrane pore gated by extracellular ATP. *J Immunol* 156:1531–1539
22. Ferrari D, Wesselborg S, Bauer MK et al (1997) Extracellular ATP activates transcription factor NF- $\kappa$ B through the P2Z purinoceptor by selectively targeting NF- $\kappa$ B p65. *J Cell Biol* 139:1635–1643
23. Sanz JM, Di Virgilio F (2000) Kinetics and mechanism of ATP-dependent IL-1 $\beta$  release from microglial cells. *J Immunol* 164:4893–4898
24. Sweitzer S, Martin D, DeLeo JA (2001) Intrathecal interleukin-1 receptor antagonist in combination with soluble tumor necrosis factor receptor exhibits an anti-allodynic action in a rat model of neuropathic pain. *Neuroscience* 103:529–539
25. Winkelstein BA, Rutkowski MD, Sweitzer SM et al (2001) Nerve injury proximal or distal to the DRG induces similar spinal glial activation and selective cytokine expression but differential behavioral responses to pharmacologic treatment. *J Comp Neurol* 439:127–139
26. Raghavendra V, Tanga F, DeLeo JA (2003) Inhibition of microglial activation attenuates the development but not existing hypersensitivity in a rat model of neuropathy. *J Pharmacol Exp Ther* 306:624–630
27. Watkins LR, Milligan ED, Maier SF (2001) Glial activation: a driving force for pathological pain. *Trends Neurosci* 24:450–455
28. Wagner R, Myers RR (1996) Endoneurial injection of TNF- $\alpha$  produces neuropathic pain behaviors. *Neuroreport* 7:2897–2901
29. Sommer C, Schafer M (1998) Painful mononeuropathy in C57BL/Wld mice with delayed wallerian degeneration: differential effects of cytokine production and nerve regeneration on thermal and mechanical hypersensitivity. *Brain Res* 784:154–162
30. Sommer C, Marziniak M, Myers RR (1998) The effect of thalidomide treatment on vascular pathology and hyperalgesia caused by chronic constriction injury of rat nerve. *Pain* 74:83–91
31. Sorkin LS, Doom CM (2000) Epineurial application of TNF elicits an acute mechanical hyperalgesia in the awake rat. *J Peripher Nerv Syst* 5:96–100
32. Schafer M, Geis C, Brors D et al (2002) Anterograde transport of tumor necrosis factor- $\alpha$  in the intact and injured rat sciatic nerve. *J Neurosci* 22:536–545
33. Hashizume H, DeLeo JA, Colburn RW et al (2000) Spinal glial activation and cytokine expression after lumbar root injury in the rat. *Spine* 25:1206–1217
34. Sommer C, Lindenlaub T, Teuteberg P et al (2001) Anti-TNF-neutralizing antibodies reduce pain-related behavior in two different mouse models of painful mononeuropathy. *Brain Res* 913:86–89

35. Honore P, Donnelly-Roberts D, Namovic MT et al (2006) A-740003 [N-(1-[(cyanoimino)(5-quinolinylamino) methyl] amino)-2,2-dimethylpropyl)-2-(3,4-dimethoxyphenyl)acetamide], a novel and selective P2X7 receptor antagonist, dose-dependently reduces neuropathic pain in the rat. *J Pharmacol Exp Ther* 319:1376–1385
36. Zhang X, Chen Y, Wang C et al (2007) Neuronal somatic ATP release triggers neuron-satellite glial cell communication in dorsal root ganglia. *Proc Natl Acad Sci U S A* 104:9864–9869
37. Haynes SE, Hollopeter G, Yang G et al (2006) The P2Y(12) receptor regulates microglial activation by extracellular nucleotides. *Nat Neurosci* 9:1512–1519
38. Tjelle TE, Lovdal T, Berg T (2000) Phagosome dynamics and function. *Bioessays* 22:255–263
39. Lauber K, Bohn E, Krober SM et al (2003) Apoptotic cells induce migration of phagocytes via caspase-3-mediated release of a lipid attraction signal. *Cell* 113:717–730
40. Davalos D, Grutzendler J, Yang G et al (2005) ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci* 8:752–758
41. Nimmerjahn A, Kirchhoff F, Helmchen F (2005) Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 308:1314–1318
42. Okada M, Nakagawa T, Minami M et al (2002) Analgesic effects of intrathecal administration of P2Y nucleotide receptor agonists UTP and UDP in normal and neuropathic pain model rats. *J Pharmacol Exp Ther* 303:66–73