

## Necrotizing fasciitis of the abdominal wall caused by *Serratia marcescens*

Naheed A. Lakhani,<sup>1</sup>  
Umesh Narsinghani,<sup>2</sup> Ritu Kumar<sup>3</sup>

<sup>1</sup>Department of Family and Preventive Medicine, Emory University School of Medicine, Dunwoody, GA; <sup>2</sup>Department of Pediatrics, Mercer University School of Medicine, Macon, GA; <sup>3</sup>Department of Internal Medicine Mercer University School of Medicine, Macon, GA, USA

### Abstract

In this article, we present the first case of necrotizing fasciitis affecting the abdominal wall caused by *Serratia marcescens* and share results of a focused review of *S. marcescens* induced necrotizing fasciitis. Our patient underwent aorto-femoral bypass grafting for advanced peripheral vascular disease and presented 3 weeks postoperatively with pain, erythema and discharge from the incision site in the left lower abdominal wall and underwent multiple debridement of the affected area. Pathology of debrided tissue indicated extensive necrosis involving the adipose tissue, fascia and skeletal muscle. Wound cultures were positive for *Serratia marcescens*. She was successfully treated with antibiotics and multiple surgical debridements. Since necrotizing fasciitis is a medical and surgical emergency, it is critical to examine infectivity trends, clinical characteristics in its causative spectrum. Using PubMed we found 17 published cases of necrotizing fasciitis caused by *Serratia marcescens*, and then analyzed patterns among those cases. *Serratia marcescens* is prominent in the community and hospital settings, and information on infection presentations, risk factors, characteristics, treatment, course, and complications as provided through this study can help identify cases earlier and mitigate poor outcomes. Patients with positive blood cultures and those patients where surgical intervention was not provided or delayed had a higher mortality. Surgical intervention is a definite way to establish the diagnosis of necrotizing infection and differentiate it from other entities.

### Introduction

Necrotizing fasciitis, also known as flesh-

eating disease, is a rare infection of the deeper layers of skin and subcutaneous tissues, and easily spreads across the fascial plane within those tissues. As bacterial toxins and the immune response cause vasoconstriction of the vasculature, the fascial spaces become avascular resulting in necrosis, which also prevents penetration of antibiotics into the tissues.<sup>1</sup> Common causes are Group A streptococcus (GAS) (*Streptococcus pyogenes*), *Staphylococcus aureus*, *Vibrio vulnificus*, *Clostridium perfringens*, and *Bacteroides fragilis*. Mortality ranges from 4.2 to 38% with improving prognosis as time to treatment decreases.<sup>2</sup>

*Serratia marcescens*, a motile bacillus, gram-negative, facultative anaerobe, is an opportunistic pathogen of increasing importance. It is part of normal colon flora, and is also found in soil, sewage, and water. This Enterobacteriaceae organism also typically colonizes the respiratory and urinary tracts and causes infections in those organ systems.<sup>3</sup> *S. marcescens* is often an opportunistic infection and may cause osteomyelitis, septic arthritis, endocarditis, and, rarely cellulitis or necrotizing fasciitis.<sup>4,6</sup> Soft tissue infections due to gram-negative organisms are relatively uncommon, and typical predisposing factors include: a history of trauma, alcoholism, peripheral vascular disease, systemic lupus erythematosus, immunosuppression, diabetes mellitus, urinary tract infection (UTI), bacteremia, pneumonia, infective arthritis, burns, and renal failure.<sup>4,7</sup> Other predisposing factors include: antibiotic use (most often first generation cephalosporins), steroid use, surgical instrumentation, urinary catheters, respiratory equipment, intravenous lines, injections, lacerations, abscesses, or ulcers.<sup>8</sup>

Common presentations of necrotizing fasciitis within 48 hours of infection include skin erythema and swelling at the affected site (97.6%), pyrexia (61.9%), hypotension (33.3%), altered consciousness (28.6%), bul- lous lesions (26.2%), and crepitus (9.5%). Hypotension, altered consciousness, ventilator support, ALT > two-fold of normal, serum creatinine >177 µmol/L, thrombocytopenia (<100×10<sup>9</sup>/L), and worsening symptoms within 48 hours of admission have been associated with higher fatality rates.<sup>9</sup> Lack of response to narrow-spectrum antibiotics, bullae formation, or a rapidly worsening clinical course, should heighten the suspicion for uncommon organisms like *S. marcescens*.

Reports of *S. marcescens* necrotizing fasciitis cases have increased in the literature. In this study, we identify characteristics, trends, and risk factors of those infections to better prepare the medical community and prevent poor outcomes.

Correspondence: Department of Family and Preventive Medicine, Emory University School of Medicine, 4500 North Shallowford Road, Dunwoody, Georgia 30338, USA.  
Tel.: +1.404.542.7238 / +1.404.778.6920  
Fax: +1.404.778.6901.  
E-mail: naheeda786@gmail.com

Key words: *S. Marcescens*, necrotizing fasciitis, community acquired infection, healthcare associated infection, surgical intervention.

Acknowledgments: we thank our case patient, the hospital, and other doctors involved in the care of this case patient.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Received for publication: 22 December 2014.

Revision received: 9 February 2015.

Accepted for publication: 10 February 2015.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright N.A. Lakhani et al., 2015  
Licensee PAGEPress, Italy  
Infectious Disease Reports 2015; 7:5774  
doi:10.4081/idr.2015.5774

### Materials and Methods

PubMed was used to search for cases of necrotizing fasciitis caused by *Serratia marcescens* published in the English language literature between 1966 and 2013. Keywords that were used included: *Serratia marcescens*, necrotizing, and fasciitis. We identified eleven additional cases of necrotizing fasciitis due to *S. marcescens* in the literature since the latest review in 2001, which are included in Table 1.<sup>4,7,10-22</sup> Cases were categorized as being healthcare-associated infections or community-acquired infections based on the Centers for Disease Control and Prevention (CDC) definition of healthcare-associated infections (HAIs) as infections that patients acquire during the course of receiving healthcare treatment for other conditions.<sup>23</sup>

### Case Report

We recently identified a case involving a 51-year-old African-American woman who pre-

sented 3 weeks postoperatively after undergoing a bifemoral bypass and left distal femoral aneurysm repair for symptomatic peripheral vascular disease. She had a long standing history of smoking and hypertension. On presentation, the patient reported having fever, chills, nausea, vomiting, and diarrhea for 6 days, redness and odorous discharge from her surgical incision site for one day. Additional history revealed that she was cleaning her wound with well water. On examination, there was erythema extending from the periumbilical area to her left groin, along with severe tenderness to palpation in that area. No crepitus was felt on palpation. Her temperature on admission was 36.9°C, blood pressure was 162/109 mmHg, respirations were 24 breaths per minute, and she was in no acute distress. Her white blood cell count was  $9.8 \times 10^9/L$ , C reactive protein (CRP) was 16.57 nmol/L, serum creatine

kinase level was 57 U/L, serum electrolytes, renal, liver functions and immunoglobulin levels were normal and blood and urine cultures were negative. The patient was started on intravenous vancomycin and meropenem and underwent multiple debridements of the abdominal wound and left groin region. Intraoperative findings confirmed the presence of deep necrotic fascia and non-adherent subcutaneous tissue. Histopathology of debrided tissue indicated acute and chronic inflammation with infiltration of granulocytes and necrosis involving the fibroadipose tissue and skeletal muscle. Intraoperative wound cultures were positive for *Serratia marcescens*. Based on sensitivities, antibiotics were changed to intravenous levofloxacin MIC <2 mcg/mL. She also received negative-pressure wound therapy using vacuum assisted closure. The patient improved clinically and was successfully dis-

charged on oral levofloxacin to complete a total of 3 weeks of treatment.

## Results

Since 1966, there have been 17 documented cases of necrotizing fasciitis caused by *S. marcescens* (Table 1).<sup>4,7,10-22</sup> Eight (47%) of those cases occurred within the past five years.<sup>4,7,15-24</sup> Among all of the identified cases, 10 (59%) impacted a leg, and 1 (5.9%) affected the forearm, the cervical spine, the axilla and chest wall, a venous access site, the oropharynx, the chest wall, or the lower abdomen. Thirteen (77%) of the cases had blood cultures that were positive for *S. marcescens*. Ten cases (59%) were community-acquired infections,

**Table 1. *Serratia marcescens* necrotizing fasciitis cases, 1966 to present.**

Author	Year	Age	Sex	Risk factors	Precipitating factor	Site of infection	Type	<i>S. marcescens</i> cultures	Treatment	Outcome
Rimailho <i>et al.</i> <sup>10</sup>	1987	74	M	Immunocompromized	Dilofenac consumption	Leg	CA	Blister and blood	None	Died
Bornstein <i>et al.</i> <sup>11</sup>	1992	37	F	Renal failure on hemodialysis	Pain during dialysis	Axilla and chest wall	HA	Wound, bullae, blood	Antibiotics and SD	Recovered
Zipper <i>et al.</i> <sup>12</sup>	1996	55	F	Diabetes	Left below-knee amputation	Leg	CA	Wound	Antibiotics	Recovered
Huang <i>et al.</i> <sup>13</sup>	1999	73	M	Nephrotic syndrome	Steroid therapy	Lower leg	HA	Necrotic tissue, blood	Antibiotics and SD	Recovered
Huang <i>et al.</i> <sup>13</sup>	1999	40	M	Uremia, peritoneal dialysis, SLE	Pneumonia with + cultures for <i>S. marcescens</i> , steroid and nabumetone	Left calf and thigh	CA	Necrotic tissue, blood	Antibiotics and SD	Recovered
Liangpunsakul <i>et al.</i> <sup>14</sup>	2001	66	F	Healthy	None	Leg	CA	Blood	Antibiotics	Died
Newton <i>et al.</i> <sup>15</sup>	2002	2	F	Healthy	Pharyngitis	Cervical spine	CA	Wound, blood	Antibiotics and SD	Died
Bachmeyer <i>et al.</i> <sup>16</sup>	2004	49	M	Small cell lung cancer, DM	Chemotherapy and cellulitis	Right leg	HA	Tissue, bullae, blood	Antibiotics	Recovered
Curtis <i>et al.</i> <sup>4</sup>	2005	51	M	ESRD, T2DM, CHF	Scraped knee on rock in river	Left leg	CA	Wound, blood	Antibiotics and SD	Died
Statham <i>et al.</i> <sup>17</sup>	2009	6	M	Immunocompetent	Suspected pharyngitis	Oro-pharynx	CA	Wound, blood	Antibiotics and SD	Recovered
Motsitsi <i>et al.</i> <sup>18</sup>	2011	37	M	Healthy	Human bite	Forearm	CA	Wound	SD	Died
Vano-Galvan <i>et al.</i> <sup>7</sup>	2012	57	F	CML, immunocompromized	Minor trauma	Right thigh	HA	Blister, blood	Antibiotics	Died
Prelog <i>et al.</i> <sup>19</sup>	2012	15	F	Acute lymphocytic leukemia	Venous access port implantation	Left axilla, venous HA access port site	Wound	Antibiotics and SD	Recovered	
Wen <i>et al.</i> <sup>20</sup>	2012	40	F	Nephrotic syndrome, cyclosporine use	Chemotherapy 10 days prior	Left leg	CA	Wound, blood	Antibiotics	Died
Rehman <i>et al.</i> <sup>21</sup>	2012	54	F	SLE, end-stage renal disease	Central venous catheter, AV fistula ligation, steroid therapy	Chest wall	HA	Wound, blood	Antibiotics and SD	Died
Present case		51	F	DM, PVD	Bifemoral bypass and left distal femoral aneurysm repair	Lower abdomen	CA	Wound, blood	Antibiotics and SD	Recovered
Cope <i>et al.</i> <sup>22</sup>	2013	97	F	Heart failure, CKD	Heart failure exacerbation	Right leg	HA	Wound (post-mortem)	Antibiotics	Died

SD, Surgical debridement; CA, community-acquired infection; HA, healthcare-associated infection; SLE, systemic lupus erythematosus; ESRD, end-stage renal disease; T2DM, Type II, diabetes mellitus; PVD, peripheral vascular disease.

and 7 cases (41%) were healthcare-associated infections. An overwhelming number of cases (81.3%) had pre-existing open wounds. Seven cases (41%) were immunocompromized, 5 cases (29%) had kidney disease, and 4 cases (24%) had diabetes. A majority (59%) of the cases were among females. Three of the cases (18%) were children.

Nine out of seventeen cases (53%) died as a result of necrotizing fasciitis and its complications. Six of the cases (67%) that died had community-acquired infections. Individuals with positive versus negative blood cultures were more likely to die (88% *vs.* 75%). Patients who did not receive surgical debridement had inferior outcomes; they had a mortality of 71% compared to 40% among those who received surgical intervention as opposed to debridement. Additionally, all patients (18%, n=3) who received only antibiotics and the one patient who received only surgical debridement died. All previously healthy patients (18%, n=3) also died. A majority of those that recovered had a surgical procedure in the hospital prior to symptom onset such as below the knee amputation or venous access port implantation (75%, n=8).

## Discussion and Conclusions

Necrotizing fasciitis is a deep infection of the subcutaneous tissue that results in progressive destruction of fascia and fat. The disease is classified as type I (polymicrobial infection), type II (monomicrobial) and type III gas gangrene, or clostridial myonecrosis. Type I infection involves anaerobic species in combination with one or more facultative anaerobic streptococci (other than group A) and members of the Enterobacteriaceae family. Type II infection is commonly caused by group A streptococci or other beta-hemolytic streptococci that are isolated alone or in combination with other species, most frequently *S. aureus*. These infections are also commonly referred as flesh eating infection. Among those with necrotizing fasciitis, the affected area is typically erythematous, swollen, warm, and exquisitely tender. The infection progresses rapidly over several days, with changes in skin color from red-purple to patches of blue-gray. Skin breakdown with bullae (containing thick pink or purple fluid) and frank cutaneous gangrene may be observed within three to five days.<sup>16</sup> The development of anesthesia may precede the appearance of skin necrosis and provide a clue that the process is necrotizing fasciitis rather than cellulitis. In advanced infection, high fever and systemic toxicity are generally observed.

Individuals with positive blood cultures typically have poorer prognosis and higher mortal-

ity rates.<sup>1,25</sup> Treatment of necrotizing infection consists of early and aggressive surgical exploration and debridement of necrotic tissue, together with broad spectrum empiric antibiotic therapy and hemodynamic support as necessary.<sup>4,26</sup> Acceptable antibiotic regimens prior to identification of the causative organism(s) include administration of a carbapenem or beta-lactam/beta-lactamase inhibitor, together with clindamycin (600 to 900 milligrams intravenously every eight hours), as well as an agent with activity against MRSA. Definitive antibiotic treatment should be tailored according to blood and or tissue gram stain, culture, and sensitivity results when available.<sup>27</sup> If *S. marcescens* is identified as the main pathogen, third-generation cephalosporins, fluoroquinolones, and imipenem/cilastatin are the antimicrobials of choice. Our patient's fever on presentation was likely masked by the pain medications she was taking, though she did give a history of having fever at home. She most likely had a subacute course and thus lacked the classic features of rapid clinical progression and fulminant toxicity. An HIV test was not performed, as the patient did not have any risk factors or any previous sexually transmitted disease. Our review noted a mortality rate of 53% among those with *S. marcescens*-induced necrotizing fasciitis, which is higher than the mortality rate reported by Elliott *et al.*<sup>28</sup> in 1996 and nearly equivalent (53% *vs.* 50%) to the mortality rate documented by Nancy *et al.* in 2002.<sup>29</sup> Patients who died were more likely to have positive blood cultures, present with a community-acquired illness, be previously healthy, experience minor trauma or sickness prior to symptom onset, and not receive both surgical debridement and antibiotic therapy. Similar observations of increased mortality among those with positive blood cultures or those not receiving appropriate antibiotic and surgical treatment were also found in other studies.<sup>4,26</sup> Increased mortality among individuals with less severe injury, lack of traditional risk factors, and those who did not have a recent surgical procedure in the hospital may have been due to less vigilance about changes in health status and the fact that *S. marcescens* is common in the outdoor environment; these individuals were also more likely to have community-acquired infections (70% *vs.* 30%). Additionally, individuals who had been hospitalized recently were more likely to have received antibiotics in the near past and also more likely to be wary of any new symptoms. Healthy patients also tend to seek care later and receive less aggressive treatment; only one of the previously healthy patients received both antibiotics and surgical debridement.

Clinicians should be wary of necrotizing fasciitis among those with soft tissue infections presenting after environmental open

skin exposure. Gram-negative bacilli, including *S. marcescens*, should especially be considered in cases of necrotizing fasciitis or cellulitis among immunocompromized persons with renal failure, steroid use, recent surgery or diabetes.<sup>29</sup> Early diagnosis is key, but unfortunately is missed in 85 to 100% of cases since necrotizing fasciitis is often confused with cellulitis, myositis, or deep-seated abscess(es). A high index of suspicion is important in view of the paucity of specific cutaneous findings early in the course of the disease. Radiographic imaging studies may be useful in determining if muscle tissue is involved but should not delay surgical intervention. Our case also underscores the importance of preoperative and more so, postoperative wound care education, which could have potentially prevented this fatal complication.<sup>30</sup> The most effective treatment regimen for suspected necrotizing fasciitis includes: early and aggressive surgical exploration and debridement, immediate broad spectrum empiric antibiotic therapy, and hemodynamic support as necessary.<sup>4,26</sup>

## References

- Childers BJ, Potyondy LD, Nachreiner R, et al. Necrotizing fasciitis: a fourteen-year retrospective study of 163 consecutive patients. *Am Surg* 2002;68:109-16.
- Bilton BD, Zibari GB, McMillan RW, et al. Aggressive surgical management of necrotizing fasciitis serves to decrease mortality: a retrospective study. *Am Surg* 1998;64:397-400.
- Hejazi A, Falkiner FR. *Serratia marcescens*. *J Med Microbiol* 1997;46:903-12.
- Curtis CE, Chock S, Henderson T, Holman MJ. A fatal case of necrotizing fasciitis caused by *Serratia marcescens*. *Am Surg* 2005;71:228-30.
- Yu VL. *Serratia marcescens*: historical perspective and clinical review. *New Engl J Med* 1979;300:887-93.
- Liangpunsakul S, Pursell K. Community-acquired necrotizing fasciitis caused by *Serratia marcescens*: case report and review. *Eur J Clin Microbiol Infect Dis* 2001;20:509-10.
- Vano-Galvan S, Alvarez-Twose I, Moreno-Martin P, Jaen P. Fulminant necrotizing fasciitis caused by *Serratia marcescens* in an immunosuppressed host. *Int J Dermatol* 2014;53:e57-8.
- Acar JF. *Serratia marcescens* infections. *Infect Control* 1986;7:273-8.
- Hung CC, Chang SC, Lin SF, et al. Clinical manifestations, microbiology and prognosis of 42 patients with necrotizing fasciitis. *J Formos Med Assoc* 1996;95:917-22.

10. Rimalho A, Riou B, Richard C, Auzepy P. Fulminant necrotizing fasciitis and nonsteroidal anti-inflammatory drugs. *J Infect Dis* 1987;155:143-6.
11. Bornstein PF, Ditto AM, Noskin GA. *Serratia marcescens* cellulitis in a patient on hemodialysis. *Am J Nephrol* 1992;12:374-6.
12. Zipper RP, Bustamante MA, Khatib R. *Serratia marcescens*: a single pathogen in necrotizing fasciitis. *Clin Infect Dis* 1996;23:648-9.
13. Huang JW, Fang CT, Hung KY, et al. Necrotizing fasciitis caused by *Serratia marcescens* in two patients receiving corticosteroid therapy. *J Formos Med Assoc* 1999;98:851-4.
14. Liangpunsakul S, Pursell K. Community-acquired necrotizing fasciitis caused by *Serratia marcescens*: case report and review. *Eur J clin microbiol infect dis* 2001;20:509-10.
15. Newton CL, deLEMOs D, Abramo TJ, et al. Cervical necrotizing fasciitis caused by *Serratia marcescens* in a 2 year old. *Pediatr Emerg Care* 2002;18:433-5.
16. Bachmeyer C, Sanguina M, Turc Y, et al. Necrotizing fasciitis due to *Serratia marcescens*. *Clin Exp Dermatol* 2004;673-4.
17. Statham MM, Vohra A, Mehta DK, et al. *Serratia marcescens* causing cervical necrotizing oropharyngitis. *Int J Pediatr Otorhinolaryngol* 2009;73:467-73.
18. Motsitsi NS. Fatal necrotizing fasciitis following human bite of the forearm. *J Hand Surg* 2011;36(605).
19. Prelog T, Jereb M, Cucek I, Jazbec J. Necrotizing fasciitis caused by *Serratia marcescens* after venous access port implantation in a child with acute lymphoblastic leukemia. *J Pediatr Hematol Oncol* 2012;34:e246-8.
20. Wen YK. Necrotizing fasciitis caused by *Serratia marcescens*: a fatal complication of nephrotic syndrome. *Renal Failure* 2012;34:649-52.
21. Rehman T, Moore TA, Seoane L. *Serratia marcescens* necrotizing fasciitis presenting as bilateral breast necrosis. *J Clin Microbiol* 2012;50:3406-8.
22. Cope TE, Cope W, Beaumont DM. A case of necrotising fasciitis caused by *Serratia marcescens*: extreme age as functional immunosuppression? *Age Ageing* 2013;42:266-8.
23. Centers for Disease Control and Prevention. Healthcare-associated infections (HAIs). Available from: <http://www.cdc.gov/HAI/surveillance/>
24. Lancerotto L, Tocco I, Salmaso R, et al. Necrotizing fasciitis: classification, diagnosis, and management. *J Trauma Acute Care Surg* 2012;72:560-6.
25. Chen IC, Li WC, Hong YC, et al. The microbiological profile and presence of bloodstream infection influence mortality rates in necrotizing fasciitis. *Crit Care* 2011;15:R152.
26. McHenry CR, Piotrowski JJ, Petrinic D, Malangoni MA. Determinants of mortality for necrotizing soft-tissue infections. *Ann Surg* 1995;221:558-65.
27. Schwartz MN. Cellulitis and subcutaneous tissue infections. In: Mandell GL, Bennett JE, Dolin R, eds. *Principles and practice of infectious diseases*. 6th ed. Philadelphia: Churchill Livingstone; 2005. p 1172.
28. Elliott DC, Kufera JA, Myers RA. Necrotizing soft tissue infections. Risk factors for mortality and strategies for management. *Ann Surg* 1996;224:672-83.
29. Nancy F, Crum MRW. *Serratia marcescens* Cellulitis: a case report and review of the literature. *Infect Dis Clin Pract* 2002;11.
30. Ronco M, Iona L, Fabbro C, et al. Patient education outcomes in surgery: a systematic review from 2004 to 2010. *Int J Evid Based Healthc* 2012;10:309-23.