

RESEARCH ARTICLE

Penile coital injuries in men decline after circumcision: Results from a prospective study of recently circumcised and uncircumcised men in western Kenya

Nelli Westercamp^{1*}, Supriya D. Mehta¹, Walter Jaoko², Timothy A. Okeyo³, Robert C. Bailey¹

1 Division of Epidemiology and Biostatistics, School of Public Health, University of Illinois at Chicago, Chicago, Illinois, United States of America, **2** Department of Medical Microbiology, University of Nairobi, Nairobi, Kenya, **3** Nyanza Reproductive Health Society, Kisumu, Kenya

* kapooosha@gmail.com



OPEN ACCESS

Citation: Westercamp N, Mehta SD, Jaoko W, Okeyo TA, Bailey RC (2017) Penile coital injuries in men decline after circumcision: Results from a prospective study of recently circumcised and uncircumcised men in western Kenya. PLoS ONE 12(10): e0185917. <https://doi.org/10.1371/journal.pone.0185917>

Editor: Robert K Hills, Cardiff University, UNITED KINGDOM

Received: May 12, 2017

Accepted: September 21, 2017

Published: October 10, 2017

Copyright: © 2017 Westercamp et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Support for this study was provided by a grant to FHI360 from the Bill and Melinda Gates Foundation (www.gatesfoundation.org) to support the Male Circumcision Consortium, a partnership between FHI360, EngenderHealth, and University of Illinois at Chicago working closely with the Nyanza Reproductive Health Society (grant

Abstract

Background

Penile coital injuries are one of the suggested mechanisms behind the increased risk of HIV among uncircumcised men. We evaluated the prevalence and correlates of self-reported penile coital injuries in a longitudinal community-based cohort of young (18–24 years old), newly circumcised and uncircumcised men in Western Kenya.

Methods

Self-reported penile coital injuries were assessed at baseline, 6, 12, 18 and 24 months of follow-up, and were defined as scratches, cuts or abrasions during sex, penile soreness during sex, and skin of the penis bleeding during sex. Associations between penile coital injuries, circumcision, sexual satisfaction, and other covariates were estimated with mixed effect models.

Results

Between November 2008 and April 2010 3,186 participants were enrolled (1,588 into circumcision group and 1,598 as age-matched controls). Among 2,106 (66%) participants sexually active at baseline, 53% reported any penile injury, including 44% scratches, cuts or abrasions; 32% penile pain/soreness; and 22% penile bleeding. In multivariable modeling, risk was lower for circumcised men than uncircumcised men for scratches, cuts and abrasions (aOR = 0.39; 95% CI 0.34–0.44); penile pain/soreness (aOR = 0.58; 95% CI 0.51–0.65), penile bleeding (aOR = 0.53; 95% CI 0.46–0.62), and any penile coital injuries (aOR = 0.47; 95%CI 0.42–0.53). Other significant risk factors included increasing age, history of STIs and genital sores, and multiple sex partners, while condom use was protective. Coital injuries were significantly associated with lower levels of sexual satisfaction in longitudinal analyses (scratches, cuts or abrasions: aOR = 0.87, 95% CI: 0.76–0.98; penile pain/

#47394, RCB). Robert C. Bailey (RCB) received support from the Chicago Developmental Center for AIDS Research, an NIH (www.nih.gov) funded program (P30 AI 082151). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

soreness: aOR = 0.82, 95% CI: 0.72–0.93; and penile bleeding: aOR = 0.65, 95% CI: 0.55–0.76).

Conclusions

Self-reported penile coital injuries were common and decreased significantly following circumcision. Improving sexual experience through the removal of a potential source of sexual discomfort may resonate with many men targeted for circumcision services. The role of penile coital injuries in sexual satisfaction, HIV, HSV-2, and as a motivator for seeking circumcision services should be explored further.

Introduction

Three randomized control trials (RCT) of male circumcision (MC) for HIV prevention in Kenya [1], Uganda [2], and South Africa [3], have demonstrated the protective effect of male circumcision against heterosexually-acquired HIV infection in men beyond any reasonable doubt [4, 5]. While the exact biological mechanism by which MC affords this protection is not known [6], there are a number of plausible explanations based on the cellular composition and environment of the inner foreskin. The earliest hypotheses concerned the gross anatomy of the uncircumcised penis, including the feasibility of potentially infectious secretions being trapped in facilitating conditions beneath the foreskin [4, 7] and the increased surface area of the inner foreskin [8, 9]. With the recognition that ulcerative STIs and other causes of genital tract inflammation increase the risk of HIV infection [10, 11], the association between these infections and circumcision offers additional possible explanation [7, 12–14]. Histologic examination and specific immune responses of the foreskin, and differences in the penile microbiome of circumcised and uncircumcised men offers another set of mechanisms [15–18]. Like most biologic mechanisms, the protective effect of circumcision almost certainly represents a complex system incorporating multiple explanatory factors [19–22].

One possible foreskin-associated HIV risk factor that is often mentioned, but that has not received attention in empirical research, is the perception that preputial mucosa is comparatively fragile and prone to injury during intercourse [7, 8, 23–25]. One difficulty in determining the role of intercourse-associated mechanical injury to the penis in HIV infection is a lack of consistent terminology or operational assessment. When discussed, mechanical penile injuries are often referred to as minor epithelial disruptions, penile trauma, traumatic lesions, or most recently as penile coital injuries, clarifying an integral sexual component [7, 12, 25–28].

Recent research in men participating in the Kenya RCT has shown that penile coital injuries are more commonly reported among uncircumcised men [27], may be an important non-STI cause of genital ulcer disease (GUD) [29], and increase the risk of *Neisseria gonorrhoeae* [30] and HIV [31]. High pre-circumcision baseline prevalence of coital injuries was observed in a circumcision cohort in the Dominican Republic, with a subsequent decline following circumcision [32]. Outside of these populations there is little information available on coital injury prevalence, associated factors, or related disease susceptibilities. Similarly, the potential association between the coital injuries and sexual satisfaction remains largely unexplored.

To increase the understanding of the prevalence and correlates of penile coital injuries and the potential effect on sexual satisfaction in a more general population, we evaluated three types of self-reported penile coital injuries in a longitudinal community-based cohort of newly circumcised and uncircumcised men in Nyanza Province, Kenya.

Methods

Study design and participants

The study took place between November 2008 and January 2012 in one urban (Kisumu East) and two rural (Nyando and Kisumu West) sub-counties in western Kenya. Additional study details, sample description and the outcomes of the behavioral risk compensation assessment have been previously described by Westercamp et al. [33]. To participate, men had to be uncircumcised, 18 to 35 years old, live within the study area, and have no plans to relocate within the next 2 years. Eligible men self-selected into the circumcision group by seeking circumcision services at a voluntary medical male circumcision (VMMC) clinic within the study area, and were recruited before risk-reduction counseling and the circumcision procedure itself were completed. A control group was recruited from the community surrounding each VMMC clinic site and frequency-matched to age and residence (community) of the circumcision group. The controls were men who declined the opportunity to become circumcised before enrollment in the study.

Participants provided written informed consent in their language of choice (English, Dholuo or Kiswahili) and were offered 200 Kenyan shillings (about \$2.50) for each study visit to cover travel expenses and loss of income. Ethical approval was obtained from the Kenyatta National Hospital Ethics and Research Committee and the Institutional Review Board of the University of Illinois at Chicago.

Study procedures

Study participants completed a detailed sexual history and behavioral questionnaire and had their circumcision status visually confirmed by specially trained research assistants at each study visit: baseline, and 6, 12, 18, and 24-month follow-up visits. All participants were uncircumcised at baseline interview. Men who intended to become circumcised proceeded through the normal VMMC clinic flow following the Kenyan government VMMC guidelines.

Study questionnaires were administered through audio computer-assisted self-interview (ACASI) modules, developed in English, Dholuo, and Kiswahili. An equivalent paper-based questionnaire was used at participant request or in cases of power outage at study facilities (~30% of all questionnaires). The questionnaire instrument included items related to socio-demographic characteristics, sexual behaviors, history of STIs, general reproductive health, and sexual function and satisfaction including a set of questions addressing penile coital injuries. All data were self-reported and no biological samples were taken.

Statistical analyses

Self-reported penile coital injuries were determined by asking men, “In the past 6 months . . . : 1) how often during sex did the skin of your penis get scratches, cuts, or abrasions? (labeled as “scratches/cuts/abrasions” in the analyses); 2) how often during sex did your penis get sore? (labeled as “pain/soreness” in the analyses); and 3) how often during or after sex did the skin of your penis bleed? (labeled as “penile bleeding” in the analyses)” The response set was dichotomized for analysis as ever (by grouping always, often, sometimes or rarely) vs. never. To facilitate comparison, the penile injury assessment was the same as used by Mehta *et al.* in their 2010 evaluation of penile coital injuries in the Kenyan RCT and, to the extent possible, variables selected for analysis were kept consistent [27]. These variables included: condom use at last sex, preference for dry sex, applying substances on penis before sex, self-reported STIs in the past 6 months, genital hygiene after sex and circumcision status. Demographic (i.e., age, marital status, education, employment and ethnicity) and behavioral (i.e., number of partners

in the past 6 months) variables were also included in the analysis. Circumcision status was treated as a time-varying covariate to accommodate crossovers. Pearson χ^2 tests for categorical variables and Kolmogorov-Smirnov two-sample test for non-normally distributed continuous variables were used to assess baseline differences between circumcision and control group participants.

Because reports of abrasions, scratches or cuts to the penis may represent a misclassification of genital ulcers related to infectious etiologies, we identified men with more chronic GUD symptoms by asking, "Have you experienced any sores on or around genitals in the past 6 months." A visual genital exam was done to confirm circumcision status, but did not assess genital health or STI signs.

Sexual satisfaction was assessed by asking participants, "In the past 6 months, how would you generally rate your satisfaction with sexual intercourse?" The responses were dichotomized for analysis as satisfied (by grouping very satisfied and satisfied) and dissatisfied (by grouping very dissatisfied, dissatisfied, and unsure).

We compared penile coital injuries reported by circumcised and uncircumcised participants using random intercept mixed-effect models for binary outcomes to account for within-subject correlation due to repeated measures. Penile coital injuries were modeled for each participant as the linear slope over time assessed in 6-month intervals between the baseline and the 24-month follow-up time. The model included circumcision group as a binary variable and a group by time interaction to allow for varying trajectories between the two groups over time. To quantify the differences between circumcised and uncircumcised men over 24 months, we estimated odds ratios (OR) for group effect through mixed-effect models by excluding the circumcision by time interaction. All analyses were restricted to men sexually active in the 6 months preceding the interview. Covariates were selected for inclusion in multi-variable models based on univariate analyses ($p < 0.05$) and previous literature. All variables, except for age, education, and ethnicity, were time-varying covariates. Final model selection was done using backwards elimination with study time and age included in all models. Sexual satisfaction was modeled for each participant as the linear slope over time assessed in 6-month intervals between the baseline and the 24-month follow-up time. The primary independent predictors were the three types of penile coital injuries, with adjustment for time, circumcision status, age, education, employment, number of partners in the past 6 months, and reports of genital sores and STIs in the past 6 months. To ensure the comparability of our results with studies using other modeling approaches, population-averaged odds ratios were calculated by transforming our subject-specific regression estimates as described by Hu et al. [34]. All presented odds ratios are population-averaged. Statistical analyses were performed using SAS v9.2 [35] with the NLMIXED procedure for mixed-effect modeling.

Results

Sample description

Between November 2008 and April 2010, 3,186 of 3,825 (97%) of eligible men agreed to participate in the study. By design, study groups were equally sized (1,588 circumcision group; 1,598 control group). Approximately 5% of both groups (79/1588: circumcision and 74/1598: controls) were lost to follow-up after the baseline assessment and were excluded from longitudinal analyses. Men not returning for any follow-up were less likely to be Luo ($p = 0.03$) and less likely to have ever had sex ($p = 0.01$). Follow-up rates were 70% (6-months), 81% (12-months), 82% (18-months), and 84% (24-months), with similar loss to follow-up for the two groups. There were differences between the self-selected groups, with men in the circumcision group

Table 1. Baseline characteristics of study participants.

	Circumcision group (N = 1 588)	Control group (N = 1 598)	Overall (N = 3 186)	p-value
Demographic characteristics				
Age, mean (IQR, range)	20 (19–24; 18–35)	20 (19–24; 18–35)	20 (19–24; 18–35)	0.08
Ethnic group				0.01
Luo	1547 (97%)	1585 (99%)	3132 (98%)	
Educational level				
				0.01
Primary and less	354 (22%)	488 (31%)	842 (26%)	
Any secondary	955 (60%)	882 (55%)	1837 (58%)	
Any post-secondary	266 (17%)	306 (13%)	472 (15%)	
Unsure / Refused to answer	13 (1%)	22 (1%)	35 (1%)	
Employment status				
				0.01
Employed	421 (27%)	584 (37%)	1005 (32%)	
Marital status				
				0.01
Single, without live-in partner	1055 (66%)	956 (60%)	2011 (63%)	
Single, with live-in partner	220 (14%)	203 (13%)	423 (13%)	
Married, living with wife	271 (17%)	401 (25%)	672 (21%)	
Married, not living with wife	42 (3%)	38 (2%)	80 (3%)	
Sexual history				
Ever had sex				
				0.13
Yes	1382 (87%)	1419 (89%)	2801 (88%)	
No	206 (13%)	179 (11%)	385 (12%)	
Age (years) at first sex (IQR; range; N)	16 (15–18; 9–30; 1380)	16 (15–18; 9–29; 1417)	16 (15–18; 9–30; 2797)	0.66
Sexual intercourse in past 6 months (sexually active only)				
				0.54
Yes	1032 (75%)	1074 (76%)	2106 (75%)	
No	350 (25%)	345 (24%)	695 (25%)	
Number of partners in past 6 months (sexually active only)				
				0.13
None	350 (25%)	345 (24%)	960 (34%)	
One	502 (36%)	564 (40%)	1008 (36%)	
2+	393 (29%)	398 (28%)	786 (28%)	
Unsure / Refused to answer	137 (10%)	112 (8%)	147 (2%)	
Lifetime number of partners	3 (2–6; 1–552; 1193)	3 (2–6; 1–122; 1270)	3 (2–6; 1–552; 2463)	0.73
Penile coital injuries in the past 6 months (sexually active in last 6 months, N = 2,045)				
Reported penile coital injuries in the past 6 months:				
Scratches/cuts/abrasions	477 (47%)	428 (41%)	905 (44%)	0.01
Penile pain/soreness	343 (34%)	321 (31%)	664 (32%)	0.15
Penile bleeding	234 (23%)	211 (20%)	445 (22%)	0.12
Any penile coital injury	562 (56%)	518 (50%)	1080 (53%)	0.01

Sample sizes vary in questions based on past or recent sexual activity. Data are median (IQR; range; n) for continuous data, or n (%) for categorical data. P values are based on Kolmogorov-Smirnov two-sample test for non-normally distributed continuous data and chi-square for categorical data.

<https://doi.org/10.1371/journal.pone.0185917.t001>

more likely to have a secondary or higher education, be unemployed, and single than the control group (Table 1); and no differences in other demographic or behavioral characteristics.

Baseline penile coital injuries

Among the 2,106 (66%) participants sexually active in the 6 months prior to baseline, 2,045 (97%) answered the questions about penile coital injuries (Table 1). In total, 1,080 (53%) reported any penile injury, including 44% reporting scratches, cuts, or abrasions; 32% penile

pain/soreness; and 22% penile bleeding. Overall, 259 (13%) of men reported all three types of coital injuries, 413 (20%) two types, and 408 (20%) reported a single type of penile coital injury.

At baseline, men who selected circumcision were significantly more likely to report a recent history of penile coital scratches, cuts, or abrasions compared to men who were initially not planning to become circumcised (47% vs. 41%, $p = 0.007$; Table 1). Reports of penile pain/soreness and penile bleeding did not differ by circumcision intent at baseline (Table 1). Combined, a greater proportion of those who selected circumcision reported at least one type of penile coital injury compared to those who selected to remain uncircumcised (56% vs. 50%; $p = 0.01$; Table 1).

Penile coital injuries by circumcision status

Over 24 months of follow-up, reports of any penile coital injuries following circumcision declined from 56% (pre-circumcision baseline) to 15%, a 73% relative decrease. Significant decline was noted for each component individually: scratches/cuts/abrasions (81% decrease), penile pain/soreness (71% decrease), and penile bleeding (87% decrease). All declines were evident at 6-months post-procedure and were sustained throughout the follow-up period (Fig 1). Among the uncircumcised group, no decline was noted in cuts, scratches, abrasions or post-coital penile soreness, while a 25% decline in coital penile bleeding from 20% of men at enrollment to 15% at 24 months was observed.

Longitudinal analysis using mixed-effect models revealed the presence of group by time interaction ($p < 0.0001$) for all measures of penile coital injuries indicating different patterns of change in injuries over time for circumcised and uncircumcised men. When stratified by




Fig 1. Observed self-reported penile coital injury by circumcision status over time. Notes: There was a significant decline with time for all measures of penile coital injuries, except for scratches/cuts/abrasions (A) and pain/soreness (B) in uncircumcised men.

<https://doi.org/10.1371/journal.pone.0185917.g001>

Table 2. Longitudinal changes in penile coital injuries over 24 months of follow up by circumcision status: the results of mixed effect models (stratified by circumcision status, unadjusted for other covariates).

	Stratified analysis					
	Circumcised			Uncircumcised		
	Estimate	SE	p	Estimate	SE	p
Scratches, cuts and abrasions						
Change over time (visit #)	-0.719	0.037	0.01	-0.031	0.026	0.24
Penile pain/soreness						
Change over time (visit #)	-0.451	0.032	0.01	-0.015	0.027	0.58
Penile bleeding						
Change over time (visit #)	-0.717	0.05	0.01	-0.121	0.032	0.01
Any penile injuries						
Change over time (visit #)	-0.609	0.031	0.01	-0.082	0.025	0.01

<https://doi.org/10.1371/journal.pone.0185917.t002>

circumcision status, decline from pre-circumcision baseline over time was significant for circumcised men across all three types of coital injuries (Table 2). For uncircumcised men, statistically significant decline of lower magnitude was observed only for penile bleeding and for the combined measure of any coital injury (Table 2).

In addition to exploring the differential changes in coital injuries in each group over time (Table 2), we reran the models without the group by time interaction term to be able to quantify the association between coital injuries and circumcision for the duration of the study. By removing the interaction term from the model, the estimates of the effect of circumcision were adjusted for time, but not for the differential dynamics of time for each group. As a result, OR comparing circumcised to uncircumcised men over 24 months of follow up was 0.49 (95% CI: 0.44–0.55) for any penile coital injury. By type, odds ratios were 0.41 (95% CI: 0.37–0.47) for scratches, cuts and abrasions, 0.59 (95% CI: 0.52–0.66) for penile pain/soreness, and 0.54 (95% CI: 0.47–0.62), for penile bleeding.

Penile coital injuries and genital sores

At baseline, 186 (9%) of sexually active men reported genital sores in the past 6 months. Compared to men not reporting sores, a higher proportion of men with genital sores were seeking circumcision, had only primary education, were employed, married, had multiple partners in the past 6 months, had unprotected sex, preferred dry sex, and reported STIs in the past 6 months. Following circumcision, the proportion of men reporting genital sores declined from 11% (baseline, among men selecting circumcision) to 2% at 24-months. Among the uncircumcised men, no change in the prevalence of genital sores was observed: 8% at baseline to 8% at 24 months. At baseline, 14% of men reporting any penile coital injuries also reported genital sores (14% of men reporting scratches, cuts, or abrasions, 19% reporting penile pain/soreness, and 14% with penile bleeding). Conversely, over 81% of men reporting a genital sore at baseline also reported a concomitant penile coital injury (Table 3), including 68% reporting scratches, cuts, or abrasions, 66% with penile pain/soreness, and 34% with bleeding.

In newly circumcised men reporting genital sores, concurrent report of any penile coital injuries did not significantly decline over follow-up (pre-circumcision baseline 81% to 79% at 24-months; $p = 0.22$). In men with no history of genital sores, however, reports of penile coital injuries declined from 53% (pre-circumcision baseline) to 14% ($p < 0.001$; ~80% relative decrease) over the 24-months following circumcision. Therefore, following circumcision we observe a decline in reports of recent genital sores and decline in penile coital injuries, but the

Table 3. Prevalence of reported coital injuries across time-varying covariates significantly associated with coital injuries in univariate analyses.

	Baseline, n(%)	6 months, n(%)	12 months, n(%)	18 months, n(%)	24 months, n(%)
Circumcision status*					
Circumcised	562/1008(56%)	200/732 (27%)	202/929(22%)	238/1071(22%)	180/1199(15%)
Uncircumcised	518/1037(50%)	401/888(45%)	390/920(42%)	386/892(43%)	375/848 (44%)
Marital status *					
Single	569/1143(50%)	270/852(32%)	295/976(30%)	271/1001(27%)	1236/1008(23%)
Married or cohabitating	511/902(57%)	331/768(43%)	297/873(34%)	353/962(37%)	319/1039(31%)
Condom use at last sex *					
No condom used	572/1001(57%)	321/736(44%)	324/861(38%)	321/861(37%)	280/912(31%)
Condom used	435/892(49%)	252/800(32%)	243/917(27%)	290/1064(27%)	263/1106(24%)
Number of partners in the past 6 months *					
One	444/1012(44%)	257/804(32%)	276/1030(27%)	311/1171(27%)	328/1353(24%)
Two or more	501/756(66%)	294/652(45%)	275/686(40%)	279/695(40%)	214/643(33%)
How long until washed penis after last time had sex *					
One hour or less	383/783(49%)	232/662(35%)	236/798(30%)	261/917(29%)	275/1085(25%)
More than one hour	697/1262(55%)	369/958(39%)	356/1051(34%)	363/1047(35%)	280/962(29%)
Applied substances to penis before sex in the past 6 months *					
No	885/1719(52%)	516/1407(37%)	507/1653(31%)	543/1776(31%)	493/1874(26%)
Yes	85/114(75%)	46/89(52%)	45/75(60%)	43/79(54%)	37/83(45%)
Preference for dry sex					
Prefers dry sex	395/749(53%)	231/614(38%)	232/662(35%)	260/727(36%)	219/695(36%)
Prefers wet sex or no opinion	575/1084(53%)	331/881(38%)	320/1066(30%)	326/1128(29%)	311/1262(25%)
STIs in the past 6 months *					
No	885/1715(52%)	525/1440(37%)	519/1675(31%)	548/1799(31%)	500/1909(26%)
Yes	88/118(75%)	38/56(68%)	33/49(67%)	37/54(69%)	29/45(64%)
Genital sores in the past 6 months *					
No	930/1859(50%)	532/1533(35%)	519/1750(30%)	540/1860(29%)	478/1950(25%)
Yes	150/186(81%)	69/87(79%)	73/99(74%)	84/104(81%)	77/97(79%)

Circumcision status at baseline is by enrollment group; at follow up by actual status;

* Significant baseline difference

<https://doi.org/10.1371/journal.pone.0185917.t003>

latter only when men have not also experienced penile sores, indicating differential trends in prevalence of coital injuries among circumcised men with and without genital sores, as confirmed by the significant interaction term ($p = 0.002$).

Considering any report of genital sores separately from that of penile coital injuries, the protective effect of circumcision on both is evident (Fig 2). Unadjusted ORs for the effect of circumcision over the duration of follow up were 0.67 (95% CI: 0.56–0.79) for recent history of genital sores. By type of coital injury, excluding those with genital sores, ORs by circumcision status were 0.43 (95%CI: 0.38–0.48) for scratches, cuts and abrasions, 0.61 (95%CI: 0.54–0.69) for penile pain/soreness, and 0.58 (95%CI: 0.50–0.67) for bleeding. Excluding men with genital sores did not greatly change (<10%) the magnitude of the association, therefore our multivariable analyses included men with and without genital sores and reports of genital sores were evaluated as a covariate for each type of penile coital injuries.

Other covariates and multivariable models

The distribution over time of penile coital injuries across time-varying covariates significant in the univariate analyses is shown in Table 3. These covariates were included in the multivariable




Fig 2. Genital sores and type of coital injury among those without genital sores over time (circumcised men only).

<https://doi.org/10.1371/journal.pone.0185917.g002>

modeling to assess their association with penile coital injuries. The statistically significant covariates were retained in the final models and are shown in Table 4. In multivariable modeling, adjusted for other covariates, circumcised men were more than 50% less likely than those uncircumcised to report any penile coital injury in the last 6 months (aOR = 0.47; 95%CI 0.42–0.53). This protection was greatest against penile cuts, scratches, or abrasions for which the difference reached 60% (aOR = 0.39; 95%CI 0.34–0.44; Table 4). Factors independently associated with increased risk of each type penile coital injury were the application of substances (lubricants) to the penis before sex (aORs ranged 1.94 to 2.29), increasing age (aOR = 1.03 for each added year), history of STI in the last 6 months (aORs ranged 1.66 to 2.48), reporting genital sores in the last 6 months (aORs ranged 2.60 to 4.27), and multiple partners in last 6-months (aORs ranged 1.38 to 1.58). Condom use (aORs ranged 0.73 to 0.75) and time remained protective for penile coital injuries. While these factors were significantly associated with penile coital injuries, they did not act as confounders to the effect of circumcision, as seen by comparing the crude and adjusted ORs for the three types of penile coital injuries, though adjusting for covariates did attenuate the magnitude of change over time. Out of the variables differentiating men selecting circumcision and controls at baseline (i.e., education, employment, ethnicity and marriage), only marriage was a significant risk factor for coital injuries in the univariate analyses, but not in the multivariable analyses. Likewise, post-coital hygiene, and preference for dry sex were not associated with any of penile coital injury measures.

Penile coital injuries and sexual satisfaction

Sexual satisfaction and its association with male circumcision are examined in depth elsewhere [36]. Here we explore the associations between male circumcision and sexual satisfaction, in

Table 4. Results of the multivariable mixed effect models for risks of penile coital injury over time (N = 2781).

	Scratches/cuts/ abrasions	Penile pain /soreness	Penile bleeding
	OR (95% CI)	OR (95% CI)	OR (95% CI)
Circumcision status*			
Uncircumcised	Reference	Reference	Reference
Circumcised	0.39 (0.34–0.44)	0.58 (0.51–0.65)	0.53 (0.46–0.62)
Age (continuous)	1.03 (1.02–1.05)	1.03 (1.01–1.04)	1.02 (1.00–1.04)
Condom use at last sex			
No condom used	Reference	Reference	Reference
Condom used	0.74 (0.67–0.83)	0.73 (0.65–0.81)	0.75 (0.65–0.86)
Number of partners in the past 6 months			
One	Reference	Reference	Reference
Two or more	1.48 (1.35–1.63)	1.38 (1.26–1.52)	1.58 (1.41–1.78)
Ever applied substances to penis before sex			
No	Reference	Reference	Reference
Yes	1.94 (1.55–2.42)	2.08 (1.66–2.60)	2.29 (1.79–2.93)
Self-reported genital sores in the past 6 months			
No	Reference	Reference	Reference
Yes	3.65 (2.99–4.46)	4.27 (5.09–6.39)	2.60 (2.10–3.20)
Self-reported STIs in the past 6 months			
No	Reference	Reference	Reference
Yes	1.66 (1.29–2.15)	2.48 (1.92–3.20)	2.16 (1.65–3.84)
Time (visit)			
Baseline	Reference	Reference	Reference
6 months	0.43 (0.37–0.49)	0.77 (0.66–0.89)	0.60 (0.50–0.71)
12 months	0.38 (0.32–0.44)	0.67 (0.57–0.78)	0.45 (0.37–0.54)
18 months	0.43 (0.37–0.49)	0.66 (0.57–0.76)	0.48 (0.40–0.58)
24 months	0.39 (0.34–0.45)	0.59 (0.51–0.69)	0.39 (0.33–0.48)

* Circumcision status at baseline is by enrollment group; at follow up by actual status.
All variables are time varying, except for age at baseline

<https://doi.org/10.1371/journal.pone.0185917.t004>

the context of penile coital injuries. At baseline, the proportion of men reporting being satisfied with sexual intercourse was lower among men reporting penile coital injuries compared to those not reporting coital injuries: 67% of men with scratches, cuts, and abrasions were satisfied with intercourse compared to 74% without scratches, cuts, and abrasions ($p = 0.001$); 65% with penile pain/soreness vs. 74% without penile pain/soreness ($p < 0.001$); and 61% with penile bleeding vs. 74% without penile bleeding ($p < 0.001$). Adjusted for time, circumcision status, age, education, employment, number of partners in the past 6 months, and reports of genital sores and STIs in the past 6 months, penile coital injuries remained significantly associated with lower levels of sexual satisfaction in longitudinal analyses (scratches, cuts and abrasions: aOR = 0.87, 95% CI: 0.76–0.98; penile pain/soreness: aOR = 0.82, 95% CI: 0.72–0.93; and penile bleeding: aOR = 0.65, 95% CI: 0.55–0.76). After adjusting for penile coital injuries, history of STIs and genital sores, and other factors, circumcision status remained marginally associated with increased sexual satisfaction (aOR = 1.13, 95% CI: 0.99–1.28).

Discussion

In our study comparing recently circumcised men and uncircumcised men in western Kenya, we confirm previous observations that circumcised men are less likely to report penile coital

injuries, with significantly decreased risk observed as early as 6 months after surgery and decreasing further over 24 months [27, 32]. Factors other than circumcision associated with penile coital injury included increasing age, increasing number of sexual partners, application of substances to the penis before sex, and self-reported history of STIs. The differences in prevalence of penile coital injuries at the baseline among otherwise similar groups indicates that it is likely that men reporting penile coital injuries were more likely to seek VMMC services. This study is the first to identify an association between self-reported penile coital injuries and decreased sexual satisfaction.

We found that coital-related scratches, cuts, and abrasions to the penis among young, uncircumcised, sexually active men were common (44%). This prevalence is comparable to the 48% observed by Mehta and colleagues in this same geographical area [27], but higher than the 34% found in a 1997 cross-sectional study in eastern Uganda [37], and lower than 61% reported in the cohort of circumcised men in the Dominican Republic before circumcision [32]. Several explanations for these differences are possible including: differences in the behavioral risk profile of the study samples, culturally specific sexual practices that increase the risk of coital injuries, circumcision status misclassification, and misclassification of GUD and injuries.

Reduction in GUD is a highly likely mechanism for at least some portion of the protective effect of MC against HIV [7, 25, 38], with a 41%-48% reduction in GUD following the procedure observed in recent studies [12, 28]. While not directly comparable due to methodological differences in the assessment of penile sores and genital ulcers, the 33% decline in the likelihood of self-reported genital sores with circumcision observed in our study is in the lower range of effect sizes reported. Moreover, men reporting genital sores were more likely to report penile coital injuries, suggesting an additional risk of injury with genital sores or some overlap between the symptoms of genital sores and of penile coital injuries. Unlike other studies, we found no significant reductions ($p = 0.22$) in penile injuries over time among men reporting genital sores after becoming circumcised. This finding could possibly be a result of unadjusted confounding due to the ambiguity in our measurements of genital sores and should be evaluated further.

Recent research has observed that approximately 40% to 60% of genital ulcers are not explained by STI etiologies [12, 28, 29]. Coital injuries may play a role in the formation of these unexplained ulcers through facilitating infections by other, non-sexually transmitted, pathogens [12, 28, 29, 38]. In stratified analysis, we observed that recent genital sores were reported by 14% of men with penile coital injuries. This leaves a great majority of penile coital injuries that are likely attributable to mechanical disruption or factors unrelated to preexisting GUD. Future studies should include questions designed to differentiate between coital injuries and GUD or include clinical examination to confirm current injuries or sores.

This is the first study to identify an association between penile coital injuries and decreased sexual satisfaction. This association remained significant over time and for all three types of coital injuries, independent of age, circumcision status, level of sexual activity or other causes of genital discomfort, such as GUD and other STIs. In our study, men who were seeking circumcision were more likely to report penile coital injuries at the baseline. As described in an in-depth analysis of male circumcision and sexual satisfaction based on the same cohort of men [36], men self-selecting into the circumcision group were also more likely to report lower levels of sexual satisfaction at baseline. While we did not assess this directly, it is possible that lower sexual satisfaction associated with coital injuries was one of the motivating factors behind their decision to become circumcised.

Our findings are subject to several limitations. Data on penile coital injuries, genital sores and STIs were based on self-report with no corresponding clinical exam. However, similarities

between our results and those from the RCT of male circumcision in Kisumu support the generalizability of our results: we used the same questions as the trial in a sample different in demographic, geographic, and other characteristics. Because study participants self-selected for enrollment and group assignment, it is possible that the motivation to become circumcised represents fundamental differences between study groups. Behavior, sexual history, and sexual satisfaction were self-reported, and they are subject to social desirability and recall biases, although we have limited these biases through the use of ACASI and experienced study staff with training in sensitive face-to-face interview techniques [39–42]. Sexual satisfaction was assessed using questions similar to those used in other studies [43], but we did not use validated instruments. Lastly, our study did not assess the likely mechanisms leading to coital injuries. This is an important aspect in determining how circumcision may be protecting men and ultimately in developing additional interventions and messaging.

Conclusion

Penile coital injuries have logical and observable associations with increased risk of HIV and STI infection [25, 38, 44, 45]. While their prevention may be important in that regard alone, the potential motivational force for circumcision may also be of value. In our study we found that men reporting penile coital injuries were more likely to seek VMMC services and observed a significant decline in coital injuries following circumcision. Further, men seeking circumcision services had consistently lower levels of pre-procedure sexual satisfaction [46] and those with penile coital injuries had lower levels of sexual satisfaction both at the baseline and across follow-up. Improvement of the sexual experience through the removal of a potential source of sexual discomfort may resonate with a significant portion of men targeted for VMMC [25, 38, 44, 45, 47]. The accumulation of evidence indicating an independent role of penile coital injuries in decreased sexual satisfaction, HIV [31], and HSV-2 [30], merits comprehensive study to clinically and etiologically define penile coital injuries for potential intervention targets.

Supporting information

S1 File. Study questionnaires (English, Dholuo, and Kiswahili).
(ZIP)

S2 File. Data repository.
(CSV)

Acknowledgments

We thank all of the participants, without whom this work would not have been possible. We are grateful to Cosam Ang'awa, Yusto Okemia, Kevine Amolloh, Kelvin Akoth, Kennedy Otieno, Danstan Ochieng', Victor Odula, George Kidi, David Ang'awa, Evans Otieno, Erik Ogutu, George Ong'eng'a, and Richard Okello for their dedication in recruitment, data collection, tracing, and overall commitment to the study; to Nicholas Obwama and Joseph Abuya for their tireless data entry and cleaning efforts; to Matthew Westercamp for his invaluable input and for reading and editing multiple versions of this manuscript; to Christine L. Mattson for inspiration; and to the entire NRHS staff for their assistance in making this study a success.

Author Contributions

Conceptualization: Nelli Westercamp, Supriya D. Mehta, Walter Jaoko, Robert C. Bailey.

Data curation: Nelli Westercamp, Timothy A. Okeyo.

Formal analysis: Nelli Westercamp.

Funding acquisition: Robert C. Bailey.

Investigation: Nelli Westercamp, Robert C. Bailey.

Methodology: Nelli Westercamp, Supriya D. Mehta, Robert C. Bailey.

Project administration: Nelli Westercamp, Timothy A. Okeyo, Robert C. Bailey.

Resources: Walter Jaoko, Robert C. Bailey.

Software: Nelli Westercamp, Timothy A. Okeyo.

Supervision: Nelli Westercamp, Robert C. Bailey.

Validation: Nelli Westercamp, Supriya D. Mehta, Timothy A. Okeyo.

Visualization: Nelli Westercamp, Supriya D. Mehta.

Writing – original draft: Nelli Westercamp.

Writing – review & editing: Nelli Westercamp, Supriya D. Mehta, Walter Jaoko, Timothy A. Okeyo, Robert C. Bailey.

References

1. Bailey RC, Moses S, Parker CB, Agot K, Maclean I, Krieger JN, et al. Male circumcision for HIV prevention in young men in Kisumu, Kenya: a randomised controlled trial. *Lancet*. 2007; 369(9562):643–56. Epub 2007/02/27. [https://doi.org/10.1016/S0140-6736\(07\)60312-2](https://doi.org/10.1016/S0140-6736(07)60312-2) PMID: 17321310.
2. Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, et al. Male circumcision for HIV prevention in men in Rakai, Uganda: a randomised trial. *Lancet*. 2007; 369(9562):657–66. Epub 2007/02/27. [https://doi.org/10.1016/S0140-6736\(07\)60313-4](https://doi.org/10.1016/S0140-6736(07)60313-4) PMID: 17321311.
3. Auvert B, Taljaard D, Lagarde E, Sobngwi-Tambekou J, Sitta R, Puren A. Randomized, controlled intervention trial of male circumcision for reduction of HIV infection risk: the ANRS 1265 Trial. *PLoS medicine*. 2005; 2(11):e298. Epub 2005/10/20. <https://doi.org/10.1371/journal.pmed.0020298> PMID: 16231970.
4. Siegfried N, Muller M, Deeks JJ, Volmink J. Male circumcision for prevention of heterosexual acquisition of HIV in men. *Cochrane database of systematic reviews (Online)*. 2009;(2):CD003362. Epub 2009/04/17. <https://doi.org/10.1002/14651858.CD003362.pub2> PMID: 19370585.
5. World Health Organization & United Nations Joint Programme on HIV/AIDS. Male circumcision: global trends and determinants of prevalence, safety and acceptability. Geneva: WHO, 2007.
6. Dinh MH, Fahrback KM, Hope TJ. The role of the foreskin in male circumcision: an evidence-based review. *American journal of reproductive immunology (New York, NY: 1989)*. 2011; 65(3):279–83. Epub 2010/12/01. <https://doi.org/10.1111/j.1600-0897.2010.00934.x> PMID: 21114567.
7. Cameron DW, Simonsen JN, D'Costa LJ, Ronald AR, Maitha GM, Gakinya MN, et al. Female to male transmission of human immunodeficiency virus type 1: risk factors for seroconversion in men. *Lancet*. 1989; 2(8660):403–7. Epub 1989/08/19. PMID: 2569597.
8. Fink AJ. A possible explanation for heterosexual male infection with AIDS. *The New England journal of medicine*. 1986; 315(18):1167. Epub 1986/10/30. PMID: 3762636.
9. Kigozi G, Wawer M, Ssettuba A, Kagaayi J, Nalugoda F, Watya S, et al. Foreskin surface area and HIV acquisition in Rakai, Uganda (size matters). *AIDS (London, England)*. 2009; 23(16):2209–13. Epub 2009/09/23. <https://doi.org/10.1097/QAD.0b013e328330eda8> PMID: 19770623.
10. Freeman EE, Weiss HA, Glynn JR, Cross PL, Whitworth JA, Hayes RJ. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *AIDS (London, England)*. 2006; 20(1):73–83. Epub 2005/12/06. PMID: 16327322.
11. Mayer KH, Venkatesh KK. Interactions of HIV, other sexually transmitted diseases, and genital tract inflammation facilitating local pathogen transmission and acquisition. *American journal of reproductive immunology (New York, NY: 1989)*. 2011; 65(3):308–16. Epub 2011/01/11. <https://doi.org/10.1111/j.1600-0897.2010.00942.x> PMID: 21214660.

12. Gray RH, Serwadda D, Tobian AA, Chen MZ, Makumbi F, Suntoke T, et al. Effects of genital ulcer disease and herpes simplex virus type 2 on the efficacy of male circumcision for HIV prevention: Analyses from the Rakai trials. *PLoS medicine*. 2009; 6(11):e1000187. Epub 2009/11/26. <https://doi.org/10.1371/journal.pmed.1000187> PMID: 19936044.
13. Weiss HA, Thomas SL, Munabi SK, Hayes RJ. Male circumcision and risk of syphilis, chancroid, and genital herpes: a systematic review and meta-analysis. *Sexually transmitted infections*. 2006; 82(2):101–9; discussion 10. Epub 2006/04/04. <https://doi.org/10.1136/sti.2005.017442> PMID: 16581731.
14. Porter WM, Bunker CB. The dysfunctional foreskin. *International journal of STD & AIDS*. 2001; 12(4):216–20. Epub 2001/04/26. <https://doi.org/10.1258/0956462011922922> PMID: 11319970.
15. Price LB, Liu CM, Johnson KE, Aziz M, Lau MK, Bowers J, et al. The effects of circumcision on the penis microbiome. *PloS one*. 2010; 5(1):e8422. Epub 2010/01/13. <https://doi.org/10.1371/journal.pone.0008422> PMID: 20066050.
16. Fahrback KM, Barry SM, Anderson MR, Hope TJ. Enhanced cellular responses and environmental sampling within inner foreskin explants: implications for the foreskin's role in HIV transmission. *Mucosal immunology*. 2010; 3(4):410–8. Epub 2010/04/23. <https://doi.org/10.1038/mi.2010.18> PMID: 20410876.
17. Ganor Y, Zhou Z, Tudor D, Schmitt A, Vacher-Lavenu MC, Gibault L, et al. Within 1 h, HIV-1 uses viral synapses to enter efficiently the inner, but not outer, foreskin mucosa and engages Langerhans-T cell conjugates. *Mucosal immunology*. 2010; 3(5):506–22. Epub 2010/06/24. <https://doi.org/10.1038/mi.2010.32> PMID: 20571487.
18. Zhou Z, Barry de Longchamps N, Schmitt A, Zerbib M, Vacher-Lavenu MC, Bomsel M, et al. HIV-1 efficient entry in inner foreskin is mediated by elevated CCL5/RANTES that recruits T cells and fuels conjugate formation with Langerhans cells. *PLoS pathogens*. 2011; 7(6):e1002100. Epub 2011/07/09. <https://doi.org/10.1371/journal.ppat.1002100> PMID: 21738469.
19. Desai K, Boily MC, Garnett GP, Masse BR, Moses S, Bailey RC. The role of sexually transmitted infections in male circumcision effectiveness against HIV—insights from clinical trial simulation. *Emerging themes in epidemiology*. 2006; 3:19. Epub 2006/12/26. <https://doi.org/10.1186/1742-7622-3-19> PMID: 17187662.
20. Boily MC, Desai K, Masse B, Gumel A. Incremental role of male circumcision on a generalised HIV epidemic through its protective effect against other sexually transmitted infections: from efficacy to effectiveness to population-level impact. *Sexually transmitted infections*. 2008; 84 Suppl 2:ii28–34. Epub 2008/10/01. <https://doi.org/10.1136/sti.2008.030346> PMID: 18799489.
21. Anderson D, Politch JA, Pudney J. HIV infection and immune defense of the penis. *American journal of reproductive immunology (New York, NY: 1989)*. 2011; 65(3):220–9. Epub 2011/01/11. <https://doi.org/10.1111/j.1600-0897.2010.00941.x> PMID: 21214659.
22. Morris BJ, Bailey RC, Klausner JD, Leibowitz A, Wamai RG, Waskett JH, et al. Review: a critical evaluation of arguments opposing male circumcision for HIV prevention in developed countries. *AIDS care*. 2012; 24(12):1565–75. Epub 2012/03/29. <https://doi.org/10.1080/09540121.2012.661836> PMID: 22452415.
23. Simonsen JN, Cameron DW, Gakinya MN, Ndinya-Achola JO, D'Costa LJ, Karasira P, et al. Human immunodeficiency virus infection among men with sexually transmitted diseases. Experience from a center in Africa. *The New England journal of medicine*. 1988; 319(5):274–8. Epub 1988/08/04. <https://doi.org/10.1056/NEJM198808043190504> PMID: 3393182.
24. Stone KM, Grimes DA, Magder LS. Primary prevention of sexually transmitted diseases. A primer for clinicians. *JAMA: the journal of the American Medical Association*. 1986; 255(13):1763–6. Epub 1986/04/04. PMID: 3512873.
25. Szabo R, Short RV. How does male circumcision protect against HIV infection? *BMJ (Clinical research ed)*. 2000; 320(7249):1592–4. Epub 2000/06/14. PMID: 10845974.
26. Halperin DT, Bailey RC. Male circumcision and HIV infection: 10 years and counting. *Lancet*. 1999; 354(9192):1813–5. Epub 1999/11/30. [https://doi.org/10.1016/S0140-6736\(99\)03421-2](https://doi.org/10.1016/S0140-6736(99)03421-2) PMID: 10577659.
27. Mehta SD, Krieger JN, Agot K, Moses S, Ndinya-Achola JO, Parker C, et al. Circumcision and reduced risk of self-reported penile coital injuries: results from a randomized controlled trial in Kisumu, Kenya. *The Journal of urology*. 2010; 184(1):203–9. Epub 2010/05/21. <https://doi.org/10.1016/j.juro.2010.03.015> PMID: 20483156.
28. Mehta SD, Moses S, Parker CB, Agot K, Maclean I, Bailey RC. Circumcision status and incident herpes simplex virus type 2 infection, genital ulcer disease, and HIV infection. *AIDS (London, England)*. 2012; 26(9):1141–9. Epub 2012/03/03. <https://doi.org/10.1097/QAD.0b013e328352d116> PMID: 22382150.
29. Mehta SD, Green SJ, Maclean I, Hu H, Bailey RC, Gillevet PM, et al. Microbial diversity of genital ulcer disease in men enrolled in a randomized trial of male circumcision in Kisumu, Kenya. *PloS one*. 2012; 7(7):e38991. Epub 2012/08/01. <https://doi.org/10.1371/journal.pone.0038991> PMID: 22848346.

30. Mehta SD, Moses S, Agot K, Parker C, Ndinya-Achola JO, Maclean I, et al. Adult male circumcision does not reduce the risk of incident *Neisseria gonorrhoeae*, *Chlamydia trachomatis*, or *Trichomonas vaginalis* infection: results from a randomized, controlled trial in Kenya. *The Journal of infectious diseases*. 2009; 200(3):370–8. Epub 2009/06/24. <https://doi.org/10.1086/600074> PMID: 19545209.
31. Mehta SD, Moses S, Agot K, Odoyo-June E, Li H, Maclean I, et al. The long term efficacy of medical male circumcision against HIV acquisition. *AIDS (London, England)*. 2013. Epub 2013/07/10. <https://doi.org/10.1097/01.aids.0000432444.30308.2d> PMID: 23835501.
32. Brito MO, Khosla S, Pananookooln S, Fleming PJ, Lerebours L, Donastorg Y, et al. Sexual Pleasure and Function, Coital Trauma, and Sex Behaviors After Voluntary Medical Male Circumcision Among Men in the Dominican Republic. *J Sex Med*. 2017; 14(4):526–34. <https://doi.org/10.1016/j.jsxm.2017.01.020> Epub Mar 1. PMID: 28258953
33. Westercamp N, Agot K, Jaoko W, Bailey RC. Risk compensation following male circumcision: results from a two-year prospective cohort study of recently circumcised and uncircumcised men in Nyanza Province, Kenya. *AIDS Behav*. 2014; 18(9):1764–75. <https://doi.org/10.1007/s10461-014-0846-4> PMID: 25047688
34. Hu FB, Goldberg J, Hedeker D, Flay BR, Pentz MA. Comparison of population-averaged and subject-specific approaches for analyzing repeated binary outcomes. *Am J Epidemiol*. 1998; 147(7):694–703. PMID: 9554609.
35. SAS Institute Inc. SAS software. Version 9.1.3 ed. Cary, NC2008.
36. Nordstrom MP, Westercamp N, Jaoko W, Okeyo T, Bailey RC. Medical Male Circumcision Is Associated With Improvements in Pain During Intercourse and Sexual Satisfaction in Kenya. *J Sex Med*. 2017; 14(4):601–12. <https://doi.org/10.1016/j.jsxm.2017.02.014> PMID: 28364982
37. Bailey RC, Neema S, Othieno R. Sexual behaviors and other HIV risk factors in circumcised and uncircumcised men in Uganda. *J Acquir Immune Defic Syndr*. 1999; 22(3):294–301. PMID: 10770351
38. O'Farrell N. Soap and water prophylaxis for limiting genital ulcer disease and HIV-1 infection in men in sub-Saharan Africa. *Genitourinary medicine*. 1993; 69(4):297–300. Epub 1993/08/01. PMID: 7721293.
39. Langhaug LF, Sherr L, Cowan FM. How to improve the validity of sexual behaviour reporting: systematic review of questionnaire delivery modes in developing countries. *Tropical medicine & international health: TM & IH*. 2010; 15(3):362–81. Epub 2010/04/23. <https://doi.org/10.1111/j.1365-3156.2009.02464.x> PMID: 20409291.
40. Mensch BS, Hewett PC, Erulkar AS. The reporting of sensitive behavior by adolescents: a methodological experiment in Kenya. *Demography*. 2003; 40(2):247–68. Epub 2003/07/09. PMID: 12846131.
41. Phillips AE, Gomez GB, Boily MC, Garnett GP. A systematic review and meta-analysis of quantitative interviewing tools to investigate self-reported HIV and STI associated behaviours in low- and middle-income countries. *International journal of epidemiology*. 2010; 39(6):1541–55. Epub 2010/07/16. <https://doi.org/10.1093/ije/dyq114> PMID: 20630991.
42. Vu LT, Nadol P, Le LC. HIV-Related Risk Behaviors Among the General Population: A Survey Using Audio Computer-Assisted Self-Interview in 3 Cities in Vietnam. *Asia-Pacific journal of public health / Asia-Pacific Academic Consortium for Public Health*. 2012. Epub 2012/06/30. <https://doi.org/10.1177/1010539512450611> PMID: 22743864.
43. Krieger JN, Mehta SD, Bailey RC, Agot K, Ndinya-Achola JO, Parker C, et al. Adult male circumcision: effects on sexual function and sexual satisfaction in Kisumu, Kenya. *J Sex Med*. 2008; 5(11):2610–22. <https://doi.org/10.1111/j.1743-6109.2008.00979.x> Epub 2008 Aug 28. PMID: 18761593
44. Figueroa JP, Brathwaite A, Morris J, Ward E, Peruga A, Blattner W, et al. Rising HIV-1 prevalence among sexually transmitted disease clinic attenders in Jamaica: traumatic sex and genital ulcers as risk factors. *J Acquir Immune Defic Syndr*. 1994; 7(3):310–6. Epub 1994/03/01. PMID: 8106971.
45. Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sexually transmitted infections*. 1999; 75(1):3–17. Epub 1999/08/17. PMID: 10448335.
46. Westercamp N, Agot K, Jaoko W, Bailey RC. Sexual function, satisfaction and penile trauma 24 months after circumcision among young men in Nyanza Province, Kenya International AIDS Conference; Washington, D.C., USA 2012.
47. Layer EH, Beckham SW, Momburi RB, Kennedy CE. Understanding the partial protection of male circumcision for HIV prevention among women in Iringa Region, Tanzania: An ethnomedical model. *AIDS care*. 2012. Epub 2012/12/12. <https://doi.org/10.1080/09540121.2012.748874> PMID: 23216512.