

Corrigendum

Clin Pediatr Endocrinol 2017; 26(3): 153–164. doi: 10.1297/cpe.26.153 PMID: PMC5537211

Original Article

Validation of auxological reference values for Japanese children with Noonan syndrome and comparison with growth in children with Turner syndrome

Tsuyoshi Isojima¹, Satoru Sakazume², Tomonobu Hasegawa³, Tsutomu Ogata⁴, Toshio Nakanishi⁵, Toshiro Nagai², and Susumu Yokoya⁶

¹ Department of Pediatrics, Teikyo University School of Medicine, Tokyo, Japan

² Department of Pediatrics, Dokkyo Medical University Koshigaya Hospital, Koshigaya, Japan

³ Department of Pediatrics, Keio University School of Medicine, Tokyo, Japan

⁴ Department of Pediatrics, Hamamatsu University School of Medicine, Hamamatsu, Japan

⁵ Department of Pediatric Cardiology, The Heart Institute, Tokyo Women's Medical University, Tokyo, Japan

⁶ Department of Medical Subspecialties, National Center for Child Health and Development, Tokyo, Japan

Received: March 10, 2017/ Accepted: March 31, 2017

Corresponding author: Susumu Yokoya, M.D., Ph.D., Department of Medical Subspecialties,

National Center for Child Health and Development, 2-10-1 Okura, Setagaya-ku, Tokyo 157-8535, Japan

E-mail: yokoya-s@fmu.ac.jp

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/4.0/>>.

The authors and publisher would like to apologize for the mistake in the following equation.

In p. 155, in lines 20–24 of the left column:

The SD score (Z-score) of each measurement (y-value) could be calculated from the L, M, and S curves using the following equation:

$$Z = \frac{[(y/M)^{L-1}] / (L \times S)}{S}, \text{ and if } L = 0, Z = \ln (y/M)/S.$$

should have been

The SD score (Z-score) of each measurement (y-value) could be calculated from the L, M, and S curves using the following equation:

$$Z = \frac{[(y/M)^{L-1}] / (L \times S)}{S}, \text{ and if } L = 0, Z = \ln (y/M)/S.$$