

RESEARCH ARTICLE

# Inducible lung epithelial resistance requires multisource reactive oxygen species generation to protect against bacterial infections

Hayden H. Ware<sup>1</sup>, Vikram V. Kulkarni<sup>1,2</sup>, Yongxing Wang<sup>1</sup>, Jezreel Pantaleón García<sup>1</sup>, Miguel Leiva Juarez<sup>1</sup>, Carson T. Kirkpatrick<sup>1</sup>, Shradha Wali<sup>1,2</sup>, Sarah Syed<sup>1</sup>, Alexander D. Kontoyiannis<sup>1</sup>, William K. A. Sikkema<sup>3</sup>, James M. Tour<sup>3</sup>, Scott E. Evans<sup>1,2\*</sup>

**1** Department of Pulmonary Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas, United States of America, **2** MD Anderson Cancer Center UT Health Graduate School of Biomedical Sciences, Houston, Texas, United States of America, **3** Smalley Institute for Nanoscale Science and Technology, Rice University, Houston, Texas, United States of America

These authors contributed equally to this work.

\* [seevans@mdanderson.org](mailto:seevans@mdanderson.org)



**OPEN ACCESS**

**Citation:** Ware HH, Kulkarni VV, Wang Y, Pantaleón García J, Leiva Juarez M, Kirkpatrick CT, et al. (2019) Inducible lung epithelial resistance requires multisource reactive oxygen species generation to protect against bacterial infections. *PLoS ONE* 14(2): e0208216. <https://doi.org/10.1371/journal.pone.0208216>

**Editor:** Samithamby Jeyaseelan, Louisiana State University, UNITED STATES

**Received:** November 12, 2018

**Accepted:** February 1, 2019

**Published:** February 22, 2019

**Copyright:** © 2019 Ware et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

**Funding:** This study was supported by US National Institutes of Health (NIH) grants R01 HL117976 and DP2 HL123229 to S.E.E. and P30 CA016672 to the MD Anderson Cancer Center. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract

Pneumonia remains a global health threat, in part due to expanding categories of susceptible individuals and increasing prevalence of antibiotic resistant pathogens. However, therapeutic stimulation of the lungs' mucosal defenses by inhaled exposure to a synergistic combination of Toll-like receptor (TLR) agonists known as Pam2-ODN promotes mouse survival of pneumonia caused by a wide array of pathogens. This inducible resistance to pneumonia relies on intact lung epithelial TLR signaling, and inducible protection against viral pathogens has recently been shown to require increased production of epithelial reactive oxygen species (ROS) from multiple epithelial ROS generators. To determine whether similar mechanisms contribute to inducible antibacterial responses, the current work investigates the role of ROS in therapeutically-stimulated protection against *Pseudomonas aeruginosa* challenges. Inhaled Pam2-ODN treatment one day before infection prevented hemorrhagic lung cytotoxicity and mouse death in a manner that correlated with reduction in bacterial burden. The bacterial killing effect of Pam2-ODN was recapitulated in isolated mouse and human lung epithelial cells, and the protection correlated with inducible epithelial generation of ROS. Scavenging or targeted blockade of ROS production from either dual oxidase or mitochondrial sources resulted in near complete loss of Pam2-ODN-induced bacterial killing, whereas deficiency of induced antimicrobial peptides had little effect. These findings support a central role for multisource epithelial ROS in inducible resistance against a bacterial pathogen and provide mechanistic insights into means to protect vulnerable patients against lethal infections.

**Competing interests:** The authors of this manuscript have read the journal's policy and have the following competing interests: S.E.E. is an author on U.S. patent 8,883,174, "Stimulation of Innate Resistance of the Lungs to Infection with Synthetic Ligands," and owns stock in Pulmotect, Inc., which holds the commercial options on these patent disclosures. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials. All other authors have declared that no competing interests exist.

## Introduction

Lower respiratory tract infections remain the leading cause of premature death and disability among both otherwise healthy and immunosuppressed people worldwide [1–5]. In an era of increasing antimicrobial resistance, human global hypermobility, proliferation of emerging and weaponized pathogens, aging populations, and ever-expanding categories of immunocompromised patients, the acute complications of pneumonia exact a staggering toll, killing an estimated 2.7 million people per year [6–10]. The 1943 introduction of penicillin for pneumonia management was a medical triumph [11], but the intervening decades have witnessed escalating age-adjusted pneumonia hospitalization rates [12–14] without survival rate improvements of corresponding magnitude [15]. In an effort to address the persisting threat of pneumonia to vulnerable populations, our laboratory has developed a program focused on manipulating the intrinsic antimicrobial capacity of the host to prevent pneumonia in susceptible populations in order to avert these catastrophic outcomes.

Based on this program, we have reported that the mucosal defenses of the lungs can be stimulated to protect mice against a wide array of otherwise lethal pneumonias, including those caused by antibiotic-resistant bacteria [16–19]. This *inducible resistance* is achieved following a single inhaled treatment comprised of a synergistic combination of Toll-like receptor (TLR) agonists: a diacylated lipopeptide ligand for TLR2/6, Pam2CSK4, and a class C unmethylated 2' = -deoxyribocytidine-phosphate-guanosine (CpG) ligand for TLR9, ODN M362, that are delivered concurrently in the same aerosolized suspension (hereafter, Pam2-ODN) [16, 17, 20].


Inducible resistance against pneumonia requires intact lung epithelial TLR signaling mechanisms, whereas no individual leukocyte populations have been identified as essential to Pam2-ODN-enhanced pneumonia survival [16, 21]. Given the epithelial requirement for inducible resistance in vivo [16, 22], we sought to determine whether epithelial cells were sufficient to act as autonomous antibacterial effector cells of therapeutically inducible protection. We recently reported that Pam2-ODN-induced antiviral protection requires therapeutic induction of reactive oxygen species (ROS) via a novel multisource mechanism [23], but it is unknown whether similar processes are required for inducible antibacterial defense.

We report here that Pam2-ODN induces active antibacterial responses from intact lungs and isolated lung epithelial cells that reduce pathogen burden, attenuate infectivity, and enhance survival. Moreover, we find that the protection requires epithelial generation of ROS via dual mechanisms, providing meaningful insights into the mechanisms underlying the novel synergistic interactions observed between the TLR ligands.

## Results

### Pam2-ODN treatment reduces pathogen burden and inflammatory injury in bacterial pneumonia

We have previously reported that a single nebulized treatment with Pam2-ODN results in improved survival of otherwise lethal pneumonias, including those caused by *P. aeruginosa* [16, 17, 20, 21]. Here, we found that the protection afforded by Pam2-ODN (Fig 1A) is associated with reduced bacterial burden immediately after infection, whether assessing burden by serial dilution culture of whole lung homogenates or culture of bronchoalveolar lavage (BAL) fluid (Fig 1B), suggesting that a Pam2-ODN-induced antimicrobial environment existed at the time of infection. No significant differences were noted in the performance of the two culture methods, in terms of precision or magnitude of induced bacterial reductions by Pam2-ODN, though the absolute bacteria per ml tended to be higher in the BAL-obtained samples than in



**Fig 1. Pam2-ODN protects against bacterial pneumonia.** (A) Survival of wildtype mice treated with Pam2-ODN or PBS (sham) by aerosol 24 h before challenge with *P. aeruginosa*. (B) Pathogen burden of mice in A immediately after challenge, as assessed by serial dilution culture of lung homogenates (left panel) or BAL fluid (right panel). (C) Gross appearance of mouse lungs 24 h after *P. aeruginosa* challenge following treatment with Pam2-ODN or sham. (D) Hematoxylin-eosin stained histology of lungs in C. Scale bar = 400  $\mu$ m left panels, 100  $\mu$ m right panels. Each panels is representative of at least three independent experiments. N = 8 mice/group for survival, N = 4 mice/group for pathogen burden. \* p < 0.0002 vs. PBS-treated; \*\* p < 0.002 vs PBS-treated.

<https://doi.org/10.1371/journal.pone.0208216.g001>

the lung homogenates. These findings are congruent with the inducible reductions in pathogen burden that were observed by quantification of microscopic lung fluorescence both immediately after infection and 24 h after infection when mice were challenged with GFP-expressing *P. aeruginosa* (S1 Fig). Interestingly, treatment-related fluorescence intensity differences immediately after infection appeared to be principally related to modulation of bacterial burden in the airways, whereas more substantial Pam2-ODN-induced fluorescence differences were observed in the lung parenchyma 24 h after infection. This observation likely reflects initial aerosolized pathogen deposition (and Pam2-ODN-induced killing) in the airways, followed by infectious progression into the more peripheral lung over the subsequent hours to days. The observation of disproportionate early airway deposition may also explain the above noted finding that the bacteria per ml of BAL fluid are higher than the bacteria per ml of lung homogenate immediately after infection, as shown in Fig 1B.

Although inhaled treatment with Pam2-ODN induces transient lung neutrophilia [16], we found here that the antimicrobial environment associated with Pam2-ODN-induced resistance also protected against inflammatory lung injury. Lungs harvested 24 h after *P. aeruginosa* challenge demonstrate severe hemorrhagic pneumonia in sham-treated mice, but there is

almost no evidence of such injury in Pam2-ODN-treated mice (Fig 1C). Similarly, histologic inspection of Pam2-ODN-treated lungs 24 h after infection demonstrate substantially less inflammatory cell infiltration and notably fewer bacteria. This is congruent with earlier studies [16] suggesting that the difference in *P. aeruginosa* continues to increase between active and sham treated groups as time elapses, indicating that the antimicrobial environment persists beyond the period of the initial challenge.

### **Pam2-ODN induces bacterial killing by isolated lung epithelial cells across a broad concentration and temporal range**

Congruent with the *in vivo* observations, we have found that treatment of isolated human or mouse lung epithelial cells results in significant reductions in culture bacterial burdens [16, 17, 21, 22]. Based on empiric *in vivo* efficacy optimization, Pam2-ODN is administered in a fixed 4:1 molar ratio [18, 20]. Fig 2A and 2B shows that, when delivered in this ratio, the antibacterial effect is inducible across treatment concentrations that extend to at least a 2 log<sub>10</sub> range. Higher Pam2-ODN treatment concentrations are expected to induce even greater bacterial killing than that shown, but when calculating the estimated deposition of the ligands in 20 μl mouse airway lining fluid [24] or in 10–30 ml of human airway lining fluid [25] after nebulization, it is unlikely that such high concentrations can be achieved *in vivo*. To avoid presenting responses that are easily detectable but not physiologically relevant to the *in vivo* model, all subsequent figures include data achieved with a lower Pam2-ODN dose (2.23 μM Pam2 and 0.56 μM ODN) that we calculate to be achievable by nebulization, except for panels explicitly labelled as dose response plots.


In this model, bacteria are inoculated into the epithelial cultures in log phase growth and there are no antibacterial leukocyte contributions. So, the antimicrobial epithelial responses must be active very early in the course of infection. To determine how quickly these responses could be induced, we tested treatment intervals prior to challenge and found that the most profound antibacterial responses seemed to be achieved with six or more hours of treatment, but significant bacterial burden reductions were achieved in a much shorter period (Fig 2C and 2D) in both mouse and human cells. In fact, antibacterial effects were even observed when Pam2-ODN was administered simultaneous to or after the infectious challenge (Fig 2E and 2F), though shorter treatment periods were associated with lesser antibacterial effects.

### **Pam2-ODN interact synergistically to induce bacterial killing**

Further substantiating the *in vitro* model as relevant to study of the *in vivo* pneumonia protection associated with Pam2-ODN treatment, we found that the antibacterial effect of the combined Pam2-ODN treatment was supra-additive to the effects of equimolar ligands delivered individually. Pam2 alone induced a modest reduction in bacterial burdens in human epithelial cells (Fig 2G). The magnitude of this effect is similar to the degree of protection we have observed *in vivo* with Pam2 alone [17]. ODN alone did not induce any significant reductions in bacterial burden in either human or mouse epithelial cultures (Fig 2H). However, in both models, the combination of Pam2 and ODN resulted in greater anti-pseudomonal effects than the combined effects of the two ligands delivered alone.

### **Pam2-ODN extends epithelial survival of Pseudomonas infections**

While Pam2-ODN induces a robust antibacterial effect, it has not been previously established whether the antimicrobial response was associated with a fitness cost to the cells themselves. For instance, it is conceivable that the microbicidal response might also be toxic to the host cells or it is possible that programmed cell death pathways contribute to bacteriostatic effects.



**Fig 2. Pam2-ODN induces antibacterial responses in isolated lung epithelial cells.** HBEC3kt (A) or MLE15 (B) cells were treated for 4 h with PBS or the indicated doses of Pam2-ODN, then challenged with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. HBEC3kt (C, E) or MLE15 (D, F) cells were treated with PBS or Pam2-ODN (middle dose used in A and B, 2.23  $\mu$ M Pam2 and 0.56  $\mu$ M ODN) for the indicated interval relative to challenge with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. HBEC3kt (G) or MLE15 (H) cells were treated for 4 h with the indicated treatments, then challenged with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. HBEC3kt (I) or MLE15 (J) cells were treated with PBS or Pam2-ODN for 6 h prior to *P. aeruginosa* challenge. Cell survival determined by Trypan blue exclusion is shown at the indicated time points. Each panel is representative of at least three independent experiments. \*  $p < 0.05$  vs. PBS-treated; \*\*  $p < 0.005$  vs. PBS-treated; †  $p < 0.05$  vs. either single ligand treatment.

<https://doi.org/10.1371/journal.pone.0208216.g002>


Indeed, we have previously reported that inducible epithelial resistance is correlated with transient but profound induction of inflammatory mediators [16, 17, 20], and here find significant induction of genes for both proinflammatory cytokines and antimicrobial peptides from lung epithelial cells treated with Pam2-ODN (S2 Fig). However, we have not found reduced survival of lung epithelial cells following Pam2-ODN treatment in the absence of infection and see dramatically improved cell survival of viral infections when the cells received Pam2-ODN pretreatment [23]. To investigate the effect of Pam2-ODN treatment on epithelial cell survival of bacterial infections, Trypan blue exclusion was used to determine cell viability following *P. aeruginosa* infection. While all epithelial cells were dead by 36 h after the infectious challenge, regardless of pretreatment, both mouse and human epithelial cells lived longer on average and had a greater percentage of cells alive at every intermediate time point if pretreated with Pam2-ODN (Fig 2I and 2J). These findings support that the antibacterial effect of the epithelial response to Pam2-ODN is more beneficial than any potential fitness cost.

### Pam2-ODN-induced antibacterial effects require DUOX-dependent ROS production

Antimicrobial peptides are well established contributors to lung epithelium-mediated antibacterial defense [22], and genes encoding antimicrobial peptides such as lipocalin 2 and acute phase serum albumin A proteins are some of the most strongly upregulated transcripts following Pam2-ODN treatments of lungs or isolated lung epithelial cells [16, 17]. However, *P. aeruginosa* challenge of mice deficient in these antibacterial molecules revealed little defect in Pam2-ODN-inducible protection, even when more than one antimicrobial peptide gene was knocked out (S2 Fig). These data suggest that, although they are robustly induced, these individual antimicrobial peptides are not essential effectors of inducible resistance, and prompted investigations into alternate effector mechanisms.

ROS are increasingly recognized to function as direct antimicrobial effector molecules, most likely through lipid peroxidation of microbial membranes and DNA damage, in addition to their well-established roles as signaling molecules [26]. We previously hypothesized that ROS contribute to Pam2-ODN-induced epithelial antibacterial effects [16, 18, 19]. More recently, we confirmed that ROS are essential to Pam2-ODN-induced antiviral responses and have published a comprehensive characterization of the epithelial ROS species induced by Pam2-ODN treatment [23]. Fig 3A and 3B confirm that Pam2-ODN induces dose-dependent production of ROS from both human and mouse lung epithelial cells, as measured by fluorescence signal from cell permeant carboxy- $H_2$ DCFDA. Our previous findings indicate that only hydrogen peroxide and superoxide are demonstrably increased by epithelial treatment [23], and there is no reason to suspect that induction of alternate species is reflected by the carboxy- $H_2$ DCFDA signal in the current studies using identical culture and treatment models.

Acting by superoxide dismutation and radical annihilation [27, 28], application of poly(ethylene glycolated) hydrophilic carbon clusters (PEG-HCCs) [28, 29] to the culture media



**Fig 3. Pam2-ODN induces antibacterial ROS from isolated lung epithelial cells.** HBEC3kt (A) or MLE15 (B) cells were exposed to CO-H<sub>2</sub>DCFDA, treated with the indicated doses of Pam2-ODN, then fluorescence intensity was measured every 5 min. HBEC3kt (C) or MLE15 (D) cells were pretreated with PEG-HCC or PBS, exposed to CO-H<sub>2</sub>DCFDA, then treated with the indicated dose of Pam2-ODN. Shown are fluorescence intensity 100 min after treatment. HBEC3kt (E) or MLE15 (F) cells were pretreated with PEG-HCC or PBS, treated for 4 h with PBS or Pam2-ODN (Pam2 2.23 μM, ODN 0.56 μM), then challenged with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. (G) HBEC3kt cells were stably transfected with scrambled (control) shRNA or shRNA targeting *DUOX1* or *DUOX2*, then treated with PBS or Pam2-ODN for 6 h prior to *P. aeruginosa* challenge. Shown are culture bacterial burdens 4 h after challenge. Each panels is representative of at least three independent experiments. \*

p < 0.005 vs no Pam2-ODN treatment; † p < 0.005 vs no PEG-HCC, same Pam2-ODN; †† p < 0.02 vs scrambled shRNA + Pam2-ODN; ‡ p < 0.003 vs *DUOX1* knockdown + Pam2-ODN.

<https://doi.org/10.1371/journal.pone.0208216.g003>

significantly reduced epithelial ROS, as demonstrated by CO-H<sub>2</sub>DCFDA fluorescence, at all Pam2-ODN doses (Fig 3C and 3D). Notably, by reducing epithelial ROS, PEG-HCC treatment also significantly impaired the Pam2-ODN-induced epithelial antibacterial effect (Fig 3E and 3F), supporting a ROS requirement for inducing the protective response from both human and mouse lung epithelial cells.

While all NADPH oxidase (NOX) isoforms are reported to be expressed by lung epithelia, the primary source of lung epithelial ROS are the dual oxidases DUOX1 and DUOX2 (sometimes called NOX6 and NOX7) [30–32]. To test the specific requirement for DUOX-derived ROS in Pam2-ODN-induced antibacterial defense, we used shRNA to stably knockdown *DUOX1* and *DUOX2* in HBEC3kt cells, then assessed the effect on Pam2-ODN-induced reductions in bacterial burden in the cells with the highest knockdown efficiency (annotated as *DUOX1*-1 and *DUOX2*-4 in S3 Fig). Knocking down the DUOX genes has no effect on short term cell survival [23]. However, Fig 3G shows that knocking down *DUOX1* moderately impairs the Pam2-ODN-induced epithelial antimicrobial response and knocking down *DUOX2* severely impairs the inducible antibacterial effect. This is congruent with prior reports that *DUOX1* produces a relatively consistent amount of ROS, though this production can be moderately enhanced by IL-4 and IL-13 exposure [33], whereas *DUOX2*-dependent ROS production can be profoundly increased by activating existing *DUOX2* and increasing *DUOX2* and *DUOXA2* transcription following exposure to cytokines such as IFN $\gamma$  [33]. Interestingly, while the *DUOX1* requirement for inducible antipseudomonal defense appears to be less substantial than the *DUOX2* requirement, the inducible protection defect observed in *DUOX1* knockdown cells is more profound than that observed in virus challenged *DUOX1* knockdown cells [23].


### Pam2-ODN-induced antibacterial effects require mitochondrial ROS production

Although we confirmed that DUOX-dependent ROS production is required for the inducible bacterial killing, there is accumulating evidence that mitochondria-derived ROS (mtROS) can also participate in antimicrobial responses of nonphagocytes [23, 34–37]. To test whether mtROS also contribute to the inducible antibacterial effect of Pam2-ODN, mtROS were selectively scavenged with mitoTEMPO prior to *P. aeruginosa* challenge with or without Pam2-ODN pretreatment. Fig 4A and 4B shows that mtROS scavenging profoundly impaired the Pam2-ODN induced bacterial killing by mouse and human epithelial cells. To address potential off-target effects or nonspecific ROS scavenging of mitoTEMPO, we tested whether we could reduce inducible mtROS production, rather than scavenging produced ROS. Fig 4C shows that combination treatment with a mitochondrial complex II inhibitor and a respiratory chain uncoupler reduces mtROS production at every tested dose of Pam2-ODN. This impaired Pam2-ODN-induced mtROS production resulted in bacterial killing defects that mirrored the mtROS scavenging experiments (Fig 4D and 4E), revealing a requirement for mtROS in Pam2-ODN-induced antibacterial responses.

### Discussion

Accumulating evidence supports the essential contributions of airway and alveolar epithelia to antimicrobial defense. In addition to their capacity to recruit and activate leukocyte-mediated





**Fig 4. Mitochondrial ROS are required for Pam2-ODN-induced antibacterial epithelial responses.** HBEC3kt (A) or MLE15 (B) cells were pretreated with MitoTEMPO or PBS, treated for 4 h with PBS or Pam2-ODN (Pam2 2.23 uM,

ODN 0.56  $\mu$ M), then challenged with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. (C) HBEC3kt cells were pretreated with FCCP-TTFA or PBS, exposed to MitoSOX, then treated with PBS or Pam2-ODN at the indicated doses. Shown are culture fluorescence intensities at 100 min after treatment. HBEC3kt (D) or MLE15 (E) cells were pretreated with FCCP-TTFA or PBS, treated for 4 h with PBS or Pam2-ODN (Pam2 2.23  $\mu$ M, ODN 0.56  $\mu$ M), then challenged with *P. aeruginosa*. Shown are culture bacterial burdens 4 h after challenge. Each panel is representative of at least three independent experiments. N = 4–5 samples/condition for all experiments. \*  $p < 0.01$  vs. PBS-treated without inhibitor/scavenger; †  $p < 0.02$  vs. Pam2-ODN-treated without inhibitor/scavenger.

<https://doi.org/10.1371/journal.pone.0208216.g004>

defenses, epithelial cells exert directly antimicrobial effects on invading pathogens [22]. Indeed, we have found that lung epithelial cells function as primary effector cells of Pam2-ODN-induced resistance to pneumonia. [16, 17, 21, 23]. The current work demonstrates that a preventative Pam2-ODN treatment promotes mouse survival of bacterial challenges by reducing pathogen burden and attenuating associated immunopathology. Similar in vivo Pam2-ODN-induced reductions in pathogen burden are demonstrated by multiple quantification techniques, suggesting that the induced antimicrobial environment eliminates bacteria from all anatomic and cellular compartments of the lungs. Similar pathogen killing is observed from Pam2-ODN-treated isolated lung epithelial cells, where the reduced pathogen burden enhances cellular survival, even in the absence of leukocytes.

Although numerous antimicrobial peptides are induced by Pam2-ODN treatment, no individual peptides have been demonstrated to be required for the enhanced mouse survival of infectious challenges nor for the intrapulmonary pathogen killing. While it was, perhaps, unanticipated that none of these highly enriched peptides would prove essential to the protection, there are a number of plausible explanations for this observation. Foremost among these is the possibility that the extreme multiplicity of induced antimicrobial molecules renders the loss of only one or two peptides largely irrelevant. An alternate consideration is that different peptide combinations may be required to protect against different challenges.

Regardless of explanation, the inability to detect any essential antimicrobial peptides underscores the robustness of the Pam2-ODN-induced protection against pneumonias and emphasizes the importance of our finding that epithelial ROS are required for inducible antiviral defense. We have established that inducible ROS are essential to protecting against challenges by orthomyxoviruses and paramyxoviruses [23]. The present study similarly finds that inducible ROS production is essential to Pam2-ODN-induced epithelial bacterial killing, notably requiring ROS from both DUOX and mitochondrial sources.

ROS are recognized to contribute to antibacterial defenses, particularly in the context of NADPH oxidase-dependent killing of bacteria in phagolysosomes of myeloid cells [26]. However, the broadly microbicidal capacity of ROS generated by a wide range of cells has been demonstrated in recent years against Gram-negative, Gram-positive, viral and fungal pathogens, even in the setting of biofilms or antibiotic resistance [38–40]. This vigorous protection may rely in part on the capacity of ROS to synergize with other antimicrobial treatments [38, 41]. Similarly, ROS can reduce the minimum inhibitory concentrations of host effector molecules, such as neutrophil proteases [26]. It has been further proposed that eliciting ROS production is an essential component of the pathogen killing mechanisms of some antimicrobial pharmaceutical agents [42, 43].

Superoxide and hydrogen peroxide are the predominant species produced by lung epithelial cells [26, 34, 44]. Our recent comprehensive assessment of ROS production following Pam2-ODN treatment revealed superoxide and hydrogen peroxide to be the only species detectably induced by Pam2-ODN [23].

The principal sources of lung epithelial hydrogen peroxide are DUOX1 and DUOX2 [30, 31, 45, 46]. Therapeutic induction of DUOX2 has been proposed as an antimicrobial therapeutic

strategy [47], and we have reported that DUOX-related genes are enriched following epithelial exposure to Pam2-ODN [17, 23]. This study confirms that DUOX-dependent ROS are required for inducible *P. aeruginosa* killing, with the protective effect profoundly attenuated when cellular ROS are annihilated by PEG-HCC or either DUOX is selectively knocked down. It is well established that the DUOX-dependent product of the lactoperoxidase/hydrogen peroxide/thiocyanate system, hypothiocyanate, exerts antimicrobial effects [45, 46, 48], however it is unlikely that the ROS dependency observed here reflects hydrogen peroxide-mediated hypothiocyanate production, as our in vivo models lack tracheobronchial seromucus glands as a lactoperoxidase source [49] and our in vitro models lack a source of thiocyanate [33]. This suggested that the DUOX-dependent ROS effects are likely achieved either through direct pathogen toxicity or through host signaling events.

mtROS have also been increasingly reported to contribute to both innate and adaptive immunity [34, 50], and increased production of mitochondria-derived species likely explains the Pam2-ODN-increased superoxide we detect [23].

mtROS are generated via electron transport chain leakage [44], resulting in superoxide that diffuses through mitochondrial membranes following dismutation to hydrogen peroxide [51]. This process is exquisitely tightly regulated by scavenging, production, and localization [34], so the substantial induction of mtROS by Pam2-ODN represents an important homeostatic perturbation. It has been reported that TLR manipulation can promote generation of both antibacterial mtROS [37] and NADPH oxidase-generated ROS [52] in macrophages, so it is plausible that TLR ligands could induce ROS from both mitochondrial and DUOX sources in epithelial cells.

The precise mechanisms underlying the requirement for dual sources of ROS have yet to be elucidated. However, this may be explained by generation of ROS-induced ROS that yield high ROS concentrations [51, 53, 54]. Alternate explanations include the hypothesis that multiple sources are required to achieve sufficiently high aggregate ROS concentrations to exert microbicidal actions or that the different sources play different signaling and pathogen killing roles. It is even possible that the mtROS regulate DUOX functions [55]. This remains an area of active investigation.

These data indicate that multisource ROS are required for Pam2-ODN-induced bacterial killing, extending the range of pathogens that are known to be susceptible to inducible epithelial ROS and highlighting the centrality of ROS generation to the protective phenomenon of inducible epithelial resistance. By advancing understanding of the mechanisms of Pam2-ODN-induced resistance, these data may facilitate development of even more efficacious resistance-inducing therapeutics and offer hope that pneumonia can be prevented in vulnerable populations.

## Materials and methods

### Animals, cells and reagents

All general reagents were obtained from Sigma-Aldrich (St Louis, MO), except as indicated. All mouse experiments were performed with 5–8 week-old C57BL/6J (The Jackson Laboratory, Bar Harbor, ME) of a single gender in accordance with the Institutional Animal Care and Use Committee of The University of Texas MD Anderson Cancer Center, protocol 00000907-RN01. Immortalized human bronchial epithelial (HBEC3kt) cells were kindly provided by Dr. John Minna. Murine lung epithelial (MLE-15) cells were kindly provided by Dr. Jeffrey Whitsett. The cell lines used were authenticated by the MD Anderson Characterized Cell Line Core Facility.

## Cell culture

HBEC3kt cells were cultured in keratinocyte serum-free media (KSFM, ThermoFisher Scientific, Grand Island, NY) supplemented with human epidermal growth factor and bovine pituitary extract. MLE-15 cells were cultured in RPMI supplemented with 10% fetal bovine serum. Cultures were maintained in the presence of penicillin and streptomycin.

## TLR treatments

For *in vivo* studies, S-[2,3-bis(palmitoyloxy)-propyl]-(R)-cysteinyl-(lysyl) 3-lysine (Pam2CSK<sub>4</sub>) and ODN M362 (InvivoGen, San Diego, CA) were reconstituted in endotoxin-free water, then diluted to the desired concentration in sterile PBS. As previously described [16], the Pam2-ODN was placed in an Aerotech II nebulizer (Biodex Medical Systems, Shirley, NY) driven by 10 l min<sup>-1</sup> air supplemented with 5% CO<sub>2</sub> for 20 min. The nebulizer was connected by polyethylene tubing to a polyethylene exposure chamber. 24 h prior to infections, 8 ml of Pam2 (4 μM) -ODN (1 μM) was delivered via nebulization to unrestrained mice for 20 minutes, and then mice were returned to normal housing. For *in vitro* studies, Pam2-ODN was added to the culture media 4 h prior to inoculation with bacteria or at the indicated time point. Pam2-ODN was given in fixed ratio, but at varying doses as indicated. For both treatments and infectious challenges, the mice are allowed to move freely and without restraint in the exposure chamber. No signs of distress have ever been observed during any of the aerosolized exposures. Aside from those that are immediately sacrificed (described below), on completion of the aerosolized exposures, mice are immediately returned to their housing cages where they have *ad lib* access to food and water.

## Infection models

As previously described [16], frozen stock of *Pseudomonas aeruginosa* strain PA103 (American Type Culture Collection, Manassas, VA) was incubated overnight in tryptic soy broth, then expanded in Luria-Bertini media to OD<sub>600</sub> 0.35. Bacterial suspensions were centrifuged, washed, re-suspended in PBS, and aerosolized over 60 min. For all bacterial challenges, a nebulized inoculum of 10 ml of ~2 × 10<sup>10</sup> CFU/ml were delivered. Immediately after bacterial challenges, some mice were anesthetized and their lungs were harvested and homogenized [16] using a Mini-Beadbeater-1 (Biospec, Bartlesville, OK). Serial dilutions of the nebulizer inoculum and lung homogenates were plated on tryptic soy agar plates (Becton Dickinson). The remaining mice were observed for 12 d to determine whether their clinical conditions met euthanasia criteria. Following infection, lab personnel coordinated with staff of the MD Anderson Department of Veterinary Medicine to ensure that the mice were evaluated a minimum of three times daily to determine whether euthanasia criteria were met. As specified in the above noted animal protocol, mice that were not submitted to anesthetic excess followed by thoracotomy with bilateral pneumonectomy for pathogen burden assessments were humanely sacrificed by inhalational exposure to approved concentrations of carbon dioxide until respiratory efforts ceased, followed by cervical dislocation as a secondary method of euthanasia, when they either achieved the end of the observation period or met the predesignated euthanasia criteria. The relevant euthanasia-triggering criteria include any evidence hypothermia, impaired mobility, respiratory distress, inability to access food or water, or any evidence of distressed behavior. Weight loss is also among the approved indications for euthanasia, but the mice that met euthanasia criteria in this model became ill or distressed within 1–2 d (before losing > 25% body weight), so no mice were euthanized due to weight loss in the current study. Despite the close observation, this same rapidity of illness resulted in up to 3 of the 56 infected mice dying spontaneously from pneumonia before being euthanized in some experimental replicates. Although meeting euthanasia criteria is the primary endpoint, the

presented “Survival (%)” in Fig 1 formally indicates mice that had not either met euthanasia criteria or spontaneously died. When mice were identified to meet criteria, they were submitted to euthanasia within 30 minutes by either lab personnel or Department of Veterinary Medicine staff. All lab personnel and Department of Veterinary Medicine staff receive formal instruction in methods to minimize stress and discomfort to experimental animals and analgesia is provided to animals that demonstrate any evidence of discomfort but do not meet euthanasia criteria. Moistened chow and gel packs are provided on the floor of cages to mice for whom there is any concern of inability to access food or water.

For the *in vitro* challenges, after the indicated treatments, confluent mouse or human epithelial cell cultures were inoculated with *P. aeruginosa* ( $20 \mu\text{l} \times 1 \times 10^5$  CFUs/ml), incubated for 6 h, then harvested and submitted to serial dilution culture.

### Lentiviral shRNA knockdown of *DUOX1* and *DUOX2*

GIPZ human *DUOX1* and *DUOX2* lentiviral shRNA clones were purchased from GE Dharmacon (Lafayette, CO). Lentiviruses bearing human *DUOX1* and *DUOX2* shRNA were produced by transfection in 293T cells per manufacturer’s instructions. Infection efficiency was enhanced by addition of 8  $\mu\text{g}/\text{ml}$  Polybrene into the culture media and centrifuging the cells at 2,000 rpm for 60 min at 32°C. Lentiviral-infected HBEC3kt cells were selected by cell sorting based on GFP expression. shRNA knockdown efficiency was determined by immunoblot analysis, as previously shown [23].

### ROS detection, scavenging and inhibition

To assess ROS generation, cells were treated with 5  $\mu\text{M}$  of each indicated detector for 1 h before exposure to Pam2-ODN or sham, as previously reported [23]. Fluorescence was continuously measured on a BioTek Synergy2 for 250 min after treatment. Excitation/emission wavelengths for ROS-detecting agents are: Carboxy-2',7'-dichlorodihydrofluorescein diacetate (CO-H<sub>2</sub>DCFDA, ThermoFisher), 490nm/525nm; and MitoSOX<sup>TM</sup> Red (ThermoFisher), 510nm/580nm.

Cellular ROS were scavenged by 1 h exposure to PEGylated hydrophilic carbon clusters (PEG-HCC, 5  $\mu\text{g}/\text{mL}$ ) prior to application of Pam2-ODN or PBS [23]. Mitochondrial ROS were scavenged by 1 h exposure to (2-(2,2,6,6-Tetramethylpiperidin-1-oxyl-4-ylamino)-2-oxoethyl) triphenylphosphonium chloride monohydrate (MitoTEMPO, 30nM, ThermoFisher) prior to treatment with Pam2-ODN or PBS [23]. Disruption of *in vitro* mitochondrial ROS production was achieved through concurrent application of trifluoromethoxy carbonyl-cyanide phenylhydrazone (FCCP, 400 nM, Cayman Chemical, Ann Arbor, MI), and 2-theyl-trifluoroacetone (TTFA, 200  $\mu\text{M}$ , Sigma) [23].

### Statistical analysis

Statistical analysis was performed using SPSS v19 (SAS Institute, Cary, NC). Student’s t-test was used to compare the lung pathogen burdens between the groups. Error bars shown in all the figures represent technical replicates within the displayed experiment, rather than aggregation of experimental replicates. Percentage of mice surviving pathogen challenges was compared using Fisher’s exact test on the final day of observation, and the log-rank test was used to compare the survival distribution estimated by the Kaplan–Meier method.

### Supporting information

**S1 Fig. Early and late bacterial burden reduction by Pam2-ODN treatment.** Wild type C57BL/6J mice were treated with Pam2-ODN or PBS (sham) 24 h prior to challenge with

GFP-expressing *P. aeruginosa*. Shown are representative micrographs of 20  $\mu\text{m}$  OCT-embedded lung sections immediately after infection (A) or 24 h after infection (B). Blue, DAPI; green, bacterial GFP; white dashed line, airway delineation; white scale bar, 200  $\mu\text{m}$ . GFP signal in the indicated conditions was then quantified. (C) Mean fluorescence intensity per 20x field. (D) Mean fluorescence intensity of airways contained within the fields examined in C. (E) Mean fluorescence intensity of five stereotyped (174  $\mu\text{m}$  x 174  $\mu\text{m}$ ), airway-free boxes from each of the fields examined in C. N = 3 mice per condition, 10 random fields measured per mouse. \*  $p < 0.05$ , \*\*  $p < 0.001$ .

(PDF)

**S2 Fig. Cytokine and antimicrobial peptide induction in Pam2-ODN-induced resistance.**

(A) HBEC3kt cells were treated with PBS (sham) or Pam2-ODN for 2 h, then submitted to RT-qPCR for the indicated transcripts. Shown are RQ values for the target transcript relative to 18s gene. Each panel is representative of at least three independent experiments. N = 4–5 samples/condition for all experiments. Wild type or mice deficient in (B) *Lcn2*, (C) *Cramp*, (D) *Lcn2* and *Cramp*, or (E) the indicated acute phase SAA genes were treated with PBS (sham) or Pam2-ODN by aerosol 24 h prior to challenge with *P. aeruginosa*. Shown are survival plots for each challenge. Each panel is representative of at least three independent experiments. N = 8–10 mice/condition. \*  $p < 0.001$  vs PBS treated. \*\*  $p < 0.007$  vs. syngeneic PBS treated. †  $p < 0.05$  vs. syngeneic PBS treated.

(PDF)

**S3 Fig. Knockdown of dual oxidase genes.** Shown are immunoblots of HBEC3kt cells transfected with scrambled shRNA (Scr) or shRNA targeting *DUOX1* (A) or *DUOX2* (B), then probed for their respective protein products. B-actin is shown as a loading control for each sample.

(PDF)

## Acknowledgments

The authors wish to thank Dr. Tak Mak and Dr. Thorsten Berger for graciously sharing *Lcn2* knockout mice. The authors wish to thank Dr. Richard Gallo for graciously sharing *Cramp* knockout mice. The authors wish to thank Dr. Maria de Beer and Dr. Frederick de Beer for graciously sharing *Saa1/2* double knockout mice. The authors wish to thank Dr. Matthew Poynter for graciously sharing *Saa3* knockout mice.

## Author Contributions

**Conceptualization:** Hayden H. Ware, Vikram V. Kulkarni, Scott E. Evans.

**Formal analysis:** Yongxing Wang, Miguel Leiva Juarez.

**Funding acquisition:** Scott E. Evans.

**Investigation:** Hayden H. Ware, Vikram V. Kulkarni, Yongxing Wang, Jezreel Pantaleón García, Miguel Leiva Juarez, Carson T. Kirkpatrick, Shradha Wali, Sarah Syed, Alexander D. Kontoyiannis, Scott E. Evans.

**Methodology:** Yongxing Wang, Miguel Leiva Juarez, Shradha Wali, Scott E. Evans.

**Resources:** Carson T. Kirkpatrick, William K. A. Sikkema, James M. Tour, Scott E. Evans.

**Supervision:** Yongxing Wang, Scott E. Evans.

**Writing – original draft:** Scott E. Evans.

**Writing – review & editing:** Hayden H. Ware, Vikram V. Kulkarni, Yongxing Wang, Jezreel Pantaleón García, Miguel Leiva Juarez, Carson T. Kirkpatrick, Shradha Wali, William K. A. Sikkema, James M. Tour.

## References

1. WHO. The World Health Report 2004—Changing History. Geneva: World Health Organization; 2004.
2. File TM. Community-acquired pneumonia. *Lancet*. 2003; 362(9400):1991–2001. [https://doi.org/10.1016/S0140-6736\(03\)15021-0](https://doi.org/10.1016/S0140-6736(03)15021-0) PMID: 14683661.
3. Joos L, Tamm M. Breakdown of pulmonary host defense in the immunocompromised host: cancer chemotherapy. *Proc Am Thorac Soc*. 2005; 2(5):445–8. <https://doi.org/10.1513/pats.200508-097JS> PMID: 16322598.
4. Mizgerd JP. Lung infection—a public health priority. *PLoS Med*. 2006; 3(2):e76. Epub 2006/01/13. 05-PLME-ES-0632R1 [pii] <https://doi.org/10.1371/journal.pmed.0030076> PMID: 16401173.
5. Collaborators GL. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory tract infections in 195 countries: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis*. 2017; 17(11):1133–61. [https://doi.org/10.1016/S1473-3099\(17\)30396-1](https://doi.org/10.1016/S1473-3099(17)30396-1) PMID: 28843578; PubMed Central PMCID: PMC5666185.
6. James JS. Empty antibiotic pipeline critically endangers public: IDSA report. *AIDS Treat News*. 2004; (404):7. PMID: 15484367.
7. Society for Healthcare Epidemiology of A, Infectious Diseases Society of A, Pediatric Infectious Diseases S. Policy statement on antimicrobial stewardship by the Society for Healthcare Epidemiology of America (SHEA), the Infectious Diseases Society of America (IDSA), and the Pediatric Infectious Diseases Society (PIDS). *Infect Control Hosp Epidemiol*. 2012; 33(4):322–7. <https://doi.org/10.1086/665010> PMID: 22418625.
8. Torres A, Peetermans WE, Viegi G, Blasi F. Risk factors for community-acquired pneumonia in adults in Europe: a literature review. *Thorax*. 2013; 68(11):1057–65. <https://doi.org/10.1136/thoraxjnl-2013-204282> PMID: 24130229; PubMed Central PMCID: PMC3812874.
9. Wong JL, Evans SE. Bacterial Pneumonia in Patients with Cancer: Novel Risk Factors and Management. *Clinics in chest medicine*. 2017; 38(2):263–77. <https://doi.org/10.1016/j.ccm.2016.12.005> PMID: 28477638; PubMed Central PMCID: PMC5424613.
10. Vinogradova Y, Hippisley-Cox J, Coupland C. Identification of new risk factors for pneumonia: population-based case-control study. *Br J Gen Pract*. 2009; 59(567):e329–38. <https://doi.org/10.3399/bjgp09X472629> PMID: 19843413; PubMed Central PMCID: PMC2751937.
11. Blake FG, Craige B Jr. Penicillin in suppurative disease of the lungs: a report of three cases. 1943. *Yale J Biol Med*. 2000; 73(1–6):103–13. PMID: 11765929.
12. Trotter CL, Stuart JM, George R, Miller E. Increasing hospital admissions for pneumonia, England. *Emerg Infect Dis*. 2008; 14(5):727–33. <https://doi.org/10.3201/eid1405.071011> PMID: 18439353; PubMed Central PMCID: PMC2600241.
13. Fry AM, Shay DK, Holman RC, Curns AT, Anderson LJ. Trends in hospitalizations for pneumonia among persons aged 65 years or older in the United States, 1988–2002. *JAMA*. 2005; 294(21):2712–9. <https://doi.org/10.1001/jama.294.21.2712> PMID: 16333006.
14. Thomsen RW, Riis A, Norgaard M, Jacobsen J, Christensen S, McDonald CJ, et al. Rising incidence and persistently high mortality of hospitalized pneumonia: a 10-year population-based study in Denmark. *J Intern Med*. 2006; 259(4):410–7. <https://doi.org/10.1111/j.1365-2796.2006.01629.x> PMID: 16594909.
15. Blasi F, Aliberti S, Pappalettera M, Tarsia P. 100 years of respiratory medicine: pneumonia. *Respir Med*. 2007; 101(5):875–81. <https://doi.org/10.1016/j.rmed.2007.02.016> PMID: 17379493.
16. Cleaver JO, You D, Michaud DR, Pruneda FA, Juarez MM, Zhang J, et al. Lung epithelial cells are essential effectors of inducible resistance to pneumonia. *Mucosal Immunol*. 2014; 7(1):78–88. <https://doi.org/10.1038/mi.2013.26> PMID: 23632328; PubMed Central PMCID: PMC3735803.
17. Duggan JM, You D, Cleaver JO, Larson DT, Garza RJ, Guzman Pruneda FA, et al. Synergistic interactions of TLR2/6 and TLR9 induce a high level of resistance to lung infection in mice. *J Immunol*. 2011; 186(10):5916–26. Epub 2011/04/13. [jimmunol.1002122 \[pii\] https://doi.org/10.1002/jimmunol.1002122](https://doi.org/10.1002/jimmunol.1002122) PMID: 21482737.
18. Evans SE, Tuvim MJ, Fox CJ, Sachdev N, Gibiansky L, Dickey BF. Inhaled innate immune ligands to prevent pneumonia. *Br J Pharmacol*. 2011; 163(1):195–206. Epub 2011/01/22. <https://doi.org/10.1111/j.1476-5381.2011.01237.x> PMID: 21250981; PubMed Central PMCID: PMC3085878.

19. Evans SE, Xu Y, Tuvim MJ, Dickey BF. Inducible innate resistance of lung epithelium to infection. *Annu Rev Physiol.* 2010; 72:413–35. Epub 2010/02/13. <https://doi.org/10.1146/annurev-physiol-021909-135909> PMID: 20148683.
20. Alfaro VY, Goldblatt DL, Valverde GR, Munsell MF, Quinton LJ, Walker AK, et al. Safety, tolerability, and biomarkers of the treatment of mice with aerosolized Toll-like receptor ligands. *Front Pharmacol.* 2014; 5:8. <https://doi.org/10.3389/fphar.2014.00008> PMID: 24567720; PubMed Central PMCID: PMC3915096.
21. Leiva-Juarez MM, Ware HH, Kulkarni VV, Zweidler-McKay PA, Tuvim MJ, Evans SE. Inducible epithelial resistance protects mice against leukemia-associated pneumonia. *Blood.* 2016; 128(7):982–92. <https://doi.org/10.1182/blood-2016-03-708511> PMID: 27317793; PubMed Central PMCID: PMC4990857.
22. Leiva-Juarez MM, Kolls JK, Evans SE. Lung epithelial cells: therapeutically inducible effectors of antimicrobial defense. *Mucosal Immunol.* 2018; 11(1):21–34. Epub 2017/08/16. <https://doi.org/10.1038/mi.2017.71> PMID: 28812547; PubMed Central PMCID: PMC5738267.
23. Kirkpatrick CT, Wang Y, Leiva Juarez MM, Shivshankar P, Pantaleon Garcia J, Plumer AK, et al. Inducible Lung Epithelial Resistance Requires Multisource Reactive Oxygen Species Generation To Protect against Viral Infections. *MBio.* 2018; 9(3). Epub 2018/05/17. <https://doi.org/10.1128/mBio.00696-18> PMID: 29764948; PubMed Central PMCID: PMC5954225.
24. Knust J, Ochs M, Gundersen HJ, Nyengaard JR. Stereological estimates of alveolar number and size and capillary length and surface area in mice lungs. *Anat Rec (Hoboken).* 2009; 292(1):113–22. <https://doi.org/10.1002/ar.20747> PMID: 19115381.
25. Patton JS, Brain JD, Davies LA, Fiegel J, Gumbleton M, Kim KJ, et al. The particle has landed—characterizing the fate of inhaled pharmaceuticals. *J Aerosol Med Pulm Drug Deliv.* 2010; 23 Suppl 2:S71–87. <https://doi.org/10.1089/jamp.2010.0836> PMID: 21133802.
26. Fang FC. Antimicrobial actions of reactive oxygen species. *MBio.* 2011; 2(5). <https://doi.org/10.1128/mBio.00141-11> PMID: 21896680; PubMed Central PMCID: PMC3171981.
27. Samuel EL, Marcano DC, Berka V, Bitner BR, Wu G, Potter A, et al. Highly efficient conversion of superoxide to oxygen using hydrophilic carbon clusters. *Proc Natl Acad Sci U S A.* 2015; 112(8):2343–8. <https://doi.org/10.1073/pnas.1417047112> PMID: 25675492; PubMed Central PMCID: PMC4345556.
28. Bitner BR, Marcano DC, Berlin JM, Fabian RH, Cherian L, Culver JC, et al. Antioxidant carbon particles improve cerebrovascular dysfunction following traumatic brain injury. *ACS Nano.* 2012; 6(9):8007–14. <https://doi.org/10.1021/nn302615f> PMID: 22866916; PubMed Central PMCID: PMC3458163.
29. Marcano DC, Bitner BR, Berlin JM, Jarjour J, Lee JM, Jacob A, et al. Design of poly(ethylene glycol)-functionalized hydrophilic carbon clusters for targeted therapy of cerebrovascular dysfunction in mild traumatic brain injury. *J Neurotrauma.* 2013; 30(9):789–96. <https://doi.org/10.1089/neu.2011.2301> PMID: 22928502.
30. van der Vliet A. NADPH oxidases in lung biology and pathology: host defense enzymes, and more. *Free Radic Biol Med.* 2008; 44(6):938–55. <https://doi.org/10.1016/j.freeradbiomed.2007.11.016> PMID: 18164271; PubMed Central PMCID: PMC3233509.
31. Grandvaux N, Mariani M, Fink K. Lung epithelial NOX/DUOX and respiratory virus infections. *Clin Sci (Lond).* 2015; 128(6):337–47. <https://doi.org/10.1042/CS20140321> PMID: 25456319.
32. Strengert M, Jennings R, Davanture S, Hayes P, Gabriel G, Knaus UG. Mucosal reactive oxygen species are required for antiviral response: role of Duox in influenza a virus infection. *Antioxid Redox Signal.* 2014; 20(17):2695–709. <https://doi.org/10.1089/ars.2013.5353> PMID: 24128054.
33. Fischer H. Mechanisms and function of DUOX in epithelia of the lung. *Antioxid Redox Signal.* 2009; 11(10):2453–65. <https://doi.org/10.1089/ARS.2009.2558> PMID: 19358684; PubMed Central PMCID: PMC32823369.
34. Sena LA, Chandel NS. Physiological roles of mitochondrial reactive oxygen species. *Mol Cell.* 2012; 48(2):158–67. <https://doi.org/10.1016/j.molcel.2012.09.025> PMID: 23102266; PubMed Central PMCID: PMC3484374.
35. Garaude J, Acin-Perez R, Martinez-Cano S, Enamorado M, Ugolini M, Nistal-Villan E, et al. Mitochondrial respiratory-chain adaptations in macrophages contribute to antibacterial host defense. *Nat Immunol.* 2016; 17(9):1037–45. Epub 2016/06/28. <https://doi.org/10.1038/ni.3509> PMID: 27348412; PubMed Central PMCID: PMC4994870.
36. Kim S, Kim MJ, Park do Y, Chung HJ, Kim CH, Yoon JH, et al. Mitochondrial reactive oxygen species modulate innate immune response to influenza A virus in human nasal epithelium. *Antiviral Res.* 2015; 119:78–83. <https://doi.org/10.1016/j.antiviral.2015.04.011> PMID: 25930096.
37. West AP, Brodsky IE, Rahner C, Woo DK, Erdjument-Bromage H, Tempst P, et al. TLR signalling augments macrophage bactericidal activity through mitochondrial ROS. *Nature.* 2011; 472(7344):476–80. <https://doi.org/10.1038/nature09973> PMID: 21525932; PubMed Central PMCID: PMC3460538.



38. Dryden M. Reactive oxygen species: a novel antimicrobial. *Int J Antimicrob Agents*. 2018; 51(3):299–303. Epub 2017/09/10. <https://doi.org/10.1016/j.ijantimicag.2017.08.029> PMID: 28887201.
39. Memar MY, Ghotaslou R, Samiei M, Adibkia K. Antimicrobial use of reactive oxygen therapy: current insights. *Infect Drug Resist*. 2018; 11:567–76. Epub 2018/05/08. <https://doi.org/10.2147/IDR.S142397> PMID: 29731645; PubMed Central PMCID: PMC5926076.
40. Vatansever F, de Melo WC, Avci P, Vecchio D, Sadasivam M, Gupta A, et al. Antimicrobial strategies centered around reactive oxygen species—bactericidal antibiotics, photodynamic therapy, and beyond. *FEMS Microbiol Rev*. 2013; 37(6):955–89. Epub 2013/06/28. <https://doi.org/10.1111/1574-6976.12026> PMID: 23802986; PubMed Central PMCID: PMC3791156.
41. Zou L, Wang J, Gao Y, Ren X, Rottenberg ME, Lu J, et al. Synergistic antibacterial activity of silver with antibiotics correlating with the upregulation of the ROS production. *Sci Rep*. 2018; 8(1):11131. Epub 2018/07/26. <https://doi.org/10.1038/s41598-018-29313-w> PMID: 30042429; PubMed Central PMCID: PMC6057937.
42. Guirao-Abad JP, Sanchez-Fresneda R, Albuquerque B, Hernandez JA, Arguelles JC. ROS formation is a differential contributory factor to the fungicidal action of Amphotericin B and Micafungin in *Candida albicans*. *Int J Med Microbiol*. 2017; 307(4–5):241–8. Epub 2017/04/17. <https://doi.org/10.1016/j.ijmm.2017.03.005> PMID: 28412040.
43. Shekhova E, Kniemeyer O, Brakhage AA. Induction of Mitochondrial Reactive Oxygen Species Production by Itraconazole, Terbinafine, and Amphotericin B as a Mode of Action against *Aspergillus fumigatus*. *Antimicrob Agents Chemother*. 2017; 61(11). Epub 2017/08/30. <https://doi.org/10.1128/AAC.00978-17> PMID: 28848005; PubMed Central PMCID: PMC5655112.
44. Zhang X, Gao F. Imaging mitochondrial reactive oxygen species with fluorescent probes: current applications and challenges. *Free Radic Res*. 2015; 49(4):374–82. <https://doi.org/10.3109/10715762.2015.1014813> PMID: 25789762.
45. Forteza R, Salathe M, Miot F, Forteza R, Conner GE. Regulated hydrogen peroxide production by Duox in human airway epithelial cells. *Am J Respir Cell Mol Biol*. 2005; 32(5):462–9. <https://doi.org/10.1165/rcmb.2004-0302OC> PMID: 15677770.
46. Klebanoff SJ, Clem WH, Luebke RG. The peroxidase-thiocyanate-hydrogen peroxide antimicrobial system. *Biochim Biophys Acta*. 1966; 117(1):63–72. Epub 1966/03/28. 0304-4165(66)90152-8 [pii]. PMID: 4380562.
47. Jeon YJ, Kim HJ. Duox2-induced innate immune responses in the respiratory epithelium and intranasal delivery of Duox2 DNA using polymer that mediates immunization. *Appl Microbiol Biotechnol*. 2018; 102(10):4339–43. Epub 2018/03/31. <https://doi.org/10.1007/s00253-018-8956-y> PMID: 29600494.
48. Cegolon L, Salata C, Piccoli E, Juarez V, Palu G, Mastrangelo G, et al. In vitro antiviral activity of hypothiocyanite against A/H1N1/2009 pandemic influenza virus. *Int J Hyg Environ Health*. 2014; 217(1):17–22. <https://doi.org/10.1016/j.ijheh.2013.03.001> PMID: 23540488.
49. Wijkstrom-Frei C, El-Chemaly S, Ali-Rachedi R, Gerson C, Cobas MA, Forteza R, et al. Lactoperoxidase and human airway host defense. *Am J Respir Cell Mol Biol*. 2003; 29(2):206–12. <https://doi.org/10.1165/rcmb.2002-0152OC> PMID: 12626341.
50. Pinegin B, Vorobjeva N, Pashenkov M, Chernyak B. The role of mitochondrial ROS in antibacterial immunity. *J Cell Physiol*. 2018; 233(5):3745–54. Epub 2017/08/05. <https://doi.org/10.1002/jcp.26117> PMID: 28771715.
51. Dikalov S. Cross talk between mitochondria and NADPH oxidases. *Free Radic Biol Med*. 2011; 51(7):1289–301. Epub 2011/07/23. <https://doi.org/10.1016/j.freeradbiomed.2011.06.033> PMID: 21777669; PubMed Central PMCID: PMC3163726.
52. Stocks CJ, Schembri MA, Sweet MJ, Kapetanovic R. For when bacterial infections persist: Toll-like receptor-inducible direct antimicrobial pathways in macrophages. *J Leukoc Biol*. 2018; 103(1):35–51. Epub 2018/01/19. <https://doi.org/10.1002/JLB.4RI0917-358R> PMID: 29345056.
53. Kim YM, Kim SJ, Tatsunami R, Yamamura H, Fukai T, Ushio-Fukai M. ROS-induced ROS release orchestrated by Nox4, Nox2, and mitochondria in VEGF signaling and angiogenesis. *Am J Physiol Cell Physiol*. 2017; 312(6):C749–C64. Epub 2017/04/21. <https://doi.org/10.1152/ajpcell.00346.2016> PMID: 28424170; PubMed Central PMCID: PMC5494593.
54. Zinkevich NS, Gutterman DD. ROS-induced ROS release in vascular biology: redox-redox signaling. *Am J Physiol Heart Circ Physiol*. 2011; 301(3):H647–53. Epub 2011/06/21. <https://doi.org/10.1152/ajpheart.01271.2010> PMID: 21685266; PubMed Central PMCID: PMC3191081.
55. Munro D, Treberg JR. A radical shift in perspective: mitochondria as regulators of reactive oxygen species. *J Exp Biol*. 2017; 220(Pt 7):1170–80. Epub 2017/03/31. <https://doi.org/10.1242/jeb.132142> PMID: 28356365.