

CORRIGENDUM

Variation in activity levels amongst dogs of different breeds: results of a large online survey of dog owners from the UK – CORRIGENDUM

Emily Pickup¹, Alexander J. German^{1,2}, Emily Blackwell³, Mark Evans⁴ and Carri Westgarth^{1,5}

¹*Institute of Veterinary Science, University of Liverpool, Neston, UK*

²*Institute of Ageing and Chronic Disease, University of Liverpool, Neston, UK*

³*School of Clinical Veterinary Science, University of Bristol, Langford, UK*

⁴*Independent Veterinary Consultant, Guildford, UK*

⁵*Institute of Infection and Global Health, University of Liverpool, Liverpool, UK*

Journal of Nutritional Science (2021), vol. 10, e93, page 1 of 1

doi:[10.1017/jns.2017.7](https://doi.org/10.1017/jns.2017.7). Published by Cambridge University Press, 17 April 2017

This article was published with errors in the first sentence of the section ‘Off-lead exercise’ on page 5. The correct sentence is below.

Off-lead exercise

Some breeds were **less** commonly allowed off the lead in public ($P < 0.001$; Table 1), including: chow chow (8, 73% **not** let off the lead); Siberian husky (67, 61%); Pyrenean mountain dog (3, 60%); Alaskan Malamute (31, 60%); and St Bernard (5, 56%).

The authors apologise for the error.

Reference

Pickup, E., German, A., Blackwell, E., Evans, M., & Westgarth, C. (2017). Variation in activity levels amongst dogs of different breeds: Results of a large online survey of dog owners from the UK. *Journal of Nutritional Science*, **6**, E10. doi:[10.1017/jns.2017.7](https://doi.org/10.1017/jns.2017.7)