Prior Abortions and Barriers to Abortion Access Reported by Pregnant Women Veterans



J Gen Intern Med 37(Suppl 3):S816–S8 DOI: 10.1007/s11606-022-07576-4 © The Author(s) 2022

INTRODUCTION

The Veterans Affairs (VA) Health Administration does not cover or provide abortion under any circumstances ¹—a policy more stringent than the federal standard, which allows for coverage in cases of rape, incest, or life endangerment. ² While Veterans obtain abortions at rates similar to women in the general US population, ³ there is little data describing abortion experiences among Veterans. We conducted telephone surveys to learn about prior abortion experiences among pregnant Veterans using VA healthcare.

METHODS

Between April 2020 and April 2021, we surveyed 299 pregnant Veterans enrolled in the Center for Maternal and Infant Outcomes Research in Translation (COMFORT) study about prior abortion experiences. COMFORT study methods are described in detail elsewhere. Participants were asked a series of multiple choice questions including whether they had a prior abortion, reasons for abortion, barriers to abortion, and if they ever discussed abortion with a VA provider (with free-response follow-up). Here we report descriptive findings and bivariate analyses (chisquared, Fisher's exact or Student's *t*-tests) of participant characteristics by any prior abortion.

RESULTS

Overall, 34 women (11%) reported at least one prior abortion (Figure 1). Veterans reporting a prior abortion were significantly more likely to disclose history of military

Prior Presentations: The data presented in this manuscript have not been presented previously.

Received May 27, 2021 Accepted April 1, 2022 Published online August 30, 2022 sexual trauma and diagnosis of post-traumatic stress disorder (Table 1).

The most common reasons for seeking abortion were not being ready to parent a(nother) child (38%) and not being a good time to have a(nother) child (26%). Five women (15%) reported needing an abortion due to concurrent intimate partner violence, and two (6%) reported that the pregnancy was a result of rape. Two women reported fetal anomalies/birth defects as the reason for their abortion, and one sought abortion due to personal health problems. Seven women (21%) reported facing barriers to abortion, including cost of travel, cost of the procedure, difficulty finding a provider or appointment, travel time, and needing childcare.

Only one participant discussed their abortion with a VA provider. She explained, "the VA discouraged the decision... [my] provider only discussed negative impacts of abortion on women."

DISCUSSION

Greater than 10% of currently pregnant Veterans report a prior abortion, with similar reasons and barriers encountered by non-Veterans. ^{5,6} Abortion is an essential component of reproductive healthcare, yet current VA policy prohibits abortion without exception.

In addition to ensuring reproductive autonomy, access to comprehensive reproductive healthcare is paramount considering that women Veterans have high rates of chronic medical and mental health conditions that may increase pregnancy-associated morbidity. Veterans who use VA healthcare are of lower socioeconomic status than Veterans with private insurance, and may therefore face greater barriers to abortion care in the private sector. Furthermore, VA serves high proportions of racial/ethnic minority women, who suffer unacceptably high rates of maternal mortality due to underlying structural and societal inequalities. While VA provides the full range of contraceptive methods at no or low-cost, pregnancy while using contraception commonly occurs and is often cited as a reason for seeking abortion.

The vast majority of abortions among US women occur in free-standing clinics (i.e., not with a patient's primary medical




Fig. 1. Participant flow from the COMFORT study cohort.

provider). ¹⁰ However, while most obstetrician-gynecologists provide direct or indirect referrals for abortion, ¹¹ VA policy explicitly forbids abortion-related counseling in addition to abortion care and coverage. ¹ This restriction directly impedes delivery of evidence-based, quality reproductive healthcare. At a minimum, it is imperative that VA policy expand to match the standard set by other federal agencies by covering abortion in cases of rape, incest, and life endangerment. Even in our small cohort, some women sought abortions due to rape or maternal health problems. However, regardless of indication, inability to obtain a desired abortion has tremendous physical, psychological, and socioeconomic conse-

quences, 12-14 and even more inclusive coverage is warranted.

This study is limited by its small sample size, inclusion of only pregnant Veterans recruited via convenience sampling, and by known under-reporting of abortion leading to underestimated prevalence. Although all COMFORT participants were enrolled in VA healthcare at the time of participation, participants were not asked whether they were enrolled in VA at the time of their abortion. These issues limit the generalizability and applicability of our results. However, this exploratory study sheds light on an important and understudied issue. Our findings highlight the potential harms of current VA abortion policy, and the need for policy change.

Table 1 Participant Characteristics at the Time of Prenatal Telephone Survey, by Report of Prior Abortion (N=299)

	No prior abortion reported (n=265)	Prior abortion reported (n=34)	P-value*
Estimated age (mean ± SD, range)	32.2 ± 4.8 (22–44)	32.5 ± 5.0 (23–43)	0.72
Marital status (N, %)			0.14
Single	69 (26.0)	14 (41.2)	
Married	177 (66.8)	17 (50.0)	
Divorced/widowed/separated	18 (6.8)	3 (8.8)	
Race† (N, %)			0.27
White	151 (57.0)	16 (47.1)	
Black	80 (30.2)	15 (44.1)	
Asian	12 (4.5)	0 (0.0)	
Native Hawaiian or Other Pacific Islander	5 (1.9)	0 (0.0)	
American Indian or Alaska Native	5 (1.9)	1 (2.9)	
Other	40 (15.1)	6 (17.6)	
Hispanic or Latino/Latina (N, %)	50 (18.9)	8 (23.5)	0.54
History of military sexual trauma $(N, \%)$			
Received uninvited and unwanted sexual attention while in the military	130 (49.1)	26 (76.5)	< 0.01
Force or the threat of force was used to have unwanted sexual contact	74 (27.9)	21 (61.8)	< 0.01
while in the military			
Self-reported mental health diagnoses, any time prior to survey $(N, \%)$			
Depression	125 (47.2)	18 (52.9)	0.53
Post-traumatic stress disorder	91 (34.3)	19 (55.9)	0.01
Anxiety disorder	124 (46.8)	18 (52.9)	0.50

^{*}P-values are from chi-squared or Fisher's tests for categorical variables and Student's t-test for continuous variables †Race categories were not mutually exclusive. "Other" races included any self-specified race not fitting the categories listed

Colleen Judge-Golden, MD, PhD^{1,2} Aimee Kroll-Desrosiers, PhD^{3,4} Kristin Mattocks, PhD, MPH^{3,4} Sonya Borrero, MD, MS^{5,6}

¹University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

²Department of Obstetrics & Gynecology, Duke University School of Medicine,

Durham, NC, USA

³Veterans Affairs Central Western Massachusetts Healthcare System,

Northampton, MA, USA

⁴University of Massachusetts Medical School,

Worcester, MA, USA

⁵Center for Health Equity, Research, and Promotion (CHERP), Veterans Affairs Pittsburgh Health Care System,

Pittsburgh, PA, USA

⁶Center for Women's Health Research and Innovation (CWHRI), University of Pittsburgh,

Pittsburgh, PA, USA

Corresponding Author: Colleen Judge-Golden, MD, PhD; Department of Obstetrics & Gynecology, Duke University School of Medicine, Durham, NC, USA (e-mail: cpj6@pitt.edu).

Funding The COMFORT study was supported by the US Department of Veterans Affairs, Veterans Health Administration, Office of Research and Development (HSR&D IIR13-081; PI: Kristin Mattocks). Colleen Judge-Golden's time was partially supported by the National Institute of General Medical Sciences of the National Institutes of Health (25T32GM008208, PI: Richard Steinman).

Declarations:

Conflict of interest: The authors declare that they do not have a conflict of interest

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended

use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/bu/4.0/.

REFERENCES

- US Department of Veterans Affairs. Medical Benefits Package. 38 CFR § 17.38(c)(1). https://ecfr.io/Title-38/Section-17.38. Accessed May 15, 2021.
- U.S. Code § 1093. Performance of abortions: restrictions. www.law. cornell.edu/uscode/text/10/1093. Accessed May 15, 2021.
- Schwarz EB, Sileanu FE, Zhao X, Mor MK, Callegari LS, Borrero S. Induced abortion among women veterans: data from the ECUUN study. Contraception. 2018;97(1):41-7.
- Mattocks KM, Baldor R, Bean-Mayberry B, et al. Factors impacting perceived access to early prenatal care among pregnant veterans enrolled in the Department of Veterans Affairs. Womens health issues. 2019;29(1):56-63.
- Increasing Access to Abortion: ACOG Committee Opinion, Number 815.
 Obstet Gynecol. 2020;136(6):e107-e15.
- Biggs MA, Gould H, Foster DG. Understanding why women seek abortions in the US. BMC Womens Health. 2013:13:29.
- Frayne S, Phibbs C, Saechao F, et al. Sourcebook: Women Veterans in the Veterans Health Administration. Volume 4: Longitudinal Trends in Sociodemographics, Utilization, Health Profile, and Geographic Distribution. Washington DC: Women's Health Evaluation Initiative, Women's Health Services, Veterans Health Administration, Department of Veterans Affairs; 2018.
- Centers for Disease Control and Prevention. Pregnancy Mortality Surveillance System. https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-mortality-surveillance-system.htm. Accessed May 15, 2021.
- Jones RK. Reported contraceptive use in the month of becoming pregnant among U.S. abortion patients in 2000 and 2014. Contraception. 2018;97(4):309-12.
- Jones RK, Witwer E, Jerman J. Abortion Incidence and Service Availability in the United States, 2017. New York: Guttmacher Institute 2019
- Daniel S, Schulkin J, Grossman D. Abortion referral practices among a national sample of obstetrician-gynecologists. Womens health issues. 2020;30(6):446-52.
- Gerdts C, Dobkin L, Foster DG, Schwarz EB. Side effects, physical health consequences, and mortality associated with abortion and birth after an unwanted pregnancy. Womens health issues. 2016;26(1):55-9.
- Biggs M, Upadhyay UD, McCulloch CE, Foster DG. Women's mental health and well-being 5 years after receiving or being denied an abortion: a prospective, longitudinal cohort study. JAMA Psychiatry. 2017;74(2):169-78.
- Foster DG, Biggs MA, Ralph L, Gerdts C, Roberts S, Glymour MM.
 Socioeconomic outcomes of women who receive and women who are denied wanted abortions in the United States. Am J Public Health. 2018;108(3):407-13.

Publisher's Note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.