



Published in final edited form as:

*Mucosal Immunol.* 2022 February ; 15(2): 362–372. doi:10.1038/s41385-021-00468-6.

## Enterocyte–innate lymphoid cell crosstalk drives early IFN- $\gamma$ -mediated control of *Cryptosporidium*

Jodi A. Gullicksrud<sup>1</sup>, Adam Sateriale<sup>1,2</sup>, Julie B. Englies<sup>3</sup>, Alexis Gibson<sup>1</sup>, Sebastian Shaw<sup>1</sup>, Zachary A. Hutchins<sup>1,4</sup>, Lindsay Martin<sup>1</sup>, David A. Christian<sup>1</sup>, Gregory A. Taylor<sup>5,6</sup>, Masahiro Yamamoto<sup>7</sup>, Daniel P. Beiting<sup>1</sup>, Boris Striepen<sup>1</sup>, Christopher A. Hunter<sup>1</sup>

<sup>1</sup>Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, 380 South University Avenue, Philadelphia, PA 19104, United States of America.

<sup>2</sup>Current: The Francis Crick Institute, London NW1 1AT, United Kingdom.

<sup>3</sup>Department of Pathobiology, New Bolton Center, School of Veterinary Medicine, University of Pennsylvania, Kennett Square, PA 19348, United States of America.

<sup>4</sup>Current: Jill Robests Institute for Research in Inflammatory Bowel Disease, Weill Cornell Medicine, New York, NY 10021, United States of America

<sup>5</sup>Departments of Medicine; Molecular Genetics and Microbiology; and Immunology; and Center for the Study of Aging and Human Development, Duke University Medical Center, Durham, North Carolina, United States of America

<sup>6</sup>Geriatric Research, Education, and Clinical Center, Durham VA Health Care System, Durham, North Carolina, United States of America

<sup>7</sup>Department of Immunoparasitology, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan.

### SUMMARY

The intestinal parasite, *Cryptosporidium*, is a major contributor to global child mortality and causes opportunistic infection in immune deficient individuals. Innate resistance to *Cryptosporidium*, which specifically invades enterocytes, is dependent on the production of IFN- $\gamma$ , yet whether enterocytes contribute to parasite control is poorly understood. In this study, utilizing a mouse-adapted strain of *C. parvum*, we show that epithelial-derived IL-18 synergized with IL-12 to stimulate innate lymphoid cell (ILC) production of IFN- $\gamma$  required for early parasite

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:[http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Corresponding Author: Christopher A. Hunter, 310F Hill Pavilion, 380 S University Ave, Philadelphia, PA 19104, Fax: 215-746-2295 Phone: 215-573-7772 [chunter@vet.upenn.edu](mailto:chunter@vet.upenn.edu).

#### AUTHOR CONTRIBUTIONS

JAG, AS, BS and CAH conceived of the study; JAG and AS conducted experiments. JBE contributed pathologist expertise and histology scoring. AG provided the sequencing analysis, with technical support from DPB. SS, ZAH, LM contributed data acquisition. DAC, GAT, and MY contributed mice and expertise. All authors approved the final version of this manuscript to be published and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

#### DISCLOSURES

The authors have no conflicts of interest to report.

control. The loss of IFN- $\gamma$ -mediated STAT1 signaling in enterocytes, but not dendritic cells or macrophages, antagonized early parasite control. Transcriptional profiling of enterocytes from infected mice identified an IFN- $\gamma$  signature and enrichment of the anti-microbial effectors IDO, GBP and IRG. Deletion experiments identified a role for *Irgm1/m3* in parasite control. Thus, enterocytes promote ILC production of IFN- $\gamma$  that acts on enterocytes to restrict the growth of *Cryptosporidium*.

---

## INTRODUCTION

The intestinal epithelium is an important site for nutrient uptake and a barrier to micro-organisms. However, this barrier can be disrupted by a diverse group of viruses, bacteria and parasites that infect the gastrointestinal tract. The enteric diseases caused by these pathogens are of public health importance, accounting for 8–10% of deaths in children worldwide (1). While many of these pathogens will disseminate from the gut, a subset is restricted to the epithelial layer, where their interactions with enterocytes are likely key determinants of disease outcome. The epithelium is composed of enterocytes, a large population of columnar epithelial cells necessary for nutrient uptake, as well as subsets of specialized epithelial cells including Paneth, goblet and tuft cells that have roles in mucosal homeostasis and immune defense (2–10).

How enterocytes participate in resistance to different types of infection is a fundamental question that is particularly relevant to epithelial-restricted pathogens such as rotavirus, norovirus, astrovirus, *Shigella*, *Cyclospora* and *Cryptosporidium*. The apicomplexan parasite, *Cryptosporidium*, is a leading cause of severe diarrhea and death in infants (11–14) and a common opportunistic infection in individuals with primary and acquired immune deficiencies (15, 16). Currently, there are no fully effective drugs or vaccines to treat or prevent cryptosporidiosis. Based on clinical experience and animal models, effective control and clearance of *Cryptosporidium* is dependent on cell-mediated immunity and the production of IFN- $\gamma$  (17–20). *Cryptosporidium* invasive stages infect the apical surface of enterocytes, where they transform into the replicative trophozoite and occupy a unique intracellular but extracytoplasmic niche (21). The restriction of this parasite to the intestinal epithelium provides a model to understand how the immune system senses enteric pathogens and an opportunity to identify enterocyte-driven pathways that promote the control of intracellular pathogens.

Most human infections are caused by the anthroponotic *Cryptosporidium hominis* and the zoonotically transmitted *C. parvum*. *C. parvum* is the species most widely used as an experimental model, however it does not robustly infect adult immune competent mice. For this reason, most studies with this organism have used neonates or immune deficient mice (such as those that lack IFN- $\gamma$  or components of adaptive immunity) to help define important pathways for resistance to *Cryptosporidium*. This approach has provided evidence that dendritic cell (DC) derived IL-12 promotes NK cell production of IFN- $\gamma$  required for innate restriction of this infection (22–26). IL-18 is another cytokine that promotes resistance to *C. parvum*, but it is unclear if this protection is due to its ability to enhance ILC production of IFN- $\gamma$  or whether IL-18 directly activates epithelial cells to limit parasite

growth (27–30). Likewise, while IFN- $\gamma$  is important for parasite control, multiple cell types in the gut can respond to this cytokine and, thus, it is unclear whether the protective effects of IFN- $\gamma$  are mediated through enterocytes, dendritic cells or macrophages.

Recent studies have developed the means to genetically modify *C. parvum* that is dependent on the ability to select for transgenic parasites while being passaged in IFN- $\gamma$ -deficient mice (31). Over time, this has resulted in parasite adaptation whereby these transgenic *C. parvum* readily infect wild type (WT) mice and provide an experimental system to dissect the events required for protective immunity in an immunocompetent setting. In this study, infection with mouse-adapted *C. parvum* (maCp) revealed that intestinal type-1 innate lymphoid cells (ILC1s) are a rapid source of IFN- $\gamma$  that limits parasite growth. This protective response was dependent on the production of IL-12 and epithelial-derived IL-18. Lineage-specific deletion of STAT1, a critical transcription factor downstream of IFN- $\gamma$  signaling, demonstrated that STAT1 was uniquely required in enterocytes—but not DCs or macrophages—to restrict parasite growth. Transcriptional profiling of enterocytes from infected mice highlighted an IFN- $\gamma$  signature, and subsequent deletion experiments identified immune-related GTPase-m1 and -m3 (*Irgm1/3*) as downstream effectors of the IFN- $\gamma$ -mediated response. Together, these studies establish that enterocytes have a central role as a source of IL-18 required to stimulate local ILC responses and are key mediators of IFN- $\gamma$ -mediated protection to an important enteric pathogen.

## RESULTS

### Localized innate IFN- $\gamma$ provides early constraint of *Cryptosporidium* infection

To assess the relationship between parasite burden and IFN- $\gamma$  production, WT mice treated with an isotype control antibody (IgG) or anti-IFN- $\gamma$  ( $\alpha$ -IFN- $\gamma$ ) were infected with a mouse-adapted *C. parvum* strain that expressed nanoluciferase (maCp-Nluc) (31, 32). At 4 days post-infection (dpi), paired biopsies were taken along the entirety of the intestine and used to quantify nanoluciferase activity or placed in culture to assess levels of secreted IFN- $\gamma$ . In control mice, parasite replication was readily detected and restricted to the distal small intestine but levels of IFN- $\gamma$  in ileal supernatants were not elevated above uninfected controls (Figure 1A). However, in mice treated with anti-IFN- $\gamma$ , there was an 80-fold increase in parasite burden and *ex vivo* IFN- $\gamma$  was readily detected and correlated with areas of the gut with the highest parasite burdens (Figure 1A, filled circles). Thus, the early production of IFN- $\gamma$  provides a mechanism of resistance to *C. parvum*, but the ability to detect the infection-induced production of IFN- $\gamma$  is dependent on the presence of a high tissue parasite burden.

To assess the contribution of innate and adaptive sources of IFN- $\gamma$  on early resistance to maCp-Nluc, WT and *Rag2*<sup>-/-</sup> mice (lacking T and B cells) were treated with  $\alpha$ -IFN- $\gamma$  or an isotype control prior to infection. No gross histological differences were detected between mouse strains at steady state (Supplemental Figure 1A). In both WT and *Rag2*<sup>-/-</sup> mice, IFN- $\gamma$  neutralization resulted in increased oocyst shedding by 4dpi, which was exacerbated at 6dpi (43-fold in WT and 263-fold in *Rag2*<sup>-/-</sup>, Figure 1B), and correlated with histological analysis that showed enhanced parasite burden in *Rag2*<sup>-/-</sup> mice treated with  $\alpha$ -IFN- $\gamma$  (Figure 1 C–D). In addition, there were significant infection-induced local changes

in mucosal architecture and enterocyte morphology by 5dpi, including reduced average villus:crypt ratio, increased crypt mitoses and a trend toward increased epithelial dysplasia (Figure 1E–G). However, quantification of these changes across multiple sections did not yield statistically significant changes between infected mice and those treated with  $\alpha$ -IFN- $\gamma$ . However, when observing ileal pathology at a later timepoint,  $\alpha$ -IFN- $\gamma$  treatment resulted in more severe pathological changes (12dpi Supplemental Figure 1B). This pathology corresponds with the sustained oocyst shedding observed after the peak parasite burden at 6dpi. Interestingly, treatment of chronically infected *Rag2*<sup>-/-</sup> mice with  $\alpha$ -IFN- $\gamma$  at 50dpi led to a marked recrudescence in parasite burden (Figure 1H). These data identify an innate mechanism of IFN- $\gamma$ -mediated resistance to *C. parvum* that operates during the acute and chronic phases of infection, but which is insufficient for parasite clearance.

### Innate lymphoid cells are required for control of *Cryptosporidium*

There are several ILC populations capable of IFN- $\gamma$  production that include: NK cells, ILC1s and ILC3s. Early studies with *C. parvum* concluded that NK cells were a major source of IFN- $\gamma$ , but the use of  $\alpha$ -asialo-GM1 (which depletes NK cells but not other ILC populations) indicated that there may be other innate sources of IFN- $\gamma$  (26, 33). Indeed, when *Rag2*<sup>-/-</sup> mice infected with maCp-Nluc were treated with  $\alpha$ -asialo-GM1, there was efficient depletion of splenic NK cells but the intestinal ILCs (CD45.2<sup>+</sup> NK1.1<sup>+</sup>NKp46<sup>+</sup>) and levels of secreted IFN- $\gamma$  were only reduced by approximately half (Supplemental Figure 1C–E). Therefore, to assess the role of ILCs in resistance to *C. parvum*, *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice, which lack all ILCs in addition to T and B cells, were compared to WT and *Rag2*<sup>-/-</sup> mice. After infection with maCp-Nluc, oocyst shedding declined in WT mice by 9dpi, and resolved by 18dpi (Figure 2A). Parasite levels in *Rag2*<sup>-/-</sup> mice were not elevated compared to WT mice at early time points (3–9dpi), but the infection failed to resolve in the absence of adaptive immune cells. In contrast, *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice demonstrated little evidence of parasite control at any time examined, with approximately 250-fold higher nanoluciferase readings by 6dpi, compared to *Rag2*<sup>-/-</sup> mice, which were sustained for the duration of the experiment (Figure 2A). In uninfected mice, there were no gross histological differences in the ileal tissues among the three strains (Supplemental Figure 1A). At 5dpi, consistent with the data in Fig 1B, WT and *Rag2*<sup>-/-</sup> mice had similar numbers of parasites and levels of infection-induced changes to the epithelium (Figure 2D–G, white and gray bars). In contrast, *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice exhibited dramatically increased parasite burden (Figure 2B–C). The absence of ILCs also resulted in increased mitoses and crypt branching, as well as more severe villus pathology, indicated by increased epithelial dysplasia and attenuation (Fig 2D–G, black bars). Thus, in the absence of adaptive immunity, ILCs are required for both early and long-term control of *Cryptosporidium* and the lack of these cells results in unrestricted parasite replication and severe intestinal pathology.

### ILC1s are a major source of innate IFN- $\gamma$ during *C. parvum* infection

To determine the innate cellular source(s) of IFN- $\gamma$  in the intestinal epithelium, mice in which the gene for the surface-expressed protein, Thy1.1, is under the transcriptional control of the *Il1ng* promoter (34) were infected with mCherry-expressing maCp-Nluc. This approach allowed the simultaneous quantification of parasite-infected (mCherry<sup>+</sup>) enterocytes (EpCAM<sup>+</sup>CD45<sup>-</sup>) and innate lymphoid cells

(CD45.2<sup>+</sup>CD3<sup>-</sup>CD19<sup>-</sup>NK1.1<sup>+</sup>NKp46<sup>+</sup>) producing IFN- $\gamma$  (Thy1.1<sup>+</sup>). Based on the data in Fig 1A, groups of mice were also treated with  $\alpha$ -IFN- $\gamma$  to increase the ability to detect cells that produce IFN- $\gamma$ . In uninfected mice, enterocytes lack mCherry, and a low basal percentage of ILCs express surface Thy1.1 regardless of  $\alpha$ -IFN- $\gamma$  treatment (Figure 3A, C). At 4dpi, a detectable portion of IECs were infected (mCherry<sup>+</sup>), and there was a small increase in the frequency of Thy1.1<sup>+</sup> ILCs. Treatment with  $\alpha$ -IFN- $\gamma$  resulted in a significant increase in parasite burden (Figure 3A–B) and a 3–5-fold increase in the proportion of Thy1.1<sup>+</sup> ILCs (Figure 3C–D). This increased production of IFN- $\gamma$  was specific to the intestinal epithelium, as few cells expressed Thy1.1 in Peyer's patches or mesenteric lymph nodes, even with IFN- $\gamma$  neutralization (Figure 3E–F). Of note, in these infected immune competent reporter mice, myeloid cells (EpCAM<sup>-</sup>CD3<sup>-</sup>CD19<sup>-</sup>NK1.1<sup>-</sup>) did not express Thy1.1 (data not shown), but a proportion of intestinal CD4<sup>+</sup> T cells and, to a lesser extent, CD4<sup>+</sup>CD8 $\alpha$ <sup>+</sup> T cells did express increased levels of Thy1.1 by 5 dpi (Supp Fig 2A).

To characterize the ILC subsets that produce IFN- $\gamma$ , expression of the transcription factors Eomes, T-bet and ROR $\gamma$ t was used to distinguish NK cells, ILC1s, and ILC3s, respectively. In uninfected reporter mice, the ILC populations in the gut were composed of 50–60% T-bet<sup>+</sup> ILC1s, 30–40% Eomes<sup>+</sup> NK cells and 5–10% were ROR $\gamma$ t<sup>+</sup> ILC3s (Supplemental Figure 2B). All three of these ILC subsets were also present at 4dpi (Figure 3G and 3I). Among ILCs not expressing IFN- $\gamma$  (Thy1.1<sup>-</sup>), the proportions of NK cells, ILC1s and ILC3s were similar to those of uninfected mice (Figure 3G, I, bottom plots and Figure 3H, J open circles). However, in infected mice, the proportion of Tbet<sup>+</sup> ILCs was significantly increased in cells expressing *Ifng* (Thy1.1<sup>+</sup>), where approximately 80% of the Thy1.1<sup>+</sup> cells expressed T-bet and not Eomes or ROR $\gamma$ t (Figure 3G–J, filled circles). In addition, while NK cells in the mesenteric lymph nodes expressed CD27 and CD49b, the Thy1.1<sup>+</sup> cells in the intestinal epithelium showed low expression of these NK cell markers (Supplemental Figure 2C–D). These data indicate that within days of *Cryptosporidium* infection, ILC1s present in the intestine become activated and are a major source of IFN- $\gamma$ .

### Epithelial-derived IL-18 synergizes with IL-12 to promote ILC production of IFN- $\gamma$

IL-12 and IL-18 have roles in innate resistance to *Cryptosporidium* sp. (22, 27, 32) and can, in other experimental systems, synergize to stimulate ILC production of IFN- $\gamma$  (35–37). To evaluate their contributions to early resistance to *C. parvum*, the level of oocyst shedding in WT, *Ifng*<sup>-/-</sup>, *Il12b*<sup>-/-</sup> (IL-12p40) and *Il18*<sup>-/-</sup> mice were compared. As expected, infection was established in WT mice, and *Ifng*<sup>-/-</sup> mice showed a rapid and marked increase in parasite burden. While the parasite levels in *Il12b*<sup>-/-</sup> mice were comparable to *Ifng*<sup>-/-</sup> mice, *Il18*<sup>-/-</sup> mice showed a phenotype that was intermediate between WT and *Ifng*<sup>-/-</sup> mice (Figure 4A). A recent study highlighted BATF3-dependent CD103<sup>+</sup> DCs as a source of IL-12 during neonatal infection with *C. parvum* (24), but the cellular source of IL-18 required to control *Cryptosporidium* was unclear. To determine the importance of the intestinal epithelium as a cellular source of IL-18 during cryptosporidiosis, oocyst shedding from mice bearing an epithelial lineage-specific deletion of IL-18 (*Villin-Cre x Il18*<sup>fl/fl</sup>, here referred to as *Il18*<sup>IEC</sup>) was compared to WT and *Il18*<sup>-/-</sup> mice. In these studies, the levels of infection in *Il18*<sup>IEC</sup> mice were comparable to whole-body *Il18*<sup>-/-</sup> mice (Figure 4B),

suggesting that the intestinal epithelium is a key source of the IL-18 that mediates early resistance to *C. parvum*.

The experiments that utilized mice with germline deletion of IL-18 or IL-12 identified an important role for these cytokines in early resistance to *C. parvum*. However, analyses of naïve *Il12b*<sup>-/-</sup> mice revealed a marked absence of ILC populations in the intestinal epithelium (Supp Fig 3A–C) and previous studies have reported that germline deletion of IL-18 impacts immune homeostasis in the gut (38). In order to control for these confounding effects, C57BL/6 and *Rag2*<sup>-/-</sup> mice were treated with antibodies against IL-12p40, IL-18, or both beginning 4 days prior to infection with maCp-Nluc. In WT and *Rag2*<sup>-/-</sup> mice, neither treatment alone dramatically impacted susceptibility to maCp-Nluc but the simultaneous blockade of IL-12 and IL-18 led to a 14.5-fold increase in parasite burden (*Rag2*<sup>-/-</sup>: Figure 4C, BL/6: Supp Fig. 3D). Moreover, despite having similar infection burdens to α-IFN-γ treated mice (Figure 4D–E), *Ifng*-Thy1.1 reporter mice treated with α-IL-12p40 plus α-IL-18 showed nearly complete loss of Thy1.1<sup>+</sup> ILC (Figure 4F–G). Collectively, these experiments demonstrate that epithelial-derived IL-18 synergizes with IL-12 to stimulate ILC production of IFN-γ required for early restriction of *Cryptosporidium* infection.

### Enterocytes are critical contributors to early IFN-γ-mediated resistance to *C. parvum*

While IFN-γ is important in acute resistance to *Cryptosporidium*, it is unclear whether it acts directly on infected cells to restrict parasite growth, or indirectly via activation and maturation of macrophages and dendritic cells (39). The transcription factor STAT1 is the major mediator of IFN signaling. *Ifng*<sup>-/-</sup> and *Stat1*<sup>-/-</sup> mice had similarly enhanced susceptibility to maCp, suggesting that IFN-γ is the main driver of STAT1 signaling required for resistance to this parasite (Supplemental Figure 4A). Therefore, lineage-specific Cre-lox mediated deletion was utilized to identify cell subsets in which STAT1 signaling was critical for early control of maCp. Loss of STAT1 only in dendritic cells (*Cd11c*-Cre x *Stat1*<sup>fl/fl</sup>; *Stat1*<sup>DC</sup>) or macrophages (*Lys2*-Cre x *Stat1*<sup>fl/fl</sup>; *Stat1*<sup>MΦ</sup>) did not enhance parasite burden over Cre<sup>-</sup> controls. In contrast, when STAT1 was deleted from enterocytes using a tamoxifen-inducible deletion (*Villin*-Cre<sup>ERT2</sup> x *Stat1*<sup>fl/fl</sup>; *Stat1*<sup>IEC</sup>), oocyst shedding was greatly increased (Figure 5A) and at a level that paralleled complete *Stat1*<sup>-/-</sup> mice (Figure 5B). This was not due to reduced levels of IFN-γ in the intestine (Supp Fig. 4B). These data indicate that IFN-γ induced STAT1-mediated activity in intestinal epithelial cells provides a cell intrinsic mechanism for the control of *Cryptosporidium*.

Because the mechanism that leads to the IFN-γ-mediated restriction of *Cryptosporidium* in enterocytes is unknown, transcriptional profiling of sort-purified enterocytes from uninfected or infected mice was performed. RNA-seq analysis of enterocytes identified a limited set of differentially expressed genes; that included several IFN-inducible GTPases: *Igtp* (encoding Irgm3), *Iigp1* (encoding Irga6), *Gm12250* (encoding Irgb10), *Gbp7*, *Tgtp1*, *Ifi47* (encoding Irg-47), and *Irgm1* (Figure 5C). Despite the limited number of differentially expressed genes, GSEA showed a strong enrichment of IFN-γ signaling in enterocytes from infected mice (Figure 5D), consistent with expression of transcripts for *Ifngr1* and *Ifngr2* in IECs (data not shown). Enrichment mapping software was used to define related gene functions and examine overlap between enriched gene sets. This analysis identified

3 main clusters: IFN- $\gamma$  signatures, mitochondrial signatures, and protein translation/ribosome signatures (Figure 5E and Supp Fig 4C–D). The mitochondrial and ribosome signatures have been associated with enterocyte stress (40, 41) and likely reflect the increased mitotic index and epithelial dysplasia associated with this infection (see Figures 1 and 2). Furthermore, while gene sets encompassing both IFN- $\gamma$  and type I IFNs were enriched, the degree of overlap was greatest among the all-encompassing “interferon signaling” and the two interferon gamma gene sets, further indicating that IFN- $\gamma$  signaling was the dominant response in infected IECs.

Given the dominant IFN- $\gamma$  signature, the expression data sets were curated for changes in IFN- $\gamma$ -induced genes associated with control of intracellular pathogens (Fig 5F). Genes strongly upregulated by maCp infection included those encoding for  $\beta 2m$  and CIITA (Class II Major Histocompatibility Complex Transactivator) which affect MHC I and MHC II expression, respectively (Figure 5F). Also markedly upregulated were transcripts encoding for indolamine dioxygenase (IDO), several Guanylate Binding Proteins (GBPs) and Immunity Related GTPases (IRGs), all of which are known to be important in non-hematopoietic cells for IFN- $\gamma$  to restrict the growth of *T. gondii* (42, 43). However, their roles in resistance to *Cryptosporidium* have not been established. To test the impact of these pathways on maCp, the course of infection in *Ido1*<sup>-/-</sup>, GBP<sup>chr3</sup> (lacking *Gbp2*, *Gbp3*, *Gbp5*, *Gbp7*, and *Gbp2ps*) and *Irgm1/m3*<sup>-/-</sup> mice was compared with that in WT mice by fecal oocyst shedding. Because *Irgm1*-deficient mice have multiple defects in immune function that are mitigated by the loss of *Irgm3*, *Irgm1/m3*<sup>-/-</sup> mice are used to study the role of *Irgm1* in resistance to infections (44, 45). While *Ido1*<sup>-/-</sup> and GBP<sup>chr3</sup> mice showed no enhanced susceptibility to maCp-Nluc, *Irgm1/m3*<sup>-/-</sup> mice demonstrated a greater than 5-fold increase in fecal oocyst shedding at 5dpi (Figure 5G). We note that parasite burden in *Irgm1/m3*<sup>-/-</sup> mice was intermediate between WT and *Ifng*<sup>-/-</sup> or *Stat1*<sup>IEC</sup> mice, which argues for additional IFN- $\gamma$ /STAT1-mediated mechanisms for parasite control. These findings provide the first evidence for a role of IRGs in the IFN- $\gamma$ -mediated mechanism of resistance to *Cryptosporidium*.

## DISCUSSION

The importance of IFN- $\gamma$  in resistance to *Cryptosporidium* is well appreciated (17, 19, 46), but significant gaps exist in our understanding of the specific cells that produce and respond to this central cytokine in order to mediate parasite control. Previous studies have focused on NK cells as a potential innate source of IFN- $\gamma$  in resistance to *C. parvum* (25, 26); however, they predate the identification of other ILC populations. As confirmed in our studies, NK cells are largely present in secondary lymphoid tissues and the circulation, whereas the other ILC subsets are predominately tissue resident (47). Although there is marked phenotypic and functional plasticity for ILCs in the intestine (48), our studies with mouse-adapted *C. parvum* suggested that ILC1s were the major early source of IFN- $\gamma$  in the small intestine, although NK cells likely also contributed. In other models of intracellular infection, NK and ILC1-mediated resistance is transient, associated with the acute phase of infection but not sufficient for long term control (49, 50). Thus, it was unexpected that although innate production of IFN- $\gamma$  was not sufficient for parasite clearance in *Rag*<sup>-/-</sup> mice, it did provide a significant level of long-term control of maCp. Because of the lack of effective therapies

to treat chronic cryptosporidiosis in patients with primary or acquired defects in T cell function (51–53), this model provides an opportunity to understand how ILC responses are maintained and whether they can be enhanced to mediate parasite clearance.

Previous studies have demonstrated that IL-12 and IL-18 are important in resistance to *C. parvum* (20, 22, 27, 29, 54). While DCs are a major source of IL-12 (24, 55, 56), a recent study reported that, *in vitro*, DCs incubated with *C. parvum* activate the inflammasome and downstream caspase-1, resulting in IL-18 secretion (30). In addition, IL-18 has been proposed to promote parasite control independent of IFN- $\gamma$  in enterocyte cell lines (29). In contrast, we found that enterocyte-intrinsic NLRP6 inflammasome activation was required for IL-18 mediated resistance to *Cryptosporidium in vivo* (32). Consistent with that observation, the studies presented here highlight enterocytes as a critical source of IL-18 which synergized with IL-12 to stimulate ILC1s to produce early IFN- $\gamma$ . During infection with *Citrobacter rodentium*, an extracellular bacteria that attaches to the luminal surface of enterocytes, NLRP3 inflammasome activation leads to enterocyte-derived IL-18, which is required for host resistance (57–59). Thus, different sensors in enterocytes allow these cells to respond to diverse pathogens but converge on the processing of IL-18. In contrast, although *Salmonella* infects enterocytes, enteric neurons (not enterocytes) are the relevant source of IL-18 required for protection (60). Intriguingly, patients with Hirschsprung disease have regions that lack distal bowel ganglia and are susceptible to *Cryptosporidium* (61, 62), but additional experiments will be required to determine whether the enteric nervous system is also a relevant source of IL-18 required for resistance to *Cryptosporidium*.

IFN- $\gamma$  is a cytokine with wide ranging effects relevant to *Cryptosporidium* that include its ability to enhance antigen presentation and activation of anti-microbial activities within non-hematopoietic cells. Here, deletion of STAT1 was utilized to target the downstream effects of IFN- $\gamma$  during maCp infection but it is important to note that STAT1 is utilized by other IFNs and cytokines such as IL-27. Indeed, there are reports that endogenous type I (63) and type III IFNs (64) contribute to early resistance to *C. parvum*. However, we have not observed a role for IL-27 or the type I IFNs in parasite control (unpublished observations). Nevertheless, in the current study, mice deficient in STAT1 or IFN- $\gamma$  showed similar susceptibility and the genes most strongly enriched in enterocytes from infected mice are more closely associated with IFN- $\gamma$ , compared to IFN- $\alpha/\beta$  or IFN- $\lambda$  (65). The finding that the loss of STAT1 in macrophage and DC populations did not impact susceptibility indicated that these populations were not important effectors of IFN- $\gamma$ -mediated parasite control. In contrast, the inducible deletion of STAT1 in enterocytes established a cell-intrinsic role for STAT1 in resistance to *C. parvum*. This pathway contrasts with *Salmonella*, where IFN- $\gamma$  is required to restrict bacterial growth in macrophages but not enterocytes (66, 67), as well as *Toxoplasma gondii*, where innate sources of IFN- $\gamma$  act specifically on cDC1s (68).

Studies using human intestinal cell lines concluded that IFN- $\gamma$  can inhibit growth of *Cryptosporidium*, but inhibitory effects *in vitro* are modest and the mechanisms that underlie parasite restriction are not understood (14, 46). The identification of a role for *Irgm1* suggests a potential overlap with mechanisms used to control other intracellular pathogens. The upregulation of *Irgm1*, *Igtp* and *Iigp1* in enterocytes from infected mice highlights anti-microbial effectors that, together with the GBPs, can intersect with autophagy pathways



involved in pathogen restriction (69). It is important to note that *Irgm1* and *Irgm3* are also expressed in other cell types, thus the use of germline deletion does not rule out additional roles for these proteins in resistance to *C. parvum*. Furthermore, while the loss of the GBPs on chromosome 3 (GBP 1, GBP2, GBP3, GBP5, and GBP7) had no detectable impact on host susceptibility, it does not rule out possible roles for the other GBPs located on chromosome 5. Of these, the expression of *Gbp6* was significantly enriched in epithelial cells from infected mice. Another possibility is that, although GBPs interact with vacuoles that contain other intracellular pathogens, perhaps the extra-cytoplasmic location of *Cryptosporidium* and the partitioning by parasite induced cellular barriers precludes this interaction (21), which would render the canonical GBP-dependent mechanisms of protection ineffective. The recent advances using enteroid-derived systems to culture *Cryptosporidium* (70, 71) should be useful to dissect how enterocytes utilize interferon-stimulated genes to directly clear or restrict growth of *Cryptosporidium* species.

The intestinal epithelium is a critical barrier where the immune system interacts with diverse microbial communities in the gut. Enterocyte interactions with the microbiome are central to homeostasis and the etiology of a variety of allergies and inflammatory conditions (5, 72, 73). There is an increased appreciation that different types of epithelial cells provide signals that help to coordinate the enteric nervous system, gut physiology and mucosal immunity in order to maintain tolerance. For extracellular pathogens, enterocytes promote clearance of the helminth *Trichuris muris* (74) and the bacteria *Clostridium difficile* (75) and *Citrobacter rodentium* (57). Less is known about how enterocytes respond to intracellular infections, although inflammasome-mediated extrusion of *Salmonella*-infected enterocytes limits bacterial spread (76). Indeed, expulsion of infected enterocytes represents a conserved mechanism of pathogen resistance that is present in insects and higher vertebrates (77–79). The studies presented here identified enterocytes as a critical source of IL-18 required for ILC activation and as key drivers of IFN- $\gamma$  mediated resistance to an important enteric pathogen. Thus, mouse-adapted *C. parvum* is a valuable model of an enteric infection that affects intestinal physiology and nutritional status. Furthermore, the ability to genetically modify this organism provides the opportunity to understand the role of enterocytes in pathogen recognition and resistance to infection.

## MATERIALS & METHODS

### Mice

C57BL/6 (stock no: B6NTac), *Rag2*<sup>-/-</sup> (stock no: RAGN12), and *Rag2*<sup>-/-</sup> *Il2rc*<sup>-/-</sup> (stock no: 4111) were purchased from Taconic. C57BL/6 (stock no: 000664), *Rag2*<sup>-/-</sup> (stock no: 008449), *Ifng*<sup>-/-</sup> (stock no: 002287), *Il12rb*<sup>-/-</sup> (stock no: 002693), *Stat1*<sup>-/-</sup> (stock no: 012606), *Il18*<sup>-/-</sup> (stock no: 004130), *Ido1*<sup>-/-</sup> (stock no:005867), *Vill*-Cre (stock no:021504), *Lys2*-Cre (stock no: 004781), and *Cd11c*-Cre (stock no. 008068) were purchased from Jackson Laboratory and maintained in-house. *Stat1*<sup>fllox</sup> mice were generated as previously described (80) and maintained in house. *Ifng*/Thy1.1 BAC-In mice were provided by Dr. Phillip Scott but originated in the laboratory of Dr. Casey Weaver (34, 81). *Il18*<sup>fllox</sup> mice were provided by Dr. Jorge Henoa-Mejia. *Vill*-Cre<sup>ERT2</sup> were provided by Dr. Lou Ghanem at Children's Hospital of Philadelphia and Dr. David Artis at Cornell University.

In-house breeding was performed to obtain all Cre-lox combinations. Unless otherwise noted, mice used in this study were males or females ranging from 7–11 weeks. We did not observe a difference in infection burden between male and female mice. All mice were age matched within individual experiments. All protocols for animal care were approved by the Institutional Animal Care and Use Committee of the University of Pennsylvania (protocol #805405 and #806292).

### Parasites and infection

Transgenic *C. parvum* expressing nanoluciferase and mCherry (32, 82) are propagated by orally infecting *Iflng*<sup>-/-</sup> mice. Oocysts are purified from fecal collections of infected mice using sucrose flotation followed by a cesium chloride gradient, as previously described (82). Mice were infected with  $5 \times 10^4$ – $1 \times 10^5$  oocysts by oral gavage. To measure parasite burden in intestinal tissue, 5mm biopsy punches were taken along murine small intestines and suspended in 0.5mL lysis buffer (50mM tris HCl (pH 7.6), 2mM DTT, 2mM EDTA, 10% glycerol, 1% TritonX in ddH<sub>2</sub>O). To quantify fecal oocyst shedding, 20mg fecal material was suspended in 1mL lysis buffer. Samples were shaken with glass beads for 5min, then combined in a 1:1 ratio with Nano-Glo® Luciferase solution (Promega, Ref N1150). A Promega GloMax plate reader was used to measure luminescence. For each experiment, comparators are described in the text and involve cage-to-cage comparisons. The numbers of mice per experimental group and experimental replicates are provided in the figure legends and no data sets were excluded. No randomization or blinding strategies were employed.

### Histology

For histological analysis of the small intestine, tissue from the distal third of the small intestine was flushed with 10% neutral buffered formalin (Sigma, St Louis, MO, USA), then ‘swiss-rolled’ and fixed overnight. Fixed samples were paraffin-embedded, sectioned, and stained with hematoxylin and eosin for detailed histologic evaluation. Slides were evaluated by a board-certified veterinary pathologist in a blinded fashion for quantitative measurements of number of parasites, villus/crypt architectural features and inflammatory infiltrates, and semi-quantitative scores for villus epithelium lesions as previously described (82).

### Cytokine neutralization and measurement

To neutralize IFN- $\gamma$ , 1mg anti-IFN- $\gamma$  (XMG1.2, BioXcell Cat #: BE0055) was given intraperitoneally (i.p.) 1 day prior and 2 days post infection with maCp. To also neutralize IL-12 or IL-18, 2mg anti-IL-12p40 (C17.8 BioXcell Cat #: BE0051) or 1mg anti-IL-18 (YIGIF74-1G7, BioXcell Cat #: BE0237) were given i.p. on days -4, -1, and 1, while anti-IFN- $\gamma$  was given on day -2 and day 2. Intestinal IFN- $\gamma$  levels were assessed from 5mm biopsy punches that were incubated in complete RPMI at 37°C for 24 hours. Clear, flat-bottom 96-well plates (Immunulon 4 HBX) were coated with 0.25 $\mu$ g/mL anti-IFN- $\gamma$  (AN-18, Invitrogen Ref #:14-7313-85) at 4°C overnight. Samples were added and IFN- $\gamma$  left to bind at 37°C for 2 hours. 0.25 $\mu$ g/mL biotinylated anti-IFN- $\gamma$  (R4-6A2, eBioscience Ref #: 13-7312-85) in PBS with 2.5% FBS and 0.05% Tween was added for 1 hour at room temperature, followed by peroxidase-labeled streptavidin for 30 minutes. Finally, KPL ABTS® peroxidase substrate (SeraCare Cat #: 5120-0041) was applied for detection.

## Flow cytometry and cell sorting

Single-cell suspensions were prepared from intestinal sections by shaking diced tissue at 37°C for 25 minutes in Hank's Balanced Salt Solution with 5 mM EDTA and 1 mM DTT. Cell pellets were then passed through 70 µm and 40 µm filters. Cells were surface stained using the following fluorochrome-conjugated Abs: anti-EpCAM (G8.8), anti-CD45.2 (104), anti-CD19 (MB19-1), anti-Thy1.1 (HIS51), anti-CD27 (LG.7F9), anti-RORγt (B2D) and anti-γδ TCR (eBioGL3) from eBioscience; anti-NK1.1 (PK136), anti-NKp46 (29A1.4), anti-CD3 (17A2), anti-CD127 (SB/199), anti-CD49a (HMA1), anti-T-bet (4B10) and anti-CD8β (YTS156.7.7) from BioLegend; anti-Eomes (Dan11mag) and Live/Dead Aqua from Invitrogen; and anti-CD49b (DX5) from BD Biosciences. Data were collected on a LSRFortessa (BD Biosciences) and analyzed with FlowJo v10 software (TreeStar). Cell sorting was performed on a BD FACS Jazz (BD Biosciences).

## RNA-seq and gene enrichment analysis

RNAseq reads were pseudo-aligned to the Ensembl *Mus musculus* reference transcriptome v79 using Kallisto v0.44.0 (83). In R, transcripts were collapsed to genes using Bioconductor tximport (84), and differentially expressed genes were identified using Limma-Voom (85, 86). Gene set enrichment analysis was performed using the GSEA software and the annotated gene sets of the Molecular Signatures Database (MSigDB) (87, 88). From the GSEA output, enrichment maps were generated to provide a visual representation of gene set overlap using Cytoscape v3.8.2 (89). Data and analyses have been deposited to the GEO repository (GSE168680). One sample was excluded for failing quality control.

## Statistics

Statistical significance was calculated using the unpaired Student's t-test for comparing 2 groups, or ANOVA followed by multiple comparisons for comparing groups of 3 or more. Analyses were performed using GraphPad Prism v.9.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGMENTS

This work was supported in part by funding from the Bill and Melinda Gates Foundation (OPP1183177) to BS and through grants from the U.S. National Institutes of Health to BS & CAH (R01AI148249), R21 AI139895 to CAH, and fellowships and career awards to JG (T32AI055400), AS (K99AI137442), and AG (T32A1055400) and a Swiss National Science Foundation fellowship (191774) to SB. We would also like to acknowledge the Commonwealth of Pennsylvania.

## References

1. Collaborators GBDCM. Global, regional, national, and selected subnational levels of stillbirths, neonatal, infant, and under-5 mortality, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1725–74. [PubMed: 27733285]

2. Birchenough GM, Johansson ME, Gustafsson JK, Bergstrom JH, Hansson GC. New developments in goblet cell mucus secretion and function. *Mucosal Immunol.* 2015;8(4):712–9. [PubMed: 25872481]
3. von Moltke J, Ji M, Liang HE, Locksley RM. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature.* 2016;529(7585):221–5. [PubMed: 26675736]
4. Adolph TE, Tomczak MF, Niederreiter L, Ko HJ, Bock J, Martinez-Naves E, et al. Paneth cells as a site of origin for intestinal inflammation. *Nature.* 2013;503(7475):272–6. [PubMed: 24089213]
5. Peterson LW, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol.* 2014;14(3):141–53. [PubMed: 24566914]
6. Cliffe LJ, Potten CS, Booth CE, Grecnis RK. An increase in epithelial cell apoptosis is associated with chronic intestinal nematode infection. *Infect Immun.* 2007;75(4):1556–64. [PubMed: 17242061]
7. Gerbe F, Sidot E, Smyth DJ, Ohmoto M, Matsumoto I, Dardalhon V, et al. Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature.* 2016;529(7585):226–30. [PubMed: 26762460]
8. Nadsombati MS, McGinty JW, Lyons-Cohen MR, Jaffe JB, DiPeso L, Schneider C, et al. Detection of Succinate by Intestinal Tuft Cells Triggers a Type 2 Innate Immune Circuit. *Immunity.* 2018;49(1):33–41 e7. [PubMed: 30021144]
9. Schneider C, O'Leary CE, von Moltke J, Liang HE, Ang QY, Turnbaugh PJ, et al. A Metabolite-Triggered Tuft Cell-ILC2 Circuit Drives Small Intestinal Remodeling. *Cell.* 2018;174(2):271–84 e14. [PubMed: 29887373]
10. Nusse YM, Savage AK, Marangoni P, Rosendahl-Huber AKM, Landman TA, de Sauvage FJ, et al. Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche. *Nature.* 2018;559(7712):109–13. [PubMed: 29950724]
11. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet.* 2013;382(9888):209–22. [PubMed: 23680352]
12. Checkley W, White AC Jr., Jaganath D, Arrowood MJ, Chalmers RM, Chen XM, et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect Dis.* 2015;15(1):85–94. [PubMed: 25278220]
13. Platts-Mills JA, Babji S, Bodhidatta L, Gratz J, Haque R, Havt A, et al. Pathogen-specific burdens of community diarrhoea in developing countries: a multisite birth cohort study (MAL-ED). *Lancet Glob Health.* 2015;3(9):e564–75. [PubMed: 26202075]
14. Khalil IA, Troeger C, Rao PC, Blacker BF, Brown A, Brewer TG, et al. Morbidity, mortality, and long-term consequences associated with diarrhoea from *Cryptosporidium* infection in children younger than 5 years: a meta-analysis study. *Lancet Glob Health.* 2018;6(7):e758–e68. [PubMed: 29903377]
15. Levy J, Espanol-Boren T, Thomas C, Fischer A, Tovo P, Bordigoni P, et al. Clinical spectrum of X-linked hyper-IgM syndrome. *J Pediatr.* 1997;131(1 Pt 1):47–54. [PubMed: 9255191]
16. Gomez Morales MA, Mele R, Ludovisi A, Bruschi F, Tosini F, Rigano R, et al. *Cryptosporidium parvum*-specific CD4 Th1 cells from sensitized donors responding to both fractionated and recombinant antigenic proteins. *Infect Immun.* 2004;72(3):1306–10. [PubMed: 14977932]
17. Hayward AR, Chmura K, Cosyns M. Interferon-gamma is required for innate immunity to *Cryptosporidium parvum* in mice. *J Infect Dis.* 2000;182(3):1001–4. [PubMed: 10950807]
18. White AC, Robinson P, Okhuysen PC, Lewis DE, Shahab I, Lahoti S, et al. Interferon-gamma expression in jejunal biopsies in experimental human cryptosporidiosis correlates with prior sensitization and control of oocyst excretion. *J Infect Dis.* 2000;181(2):701–9. [PubMed: 10669358]
19. Leav BA, Yoshida M, Rogers K, Cohen S, Godiwala N, Blumberg RS, et al. An early intestinal mucosal source of gamma interferon is associated with resistance to and control of *Cryptosporidium parvum* infection in mice. *Infect Immun.* 2005;73(12):8425–8. [PubMed: 16299343]


20. Tessema TS, Dauber E, Petry F. Adoptive transfer of protective immunity from *Cryptosporidium parvum*-infected interferon-gamma and interleukin-12-deficient mice to naive recipients. *Vaccine*. 2009;27(47):6575–81. [PubMed: 19717136]
21. Guerin A, Striepen B. The Biology of the Intestinal Intracellular Parasite *Cryptosporidium*. *Cell Host Microbe*. 2020;28(4):509–15. [PubMed: 33031769]
22. Ehigiator HN, McNair N, Mead JR. *Cryptosporidium parvum*: the contribution of Th1-inducing pathways to the resolution of infection in mice. *Exp Parasitol*. 2007;115(2):107–13. [PubMed: 16920103]
23. Bedi B, McNair NN, Mead JR. Dendritic cells play a role in host susceptibility to *Cryptosporidium parvum* infection. *Immunol Lett*. 2014;158(1–2):42–51. [PubMed: 24295591]
24. Potiron L, Lacroix-Lamande S, Marquis M, Levern Y, Fort G, Franceschini I, et al. Batf3-Dependent Intestinal Dendritic Cells Play a Critical Role in the Control of *Cryptosporidium parvum* Infection. *J Infect Dis*. 2019;219(6):925–35. [PubMed: 30203075]
25. Barakat FM, McDonald V, Di Santo JP, Korbel DS. Roles for NK cells and an NK cell-independent source of intestinal gamma interferon for innate immunity to *Cryptosporidium parvum* infection. *Infect Immun*. 2009;77(11):5044–9. [PubMed: 19687195]
26. Rohlman VC, Kuhls TL, Mosier DA, Crawford DL, Greenfield RA. *Cryptosporidium parvum* infection after abrogation of natural killer cell activity in normal and severe combined immunodeficiency mice. *J Parasitol*. 1993;79(2):295–7. [PubMed: 8459345]
27. Bedi B, McNair NN, Forster I, Mead JR. IL-18 cytokine levels modulate innate immune responses and cryptosporidiosis in mice. *J Eukaryot Microbiol*. 2015;62(1):44–50. [PubMed: 25155632]
28. Choudhry N, Petry F, van Rooijen N, McDonald V. A protective role for interleukin 18 in interferon gamma-mediated innate immunity to *Cryptosporidium parvum* that is independent of natural killer cells. *J Infect Dis*. 2012;206(1):117–24. [PubMed: 22517912]
29. McDonald V, Pollok RC, Dhaliwal W, Naik S, Farthing MJ, Bajaj-Elliott M. A potential role for interleukin-18 in inhibition of the development of *Cryptosporidium parvum*. *Clin Exp Immunol*. 2006;145(3):555–62. [PubMed: 16907926]
30. McNair NN, Bedi C, Shayakhmetov DM, Arrowood MJ, Mead JR. Inflammasome components caspase-1 and adaptor protein apoptosis-associated speck-like proteins are important in resistance to *Cryptosporidium parvum*. *Microbes Infect*. 2018;20(6):369–75. [PubMed: 29842985]
31. Vinayak S, Pawlowic MC, Sateriale A, Brooks CF, Studstill CJ, Bar-Peled Y, et al. Genetic modification of the diarrhoeal pathogen *Cryptosporidium parvum*. *Nature*. 2015;523(7561):477–80. [PubMed: 26176919]
32. Sateriale A, Gullicksrud JA, Engiles JB, McLeod BI, Kugler EM, Henao-Mejia J, et al. The intestinal parasite *Cryptosporidium* is controlled by an enterocyte intrinsic inflammasome that depends on NLRP6. *Proc Natl Acad Sci U S A*. 2021;118(2).
33. Ungar BL, Kao TC, Burris JA, Finkelman FD. *Cryptosporidium* infection in an adult mouse model. Independent roles for IFN-gamma and CD4+ T lymphocytes in protective immunity. *J Immunol*. 1991;147(3):1014–22. [PubMed: 1677668]
34. Harrington LE, Janowski KM, Oliver JR, Zajac AJ, Weaver CT. Memory CD4 T cells emerge from effector T-cell progenitors. *Nature*. 2008;452(7185):356–60. [PubMed: 18322463]
35. Takeda K, Tsutsui H, Yoshimoto T, Adachi O, Yoshida N, Kishimoto T, et al. Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity*. 1998;8(3):383–90. [PubMed: 9529155]
36. Fuchs A, Vermi W, Lee JS, Lonardi S, Gilfillan S, Newberry RD, et al. Intraepithelial type 1 innate lymphoid cells are a unique subset of IL-12- and IL-15-responsive IFN-gamma-producing cells. *Immunity*. 2013;38(4):769–81. [PubMed: 23453631]
37. Hunter CA, Timans J, Pisacane P, Menon S, Cai G, Walker W, et al. Comparison of the effects of interleukin-1 alpha, interleukin-1 beta and interferon-gamma-inducing factor on the production of interferon-gamma by natural killer. *Eur J Immunol*. 1997;27(11):2787–92. [PubMed: 9394800]
38. Harrison OJ, Srinivasan N, Pott J, Schiering C, Krausgruber T, Ilott NE, et al. Epithelial-derived IL-18 regulates Th17 cell differentiation and Foxp3(+) Treg cell function in the intestine. *Mucosal Immunol*. 2015;8(6):1226–36. [PubMed: 25736457]

39. Laurent F, Lacroix-Lamande S. Innate immune responses play a key role in controlling infection of the intestinal epithelium by *Cryptosporidium*. *Int J Parasitol*. 2017;47(12):711–21. [PubMed: 28893638]
40. Rath E, Moschetta A, Haller D. Mitochondrial function - gatekeeper of intestinal epithelial cell homeostasis. *Nat Rev Gastroenterol Hepatol*. 2018;15(8):497–516. [PubMed: 29844587]
41. Moon Y. Mucosal injuries due to ribosome-inactivating stress and the compensatory responses of the intestinal epithelial barrier. *Toxins (Basel)*. 2011;3(10):1263–77. [PubMed: 22069695]
42. Saeij JP, Frickel EM. Exposing *Toxoplasma gondii* hiding inside the vacuole: a role for GBPs, autophagy and host cell death. *Curr Opin Microbiol*. 2017;40:72–80. [PubMed: 29141239]
43. Yamamoto M, Okuyama M, Ma JS, Kimura T, Kamiyama N, Saiga H, et al. A cluster of interferon-gamma-inducible p65 GTPases plays a critical role in host defense against *Toxoplasma gondii*. *Immunity*. 2012;37(2):302–13. [PubMed: 22795875]
44. Liu B, Gulati AS, Cantillana V, Henry SC, Schmidt EA, Daniell X, et al. *Irgm1*-deficient mice exhibit Paneth cell abnormalities and increased susceptibility to acute intestinal inflammation. *Am J Physiol Gastrointest Liver Physiol*. 2013;305(8):G573–84. [PubMed: 23989005]
45. Maric-Biresev J, Hunn JP, Krut O, Helms JB, Martens S, Howard JC. Loss of the interferon-gamma-inducible regulatory immunity-related GTPase (IRG), *Irgm1*, causes activation of effector IRG proteins on lysosomes, damaging lysosomal function and predicting the dramatic susceptibility of *Irgm1*-deficient mice to infection. *BMC Biol*. 2016;14:33. [PubMed: 27098192]
46. Pollok RC, Farthing MJ, Bajaj-Elliott M, Sanderson IR, McDonald V. Interferon gamma induces enterocyte resistance against infection by the intracellular pathogen *Cryptosporidium parvum*. *Gastroenterology*. 2001;120(1):99–107. [PubMed: 11208718]
47. Kim CH, Hashimoto-Hill S, Kim M. Migration and Tissue Tropism of Innate Lymphoid Cells. *Trends Immunol*. 2016;37(1):68–79. [PubMed: 26708278]
48. Gury-BenAri M, Thaiss CA, Serafini N, Winter DR, Giladi A, Lara-Astiaso D, et al. The Spectrum and Regulatory Landscape of Intestinal Innate Lymphoid Cells Are Shaped by the Microbiome. *Cell*. 2016;166(5):1231–46 e13. [PubMed: 27545347]
49. Park E, Patel S, Wang Q, Andhey P, Zaitsev K, Porter S, et al. *Toxoplasma gondii* infection drives conversion of NK cells into ILC1-like cells. *Elife*. 2019;8.
50. Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C, et al. ILC1 Confer Early Host Protection at Initial Sites of Viral Infection. *Cell*. 2017;171(4):795–808 e12. [PubMed: 29056343]
51. Flanigan T, Whalen C, Turner J, Soave R, Toerner J, Havlir D, et al. *Cryptosporidium* infection and CD4 counts. *Ann Intern Med*. 1992;116(10):840–2. [PubMed: 1348918]
52. Navin TR, Weber R, Vugia DJ, Rimland D, Roberts JM, Addiss DG, et al. Declining CD4+ T-lymphocyte counts are associated with increased risk of enteric parasitosis and chronic diarrhea: results of a 3-year longitudinal study. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1999;20(2):154–9. [PubMed: 10048902]
53. O'Hara SP, Small AJ, Nelson JB, Badley AD, Chen XM, Gores GJ, et al. The human immunodeficiency virus type 1 tat protein enhances *Cryptosporidium parvum*-induced apoptosis in cholangiocytes via a Fas ligand-dependent mechanism. *Infect Immun*. 2007;75(2):684–96. [PubMed: 17118988]
54. Urban JF Jr., Fayer R, Chen SJ, Gause WC, Gatley MK, Finkelman FD. IL-12 protects immunocompetent and immunodeficient neonatal mice against infection with *Cryptosporidium parvum*. *J Immunol*. 1996;156(1):263–8. [PubMed: 8598471]
55. Martinez-Lopez M, Iborra S, Conde-Garrosa R, Sancho D. *Batf3*-dependent CD103+ dendritic cells are major producers of IL-12 that drive local Th1 immunity against *Leishmania major* infection in mice. *Eur J Immunol*. 2015;45(1):119–29. [PubMed: 25312824]
56. Mashayekhi M, Sandau MM, Dunay IR, Frickel EM, Khan A, Goldszmid RS, et al. CD8alpha(+) dendritic cells are the critical source of interleukin-12 that controls acute infection by *Toxoplasma gondii* tachyzoites. *Immunity*. 2011;35(2):249–59. [PubMed: 21867928]

57. Navabi N, Whitt J, Wu SE, Woo V, Moncivaiz J, Jordan MB, et al. Epithelial Histone Deacetylase 3 Instructs Intestinal Immunity by Coordinating Local Lymphocyte Activation. *Cell Rep.* 2017;19(6):1165–75. [PubMed: 28494866]
58. Liu Z, Zaki MH, Vogel P, Gurung P, Finlay BB, Deng W, et al. Role of inflammasomes in host defense against *Citrobacter rodentium* infection. *J Biol Chem.* 2012;287(20):16955–64. [PubMed: 22461621]
59. Munoz M, Eidenschenk C, Ota N, Wong K, Lohmann U, Kuhl AA, et al. Interleukin-22 induces interleukin-18 expression from epithelial cells during intestinal infection. *Immunity.* 2015;42(2):321–31. [PubMed: 25680273]
60. Jarret A, Jackson R, Duizer C, Healy ME, Zhao J, Rone JM, et al. Enteric Nervous System-Derived IL-18 Orchestrates Mucosal Barrier Immunity. *Cell.* 2020;180(1):50–63 e12. [PubMed: 31923399]
61. Sellers M, Udaondo C, Moreno B, Martinez-Ales G, Diez J, Martinez L, et al. [Hirschsprung-associated enterocolitis: Observational study in a paediatric emergency care unit]. *An Pediatr (Barc).* 2018;88(6):329–34.
62. Teitelbaum DH, Caniano DA, Qualman SJ. The pathophysiology of Hirschsprung's-associated enterocolitis: importance of histologic correlates. *J Pediatr Surg.* 1989;24(12):1271–7. [PubMed: 2593059]
63. Barakat FM, McDonald V, Foster GR, Tovey MG, Korbel DS. *Cryptosporidium parvum* infection rapidly induces a protective innate immune response involving type I interferon. *J Infect Dis.* 2009;200(10):1548–55. [PubMed: 19821721]
64. Ferguson SH, Foster DM, Sherry B, Magness ST, Nielsen DM, Gookin JL. Interferon-lambda3 promotes epithelial defense and barrier function against *Cryptosporidium parvum* infection. *Cell Mol Gastroenterol Hepatol.* 2019.
65. Tretina K, Park ES, Maminska A, MacMicking JD. Interferon-induced guanylate-binding proteins: Guardians of host defense in health and disease. *J Exp Med.* 2019;216(3):482–500. [PubMed: 30755454]
66. Songhet P, Barthel M, Stecher B, Muller AJ, Kremer M, Hansson GC, et al. Stromal IFN-gammaR-signaling modulates goblet cell function during *Salmonella Typhimurium* infection. *PLoS One.* 2011;6(7):e22459. [PubMed: 21829463]
67. Monack DM, Bouley DM, Falkow S. *Salmonella typhimurium* persists within macrophages in the mesenteric lymph nodes of chronically infected *Nramp1*<sup>+/+</sup> mice and can be reactivated by IFN-gamma neutralization. *J Exp Med.* 2004;199(2):231–41. [PubMed: 14734525]
68. Lopez-Yglesias AH, Burger E, Camanzo E, Martin AT, Araujo AM, Kwok SF, et al. T-bet-dependent ILC1- and NK cell-derived IFN-gamma mediates cDC1-dependent host resistance against *Toxoplasma gondii*. *PLoS Pathog.* 2021;17(1):e1008299. [PubMed: 33465134]
69. Coers J, Brown HM, Hwang S, Taylor GA. Partners in anti-crime: how interferon-inducible GTPases and autophagy proteins team up in cell-intrinsic host defense. *Curr Opin Immunol.* 2018;54:93–101. [PubMed: 29986303]
70. Wilke G, Funkhouser-Jones LJ, Wang Y, Ravindran S, Wang Q, Beatty WL, et al. A Stem-Cell-Derived Platform Enables Complete *Cryptosporidium* Development In Vitro and Genetic Tractability. *Cell Host Microbe.* 2019.
71. Heo I, Dutta D, Schaefer DA, Iakobachvili N, Artegiani B, Sachs N, et al. Modelling *Cryptosporidium* infection in human small intestinal and lung organoids. *Nat Microbiol.* 2018;3(7):814–23. [PubMed: 29946163]
72. Dahan S, Roth-Walter F, Arnaboldi P, Agarwal S, Mayer L. Epithelia: lymphocyte interactions in the gut. *Immunol Rev.* 2007;215:243–53. [PubMed: 17291293]
73. Eberl G, Lochner M. The development of intestinal lymphoid tissues at the interface of self and microbiota. *Mucosal Immunol.* 2009;2(6):478–85. [PubMed: 19741595]
74. Zaph C, Troy AE, Taylor BC, Berman-Booty LD, Guild KJ, Du Y, et al. Epithelial-cell-intrinsic IKK-beta expression regulates intestinal immune homeostasis. *Nature.* 2007;446(7135):552–6. [PubMed: 17322906]
75. Mamareli P, Kruse F, Friedrich C, Smit N, Strowig T, Sparwasser T, et al. Epithelium-specific MyD88 signaling, but not DCs or macrophages, control acute intestinal infection with *Clostridium difficile*. *Eur J Immunol.* 2019;49(5):747–57. [PubMed: 30802297]

76. Rauch I, Deets KA, Ji DX, von Moltke J, Tenthorey JL, Lee AY, et al. NAIP-NLRC4 Inflammasomes Coordinate Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of Caspase-1 and -8. *Immunity*. 2017;46(4):649–59. [PubMed: 28410991]
77. Chatterjee M, Ip YT. Pathogenic stimulation of intestinal stem cell response in *Drosophila*. *J Cell Physiol*. 2009;220(3):664–71. [PubMed: 19452446]
78. Lee KZ, Lestrade M, Socha C, Schirmeier S, Schmitz A, Spenle C, et al. Enterocyte Purge and Rapid Recovery Is a Resilience Reaction of the Gut Epithelium to Pore-Forming Toxin Attack. *Cell Host Microbe*. 2016;20(6):716–30. [PubMed: 27889464]
79. Ayyaz A, Jasper H. Intestinal inflammation and stem cell homeostasis in aging *Drosophila melanogaster*. *Front Cell Infect Microbiol*. 2013;3:98. [PubMed: 24380076]
80. Klover PJ, Muller WJ, Robinson GW, Pfeiffer RM, Yamaji D, Hennighausen L. Loss of STAT1 from mouse mammary epithelium results in an increased Neu-induced tumor burden. *Neoplasia*. 2010;12(11):899–905. [PubMed: 21076615]
81. Hatton RD, Harrington LE, Luther RJ, Wakefield T, Janowski KM, Oliver JR, et al. A distal conserved sequence element controls *Irfng* gene expression by T cells and NK cells. *Immunity*. 2006;25(5):717–29. [PubMed: 17070076]
82. Sateriale A, Slapeta J, Baptista R, Engiles JB, Gullicksrud JA, Herbert GT, et al. A Genetically Tractable, Natural Mouse Model of Cryptosporidiosis Offers Insights into Host Protective Immunity. *Cell Host Microbe*. 2019;26(1):135–46 e5. [PubMed: 31231045]
83. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol*. 2016;34(5):525–7. [PubMed: 27043002]
84. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139–40. [PubMed: 19910308]
85. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-seq and microarray studies. *Nucleic Acids Res*. 2015;43(7):e47. [PubMed: 25605792]
86. Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*. 2014;15(2):R29. [PubMed: 24485249]
87. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, et al. PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet*. 2003;34(3):267–73. [PubMed: 12808457]
88. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545–50. [PubMed: 16199517]
89. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*. 2003;13(11):2498–504. [PubMed: 14597658]





**Figure 1. Localized innate IFN- $\gamma$  provides early protection from mouse-adapted *C. parvum* and pathology.**

WT mice were infected with  $10^5$  maCp-Nluc oocysts and treated with either  $\alpha$ -IFN- $\gamma$  or isotype control. (A) At 4dpi, two adjacent 5mm biopsies were taken along the length of the small and large intestines (D=duodenum, J=jejunum, I=ileum, Ce=cecum, C=colon). One biopsy was evaluated for nanoluciferase (top), the other was incubated at 37°C for 24h and IFN- $\gamma$  was measured from the supernatant. Graph depicts 1 representative mouse from n=3 for each group. (B) WT C57BL/6 and *Rag2*<sup>-/-</sup> mice were treated with isotype or  $\alpha$ -IFN- $\gamma$  and nanoluciferase was used to measure fecal oocyst shedding, n=4 per group. Similar results were observed in 2 additional experiments. (C) Representative mucosa


and (D–G) cumulative histology scoring from *Rag2*<sup>-/-</sup> mice 5dpi that shows a marked increase in cryptosporidia organisms (black arrows) infecting the villus enterocytes in treated mice versus controls with a progressive increase in epithelial dysplasia (arrowheads), reduced villus:crypt ratios with increased crypt depth (double-headed arrows) and crypt mitoses (white arrows); scale bars = 20µm. (H) *Rag2*<sup>-/-</sup> mice were infected with 10<sup>5</sup> maCp-Nluc oocysts and treated with α-IFN-γ at 50, 53, and 56dpi. Oocysts shedding was monitored throughout by nanoluciferase. n=4, representative from 3 experimental replicates. Bars denote mean ± SD. ANOVA followed by multiple comparisons were performed on cumulative pathology scores, \*\**p* 0.01, \*\*\**p* 0.001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 2. Innate lymphoid cells are critical components of protection against *C. parvum*.** C57BL/6 WT, *Rag2*<sup>-/-</sup> and *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice were infected and used for histological analysis or oocyst shedding. (A) Kinetics of fecal oocyst shedding; n=3–5, representative of 3 replicates comparing *Rag2*<sup>-/-</sup> and *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup>. (B–G) H&E staining of representative villi (upper panels) and crypts (lower panels), and cumulative histology scoring from WT, *Rag2*<sup>-/-</sup> and *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice at 5dpi that show a marked increase in *Cryptosporidium* organisms (black arrows) infecting the villus enterocytes in *Rag2*<sup>-/-</sup>*Il2rg*<sup>-/-</sup> mice versus WT or *Rag2*<sup>-/-</sup> mice as well as progressive villus epithelial dysplasia with attenuation (arrowheads), increases in crypt branching (double arrows) and crypt mitoses (blue arrows); scale bars = 20µm; n = 4–8 where each symbol denotes 1 mouse cumulative of 3 experiments. Bars indicate mean ± SD. n = 4–8 where each symbol denotes 1 mouse, cumulative of 3 experiments. Bars indicate mean ± SD. For histology scoring, t-tests were


used to compare uninfected and infected within each mouse strain, \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.0001$ .

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Figure 3. ILC1s are a critical source of innate IFN- $\gamma$ .**

*Ifng*-Thy1.1 KI mice were infected with mCherry-expressing maCp-Nluc and cells from the ileal epithelium were assessed by flow cytometry at 4dpi. (A) Representative flow plots and (B) summary bar graph of maCp-infected (mCherry<sup>+</sup>) intestinal epithelial cells, gated on Live CD45.2<sup>-</sup>EpCAM<sup>+</sup> cells. (C) Representative flow plots and (D) summary bar graph of *Ifng*<sup>+</sup> (Thy1.1<sup>+</sup>) innate lymphoid cells (NK1.1<sup>+</sup>NKp46<sup>+</sup>),  $n = 7-24$  from 4 experiments. (E) Representative flow plots and (F) summary bar graph of frequency of *Ifng*<sup>+</sup> (Thy1.1) cells from ILCs (seeing gating strategy from (C)) taken from intestinal epithelium (IE), Peyer's patches (PP), or mesenteric lymph nodes (mesLN). mesLN and PP were pooled from 2-4 mice, representative of 3 experiments. (G) Representative flow plots and (H) summary bar graph of T-bet and Eomes expression on *Ifng*<sup>+</sup> (Thy1.1<sup>+</sup>) or *Ifng*<sup>-</sup> (Thy1.1<sup>-</sup>) ILCs. (I) Representative flow plots and (J) summary bar graph of T-bet and ROR $\gamma$ t expression on *Ifng*<sup>+</sup> (Thy1.1<sup>+</sup>) or *Ifng*<sup>-</sup> (Thy1.1<sup>-</sup>) ILCs. All ILCs were gated on


Live CD45.2<sup>+</sup>CD3<sup>-</sup>CD19<sup>-</sup>NKp46<sup>+</sup>NK1.1<sup>+</sup> cells. Statistical significance between  $\alpha$ -IFN- $\gamma$  treated or untreated in B and D was determined by Student's t-test. Statistical significance in F was determined by one-way ANOVA and multiple comparisons. Student's t-tests were used to compare Thy1.1<sup>+</sup> and Thy1.1<sup>-</sup> in H and J. ns = not significant ( $p > 0.05$ ), \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\*\* $p < 0.0001$ .

Author Manuscript

Author Manuscript


Author Manuscript

Author Manuscript



**Figure 4. IL-12 and IL-18 stimulate IFN- $\gamma$  production by ILCs.**

(A) WT C57BL/6 mice or mice lacking IL-18 (*Il18*<sup>-/-</sup>), IL-12p40 (*Il12b*<sup>-/-</sup>) or IFN- $\gamma$  (*Ifng*<sup>-/-</sup>) were infected with maCp-Nluc and oocyst shedding in feces was monitored. Representative of 3 experimental replicates. (B) Control (Cre<sup>-</sup>*Il18*<sup>fl/fl</sup>), epithelial-specific IL-18-deficient (Villin<sup>Cre</sup> x *Il18*<sup>fl/fl</sup>; *Il18*<sup>IEC</sup>) and *Il18*<sup>-/-</sup> mice were infected with maCp-Nluc and oocyst shedding in feces was monitored. (C-G) *Rag2*<sup>-/-</sup> mice (C) or *Ifng*-Thy1.1 KI mice (D-G) were treated with  $\alpha$ -IL-18,  $\alpha$ -IL-12p40, or both and infected with maCp-Nluc. (C) Oocyst shedding in feces was monitored. (D) Representative flow plots and (E) summary bar graph of maCp-infected (mCherry<sup>+</sup>) IECs 4dpi. (F) Representative flow plots and (G) summary bar graph of *Ifng*<sup>+</sup> (Thy1.1<sup>+</sup>) ILCs 4dpi (gating same as Figure 3). n = 8–11 from 3 experiments. Statistical significance was determined by one-way ANOVA and multiple comparisons; ns = not significant ( $p > 0.05$ ) \*\*\*\* $p < 0.0001$ .



**Figure 5. IFN- $\gamma$ -mediated protection is dependent on enterocyte expression of STAT1.** (A–B)  $Cre^{-}Stat1^{fl/fl}$  (Control),  $Cd11c^{Cre}Stat1^{fl/fl}$  ( $Stat1^{DC}$ ) and  $Lyz2^{Cre}Stat1^{fl/fl}$  ( $Stat1^{MΦ}$ ),  $Villin^{CreERT2-Stat1^{fl/fl}}$  ( $Stat1^{IEC}$ ) and  $Stat1^{-/-}$  mice were infected with maCp-Nluc and oocyst shedding was monitored in feces. (C–F) IEC from naïve mice or mice infected with maCp-mCherry for 3 or 5 days were FACS sorted and used for RNA-sequencing analysis. (C) Volcano plot with IFN signature genes marked in red. DEGs that were significantly upregulated ( $p = 0.05$ ) in IECs from maCp-mCherry-infected mice are labeled. (D–E) Gene set enrichment analysis highlighting an IFN- $\gamma$  signature (D), a cluster of IFN pathways (E). (F) Heatmap of pertinent IFN-stimulated genes. (G) WT C57BL/6,  $Ido1^{-/-}$ ,  $GBP^{chr3}$ , and  $Irgm1/m3^{-/-}$  mice were infected with  $5 \times 10^4$  maCp-Nluc oocysts and fecal shedding of oocysts was monitored. Representative of 3 experimental replicates.