



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

REFERENCES

1. Xierali IM, Castillo-Page L, Zhang K, Gampfer KR, Nivet MA. AMlast page: the urgency of physician workforce diversity. *Acad Med* 2014;89:1192.
2. Amalba A, Abantanga FA, Scherpier AJ, van Mook WN. Community-based education: the influence of role modeling on career choice and practice location. *Med Teach* 2017;39:174–80.
3. Association of American Medical Colleges. Faculty roster: U.S. medical school faculty. Available at: <https://www.aamc.org/data-reports/faculty-institutions/report/famous>. Accessed May 10, 2021.
4. Accreditation Council for Graduate Medical Education. Data resource book, academic year 2020-2021. 2020. Available at: https://www.acgme.org/globalassets/pfassets/publicationsbooks/2020-2021_acgme_databook_document.pdf. Accessed October 24, 2021.

5. United States Census Bureau. The 2019 American community survey 5-year Public Use Microdata Sample (PUMS) files. Available at: <https://census.gov/data/developers/data-sets/acs-5year.html>. Accessed July 11, 2021.
6. Rayburn WF, Liu CQ, Elwell EC, Rogers RG. Diversity of physician faculty in obstetrics and gynecology. *J Reprod Med* 2016;61:22–6.

© 2022 Elsevier Inc. All rights reserved. <https://doi.org/10.1016/j.ajog.2022.02.007>

Impact of the COVID-19 pandemic on the incidence of prematurity: critical role of gestational age and environment



OBJECTIVE: Data on the incidence of prematurity during the COVID-19 pandemic are contradictory, with some reports showing decreased preterm births and others showing no effect.^{1,2} We propose that multiple biologic effects of SARS-CoV-2 infection and environmental changes during the pandemic exert competing effects on the preterm birth rate.

SARS-CoV-2 infection may increase preterm births by increasing preeclampsia and medically indicated preterm births,³ but its effects on spontaneous preterm births are unknown. Infection is the most commonly identified etiologic contributor to spontaneous preterm births at <28 weeks' gestation.⁴ Our preliminary data demonstrated that SARS-CoV-2 placental infection increases the expression of placenta-specific microRNA 519c, which protects against infection-induced preterm birth.⁵ Furthermore, environmental effects of the COVID-19 pandemic (lockdowns, quarantine, decreased travel) may decrease pathogen exposures that can trigger infection-

induced preterm birth.⁶ Therefore, the effects of the COVID-19 pandemic on the rates of prematurity likely reflect a balance between increased preeclampsia-induced preterm births and decreased infection-induced spontaneous preterm births.

New York was the first epicenter of the COVID-19 pandemic in the United States. The prevalence of SARS-CoV-2 infection exploded in March 2020, with a surge in the urban center. Lockdowns were initiated in all jurisdictions by mid-March. We compared the incidence of extreme prematurity (with and without preeclampsia) in 2020 with that of 2019 in a large cross-sectional study of hospitals in the New York City area.

STUDY DESIGN: Data were collected from urban medical centers (New York University Tisch Hospital, Lenox Hill Hospital, Montefiore Medical Center, Mount Sinai Hospital) and suburban medical centers (New York University Langone Hospital—Long Island, Long Island Jewish Medical Center,

TABLE 1
Number and rates of preterm deliveries in 2019 and 2020

Variables	2019		2020		Difference (95% CI)	P value
	N	Rate/1000	N	Rate/1000		
Deliveries	63,327	—	62,020	—	–1307	.0002
Delivery <35 wk	2255	35.6	2153	34.7	–0.9 (–1.9 to 0.1)	.08
Delivery 28–35 wk	1900	30.0	1864	30.1	0.1 (–0.9 to 1.0)	.9
Delivery <28 wk	355	5.6	289	4.7	–0.9 (–1.3 to –0.6)	<.0001
With preeclampsia	44	123.9	52	179.9	56.0 (28.0–83.0)	<.0008
COVID-19 positive	—	—	1870	30.15	—	—

CI, confidence interval.

Weinberger. Impact of the COVID-19 pandemic on the incidence of prematurity. *Am J Obstet Gynecol* 2022.

TABLE 2
Number and rate of preterm deliveries at urban and suburban hospitals

Variables	Urban		Suburban		Difference (95% CI)	P value
	2019	2020	2019	2020		
	N	Rate/1000	N	Rate/1000		
Total deliveries	35,466	—	33,471	—	—	—
Delivery <35 wk	1311	37.0	1269	37.9	0.9 (−0.5 to 2.4)	.2
Delivery 28–35 wk	1103	31.1	1091	32.6	1.5 (0.2–2.8)	.03
Delivery <28 wk	208	5.9	178	5.3	−0.5 (−1.1 to 0.0)	.06
With preeclampsia	21	101.0	28	157.3	56.3 (22.2–88.6)	.001
COVID-19 positive	—	—	1344	40.2	—	—
Suburban						
Total deliveries	27,861	—	28,549	—	—	—
Delivery <35 wk	944	33.9	884	30.9	−2.9 (−4.4 to −1.5)	.0001
Delivery 28–35 wk	847	30.4	812	28.4	−2.0 (−3.4 to −0.6)	.06
Delivery <28 wk	97	3.5	72	2.5	−1.4 (−2.0 to −0.8)	<.0001
With preeclampsia	23	156.5	24	216.2	59.8 (11.5–105.7)	.01163
COVID-19 positive	—	—	526	18.4	—	—

CI, confidence interval.

Weinberger. Impact of the COVID-19 pandemic on the incidence of prematurity. *Am J Obstet Gynecol* 2022.

North Shore University Hospital, Westchester Medical Center, and Stony Brook University Hospital) in New York. Rates of prematurity with and without preeclampsia were compared using the inverted skew-corrected score test for binomial distributions and Poisson regression.

RESULTS: Total deliveries decreased from 63,327 in 2019 to 62,020 in 2020, with a SARS-CoV-2 positivity rate of 30.1 per 1000 (Table 1). Extreme prematurity (<28 weeks' gestation) decreased from 5.6 to 4.7 per 1000 deliveries in 2020 ($P<.0001$), but the rate of moderate prematurity (28–35 weeks' gestation) did not change. Preeclampsia increased in 2020, from 123.9 to 179.9 per 1000 in mothers who delivered at <28 weeks' gestation.

At the suburban sites, overall prematurity dropped significantly (33.9 to 30.9/1000, $P=.0001$), primarily driven by decreased rates of extreme prematurity (3.5 to 2.5/1000, $P<.0001$). In contrast, at urban sites, total and extreme prematurity did not significantly decrease, and moderate prematurity increased (31.1 to 32.6/1000, $P=.03$). The rates of SARS-CoV-2 positivity were significantly higher in urban than in suburban sites (40.2 vs 18.4/1000, $P<.0001$) (Table 2).

CONCLUSION: In 2020, the COVID-19 pandemic in New York was associated with an overall decreased rate of extreme preterm births (<28 weeks' gestation) despite an increased rate of preeclampsia. This is consistent with the fact that extreme preterm births usually occur spontaneously

secondary to infection. Decreased intrapartum infection was likely related to both the environmental effects of lockdowns and possible biologic effects of SARS-CoV-2 infection on the placenta. In contrast, the rates of moderate prematurity (>28 weeks' gestation) did not decrease. This is most likely related to the relative impact of increased preeclampsia and other medical indications for preterm delivery. Conflicting results in earlier reports may be related to the diversity in the gestational ages and the contributing etiologic factors for preterm birth in the studied population.

The effects of the pandemic in New York were different in urban and suburban populations. Total prematurity decreased at suburban sites, driven by significant decreases in extreme prematurity. In contrast, at urban sites, total and extreme prematurity remained unchanged, and moderate prematurity increased. It is possible that lockdowns were less effective in the urban locations, as demonstrated by the increased prevalence of COVID-19 infection. Our findings highlight the importance of providing socioeconomic data in reports on the population effects of the pandemic. ■

Barry Weinberger, MD
 Division of Neonatology
 Department of Pediatrics
 Cohen Children's Medical Center, Northwell Health
 Donald and Barbara Zucker School of Medicine at Hofstra/
 Northwell
 New Hyde Park, NY

Jasmin Divers, PhD
Division of Health Services Research
Department of Foundations of Medicine
NYU Long Island School of Medicine
Mineola, NY

Deborah Campbell, MD, FAAP
Division of Neonatology
Department of Pediatrics
Children's Hospital at Montefiore
Albert Einstein College of Medicine
Weiler Hospital
Bronx, NY

Steven Ham, MD, MPH
Division of Neonatology
Department of Pediatrics
Cohen Children's Medical Center, Northwell Health
Donald and Barbara Zucker School of Medicine at Hofstra/
Northwell
New Hyde Park, NY

Courtney Juliano, MD
Division of Neonatology
Department of Pediatrics
Icahn School of Medicine at Mount Sinai Hospital
New York, NY

Dalibor Kurepa, MD
Division of Neonatology
Department of Pediatrics
Cohen Children's Medical Center, Northwell Health
Donald and Barbara Zucker School of Medicine at Hofstra/
Northwell
New Hyde Park, NY

Edmund Lagamma, MD
Division of Newborn Medicine
Department of Pediatrics
Maria Fareri Children's Hospital
Westchester Medical Center
New York Medical College
Valhalla, NY

Pradeep Mally, MD
Division of Neonatology
Department of Pediatrics
Hassenfeld Children's Hospital at NYU Langone
NYU Grossman School of Medicine
New York, NY

Suhas Nafday, MD
Division of Neonatology
Department of Pediatrics
Children's Hospital at Montefiore-Weiler
Bronx, NY

Sheri Nemerofsky, MD
Division of Neonatology

Department of Pediatrics
Children's Hospital at Montefiore
Albert Einstein College of Medicine
Wakefield Division
Bronx, NY

Shanthy Sridhar, MD
Division of Neonatology
Department of Pediatrics
Stony Brook Children's Hospital
Renaissance School of Medicine at Stony Brook University
Stony Brook, NY

Kim Williams, MBBS
Division of Neonatology
Department of Pediatrics
Lenox Hill Hospital, Northwell Health
Donald and Barbara Zucker School of Medicine at Hofstra/
Northwell
New York, NY

Nazeeh Hanna, MD
Division of Neonatology
Department of Pediatrics
NYU Langone Hospital—Long Island
New York University Long Island School of Medicine
259 First St.
Mineola, NY 11501
nazeehanna@gmail.com

The authors report no conflict of interest.

This work was supported partially by the the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, under grant number R21 HD100653-01.

REFERENCES

1. Harvey EM, McNeer E, McDonald MF, et al. Association of preterm birth rate with COVID-19 statewide stay-at-home orders in Tennessee. *JAMA Pediatr* 2021;175:635–7.
2. Handley SC, Mullin AM, Elovitz MA, et al. Changes in preterm birth phenotypes and stillbirth at 2 Philadelphia hospitals during the SARS-CoV-2 pandemic, March–June 2020. *JAMA* 2021;325:87–9.
3. Conde-Agudelo A, Romero R. SARS-CoV-2 infection during pregnancy and risk of preeclampsia: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2022;226:68–89.e3.
4. Goldenberg RL, Culhane JF, Iams JD, Romero R. Epidemiology and causes of preterm birth. *Lancet* 2008;371:75–84.
5. Tiozzo C, Bustoros M, Lin X, et al. Placental extracellular vesicles-associated microRNA-519c mediates endotoxin adaptation in pregnancy. *Am J Obstet Gynecol* 2021;225:681.e1–20.
6. Racicot K, Mor G. Risks associated with viral infections during pregnancy. *J Clin Invest* 2017;127:1591–9.

© 2022 Elsevier Inc. All rights reserved. <https://doi.org/10.1016/j.ajog.2022.02.028>