



# Glycine receptors caught between genome and proteome – functional implications of RNA editing and splicing

Pascal Legendre<sup>1,2,3</sup>, Benjamin Förster<sup>4,5</sup>, Rene Jüttner<sup>6</sup> and Jochen C. Meier<sup>5\*</sup>

<sup>1</sup> INSERM U952, Paris, France

<sup>2</sup> CNRS UMR 7224, Paris, France

<sup>3</sup> Université Pierre et Marie Curie, Paris, France

<sup>4</sup> Fachbereich Biologie, Chemie, Pharmazie, Freie Universität Berlin, Berlin, Germany

<sup>5</sup> RNA Editing and Hyperexcitability Disorders, Max Delbrück Center for Molecular Medicine, Berlin-Buch, Germany

<sup>6</sup> Developmental Neurobiology, Max Delbrück Center for Molecular Medicine, Berlin-Buch, Germany

## Edited by:

Jean-Michel Rigo, Université Hasselt, Belgium

## Reviewed by:

Hans-Georg Breiteringer, The German University in Cairo, Egypt

Jean-Michel Rigo, Université Hasselt, Belgium

## \*Correspondence:

Jochen C. Meier, Max Delbrück Center for Molecular Medicine, Robert-Rössle-Strasse 10, 13092 Berlin, Germany.  
e-mail: jochen.meier@mdc-berlin.de

Information processing in the brain requires a delicate balance between excitation and inhibition. Glycine receptors (GlyR) are involved in inhibitory mechanisms mainly at a synaptic level, but potential novel roles for these receptors recently emerged due to the discovery of posttranscriptional processing. *GLR* transcripts are edited through enzymatic modification of a single nucleotide leading to amino acid substitution within the neurotransmitter binding domain. RNA editing produces gain-of-function receptors well suited for generation and maintenance of tonic inhibition of neuronal excitability. As neuronal activity deprivation in early stages of development or in epileptic tissue is detrimental to neurons and because RNA editing of GlyR is up-regulated in temporal lobe epilepsy patients with a severe course of disease a pathophysiological role of these receptors emerges. This review contains a state-of-the-art discussion of (patho)physiological implications of GlyR RNA editing.

**Keywords:** RNA editing, RNA splicing, hippocampus, epilepsy, GABA

## GLYCINE RECEPTOR GENES AND SUBUNITS

Glycine receptors (GlyRs) are pentameric chloride (Cl<sup>-</sup>) channels. They contribute to inhibition of neuron firing in brain. Expression of GlyR genes (*GLRA1-4*) produces four different  $\alpha$  subunits that are capable of forming functional, homomeric channels (Betz and Laube, 2006; Lynch, 2009), while a single gene (*GLRB*) delivers neurons with GlyR  $\beta$  subunit mRNA. Besides modulation of ligand binding (Grudzinska et al., 2005) the GlyR  $\beta$  subunit mediates receptor stabilization at postsynaptic sites (Kirsch et al., 1991; Meyer et al., 1995; Meier et al., 2000, 2001; Meier and Grantyn, 2004). GlyR genes are located on several chromosomes in humans, with *GLRA1* on chromosome 5 (5q32), *GLRA2* on chromosome X (Xp22.1-p21.3) and *GLRA3* as well as *GLRB* on chromosome 4 (4q33-q34 and 4q31.3, respectively). *GLRA4* is a pseudogene (Simon et al., 2004) located on the X-chromosome, position Xq22.2. Expression of all other genes occurs in a wide range of brain regions (Lynch, 2009), and generally it is thought that GlyR  $\alpha 2$  expression predominates in the juvenile brain and declines with development (Lynch, 2009). However, at least in retina (Haverkamp et al., 2004) and in the hippocampus (Eichler et al., 2008) GlyR  $\alpha 2$  expression persists throughout development.

## FOR WHAT PURPOSE DO WE NEED RNA EDITING?

Through RNA editing the genetically encoded information can be modified. Enzymatic deamination of adenosine and cytidine is mediated by adenosine deaminases acting on RNA (ADAR) and apolipoprotein B mRNA editing complex (APOBEC) or activation induced deaminase (AID), respectively (Anant and Davidson, 2001; Seeburg and Hartner, 2003; Honjo et al., 2005). The resulting inosine (equivalent to guanosine) or uracil may then lead to amino acid substitutions in corresponding proteins, provided that the resulting

amino acid triplet codons engage different transfer RNAs. These enzyme machineries thus allow for diversification of the proteome or even correction of the genome code. Although the role of diversification by cytidine-to-uracil (C-to-U) RNA editing in generation of immunoglobulin variability is under debate (Honjo et al., 2005) the critical impact of RNA editing on correction of genome codes is well established. For example, the permeability of glutamate receptors for calcium ions increases when adenosine-to-inosine (A-to-I) RNA editing is prohibited, resulting in severe epilepsy (Brusa et al., 1995). Therefore in this case, the A-to-I RNA editing machinery is required for maintenance of physiological brain state through correction of unwanted genome codes (Seeburg and Hartner, 2003). The number of identified RNA-edited transcripts steadily increases, and for example serotonin receptors, potassium channels, GABA and glycine receptors were recently included in this register (Gurevich et al., 2002; Bhalla et al., 2004; Buckingham et al., 2005; Meier et al., 2005).

## GlyR C-TO-U RNA EDITING IS NOT SPECIES-SPECIFIC

A cDNA clone corresponding to GlyR  $\alpha 3^{185L}$  was originally isolated from *Sprague Dawley* rat brain, and molecular analysis revealed the involvement of C-to-U RNA editing in proline-to-leucine substitution at position 185 of the mature GlyR  $\alpha 3$  protein (Meier et al., 2005). Molecular analysis of hippocampi from pharmacoresistant temporal lobe epilepsy (TLE) patients further revealed expression of mRNAs coding for the high affinity GlyR  $\alpha 3^{185L}$  variant (Eichler et al., 2008). In addition, *GLRA2* transcripts of GlyR  $\alpha 2^{192L}$  were found in these patients. Sequencing of corresponding exons demonstrated lack of genomic  $\alpha 2^{192L}$  codons, again supporting the involvement of C-to-U RNA editing in proline-to-leucine substitution in

TLE patients (Eichler et al., 2008). However, the amount of RNA-edited *GLR* transcripts was not constant between TLE patients, but increased according to the frequency of secondarily generalized tonic-clonic seizures or the degree of hippocampal sclerosis. That hippocampal sclerosis was associated with increased amounts of RNA-edited *GLRA2/3* transcripts indicates a pathophysiological role of high affinity GlyRs in human hippocampus, which already was suggested by our data on experimentally induced brain lesion (Meier et al., 2005). Consequently, in TLE patients without hippocampal sclerosis the amount of RNA-edited *GLR* transcripts was very low, particularly in case of GlyR  $\alpha 3$  (below 1% of *GLRA3* transcripts). This renders quantification of RNA-edited transcripts rather difficult (Nakae et al., 2008). However, our methods for quantification were recently shown to be suitable for detection of RNA-edited messenger fractions below 1% (Eichler et al., 2008).

### LEUCINE AT POSITIONS 192 ( $\alpha 2$ ) AND 185 ( $\alpha 1$ AND $\alpha 3$ ) IS RESPONSIBLE FOR HIGH APPARENT GlyR AGONIST AFFINITY


GlyRs are members of the ligand-gated ion channel (LGIC) superfamily, which among others comprises A- and C-type GABA receptors ( $GABA_{A/C}R$ ) and the nicotinic acetylcholine receptor (nAChR). All members share common structural features, particularly in the ligand binding domain (Celie et al., 2005). In fact, all GlyR subunits and  $GABA_A \beta 1-3$  contain proline at the position corresponding to amino acid 185 of the mature signal-peptide cleaved  $\alpha 3$  protein.

However,  $GABA_A$ Rs that require a high agonist affinity because of their involvement in tonic inhibition of neuronal excitability don't have proline at this position. Instead,  $GABA_A$ Rs containing  $\alpha 4$ ,  $\alpha 6$  or  $\delta$  subunits are equipped with aliphatic hydrophobic neutral amino acids (such as alanine, leucine or valine) (Meier et al., 2005), and these receptors are well known to respond to low neuronal ambient GABA with tonic chloride currents (Mody, 2001; Wisden et al., 2002; Caraiscos et al., 2004; Mody and Pearce, 2004). Therefore, amino acid substitution at this position of the ligand binding domain should increase the apparent GlyR agonist affinity. Indeed, proline-to-leucine substitution by C-to-U editing of *GLRA3* transcripts was shown to increase the apparent agonist affinity (Meier et al., 2005). Similarly, editing of *GLRA2* transcripts at the corresponding position (192) produces receptors with increased apparent agonist affinity (Eichler et al., 2008).

To further emphasize the critical role of position 185/192 in determining apparent receptor agonist affinity dose-response curves were obtained from HEK293 cells expressing GlyR  $\alpha 1^{185P}$  or  $\alpha 1^{185L}$  (Figure 1). The short GlyR  $\alpha 1$  splice variant lacking insert SPMLNLFQ (Malosio et al., 1991) was used throughout. Again, compared with GlyR  $\alpha 1^{185P}$ , the apparent affinity of  $\alpha 1^{185L}$  channels was increased several fold [ $EC_{50(\text{Glycine})}$  ( $\mu\text{M}$ )  $\alpha 1^{185L}$ :  $17.7 \pm 1.5$  vs.  $\alpha 1^{185P}$ :  $85.3 \pm 4.4$ ,  $EC_{50(\text{Taurine})}$  ( $\mu\text{M}$ )  $\alpha 1^{185L}$ :  $67.0 \pm 6.8$  vs.  $\alpha 1^{185P}$ :  $378 \pm 33$ , mean  $\pm$  SD].

It should be noted here that the apparent affinity of GlyRs does not seem to be influenced by RNA splicing because GlyR  $\alpha 1$  with insertion ( $\alpha 1^{\text{ins}}$ ) (Malosio et al., 1991) exhibited similar current profiles (not shown). In addition, compared with  $\alpha 3K$ , the long splice variant of  $\alpha 3$ -GlyRs ( $\alpha 3L$ ) (Nikolic et al., 1998) also had increased apparent glycine affinity when RNA-edited (Figure 2).


Altogether, proline-to-leucine substitution at positions 192 ( $\alpha 2$ ) and 185 ( $\alpha 1$  and  $\alpha 3$ ) confers high apparent agonist affinities to GlyRs. On the contrary, GlyR apparent agonist affinity is not regulated by



RNA splicing, except for GlyR  $\alpha 2$  where changes in the apparent affinity were observed upon RNA splicing in the region coding for the N-terminal domain (Miller et al., 2004; Eichler et al., 2008).

### HOW RNA EDITING CAN CHANGE RECEPTOR-CHANNEL KINETICS, A PERSPECTIVE


RNA editing involves a single amino acid substitution in the N-terminal GlyR domain. The main consequence of RNA editing is an increase in the apparent affinities ( $EC_{50}$ ) of GlyRs for glycine and taurine. RNA editing does not modify GlyR desensitization kinetics (Meier et al., 2005), which is consistent with previous studies showing that receptor desensitization kinetics are mainly affected by change in amino acid sequences in the TM3-4 loop (Legendre, 2001; Breitinger and Becker, 2002; Breitinger et al., 2002). A single amino acid substitution in the N-terminal GlyR domain at positions 192 ( $\alpha 2$ ) or 185 ( $\alpha 1$  and  $\alpha 3$ ) could result in a change in the transduction process between agonist binding and channel gating, but it cannot be excluded at this stage that proline-to-leucine



substitution modifies the conformation of the N-terminal domain and hence the access of the ligand to the binding pocket. Changes in  $EC_{50}$  values do therefore not necessarily imply changes in the affinity of the receptor for its agonist. Effectively, in the simplest case, the  $EC_{50}$  depends both on the agonist dissociation constant ( $K_d$ ) and on the receptor-channel efficacy ( $E$ ), and  $EC_{50} = K_d / (1 + E)$  [for a more detailed explanation see (Colquhoun, 1998; Legendre, 2001)]. Several models have been proposed to describe GlyR activation kinetics, e.g. the “reluctant gating model” (Legendre, 1998) and the “flip model” (Burzomato et al., 2004). The flip model was recently extended to nAChR (Lape et al., 2008), although it has partially been challenged recently (Mukhtasimova et al., 2009). In any case, these studies demonstrate that receptor channels can have intermediate conformational states between binding steps and opening states, which must be taken into account when interpreting changes in  $EC_{50}$ . Accordingly, the best way to determine how GlyR RNA editing influences their apparent glycine affinity will be to compare activation kinetics of regular and RNA-edited GlyRs at the single channel level, as was already done in case of the hyperkplexia amino acid substitution  $\alpha 1^{K276E}$  (Lewis et al., 1998), the hyperkplexia amino acid substitution  $\alpha 1^{Q266H}$  (Moorhouse et al., 1999) and the spasmodic amino acid substitution  $\alpha 1^{A52S}$  (Plested et al., 2007). Nonetheless, the net outcome of proline-to-leucine substitution is a gain-of-function GlyR.

### CONFORMATIONAL IMPLICATIONS OF GlyR RNA EDITING

The crystal structure of the acetylcholine binding protein (AChBP) was recently determined at high resolution (Celie et al., 2005). Because members of the LGIC superfamily have a common ancestor (Celie et al., 2005) we have used the AChBP structure as a model for the ligand binding domain of GlyR. At least from the point of view of intramolecular location of  $\beta$ -sheets a projection



of GlyR sequences on AChBP protein structure can be made (Grenningloh et al., 1987). According to these thoughts proline 185/192 of GlyR  $\alpha$  subunits is located directly at the beginning of  $\beta$ -sheet 9 (Figure 3,  $\beta 9$ ), a position that is close to the plasma membrane where these receptors enter the cytosol through transmembrane domain (TM) 1 (Figure 3, C-terminus). The glycine binding pocket (Figure 3) is formed by two distant amino acid groups (Legendre, 2001), namely GY (160 and 161, respectively) and KYT (200, 202 and 204, respectively). These two domains are separated by  $\beta$ -sheets 7 and 10, and prolines 185/192 are part of the  $\beta 8/\beta 9$ -flanked connecting F-loop (Figure 3).

Further pursuing these considerations we have arguments that support a conformational role of proline-to-leucine substitution. First, leucine residues have bifurcated non-polar side chains that can mediate interaction with membrane lipids, while proline is frequently found in regions where the protein backbone has to make a sharp turn. In fact, increasing local hydrophobicity at position 250 (located between TM1 and TM2 close to the intracellular side of the plasma membrane) was shown to impact on receptor apparent affinity (Breitinger et al., 2001; Breitinger and Becker, 2002). In this case however, increased hydrophobicity decreased apparent GlyR affinity and favored rapid desensitization. Second, the disulfide bridge that connects  $\beta$ -sheets 6' and 7 (Figure 3) could serve as a hinge (Laube et al., 1993), potentially transmitting conformational rearrangement by proline-to-leucine substitution to the glycine binding pocket. Third, the glycine binding sequence KYT is located within the C-loop (Figure 3), which was shown

to undergo conformational change upon AChR agonist binding (Unwin et al., 2002), and fourth, F-loop dynamics are involved in benzodiazepine mediated GABA<sub>A</sub>R activation (Padgett and Lummis, 2008). More recently, the F-loop was involved in sensing conformational rearrangements upon ligand docking (Pless and Lynch, 2009). Therefore, we put forward the idea that structural rearrangement by proline-to-leucine substitution will impact on the accessibility of GlyR agonists to the C-loop binding pocket.

### GABA ACTIVATES RNA-EDITED GlyRs

Because proline-to-leucine substitution can cause structural rearrangement and therefore impact on accessibility of GlyR agonists to the C-loop binding pocket we have considered possible GABA responsiveness of RNA-edited GlyRs. Non-edited GlyRs can be activated by GABA at concentrations in the millimolar range (De Saint et al., 2001), which apparently precluded any physiological role for GABA on GlyRs. However, an elegant study recently demonstrated the opposite. Lu and colleagues (Lu et al., 2008) show that co-release of GABA and glycine accelerates glycinergic transmission by acting directly on GlyRs. This effect is explained by competition of GABA and glycine with the same binding site. The deactivation phase time constant of a synaptic event reflects the duration of bursts of channel openings (Legendre, 2001). Since burst duration partly depends on the dissociation rate constant of the agonist, a weak partial agonist will evoke bursts of short duration, which will in turn result in a decrease in the deactivation time constant of the synaptic response. It is tempting to speculate that postsynaptic GlyRs facing GABAergic nerve endings as observed in some CNS area such as hippocampus, and brain stem and spinal cord during development, could be activated by synaptic GABA release (Geiman et al., 2002; Levi et al., 2004; Meier and Grantyn, 2004; Muller et al., 2006). However, the


concentration of GABA, released into the synaptic cleft, must be high enough (>1 mM) to activate GlyRs. This limitation can be overcome as GlyRs acquire a higher agonist affinity if RNA-edited (Meier et al., 2005; Eichler et al., 2008). As GABA is a weak competitive partial agonist on GlyRs it is likely that an increase in the apparent affinity for glycine will also result in an increase in the apparent affinity for GABA. This is the case (Figure 4).

As shown in Figure 4 the threshold for GlyR activation by GABA is >0.3 mM in case of GlyR  $\alpha 3K^{185P}$ , whereas it is close to 0.1 mM for RNA-edited GlyR  $\alpha 3K^{185L}$ . Interestingly, application of 10 mM GABA (Figure 4B) on outside-out patches containing GlyR  $\alpha 3K^{185L}$  evoked responses with amplitudes representing ~80% of current amplitudes obtained with saturating glycine concentration (0.1 mM) (Meier et al., 2005). We did not test the effect of GABA concentrations above 10 mM, but according to the amplitude of GABA-evoked responses relative to current amplitudes obtained with 0.1 mM glycine, the apparent affinity of GlyR  $\alpha 3K^{185L}$  for GABA ( $EC_{50}$ ) is in the range of 2–3 mM (Figure 4B).

At GABAergic synapses the peak concentration of GABA molecules released per vesicle is likely to range from 0.5 to 1 mM (Jones and Westbrook, 1995). Accordingly, GlyR  $\alpha 3K^{185L}$  could be partially activated if located vis-à-vis GABAergic nerve endings. In fact, we have shown recently that GlyR  $\alpha 3K$  preferentially associates with GABAergic hippocampal nerve endings (Eichler et al., 2009). However, it still remains to be shown that RNA-edited  $\alpha 3K$  behaves similarly.

### A ROLE FOR $\alpha 3$ -GlyRs AT GLUTAMATERGIC SYNAPSES?

In addition to these findings, recent analysis of  $\alpha 3$ -GlyR distribution revealed an unexpected location of the long splice variant of these receptors ( $\alpha 3L$ ) adjacent to glutamatergic nerve endings





in hippocampal principle cell layers in adult mice (Eichler et al., 2009). It therefore is conceivable that activation of GlyRs can evoke local chloride currents at glutamatergic synapses, which will reduce the efficacy of these synapses. However, these GlyRs must be activated at the level of glutamatergic postsynaptic domains. Presence of glycine transporter 1 at hippocampal glutamatergic synapses (Cubelos et al., 2005) together with the observation of postsynaptic glutamatergic responses with recurrent small strychnine-sensitive components (Müller et al., 2007) can support this hypothesis, particularly in case of RNA-edited receptors (Meier et al., 2005; Eichler et al., 2008). Again, it remains to be shown whether RNA editing impacts on the subcellular distribution of these receptors.

### HIGH AFFINITY GlyR EFFECTIVELY INHIBITS NEURONAL ACTIVITY THROUGH SHUNT INHIBITION

It is now well established that GlyRs are highly mobile entities of the neuronal plasma membrane (Meier et al., 2001; Dahan et al., 2003), which implies that receptors have a certain probability to be encountered at non-synaptic locations.

In the hippocampus, functional glycinergic synapses have not been described, and the majority of  $\alpha 3$ -GlyR immunoreactivity indeed is found at non-synaptic places (Eichler et al., 2009). Furthermore, it has recently been shown that glycine uptake can regulate hippocampal network activity via GlyR-mediated non-synaptic tonic inhibition (Zhang et al., 2008), and the source of extracellular glycine was suggested to be glial cells (Zhang et al., 2008). A continuous activation of non-synaptic receptors requires a sufficient amount of extracellular glycine or taurine. However, in the central nervous system, the concentration of glycine is tightly regulated by glycine transporters (Gomez et al., 2003), and the residual glycine concentration in the extracellular space is likely to be close to 0.1  $\mu\text{M}$  in normal conditions (Roux and Supplisson, 2000).

Extracellular glycine can increase temporarily in case of sustained presynaptic activity (Roux and Supplisson, 2000; Supplisson and Roux, 2002), as for example in the epileptic tissue where hyper-synchronous high frequency network activity occurs (Fisher et al., 1992; Bragin et al., 2007). Accordingly, this might have important functional consequences, because non-synaptic slightly desensitizing GlyRs could be continuously activated as a result of temporarily altered GlyT activity (e.g. reverse transport) or even due to glycine spillover. Remarkably, the embryonic form of GlyRs, which is composed of five  $\alpha 2$  subunits and expressed before synaptogenesis displays functional properties adapted to non-synaptic release of the agonist (Mangin et al., 2003). Therefore, it can sustain a long lasting activation state when the extracellular glycine concentration increases above its activation threshold (Mangin et al., 2003).

Irrespective of the subunit, remarkable features of RNA-edited GlyRs are their high apparent affinities for glycine (Meier et al., 2005; Eichler et al., 2008). Effectively, these receptors can be activated by very low glycine concentrations. In case of GlyR  $\alpha 3^{185\text{L}}$ , 300 nM glycine already is sufficient for receptor activation, which in turn results in the occurrence of single channel openings. Independently of the chloride equilibrium potential, the activation of several GlyR  $\alpha 3^{185\text{L}}$  can evoke an inhibition of the neuronal activity. Effectively, the opening of several channels can lead to a decrease in the cell

input resistance, which will reduce the propagation of synaptic events and of action potentials, resulting in an inhibitory shunting process (Eichler et al., 2008).

### HIGH AFFINITY GlyRs ARE HARMFUL FOR HIPPOCAMPAL NEURONS WITH HIGH CHLORIDE EQUILIBRIUM POTENTIAL

Although GlyRs were reported to have anticonvulsant effects (Kirchner et al., 2003; Song et al., 2006) results obtained more recently (Eichler et al., 2008) point to a pathophysiological role of enhanced GlyR-mediated signaling, particularly in the context of high chloride equilibrium potential as is the case in TLE (Palma et al., 2006). Under these circumstances, neuron silencing triggers up-regulation of the ratio between glutamatergic and GABAergic nerve terminals and is associated with abnormal dendrite morphology (Lohmann et al., 2005; Eichler et al., 2008), both of which phenomena are reminiscent of the TLE histopathology (Loup et al., 2000; Stief et al., 2007). Most importantly, silenced hippocampal neurons were prone to die due to glutamate excitotoxicity, but amongst others could be rescued by expression of the potassium chloride cotransporter type 2 (KCC2) (Eichler et al., 2008). KCC2 expression provides neurons with an efficient chloride extrusion mechanism that decreases their chloride equilibrium potential (Lee et al., 2005). Although the mechanisms underlying this neuroprotective effect of KCC2 remained obscure our results support the previously postulated developmentally dichotomous effects of single cell activity deprivation on neuron integrity within an otherwise active neuronal network (Tao and Poo, 2005).

Furthermore considering altered *GLRA3* splicing in the hippocampus of TLE patients, favouring association of  $\alpha 3$ -GlyRs with hippocampal GABAergic synapses, the net outcome of anomalous posttranscriptional *GLRA3* processing is predicted to add to the pathophysiology of TLE. In a worst case scenario, provided that the chloride equilibrium potential is high enough (Rivera et al., 2004; Palma et al., 2006; Eichler et al., 2008), up-regulation of RNA-edited  $\alpha 3\text{K}$ -GlyRs (associated preferentially with GABAergic synapses) accompanied by down-regulation of RNA-edited  $\alpha 3\text{L}$ -GlyRs (associated preferentially with glutamatergic synapses) could increase glycinergic signalling at depolarizing GABAergic synapses while reducing inhibitory effects on glutamatergic postsynaptic currents.

### EXPERIMENTAL STRATEGIES FOR DETECTION OF RNA-EDITED GlyR PROTEIN

In any case, it is necessary to prove the existence of RNA-edited GlyR protein. However, this figured out to be rather difficult. Because of the position of leucine 185/192 at the interior of the channel pore close to the plasma membrane antibody accessibility is predicted to be low. Indeed, a potentially GlyR  $\alpha 3^{185\text{L}}$ -specific antibody obtained by immunization of guinea pigs with peptide EGLTLL**Q**FLLK (185L, underlined and bold) and cross-absorption against peptide EGLTLP**Q**FLLK (185P, underlined and bold) was not able to visualize RNA-edited GlyR  $\alpha 3^{185\text{L}}$  in transfected HEK293 (not shown). Even variation of fixation protocols and reagents did not ameliorate cellular protein detection. Therefore, it will be necessary to isolate a particular single channel conductance specific to RNA-edited GlyRs or any other differentiating factor in order to validate existence of these proteins

in neuronal tissue. It is conceivable to screen ligand databases for the presence of agonists or antagonists specific to RNA-edited GlyRs. So far, however, poor availability of reliable chloride indicators applicable to living cells was a rather rate limiting step in identification of novel ligands.

## CONCLUSION

RNA-edited GlyRs may open avenues to medication of excitability disorders (Eichler and Meier, 2008) because they could provide researchers and clinicians with a battery of new therapeutically relevant drugs, yet these enigmatic receptors still have to be captured at the protein level of investigation. In particular, a possible cell-type specific hippocampal expression has to be determined.

## SUPPORTING METHODS

### WHOLE-CELL PATCH-CLAMP RECORDINGS

Whole-cell patch-clamp recordings of HEK293 cells were performed 2–3 days after  $\text{Ca}^{2+}$ -phosphate transfection with  $\alpha 1$ -GlyR constructs. For electrophysiological recordings, the growth medium was replaced with a bath solution containing (in mM): 145 NaCl, 2.5 KCl, 20 HEPES, 10 glucose, 2  $\text{CaCl}_2$ , 1  $\text{MgCl}_2$ , adjusted to pH 7.3. All recordings were carried out at room temperature using patch-clamp amplifier EPC-9 (HEKA Elektronik, Lambrecht, Germany). Patch electrodes were fabricated from borosilicate glass capillaries tubing using a P-97 puller (Sutter Instruments, Novato, CA, USA). The pipette solution contained (in mM): 4 NaCl, 130 KCl, 5 EGTA, 10 HEPES, 10 glucose, 0.5  $\text{CaCl}_2$ , 4  $\text{MgCl}_2$ , adjusted to pH 7.3. The pipette-to-bath DC resistance of patch electrodes ranged from 2–4 M $\Omega$ . Recordings were performed at a holding potential of  $-70$  mV. Glycine or taurine was locally applied for 3 s via a gravity-driven 5-channel superfusion system. An interval of at least 60 s was left between successive agonist concentrations to allow for recovery from receptor desensitization. Different agonist concentrations were applied randomly. Electrophysiological signals were sampled at a rate of 5 kHz, filtered at 1 kHz and analyzed off-line using software TIDA 5.1 (HEKA Electronics, Lambrecht, Germany). To determine the  $\text{EC}_{50}$  value for agonists, peak currents obtained with different glycine or taurine concentrations were plotted and fitted to the Hill equation using software Origin 7.0 (OriginLab, Northampton, USA).

## REFERENCES

- Anant, S., and Davidson, N. O. (2001). Molecular mechanisms of apolipoprotein B mRNA editing. *Curr. Opin. Lipidol.* 12, 159–165.
- Betz, H., and Laube, B. (2006). Glycine receptors: recent insights into their structural organization and functional diversity. *J. Neurochem.* 97, 1600–1610.
- Bhalla, T., Rosenthal, J. J., Holmgren, M., and Reenan, R. (2004). Control of human potassium channel inactivation by editing of a small mRNA hairpin. *Nat. Struct. Mol. Biol.* 11, 950–956.
- Bragin, A., Claeys, P., Vonck, K., Van, R. D., Wilson, C., Boon, P., and Engel, J. Jr. (2007). Analysis of initial slow waves (ISWs) at the seizure onset in patients with drug resistant temporal lobe epilepsy. *Epilepsia* 48, 1883–1894.
- Breitinger, H. G., and Becker, C. M. (2002). The inhibitory glycine receptor—simple views of a complicated channel. *Chembiochem* 3, 1042–1052.
- Breitinger, H. G., Villmann, C., Becker, K., and Becker, C. M. (2001). Opposing effects of molecular volume and charge at the hyperekplexia site alpha 1 (P250) govern glycine receptor activation and desensitization. *J. Biol. Chem.* 276, 29657–29663.
- Breitinger, H. G., Villmann, C., Rennett, J., Ballhausen, D., and Becker, C. M. (2002). Hydroxylated residues influence desensitization behaviour of recombinant alpha3 glycine receptor channels. *J. Neurochem.* 83, 30–36.
- Brusa, R., Zimmermann, F., Koh, D. S., Feldmeyer, D., Gass, P., Seeburg, P. H., and Sprengel, R. (1995). Early-onset epilepsy and postnatal lethality associated with an editing-deficient GluR-B allele in mice. *Science* 270, 1677–1680.
- Buckingham, S. D., Biggin, P. C., Sattelle, B. M., Brown, L. A., and Sattelle, D. B. (2005). Insect GABA receptors: splicing, editing, and targeting by antiparasitics and insecticides. *Mol. Pharmacol.* 68, 942–951.
- Burzomato, V., Beato, M., Groot-Kormelink, P. J., Colquhoun, D., and Sivillotti, L. G. (2004). Single-channel behavior of heteromeric alpha1beta glycine receptors: an attempt to detect a conformational change before the channel opens. *J. Neurosci.* 24, 10924–10940.
- Caraiscos, V. B., Elliott, E. M., You, T., Cheng, V. Y., Belelli, D., Newell, J. G., Jackson, M. F., Lambert, J. J., Rosahl, T. W., Wafford, K. A., MacDonald, J. F., and Orser, B. A. (2004). Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by alpha5 subunit-containing gamma-aminobutyric acid

## OUTSIDE-OUT PATCH-CLAMP RECORDINGS

Outside-out patch-clamp recordings of HEK293 cells were performed 2 days after  $\text{Ca}^{2+}$ -phosphate transfection with  $\alpha 3\text{K}^{185\text{P}}$  or  $\alpha 3\text{K}^{185\text{L}}$  GlyR constructs. Cells were continuously perfused at room temperature (20–22°C) with bathing solution (2 ml/min) containing (in mM): NaCl 147, KCl 2.4,  $\text{CaCl}_2$  2,  $\text{MgCl}_2$  2, HEPES 10, glucose 10 (pH 7.2, osmolarity 320 mOsm). Patch-clamp electrodes (5–10 M $\Omega$ ) were pulled from thick-wall borosilicate glass (Harvard Apparatus, Kent, UK) using a Brown-Flaming puller (Sutter Instrument Co., Navato, USA). They were fire-polished and filled with (in mM): CsCl 130,  $\text{MgCl}_2$  4,  $\text{Na}_2\text{ATP}$  4, EGTA 10, HEPES 10 (pH 7.2, osmolarity 290 mOsm). Currents were recorded using an Axopatch 1D amplifier (Axon Instruments, Foster City, USA). Recordings were filtered at 10 kHz using an eight-pole Bessel filter (Frequency Devices, Haverhill, USA), sampled at 50 kHz and stored on a PC computer using pClamp software 10.1. (Axon Instruments, Foster City, USA). The membrane potential was held at  $-50$  mV throughout the experiment. Outside-out single-channel currents were evoked using a fast-flow operating system (Legendre, 1998). Control and drug solution were gravity-fed into two channels of a thin-wall glass theta tube (2-mm outer diameter; Hilgenberg, Malsfeld, Germany) pulled and broken to obtain a 200- $\mu\text{m}$  tip diameter. The outside-out patch was positioned 100  $\mu\text{m}$  away from the glass theta tube, one lumen of the application pipette was connected to reservoirs filled with solutions containing glycine or GABA. The solution exchange was performed by rapidly moving the solution interface across the tip of the patch pipette, using a piezo-electric translator (model P245.30, Physik Instrument, Waldbronn, Germany). Concentration steps of glycine or GABA were applied with an interval of  $\geq 10$  s. Exchange time solution ( $< 100$   $\mu\text{s}$ ) was determined before each set of experiments by monitoring the change in the liquid junction potential evoked by the application of a 10%-diluted control solution to the open tip of the patch pipette.

## ACKNOWLEDGMENTS

This study was supported by the DFG (ME2075/3-1 to JCM and RA424/5-1 to RJ, and Sonderforschungsbereich SFB-TR3 B5 to Jochen C. Meier) and the Helmholtz Association (VH-NG-246 to Jochen C. Meier), and Institut National de la Santé Et de la Recherche Médicale (INSERM to Pascal Legendre), and Centre de la Recherche Scientifique (CNRS to Pascal Legendre).

- type A receptors. *Proc. Natl. Acad. Sci. U.S.A.* 101, 3662–3667.
- Celie, P. H., Klaassen, R. V., Rossum-Fikkert, S. E., van Elk, R., van Nierop, P., Smit, A. B., and Sixma, T. K. (2005). Crystal structure of AChBP from *Bulinus truncatus* reveals the conserved structural scaffold and sites of variation in nicotinic acetylcholine receptors. *J. Biol. Chem.* 280, 26457–26466.
- Colquhoun, D. (1998). Binding, gating, affinity and efficacy: the interpretation of structure-activity relationships for agonists and of the effects of mutating receptors. *Br. J. Pharmacol.* 125, 924–947.
- Cubelos, B., Gimenez, C., and Zafra, F. (2005). Localization of the GLYT1 glycine transporter at glutamatergic synapses in the rat brain. *Cereb. Cortex* 15, 448–459.
- Dahan, M., Levi, S., Luccardini, C., Rostaing, P., Riveau, B., and Triller, A. (2003). Diffusion dynamics of glycine receptors revealed by single-quantum dot tracking. *Science* 302, 442–445.
- De Saint, J., David-Watine, B., Korn, H., and Gregostovski, P. (2001). Activation of human  $\alpha 1$  and  $\alpha 2$  homomeric glycine receptors by taurine and GABA. *J. Physiol.* 535, 741–755.
- Eichler, S. A., Forstera, B., Smolinsky, B., Jüttner, R., Lehmann, T. N., Fahling, M., Schwarz, G., Legendre, P., and Meier, J. C. (2009). Splice-specific roles of glycine receptor  $\alpha 3$  in the hippocampus. *Eur. J. Neurosci.* 30, 1077–1091.
- Eichler, S. A., Kirischuk, S., Jüttner, R., Schäfermeier, P. K., Legendre, P., Lehmann, T. N., Gloveli, T., Grantyn, R., and Meier, J. C. (2008). Glycinergic tonic inhibition of hippocampal neurons with depolarising GABAergic transmission elicits histopathological signs of temporal lobe epilepsy. *J. Cell. Mol. Med.* 12, 2848–2866.
- Eichler, S. A., and Meier, J. C. (2008). E-I balance and human diseases – from molecules to networking. *Front. Mol. Neurosci.* 1, 2. doi: 10.3389/neuro.02.002.2008.
- Fisher, R. S., Webber, W. R., Lesser, R. P., Arroyo, S., and Uematsu, S. (1992). High-frequency EEG activity at the start of seizures. *J. Clin. Neurophysiol.* 9, 441–448.
- Geiman, E. J., Zheng, W., Fritschy, J. M., and Alvarez, F. J. (2002). Glycine and GABA(A) receptor subunits on Renshaw cells: relationship with presynaptic neurotransmitters and postsynaptic gephyrin clusters. *J. Comp. Neurol.* 444, 275–289.
- Gomez, J., Ohno, K., and Betz, H. (2003). Glycine transporter isoforms in the mammalian central nervous system: structures, functions and therapeutic promises. *Curr. Opin. Drug Discov. Devel.* 6, 675–682.
- Grenningloh, G., Rienitz, A., Schmitt, B., Methfessel, C., Zensen, M., Beyreuther, K., Gundelfinger, E. D., and Betz, H. (1987). The strychnine-binding subunit of the glycine receptor shows homology with nicotinic acetylcholine receptors. *Nature* 328, 215–220.
- Grudzinska, J., Schemm, R., Haeger, S., Nicke, A., Schmalzing, G., Betz, H., and Laube, B. (2005). The Beta subunit determines the ligand binding properties of synaptic glycine receptors. *Neuron* 45, 727–739.
- Gurevich, I., Englander, M. T., Adlersberg, M., Siegal, N. B., and Schmauss, C. (2002). Modulation of serotonin 2C receptor editing by sustained changes in serotonergic neurotransmission. *J. Neurosci.* 22, 10529–10532.
- Haverkamp, S., Muller, U., Zeilhofer, H. U., Harvey, R. J., and Wässle, H. (2004). Diversity of glycine receptors in the mouse retina: localization of the  $\alpha 2$  subunit. *J. Comp. Neurol.* 477, 399–411.
- Honjo, T., Nagaoka, H., Shinkura, R., and Muramatsu, M. (2005). AID to overcome the limitations of genomic information. *Nat. Immunol.* 6, 655–661.
- Jones, M. V., and Westbrook, G. L. (1995). Desensitized states prolong GABA channel responses to brief agonist pulses. *Neuron* 15, 181–191.
- Kirchner, A., Breustedt, J., Rosche, B., Heinemann, U. F., and Schmieden, V. (2003). Effects of taurine and glycine on epileptiform activity induced by removal of Mg<sup>2+</sup> in combined rat entorhinal cortex-hippocampal slices. *Epilepsia* 44, 1145–1152.
- Kirsch, J., Langosch, D., Prior, P., Littauer, U. Z., Schmitt, B., and Betz, H. (1991). The 93-kDa glycine receptor-associated protein binds to tubulin. *J. Biol. Chem.* 266, 22242–22245.
- Lape, R., Colquhoun, D., and Sivilotti, L. G. (2008). On the nature of partial agonism in the nicotinic receptor superfamily. *Nature* 454, 722–727.
- Laube, B., Kuryatov, A., Kuhse, J., and Betz, H. (1993). Glycine-glutamate interactions at the NMDA receptor: role of cysteine residues. *FEBS Lett.* 335, 331–334.
- Lee, H., Chen, C. X., Liu, Y. J., Aizenman, E., and Kandler, K. (2005). KCC2 expression in immature rat cortical neurons is sufficient to switch the polarity of GABA responses. *Eur. J. Neurosci.* 21, 2593–2599.
- Legendre, P. (1998). A reluctant gating mode of glycine receptor channels determines the time course of inhibitory miniature synaptic events in zebrafish hindbrain neurons. *J. Neurosci.* 18, 2856–2870.
- Legendre, P. (2001). The glycinergic inhibitory synapse. *Cell Mol. Life Sci.* 58, 760–793.
- Levi, S., Logan, S. M., Tovar, K. R., and Craig, A. M. (2004). Gephyrin is critical for glycine receptor clustering but not for the formation of functional GABAergic synapses in hippocampal neurons. *J. Neurosci.* 24, 207–217.
- Lewis, T. M., Sivilotti, L. G., Colquhoun, D., Gardiner, R. M., Schoepfer, R., and Rees, M. (1998). Properties of human glycine receptors containing the hyperekplexia mutation  $\alpha 1(K276E)$ , expressed in *Xenopus* oocytes. *J. Physiol.* 507, 25–40.
- Lohmann, C., Finski, A., and Bonhoeffer, T. (2005). Local calcium transients regulate the spontaneous motility of dendritic filopodia. *Nat. Neurosci.* 8, 305–312.
- Loup, F., Wieser, H. G., Yonekawa, Y., Aguzzi, A., and Fritschy, J. M. (2000). Selective alterations in GABA<sub>A</sub> receptor subtypes in human temporal lobe epilepsy. *J. Neurosci.* 20, 5401–5419.
- Lu, T., Rubio, M. E., and Trussell, L. O. (2008). Glycinergic transmission shaped by the corelease of GABA in a mammalian auditory synapse. *Neuron* 57, 524–535.
- Lynch, J. W. (2009). Native glycine receptor subtypes and their physiological roles. *Neuropharmacology* 56, 303–309.
- Malosio, M. L., Grenningloh, G., Kuhse, J., Schmieden, V., Schmitt, B., Prior, P., and Betz, H. (1991). Alternative splicing generates two variants of the  $\alpha 1$  subunit of the inhibitory glycine receptor. *J. Biol. Chem.* 266, 2048–2053.
- Mangin, J. M., Baloul, M., Prado, D. C., Rogister, B., Rigo, J. M., and Legendre, P. (2003). Kinetics properties of the  $\alpha 2$  homo-oligomeric glycine receptor impairs a proper synaptic functioning. *J. Physiol. (Lond.)* 553, 369–386.
- Meier, J., and Grantyn, R. (2004). A gephyrin-related mechanism restraining glycine receptor anchoring at GABAergic synapses. *J. Neurosci.* 24, 1398–1405.
- Meier, J., Meunier-Durmort, C., Forest, C., Triller, A., and Vannier, C. (2000). Formation of glycine receptor clusters and their accumulation at synapses. *J. Cell Sci.* 113, 2783–2795.
- Meier, J., Vannier, C., Serge, A., Triller, A., and Choquet, D. (2001). Fast and reversible trapping of surface glycine receptors by gephyrin. *Nat. Neurosci.* 4, 253–260.
- Meier, J. C., Henneberger, C., Melnick, I., Racca, C., Harvey, R. J., Heinemann, U., Schmieden, V., and Grantyn, R. (2005). RNA editing produces glycine receptor  $\alpha 3P185L$  resulting in high agonist potency. *Nat. Neurosci.* 8, 736–744.
- Meyer, G., Kirsch, J., Betz, H., and Langosch, D. (1995). Identification of a gephyrin binding motif on the glycine receptor beta subunit. *Neuron* 15, 563–572.
- Miller, P. S., Harvey, R. J., and Smart, T. G. (2004). Differential agonist sensitivity of glycine receptor  $\alpha 2$  subunit splice variants. *Br. J. Pharmacol.* 143, 19–26.
- Mody, I. (2001). Distinguishing between GABA(A) receptors responsible for tonic and phasic conductances. *Neurochem. Res.* 26, 907–913.
- Mody, I., and Pearce, R. A. (2004). Diversity of inhibitory neurotransmission through GABA(A) receptors. *Trends Neurosci.* 27, 569–575.
- Moorhouse, A. J., Jacques, P., Barry, P. H., and Schofield, P. R. (1999). The startle disease mutation Q266H, in the second transmembrane domain of the human glycine receptor, impairs channel gating. *Mol. Pharmacol.* 55, 386–395.
- Mukhtasimova, N., Lee, W. Y., Wang, H. L., and Sine, S. M. (2009). Detection and trapping of intermediate states priming nicotinic receptor channel opening. *Nature* 459, 451–454.
- Müller, E., Bakkar, W., Legendre, P., Bergeron, R., and Martina, M. (2007). Site and mechanism of glycine release at glutamatergic synapse in mouse hippocampus. (Program No. 358.12/J20). San Diego, CA: Society for Neuroscience.
- Muller, E., Le, C. H., Triller, A., and Legendre, P. (2006). Developmental dissociation of presynaptic inhibitory neurotransmitter and postsynaptic receptor clustering in the hypoglossal nucleus. *Mol. Cell. Neurosci.* 32, 254–273.
- Nakae, A., Tanaka, T., Miyake, K., Hase, M., and Mashimo, T. (2008). Comparing methods of detection and quantitation of RNA editing of rat glycine receptor  $\alpha 3$ . *Int. J. Biol. Sci.* 4, 397–405.
- Nikolic, Z., Laube, B., Weber, R. G., Lichter, P., Kioschis, P., Poustka, A., Mulhardt, C., and Becker, C. M. (1998). The human glycine receptor subunit  $\alpha 3$ . Glra3 gene structure, chromosomal localization, and functional characterization of alternative transcripts. *J. Biol. Chem.* 273, 19708–19714.
- Padgett, C. L., and Lummis, S. C. (2008). The F-loop of the GABA A receptor  $\gamma 2$  subunit contributes to benzodiazepine modulation. *J. Biol. Chem.* 283, 2702–2708.
- Palma, E., Amici, M., Sobrero, F., Spinelli, G., Di, A. S., Ragozzino, D., Mascia, A., Scopetta, C., Esposito, V., Miledi, R., and Eusebi, F. (2006). Anomalous levels of Cl<sup>-</sup> transporters

- in the hippocampal subiculum from temporal lobe epilepsy patients make GABA excitatory. *Proc. Natl. Acad. Sci. U.S.A.* 103, 8465–8468.
- Pless, S. A., and Lynch, J. W. (2009). Ligand-specific conformational changes in the alpha1 glycine receptor ligand-binding domain. *J. Biol. Chem.* 284, 15847–15856.
- Plested, A. J., Groot-Kormelink, P. J., Colquhoun, D., and Sivilotti, L. G. (2007). Single-channel study of the spasmodic mutation alpha1A52S in recombinant rat glycine receptors. *J. Physiol.* 581, 51–73.
- Rivera, C., Voipio, J., Thomas-Crusells, J., Li, H., Emri, Z., Sipila, S., Payne, J. A., Minichiello, L., Saarma, M., and Kaila, K. (2004). Mechanism of activity-dependent downregulation of the neuron-specific K-Cl cotransporter KCC2. *J. Neurosci.* 24, 4683–4691.
- Roux, M. J., and Supplisson, S. (2000). Neuronal and glial glycine transporters have different stoichiometries. *Neuron* 25, 373–383.
- Seeburg, P. H., and Hartner, J. (2003). Regulation of ion channel/neurotransmitter receptor function by RNA editing. *Curr. Opin. Neurobiol.* 13, 279–283.
- Simon, J., Wakimoto, H., Fujita, N., Lalande, M., and Barnard, E. A. (2004). Analysis of the set of GABA(A) receptor genes in the human genome. *J. Biol. Chem.* 279, 41422–41435.
- Song, W., Chattipakorn, S. C., and McMahon, L. L. (2006). Glycine-gated chloride channels depress synaptic transmission in rat hippocampus. *J. Neurophysiol.* 95, 2366–2379.
- Stief, F., Zuschratter, W., Hartmann, K., Schmitz, D., and Draguhn, A. (2007). Enhanced synaptic excitation-inhibition ratio in hippocampal interneurons of rats with temporal lobe epilepsy. *Eur. J. Neurosci.* 25, 519–528.
- Supplisson, S., and Roux, M. J. (2002). Why glycine transporters have different stoichiometries. *FEBS Lett.* 529, 93–101.
- Tao, H. W., and Poo, M. M. (2005). Activity-dependent matching of excitatory and inhibitory inputs during refinement of visual receptive fields. *Neuron* 45, 829–836.
- Unwin, N., Miyazawa, A., Li, J., and Fujiyoshi, Y. (2002). Activation of the Nicotinic Acetylcholine Receptor Involves a Switch in Conformation of the alpha Subunits. *J. Mol. Biol.* 319, 1165–1176.
- Wisden, W., Cope, D., Klausberger, T., Hauer, B., Sinkkonen, S. T., Tretter, V., Lujan, R., Jones, A., Korpi, E. R., Mody, I., Sieghart, W., and Somogyi, P. (2002). Ectopic expression of the GABA(A) receptor alpha6 subunit in hippocampal pyramidal neurons produces extrasynaptic receptors and an increased tonic inhibition. *Neuropharmacology* 43, 530–549.
- Zhang, L. H., Gong, N., Fei, D., Xu, L., and Xu, T. L. (2008). Glycine uptake regulates hippocampal network activity via glycine receptor-mediated tonic inhibition. *Neuropsychopharmacology* 33, 701–711.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationship that could be construed as a potential conflict of interest.

Received: 17 July 2009; paper pending published: 30 September 2009; accepted: 13 October 2009; published online: 09 November 2009.

Citation: Legendre P, Förstera B, Jüttner R and Meier JC (2009) Glycine receptors caught between genome and proteome-functional implications of RNA editing and splicing. *Front. Mol. Neurosci.* 2:23. doi: 10.3389/neuro.02.023.2009

Copyright © 2009 Legendre, Förstera, Jüttner and Meier. This is an open-access article subject to an exclusive license agreement between the authors and the Frontiers Research Foundation, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.