



POSTER PRESENTATION

Open Access

hMENA splicing program impacts the clinical outcome of early stage lung cancer patients. How and why?

Paolo Visca¹, Sheila Spada¹, Francesca Di Modugno¹, Emilio Bria², Isabella Sperduti¹, Barbara Antoniani¹, Gabriele Alessandrini¹, Belinda Palermo¹, Vienna Ludovini³, Lucio Crinò³, Francesco Facciolo¹, Michele Milella¹, Marcella Mottolese¹, Paola Nisticò^{1*}

From Melanoma Bridge meeting 2013
Naples, Italy. 5-8 December 2013

Background

In lung cancer, reliable prognostic indicators of the risk of recurrence are still not available. Alternative splicing represents a potential biomarker of diagnosis, prognosis, invasiveness, and response to therapy in different tumors [1], including lung cancer [2].

Human MENA (hMENA) is an actin regulatory protein that modulates cell adhesion and migration [3]. We have isolated three hMENA splice variants, namely hMENA, hMENA^{11a} and hMENAΔv6, impacting differently cell shape and function. hMENA^{11a} expression ensures the integrity of cell-cell adhesion and is associated with an epithelial phenotype, whereas hMENAΔv6 is related to a mesenchymal invasive phenotype. The splicing of hMENA, relevant to epithelial mesenchymal transition, is also regulated by microenvironmental cues [4].

The dynamic reciprocity between tumor and stroma influences the tumor tissue architecture including the T cell localization. This, proposed as a prognostic marker [5], is a prerequisite for antitumor immune surveillance and recently the antibody blockade of immune checkpoints is a new reality in lung cancer treatment [6,7].

Materials and methods

Pan-hMENA and specific hMENA^{11a} Abs were tested by immunohistochemistry on duplicate TMA from 248 N0 NSCLC, and clinical factors (sex, age, histology, grading, T-size, number of resected nodes, RN) were correlated to 3-yr disease-free (DFS), cancer-specific (CSS), and overall

survival (OS) using a Cox model. ROC analysis provided optimal cut-off values and model validation. A logistic equation including regression analysis coefficients was constructed to estimate individual patients' probability (IPP) of relapse. Internal cross-validation (100 simulations with 80% of the dataset) and external validation was accomplished.

A panel of antibodies recognizing CD3, CD4, CD8, CD20 molecules has been employed for the characterization and localization of lymphocytes, by immunohistochemistry.

Results

In the series of N0 NSCLC patients (median follow-up: 36 months, range 1-96), Pan-hMENA and hMENA^{11a} were the only biological variables displaying significant correlation with outcome(s), confirmed by the cross-validation (replication rate: 78%, 83%), with a prognostic model accuracy of 61% (standard error 0.04, p=0.0001). The subgroup of patients with High Pan-hMENA/Low hMENA^{11a} relative expression fared significantly better than the other 3 groups (p≤0.002 for all outcomes). On the basis of the combination between this molecular hybrid variable and T-size and RN, a 3-risk class stratification model was generated, discriminating between patients at different risk of relapse, cancer-related death, and death for any cause, with a prognostic accuracy of 61% (standard error 0.03, p=0.01), according to ROC analysis and validated in an independent dataset of 133 patients.

The correlation between hMENA isoforms and the pattern of expression and localization of lymphocytes in the different groups of risk of relapse identified is under evaluation.

* Correspondence: nistico@ifo.it

¹Regina Elena National Cancer Institute, Rome, Italy

Full list of author information is available at the end of the article

Conclusions

The hMENA splicing program is an early prognostic marker of NSCLC patients and may represent a surrogate marker of a permissive or not tumor microenvironment for lymphocyte recruitment.

Authors' details

¹Regina Elena National Cancer Institute, Rome, Italy. ²Medical Oncology, Azienda Ospedaliera Universitaria Integrata, University of Verona, Verona, Italy. ³Medical Oncology, S. Maria della Misericordia Hospital, Perugia, Italy.

Published: 6 May 2014

References

1. Pal S, Gupta R, Davuluri RV: Alternative transcription and alternative splicing in cancer. *Pharmacol Ther* 2012, **136**(3):283-94.
2. Stallings-Mann ML, Waldmann J, Zhang Y, Miller E, Gauthier ML, Visscher DW, Downey GP, Radisky ES, Fields AP, Radisky DC: Matrix metalloproteinase induction of Rac1b, a key effector of lung cancer progression. *Sci Transl Med* 2012, **4**(142):142ra95.
3. Di Modugno F, Bronzi G, Scanlan MJ, Del Bello D, Cascioli S, Venturo I, Botti C, Nicotra MR, Mottolese M, Natali PG, Santoni A, Jager E, Nisticò P: Human Mena protein, a serex-defined antigen overexpressed in breast cancer eliciting both humoral and CD8+ T-cell immune response. *Int J Cancer* 2004, **109**(6):909-18.
4. Di Modugno F, Iapicca P, Boudreau A, Mottolese M, Terrenato I, Perracchio L, Carstens RP, Santoni A, Bissell MJ, Nisticò P: Splicing program of human Mena produces a previously undescribed isoform associated with invasive, mesenchymal-like breast tumors. *Proc Natl Acad Sci U S A* 2012, **109**(47):19280-5.
5. Fridman WH, Pagès F, Sautès-Fridman C, Galon J: The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer* 2012, **12**(4):298-306.
6. Pardoll DM: The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012, **12**(4):252-64.
7. Brahmer JR, Tykodi SS, Chow LQ, Hwu WJ, Topalian SL, Hwu P, Drake CG, Camacho LH, Kauh J, Odunsi K, Pitot HC, Hamid O, Bhatia S, Martins R, Eaton K, Chen S, Salay TM, Alaparthy S, Gross JF, Korman AJ, Parker SM, Agrawal S, Goldberg SM, Pardoll DM, Gupta A, Wigginton JM: Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med* 2012, **366**(26):2455-65.

doi:10.1186/1479-5876-12-S1-P12

Cite this article as: Visca et al.: hMENA splicing program impacts the clinical outcome of early stage lung cancer patients. How and why? *Journal of Translational Medicine* 2014 **12**(Suppl 1):P12.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

