

Pyogenic Liver Abscess and Endogenous Endophthalmitis Due to K64-ST1764 Hypervirulent *Klebsiella pneumoniae*: A Case Report

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Bo Zhao^{1,*}
Renjing Hu^{1,2,*}
Lei Gong¹
Xiaoyun Wang¹
Yingwei Zhu¹
Gaojue Wu¹

¹Department of Gastroenterology, The Affiliated Wuxi Second People's Hospital of Nanjing Medical University, Wuxi, People's Republic of China; ²Department of Clinical Laboratory, The Affiliated Wuxi Second People's Hospital of Nanjing Medical University, Wuxi, People's Republic of China

*These authors contributed equally to this work

Rationale: *Klebsiella pneumoniae* (*K. pneumoniae*, KP) are divided into two types: classic *K. pneumoniae* (cKP) and hypervirulent *K. pneumoniae* (hvKP). hvKP causes liver abscess and metastatic infection. Here, we report one case with pyogenic liver abscess (PLA) and endogenous endophthalmitis (EE) due to a relatively rarely reported serotype of *K. pneumoniae* in China.

Patient Concerns: An 80-year old man presented with nausea, vomiting, and epigastric discomfort for 2 weeks.

Diagnoses: PLA was identified by CT scan and abdominal ultrasound. Urgent ophthalmologic consultation was performed. B-scan ocular ultrasound was done and he was diagnosed as EE.

Interventions: Antibiotic treatment, intravitreal injection of eyes and eye drops were given. Percutaneous needle aspiration, evisceration, and drainage of the right eye were performed.

Outcomes: Cultures of the blood, the aspirated pus from the liver abscess, and the contents of the eyeball all yielded *K. pneumoniae* with a positive string test. The capsular serotype was K64. According to the existence of multiple virulence genes and the severe invasive clinical manifestation, this strain is regarded as a hvKp strain. Multilocus sequence typing (MLST) revealed the sequence type (ST) of this strain was K64-ST1764. Antimicrobial resistance genes, *bla*_{NDM-1} and *bla*_{KPC-2}, were not detected in the genome. The patient lost his eyesight but his symptoms subsided. During 15 months follow-up, the result was satisfactory.

Lessons: Here, we report one case with PLA due to a relatively rarely reported serotype of *K. pneumoniae* in China. This K64 *K. pneumoniae* strain is confirmed as hvKp by multiple methods. It is noteworthy that the sequence type is K64-ST1764 instead of the commonest ST11. Moreover, this strain is not considered a *K. pneumoniae* carbapenemase-producing *K. pneumoniae* (KPC-Kp) or a carbapenem-resistant *K. pneumoniae* (CRKP) as it is usually. Further follow-up and research are required to investigate this strain.

Keywords: case report, hypervirulent *Klebsiella pneumoniae*, pyogenic liver abscess, antibiotic resistance, ultrasonography guided puncture aspiration

Correspondence: Xiaoyun Wang
Department of Gastroenterology, The Affiliated Wuxi Second People's Hospital of Nanjing Medical University, Wuxi, People's Republic of China
Tel +86 15061857125
Fax +86 510 685662052
Email xiaoyunwang68@aliyun.com

Introduction

PLA is a potentially life-threatening disease which is mainly caused by bacteria. Along with the increasing prevalence of hvKp, PLA due to *K. pneumoniae* has been steadily increasing. *K. pneumoniae* are divided into two types: cKP and hvKP.¹ Ckp is an

opportunistic pathogen which mainly infects immunodeficient patients or the immunocompromised population. HvKP can infect young healthy people without underlying diseases.² Increased virulence of hvkp is related to its hypermucoviscous phenotype and stronger ability of iron acquisition.^{3,4} HvKP often causes primary liver abscess and metastatic infections like endophthalmitis, meningitis, and so on.⁵ HvKP infections are more common in the Asian Pacific but now are occurring globally.

Here we report one case who presented in China with PLA and EE due to a relatively rarely reported serotype of hvKP. Different from previous reports, the ST of this strain was K64-ST1764 instead of the commonest ST11. It is especially noteworthy that antimicrobial susceptibility testing showed that this strain was sensitive to all antibiotics, which means it is not a KPC-Kp or a CRKP.

Case Presentation

An 80-year old Chinese male patient presented to the emergency department with nausea, vomiting, and epigastric discomfort for 2 weeks without treatment. This patient had a history of Type II diabetes mellitus, hypertension, and coronary heart disease. There was no relevant family history or history of genetic disease. Laboratory investigations revealed a white cell count 14.42×10^9 cells/L (normal range = $3.5\text{--}9.5 \times 10^9$ cells/L), 85% neutrophils (normal range = 40–75%), aspartate transaminase (AST) 10^9 U/L (normal range < 50 U/L), lactate dehydrogenase (LDH) 349 U/L (normal range = 109–245 U/L), random blood glucose 19.96 mmol/L (normal range = 3.8–6.2 mmol/L), blood urea 11.3 mmol/L (normal range = 2.1–8.6 mmol/L), creatinine $141.7 \mu\text{mol/L}$ (normal range = $35.2\text{--}97.5 \mu\text{mol/L}$). Severe infection and ketoacidosis were suspected so emergency doctors gave him antibiotics (imipenem combined with ornidazole), rabeprazole, antiemetic drug, and fluid infusion, but his condition worsened. Fever appeared and the highest armpit temperature was 39°C.

He was admitted to the Department of Gastroenterology for further treatment. Laboratory investigations revealed C-reactive protein 439.43 mg/L (normal range < 10 mg/L), serum sodium (Na) 133.9 mmol/L (normal range = 136–145 mmol/L), serum potassium (K) 3.71 mmol/L (normal range = 3.5–5.3 mmol/L), total bilirubin $21 \mu\text{mol/L}$ (normal range = 3.4–20.6 $\mu\text{mol/L}$), and alanine aminotransferase (ALT) 88.6 U/L (normal range < 50 U/L). Physical examination revealed mild epigastric tenderness. A right lobe liver abscess was identified on CT scan (Figure 1). Abdominal



Figure 1 Abdominal CT scan showed a right lobe liver abscess (rounded, low density areas).

ultrasound showed a right lobe liver abscess (mixed echoes) with the size of 3.5×2.7 cm in the right liver lobe (Figure 2).

The patient was treated by anti-infection treatment (biapenem and morinidazole by intravenous injection), liver protection (polyene phosphatidylcholine), decreasing blood sugar, controlling blood pressure, and fluid infusion. Ultrasonography guided puncture aspiration for hepatic abscess was proposed but he refused. Both his white cell count and C-reactive protein were dropped after treatment but he was still feverish. Bacteria culturing and antimicrobial susceptibility testing were performed by using the VITEK-II Compact automated microbiological system and confirmed by using the broth microdilution method. Blood cultures, from both aerobic and anaerobic bottles, showed *K. pneumoniae* which was sensitive to all antibiotics.

The vision of this patient became blurred so urgent ophthalmologic consultation was performed. A B-scan ocular ultrasound was done and he was diagnosed as EE with bad prognosis. Intravitreal injection (vancomycin and cefoperazone) of eyes and eye drops were given. Finally, after obtaining the patient's consent, ultrasonography guided puncture aspiration was performed, which resulted in lowering the temperature and improving the inflammatory marker levels. Liver aspirate culture was also positive for *K. pneumoniae* and sensitive to all antibiotics (Table 1). The patient's eyes were swollen and empyema and the light perception of both eyes was basically lost. The ophthalmologist performed evisceration and drainage of the right eye, as well as repeated




Figure 2 Abdominal ultrasound showed a right lobe liver abscess (mixed echoes, with a size of 3.5cm×2.7 cm).

intravitreal injection of the left eye. Abdominal ultrasonography was performed 6 days after the first therapeutic puncture. The abscess cavity decreased conspicuously

Table I Antimicrobial Susceptibility Testing Results for *K. pneumoniae* Isolate, Using Reference Broth Microdilution According to Clinical and Laboratory Standards Institute (CLSI) Guidelines

Antimicrobial Agent	Minimum Inhibitory Concentration (µg/mL)	Interpretation*
Amikacin	≤2	S
Ampicillin	>32	S
Aztreonam	≤1	S
Ceftriaxone	≤1	S
Ceftazidime	≤1	S
Cefepime	≤1	S
Ciprofloxacin	≤0.25	S
Gentamicin	≤1	S
Imipenem	≤1	S
Levofloxacin	≤0.25	S
Ertapenem	≤0.5	S
Piperacillin-Tazobactam	≤4	S
Tigecycline	0.5	S

Notes: *Interpretative criteria were applied according to CLSI document M100 (Performance Standards for Antimicrobial Susceptibility Testing, 28th ed. 2018) except for tigecycline where criteria established by the US Food and Drug Administration were used.

Abbreviations: R, resistant; S, susceptible.

and the second percutaneous needle aspiration was given. After treatment, the patient's eye swelling and the empyema was significantly improved and he became afebrile. He was given ceftriaxon 2 g by intravenous injection once a day for 4 weeks. Repeated blood cultures were negative and ultrasound reexamination showed the abscess cavity had disappeared. He was discharged and referred to the gastroenterology clinic for follow-up. During 15 months follow-up, the result was satisfactory.

Isolates were retested by 16S rRNA gene sequences as previously reported⁶ and *K. pneumoniae* was identified and reconfirmed.

String test was performed as previously reported.² The colony was grown at 37°C overnight on a sheep blood agar plate and then stretched by a bacteriology inoculation loop. A positive result is defined as the formation of a string >5 mm in length. Cultures of the blood, the aspirated pus from the liver abscess, and the contents of the eyeball all yielded *K. pneumoniae* with a positive string test (Figure 3).

Capsular serotype of each isolate was performed by polymerase chain reaction (PCR) as previously reported.⁷ PCR analysis indicated that the capsular serotype of the isolated *K. pneumoniae* was K64, which was validated by wzi sequencing as described.⁸



Figure 3 All cultures yielded *K. pneumoniae* with a positive string test (>5mm in string length).

MLST was performed by PCR amplification as previously reported⁹ and this strain belongs to ST1764.

Multiple biomarkers, including *peg-344* (+), *iroB* (+), *iucA* (+), *p_rmpA* (+), *p_rmpA2* (+), *c_rmpA* (-), *ureA* (+), *uge* (+), *wabG* (+), *allS* (+), *mrkD* (+), and *fimH* (+) were detected by PCR as described¹⁰ and multiple virulence genes were found in all the above isolates.

Screening for carbapenemase genes was performed by PCR as described¹¹ and *bla*_{NDM-1} and *bla*_{KPC-2} were not found in this K64 strain.

We evaluated the virulence of this *K. pneumoniae* strain in a *Galleria mellonella* infection model.¹² We infected the *Galleria mellonella* larvae with different concentrations (10^2 – 10^8 colony-forming units/mL) of *K. pneumoniae* and recorded the survival rate (Figure 4). This strain showed a strong concentration-dependent lethality to the larvae.

Discussion and Conclusions

The first clinical report about hvKp was in Taiwan in 1986.¹³ Although it is more common in the Asia-Pacific region, infections due to hvKp have gradually become global.^{14,15} Because hvKp was considered to have a hypermucoviscosity phenotype, the string test has been used to defining hvKP.³ However, more and more studies have shown that hypermucoviscosity is not pathognomonic for hvKp since this phenotype also can be observed in cKp strains.¹⁶ Recently, several virulence genes, like *iuc*, *peg-344*, *rmpA*, and *rmpA2*, have been shown to most accurately predict hvKp strains.¹⁶




Figure 4 Survival curves for *Galleria mellonella* larvae inoculated with different concentrations (10^2 – 10^8 colony-forming units/mL) of the *K. pneumoniae* strain.

Since the 1980s, PLA due to *K. pneumoniae* has been steadily increasing. Especially in the Asia-Pacific region, hvKP has already become the predominant pathogen of PLA.^{17–19} Furthermore, these infections are occurring globally. A study in China has reported that among all PLA cases, *K pneumoniae* was the most common pathogen and a higher incidence of diabetes mellitus was shown in patients with *K. pneumoniae*-induced PLA.²⁰ Another study has reported that most of the liver abscesses were caused by *K. pneumoniae* and mostly occurred in patients with diabetes mellitus.²¹ The major clinical features of KP-PLA are fever, nausea, vomiting, abdominal discomfort, and laboratory inspection shows elevated white blood cell count and liver dysfunction. In this patient, all the above features can be found.

There are mainly three treatment methods: appropriate early antibiotic treatment, ultrasonography guided puncture aspiration or draining treatment, and open surgical drainage or surgical resection of the liver abscesses. We initially treated the patient with carbapenems and Morinidazole. After the condition stabilized, the antibiotics were changed to ceftriaxone for another 4 weeks. Some studies have suggested that the majority of PLA cases require drainage.²² A report showed that antimicrobial therapy alone was effective for small abscesses, while percutaneous needle aspiration or draining should be chosen for larger abscesses.²³ A study reported that liver abscesses due to *K. pneumoniae* that were >5 cm were likely to have a delayed response to therapy.²⁴ In the present study, the liver abscess was <5 cm (35×27 mm) and this hvKP strain was sensitive to all antibiotics. Even so, early antimicrobial therapy alone was ineffective. Once aspiration was performed, clinical improvement was observed. This suggested that, even for small abscesses,

puncture or drainage should be performed as early as possible when the effect of drug therapy is unsatisfactory.

In recent years, the incidence of EE has gradually increased. Gram-positive bacteria and fungal organisms account for the majority of the cases in the developed world, whereas gram-negative organisms like *K. pneumoniae* are more common in the Asia-Pacific region.²⁵ Now it has become a frequent complication: approximately 5% of individuals with hvKp bacteremia ultimately develop EE.²⁶

EE is a common metastatic complication of *K. pneumoniae* liver abscess.²⁷ Diabetes mellitus was considered to be a significant risk factor. The prognosis of EE is generally poor even with prompt diagnosis and aggressive treatment given. The final visual outcome was only light perception or worse in 89% of patients with EE associated with *K. pneumoniae*-induced PLA.²⁷ Diabetes mellitus and poor initial visual acuity were associated with poor visual outcome.²⁸ Reports showed that the main prognostic factor in *Klebsiella* EE is the presence of hypopyon which can be found in our patient.²⁹ This patient lost his eyesight.

It has long been recognized that almost all patients with severe infection with liver abscess and extrahepatic infections are infected exclusively with *K. pneumoniae* serotypes K1 or K2.² However, it is clear now that non-K1/K2 hvKp strains can express virulence genes and are capable of causing liver abscess and metastatic spread.^{16,30} The serotype of the isolated *K. pneumoniae* strain in our patient was K64, a relatively rarely reported serotype. This serotype has always been considered a cKP. But it is recently recognized that K64 is predicted to be observed for hvKp strains.^{31,32} Recent research finds that some cKp strains can acquire the hvKp virulence plasmid and evolve into hvKp strains.^{12,32} Our K64 *K. pneumoniae* strain showed strong virulence in the *Galleria mellonella* infection model. According to positive string test, the existence of multiple virulence genes, the strong lethality to the larvae and the severe invasive clinical manifestation of its infection, we can regard this K64 strain as a hvKp strain. Compared to the classification based on string test alone, our method is more reasonable.

In particular, ST was determined by MLST and the result was K64-ST1764 instead of the commonest ST11, which was considered to be predominant in Extended-Spectrum β -lactamases (ESBL)-producing strains and was prevalent in CRKP strains.^{33,34} However, previous reports showed that ST11 was predominant in K64 capsular type, hence K64 was considered as a prevalent type of CRKP.^{35–38} Furthermore, *bla*_{NDM-1} and *bla*_{KPC-2}, two important carbapenemase genes,

were not found in this K64 strain and antimicrobial susceptibility testing showed that it was sensitive to all antibiotics. Contrary to previous studies,^{31,32,36,39} these findings indicated that this K64 strain is not a KPC-Kp or a CRKP strain.

In conclusion, here we report one patient with PLA and EE due to a relatively rarely reported serotype of *K. pneumoniae*: K64. As the existence of multiple virulence genes, the strong lethality to the larvae and the severe invasive clinical manifestation and positive string test, this K64 *K. pneumoniae* can be regarded as a hvKp strain. Different from previous reports, the ST of this strain was K64-ST1764 instead of the commonest ST11. It is not a KPC-Kp or a CRKP, either. Further follow-up and research are required to investigate this strain.

Abbreviations

CRKP, carbapenem-resistant *K. pneumoniae*; cKP, classic *K. pneumoniae*; ESBL, extended-spectrum β -lactamases; hvKp, hypervirulent *K. pneumoniae*; KPC-Kp, *K. pneumoniae* carbapenemase (KPC)-producing; MLST, multilocus sequence typing; PLA, pyogenic liver abscess; ST, sequence type.

Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics Approval

This case report was approved by the ethics committee of The Affiliated Wuxi Second People's Hospital of Nanjing Medical University.

Consent for Publication

Written informed consent for publication of his clinical details and clinical images was obtained from the patient.

Acknowledgments

The key discipline of Wuxi No.ZDXK002 Workstation of academician Fan Daiming No.CYR1705

Author Contributions

Bo Zhao and Renjing Hu have contributed equally to this work and they are co-first authors. Conceived the study: Xiaoyun Wang, Bo Zhao; Acquired data, analyzed data: Bo Zhao, Renjing Hu; Performed the microbiological analysis: Renjing Hu; Wrote the manuscript: Bo Zhao; Contributed reagents/materials/analysis tools: Lei Gong,

Xiaoyun Wang; and Revised the manuscript: Xiaoyun Wang, Yingwei Zhu, Gaojue Wu.

All authors made substantial contributions to the conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the National Natural Science Foundation of China No.81500467, Foundation for Key Medical Talents in Jiangsu Province No.QNRC2016151, Six Talent Peaks Project of Jiangsu Province No.WSN-220, Six One Top-notch Talent Project of Jiangsu Province No.LGY201701. The funders had no role in the study design, conduct, analysis, interpretation of study results, decision to publish, or preparation of the manuscript. No additional external funding was received for this study.

Disclosure

The authors declare that they have no conflicts of interest to this work.

References

- Shon AS, Bajwa RP, Russo TA. Hypervirulent (hyper-mucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. *Virulence*. 2013;4(2):107–118. doi:10.4161/viru.22718
- Siu LK, Yeh K-M, Lin J-C, Fung C-P, Chang F-Y. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. *Lancet Infect Dis*. 2012;12(11):881–887. doi:10.1016/S1473-3099(12)70205-0
- Fang CT, Chuang YP, Shun CT, Chang SC, Wang JT. A novel virulence gene in *Klebsiella pneumoniae* strains causing primary liver abscess and septic metastatic complications. *J Exp Med*. 2004;199(5):697–705. doi:10.1084/jem.20030857
- Huang YH, Chou SH, Liang SW, et al. Emergence of an XDR and carbapenemase-producing hypervirulent *Klebsiella pneumoniae* strain in Taiwan. *J Antimicrob Chemother*. 2018;73(8):2039–2046. doi:10.1093/jac/dky164
- Shon AS, Russo TA. Hypervirulent *Klebsiella pneumoniae*: the next superbug? *Future Microbiol*. 2012;7(6):669–671. doi:10.2217/fmb.12.43
- He Y, Guo X, Xiang S, et al. Comparative analyses of phenotypic methods and 16S rRNA, *khe*, *rpoB* genes sequencing for identification of clinical isolates of *Klebsiella pneumoniae*. *Antonie Van Leeuwenhoek*. 2016;109(7):1029–1040. doi:10.1007/s10482-016-0702-9
- Turton J, Perry C, Elgohari S, Hampton C. PCR characterization and typing of *Klebsiella pneumoniae* using capsular type-specific, variable number tandem repeat and virulence gene targets. *J Med Microbiol*. 2010;59:541–547. doi:10.1099/jmm.0.015198-0
- Brisse S, Passet V, Haugaard A, et al. *wzi* Gene sequencing, a rapid method for determination of capsular type for *Klebsiella* strains. *J Clin Microbiol*. 2013;51(12):4073–4078. doi:10.1128/JCM.01924-13
- Diancourt L, Passet V, Verhoef J, Grimont P, Brisse S. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol*. 2005;43(8):4178–4182. doi:10.1128/JCM.43.8.4178-4182.2005
- Yan Q, Zhou M, Zou M, Liu W. Hypervirulent *Klebsiella pneumoniae* induced ventilator-associated pneumonia in mechanically ventilated patients in China. *European J Clin Microbiol Infectious Diseases*. 2016;35(3):387–396. doi:10.1007/s10096-015-2551-2
- Poirer L, Walsh T, Cuvillier V, Nordmann P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis*. 2011;70(1):119–123. doi:10.1016/j.diagmicrobio.2010.12.002
- Gu D, Dong N, Zheng Z, et al. A fatal outbreak of ST11 carbapenem-resistant hypervirulent *Klebsiella pneumoniae* in a Chinese hospital: a molecular epidemiological study. *Lancet Infect Dis*. 2018;18(1):37–46. doi:10.1016/S1473-3099(17)30489-9
- Liu YC, Cheng DL, Lin CL. *Klebsiella pneumoniae* liver abscess associated with septic endophthalmitis. *Arch Intern Med*. 1986;146(10):1913–1916. doi:10.1001/archinte.1986.00360220057011
- Rossi B, Gasperini ML, Leflon-Guibout V, et al. Hypervirulent *Klebsiella pneumoniae* in cryptogenic liver abscesses, Paris, France. *Emerg Infect Dis*. 2018;24(2):221–229. doi:10.3201/eid2402.170957
- Fazili T, Sharnogoe C, Endy T, Kiska D, Javaid W, Polhemus M. *Klebsiella pneumoniae* liver abscess: an emerging disease. *Am J Med Sci*. 2016;351(3):297–304. doi:10.1016/j.amjms.2015.12.018
- Russo TA, Olson R, Fang CT, et al. Identification of biomarkers for differentiation of hypervirulent *Klebsiella pneumoniae* from classical *K. pneumoniae*. *J Clin Microbiol*. 2018;56:9. doi:10.1128/JCM.00776-18
- Cheng DL, Liu YC, Yen MY, Liu CY, Shi FW, Wang LS. Causal bacteria of pyogenic liver abscess. *Taiwan Yi Xue Hui Za Zhi*. 1989;88(10):1008–1011.
- Chang FY, Chou MY, Fan RL, Shaio MF. A clinical study of *Klebsiella* liver abscess. *Taiwan Yi Xue Hui Za Zhi*. 1988;87(3):282–287.
- Lee TY, Wan YL, Tsai CC. Gas-containing liver abscess: radiological findings and clinical significance. *Abdom Imaging*. 1994;19(1):47–52. doi:10.1007/BF02165861
- Kong H, Yu F, Zhang W, Li X. Clinical and microbiological characteristics of pyogenic liver abscess in a tertiary hospital in East China. *Medicine*. 2017;96(37):e8050. doi:10.1097/MD.00000000000008050
- Wang WJ, Tao Z, Wu HL. Etiology and clinical manifestations of bacterial liver abscess: A study of 102 cases. *Medicine*. 2018;97(38):e12326. doi:10.1097/MD.00000000000012326
- Waghmare M, Shah H, Tiwari C, Khedkar K, Gandhi S. Management of liver abscess in children: our experience. *Euroasian J Hepatogastroenterol*. 2017;7(1):23–26. doi:10.5005/jp-journals-10018-1206
- Cheema HA, Saeed A. Etiology, presentation and management of liver abscesses at the children's hospital lahore. *Annals King Edward Medical University*. 2008;14(4):148.
- Chan DS, Archuleta S, Llorin RM, Lye DC, Fisher D. Standardized outpatient management of *Klebsiella pneumoniae* liver abscesses. *Int J Infect Dis*. 2013;17(3):e185–188. doi:10.1016/j.ijid.2012.10.002
- Sadiq MA, Hassan M, Agarwal A, et al. Endogenous endophthalmitis: diagnosis, management, and prognosis. *J Ophthalmic Inflamm Infect*. 2015;5(1):32. doi:10.1186/s12348-015-0063-y
- Lee HC, Chuang YC, Yu WL, et al. Clinical implications of hyper-mucoviscosity phenotype in *Klebsiella pneumoniae* isolates: association with invasive syndrome in patients with community-acquired bacteraemia. *J Intern Med*. 2006;259(6):606–614. doi:10.1111/j.1365-2796.2006.01641.x
- Yang CS, Tsai HY, Sung CS, Lin KH, Lee FL, Hsu WM. Endogenous *Klebsiella* endophthalmitis associated with pyogenic liver abscess. *Ophthalmology*. 2007;114(5):876–880. doi:10.1016/j.optha.2006.12.035

28. Sheu SJ, Kung YH, Wu TT, Chang FP, Horng YH. Risk factors for endogenous endophthalmitis secondary to klebsiella pneumoniae liver abscess: 20-year experience in Southern Taiwan. *Retina*. 2011;31(10):2026–2031. doi:10.1097/IAE.0b013e31820d3f9e
29. Ang M, Jap A, Chee SP. Prognostic factors and outcomes in endogenous Klebsiella pneumoniae endophthalmitis. *Am J Ophthalmol*. 2011;151(2):338–344.e332. doi:10.1016/j.ajo.2010.08.036
30. Gu DX, Huang YL, Ma JH, et al. Detection of colistin resistance gene mcr-1 in hypervirulent klebsiella pneumoniae and escherichia coli isolates from an infant with diarrhea in China. *Antimicrob Agents Chemother*. 2016;60(8):5099–5100. doi:10.1128/AAC.00476-16
31. Liu Y, Liu PP, Wang LH, Wei DD, Wan LG, Zhang W. Capsular polysaccharide types and virulence-related traits of epidemic kpc-producing klebsiella pneumoniae isolates in a Chinese University Hospital. *Microb Drug Resist*. 2017;23(7):901–907. doi:10.1089/mdr.2016.0222
32. Yu F, Lv J, Niu S, et al. In vitro activity of ceftazidime-avibactam against carbapenem-resistant and hypervirulent klebsiella pneumoniae isolates. *Antimicrob Agents Chemother*. 2018;62:8. doi:10.1128/AAC.01031-18
33. Rhee JY, Park YK, Shin JY, et al. KPC-producing extreme drug-resistant Klebsiella pneumoniae isolate from a patient with diabetes mellitus and chronic renal failure on hemodialysis in South Korea. *Antimicrob Agents Chemother*. 2010;54(5):2278–2279. doi:10.1128/AAC.00011-10
34. Lee MY, Ko KS, Kang CI, Chung DR, Peck KR, Song JH. High prevalence of CTX-M-15-producing Klebsiella pneumoniae isolates in Asian countries: diverse clones and clonal dissemination. *Int J Antimicrob Agents*. 2011;38(2):160–163. doi:10.1016/j.ijantimicag.2011.03.020
35. Pan YJ, Lin TL, Lin YT, et al. Identification of capsular types in carbapenem-resistant Klebsiella pneumoniae strains by wzc sequencing and implications for capsule depolymerase treatment. *Antimicrob Agents Chemother*. 2015;59(2):1038–1047. doi:10.1128/AAC.03560-14
36. Lu MC, Tang HL, Chiou CS, Wang YC, Chiang MK, Lai YC. Clonal dissemination of carbapenemase-producing Klebsiella pneumoniae: two distinct sub-lineages of Sequence Type 11 carrying blaKPC-2 and blaOXA-48. *Int J Antimicrob Agents*. 2018;52(5):658–662. doi:10.1016/j.ijantimicag.2018.04.023
37. Cheng YH, Lin TL, Pan YJ, Wang YP, Lin YT, Wang JT. Colistin resistance mechanisms in Klebsiella pneumoniae strains from Taiwan. *Antimicrob Agents Chemother*. 2015;59(5):2909–2913. doi:10.1128/AAC.04763-14
38. Yu F, Lv J, Niu S, et al. Multiplex pcr analysis for rapid detection of klebsiella pneumoniae carbapenem-resistant (Sequence Type 258 [ST258] and ST11) and Hypervirulent (ST23, ST65, ST86, and ST375) Strains. *J Clin Microbiol*. 2018;56:9.
39. Dong N, Zhang R, Liu L, et al. Genome analysis of clinical multi-locus sequence Type 11 Klebsiella pneumoniae from China. *Microb Genom*. 2018.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>