

POSTER PRESENTATION

Open Access

Theoretical studies on cycloaddition reactions

Lydia Rhyman^{1*}, Ponnadurai Ramasami¹, Luis R Domingo²

From 9th German Conference on Chemoinformatics
Fulda, Germany. 10-12 November 2013

Cycloaddition reactions represent one of the most powerful processes in organic chemistry. The most common types of cycloaddition reactions are the Diels-Alder (DA) and the 1,3-dipolar cycloaddition reactions (1,3-DCs) which lead to five and six membered rings, respectively.

In our ongoing efforts to contribute to the understanding of DA and 1,3-DCs; we studied the following using the B3LYP/6-31G(d) level of theory:

1. The 1,3-DCs of the pyridinium-3-olates and pyrazinium-3-olates with methyl acrylate and methyl methacrylate [1,2].
2. The competitive hetero-DA and 1,3-DCs of methyl glyoxylate oxime and its tautomeric nitrone with cyclopentadiene in the absence and in the presence of BF_3 as a Lewis acid catalyst [3].
3. A systematic study on the 1,3-DCs of C_{60} with substituted nitrile oxides (RCNO ; R = F, Cl, Br, NC, CN and NO_2) [4].

Among the outcomes of our investigations is the successful use of theoretical methods to understand the regio- and stereoselectivity of the reactions considered. It is expected that experimentalists find the results useful for synthesis involving these moieties and cycloaddition reactions. This presentation will overview our ongoing research program to have more understanding of these cycloaddition reactions.

Authors' details

¹Computational Chemistry Group, Department of Chemistry, Faculty of Science, University of Mauritius, Réduit, Mauritius. ²Departamento de Química Orgánica, Universidad de Valencia, Dr. Moliner 50, 46100 Burjassot, Valencia, Spain.

Published: 11 March 2014

* Correspondence: lyd.rhyman@gmail.com

¹Computational Chemistry Group, Department of Chemistry, Faculty of Science, University of Mauritius, Réduit, Mauritius
Full list of author information is available at the end of the article

References

1. Rhyman L, Abdallah HH, Jhaumeer-Laulloo S, Domingo LR, Joule JA, Ramasami P: The 1,3-Dipolar Cycloaddition of 1*H*-pyridinium-3-olate and 1-Methylpyridinium-3-olate with Methyl Acrylate: A Density Functional Theory Study. *Tetrahedron* 2010, **66**:9187-9193.
2. Domingo LR, Sáez JA, Joule JA, Rhyman L, Ramasami P: A DFT Study of the [3+2] versus [4+2] Cycloaddition Reactions of 1,5,6-Trimethylpyrazinium-3-olate with Methyl Methacrylate. *J Org Chem* 2013, **78**:1621-1629.
3. Rhyman L, Ramasami P, Joule JA, Sáez JA, Domingo LR: A DFT Study of the Regio- and Stereoselectivity of the 1,3-Dipolar Cycloaddition of C-Methyl Substituted Pyrazinium-3-olates with Methyl Acrylate and Methyl Methacrylate. *RSC Adv* 2013, **3**:447-457.
4. Rhyman L, Jhaumeer-Laulloo S, Domingo LR, Joule JA, Ramasami P: Computational Assessment of 1,3-Dipolar Cycloaddition of nitrile oxides with ethene and [60]fullerene. *Heterocycles* 2012, **84**:719-735.

doi:10.1186/1758-2946-6-S1-P2

Cite this article as: Rhyman et al.: Theoretical studies on cycloaddition reactions. *Journal of Cheminformatics* 2014 **6**(Suppl 1):P2.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

"Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge."

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>

